

World Journal of *Clinical Cases*

World J Clin Cases 2021 November 6; 9(31): 9320-9698



FRONTIER

- 9320 Gut-liver axis in cirrhosis: Are hemodynamic changes a missing link?
Maslennikov R, Ivashkin V, Efremova I, Poluektova E, Shirokova E

REVIEW

- 9333 Pharmaconutrition strategy to resolve SARS-CoV-2-induced inflammatory cytokine storm in non-alcoholic fatty liver disease: Omega-3 long-chain polyunsaturated fatty acids
Jeyakumar SM, Vajreswari A
- 9350 Major depressive disorder: Validated treatments and future challenges
Karrouri R, Hammani Z, Benjelloun R, Otheman Y

MINIREVIEWS

- 9368 Gene × environment interaction in major depressive disorder
Zhao MZ, Song XS, Ma JS
- 9376 Deep learning driven colorectal lesion detection in gastrointestinal endoscopic and pathological imaging
Cai YW, Dong FF, Shi YH, Lu LY, Chen C, Lin P, Xue YS, Chen JH, Chen SY, Luo XB

ORIGINAL ARTICLE**Case Control Study**

- 9386 Cognitive behavioral therapy on personality characteristics of cancer patients
Yuan XH, Peng J, Hu SW, Yang Y, Bai YJ

Retrospective Cohort Study

- 9395 Extrapneumonic necrosis volume: A new tool in acute pancreatitis severity assessment?
Cucuteanu B, Negru D, Gavrilescu O, Popa IV, Floria M, Mihai C, Cijevschi Prelipcean C, Dranga M
- 9406 Establishment of a risk assessment score for deep vein thrombosis after artificial liver support system treatment
Ye Y, Li X, Zhu L, Yang C, Tan YW

Retrospective Study

- 9417 Clinical management and susceptibility of primary hepatic lymphoma: A cases-based retrospective study
Hai T, Zou LQ
- 9431 Association of serum pepsinogen with degree of gastric mucosal atrophy in an asymptomatic population
Cai HL, Tong YL

- 9440** Risk factors for relapse and nomogram for relapse probability prediction in patients with minor ischemic stroke

Yu XF, Yin WW, Huang CJ, Yuan X, Xia Y, Zhang W, Zhou X, Sun ZW

- 9452** Incidence, prognosis, and risk factors of sepsis-induced cardiomyopathy

Liang YW, Zhu YF, Zhang R, Zhang M, Ye XL, Wei JR

- 9469** Associations with pancreatic exocrine insufficiency: An United Kingdom single-centre study

Shandro BM, Chen J, Ritehnia J, Poullis A

- 9481** Retrospective analysis of influencing factors on the efficacy of mechanical ventilation in severe and critical COVID-19 patients

Zeng J, Qi XX, Cai WW, Pan YP, Xie Y

Observational Study

- 9491** Vitamin D deficiency, functional status, and balance in older adults with osteoarthritis

Montemor CN, Fernandes MTP, Marquez AS, Poli-Frederico RC, da Silva RA, Fernandes KBP

- 9500** Psychological impact of the COVID-19 pandemic on Chinese population: An online survey

Shah T, Shah Z, Yasmeen N, Ma ZR

- 9509** Outcomes of different minimally invasive surgical treatments for vertebral compression fractures: An observational study

Yeh KL, Wu SH, Liaw CK, Hou SM, Wu SS

META-ANALYSIS

- 9520** Glycated albumin as a biomarker for diagnosis of diabetes mellitus: A systematic review and meta-analysis

Xiong JY, Wang JM, Zhao XL, Yang C, Jiang XS, Chen YM, Chen CQ, Li ZY

CASE REPORT

- 9535** Rapid response to radiotherapy in unresectable tracheal adenoid cystic carcinoma: A case report

Wu Q, Xu F

- 9542** Clinical observation of pediatric-type follicular lymphomas in adult: Two case reports

Liu Y, Xing H, Liu YP

- 9549** Malignant adenomyoepithelioma of the breast: Two case reports and review of the literature

Zhai DY, Zhen TT, Zhang XL, Luo J, Shi HJ, Shi YW, Shao N

- 9557** Validation of diagnostic strategies of autoimmune atrophic gastritis: A case report

Sun WJ, Ma Q, Liang RZ, Ran YM, Zhang L, Xiao J, Peng YM, Zhan B

- 9564** Characteristics of primary giant cell tumor in soft tissue on magnetic resonance imaging: A case report

Kang JY, Zhang K, Liu AL, Wang HL, Zhang LN, Liu WV

- 9571** Acute esophageal necrosis as a complication of diabetic ketoacidosis: A case report
Moss K, Mahmood T, Spaziani R
- 9577** Simultaneous embolization of a spontaneous porto-systemic shunt and intrahepatic arterioportal fistula: A case report
Liu GF, Wang XZ, Luo XF
- 9584** Ureteroscopic holmium laser to transect the greater omentum to remove an abdominal drain: Four case reports
Liu HM, Luo GH, Yang XF, Chu ZG, Ye T, Su ZY, Kai L, Yang XS, Wang Z
- 9592** Forearm compartment syndrome due to acquired hemophilia that required massive blood transfusions after fasciotomy: A case report
Kameda T, Yokota T, Ejiri S, Konno SI
- 9598** Transforaminal endoscopic excision of bi-segmental non-communicating spinal extradural arachnoid cysts: A case report and literature review
Yun ZH, Zhang J, Wu JP, Yu T, Liu QY
- 9607** T-cell lymphoblastic lymphoma with extensive thrombi and cardiac thrombosis: A case report and review of literature
Ma YY, Zhang QC, Tan X, Zhang X, Zhang C
- 9617** Perfect pair, scopes unite – laparoscopic-assisted transumbilical gastroscopy for gallbladder-preserving polypectomy: A case report
Zheng Q, Zhang G, Yu XH, Zhao ZF, Lu L, Han J, Zhang JZ, Zhang JK, Xiong Y
- 9623** Bilateral hematoma after tubeless percutaneous nephrolithotomy for unilateral horseshoe kidney stones: A case report
Zhou C, Yan ZJ, Cheng Y, Jiang JH
- 9629** Atypical endometrial hyperplasia in a 35-year-old woman: A case report and literature review
Wu X, Luo J, Wu F, Li N, Tang AQ, Li A, Tang XL, Chen M
- 9635** Clinical features and literature review related to the material differences in thread rhinoplasty: Two case reports
Lee DW, Ryu H, Jang SH, Kim JH
- 9645** Concurrent tuberculous transverse myelitis and asymptomatic neurosyphilis: A case report
Gu LY, Tian J, Yan YP
- 9652** Diagnostic value of contrast-enhanced ultrasonography in mediastinal leiomyosarcoma mimicking aortic hematoma: A case report and review of literature
Xie XJ, Jiang TA, Zhao QY
- 9662** Misidentification of hepatic tuberculosis as cholangiocarcinoma: A case report
Li W, Tang YF, Yang XF, Huang XY

- 9670** Brunner's gland hyperplasia associated with lipomatous pseudohypertrophy of the pancreas presenting with gastrointestinal bleeding: A case report
Nguyen LC, Vu KT, Vo TTT, Trinh CH, Do TD, Pham NTV, Pham TV, Nguyen TT, Nguyen HC, Byeon JS
- 9680** Metachronous squamous cell carcinoma of pancreas and stomach in an elderly female patient: A case report
Kim JH, Kang CD, Lee K, Lim KH
- 9686** Iatrogenic giant pseudomeningocele of the cervical spine: A case report
Kim KW, Cho JH
- 9691** Traditional Chinese medicine for gait disturbance in adrenoleukodystrophy: A case report and review of literature
Kim H, Kim T, Cho W, Chang H, Chung WS

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Takeo Furuya, MD, PhD, Assistant Professor, Department of Orthopaedic Surgery, Chiba University Graduate School of Medicine, Chiba 2608670, Japan. furuya-takeo@chiba-u.jp

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases (WJCC, World J Clin Cases)* is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for *WJCC* as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The *WJCC*'s CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yan-Xia Xing; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Sandro Vento, Bao-Gan Peng

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

November 6, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Observational Study

Vitamin D deficiency, functional status, and balance in older adults with osteoarthritis

Cláudia N Montemor, Marcos Tadeu P Fernandes, Audrey S Marquez, Regina Célia Poli-Frederico, Rubens Alexandre da Silva, Karen B P Fernandes

ORCID number: Cláudia N Montemor 0000-0002-5180-3108; Marcos Tadeu P Fernandes 0000-0003-4494-0187; Audrey S Marquez 0000-0001-7097-4204; Regina Célia Poli-Frederico 0000-0003-4631-4606; Rubens Alexandre da Silva 0000-0001-6879-436X; Karen B P Fernandes 0000-0002-1276-4900.

Author contributions: Montemor CN, Fernandes MTP, da Silva RA, and Fernandes KBP collected the patients' clinical data; Fernandes MTP and Fernandes KBP organized the collected clinical data; Montemor CN, Marquez AS, and Fernandes KBP analyzed the data; Montemor CN and Fernandes MTP wrote the paper; Marquez AS, Poli-Frederico RC, da Silva RA, and Fernandes KBP revised the paper; Poli-Frederico RC and da Silva RA made important contributions to the study performance; Fernandes KBP designed the project and the study.

Supported by National Foundation for Development of Private Higher Education Institutions (FUNADESP)/University Pitágoras UNOPAR, No. 049/15.

Institutional review board statement: This study was approved by the Ethics Committee of the University Pitágoras Unopar

Cláudia N Montemor, Rubens Alexandre da Silva, Karen B P Fernandes, Laboratory of Rehabilitation Research, Doctoral Program of Rehabilitation Sciences, University Pitágoras Unopar (UNOPAR), Londrina 86041-140, Parana, Brazil

Marcos Tadeu P Fernandes, Department of Anesthesiology, Irmandade da Santa Casa de Londrina, Londrina 86010-160, Parana, Brazil

Audrey S Marquez, Health Sciences Research Center, University Pitágoras Unopar (UNOPAR), Londrina 86041-140, Parana, Brazil

Regina Célia Poli-Frederico, Doctoral Program of Rehabilitation Sciences, University Pitágoras Unopar (UNOPAR), Londrina 86041-140, Parana, Brazil

Rubens Alexandre da Silva, Karen B P Fernandes, Département des Sciences de la Santé, Centre intersectoriel en santé durable, Laboratoire de recherche BioNR, Université du Québec à Chicoutimi (UQAC), Saguenay, G7H 2B1, Québec, Canada

Rubens Alexandre da Silva, Centre intégré de santé et services sociaux du Saguenay-Lac-Saint-Jean (CIUSSS SLSJ), Hôpital de La Baie - Services gériatriques spécialisés, Saguenay, Québec, G7H 7K9, Canada

Corresponding author: Karen B P Fernandes, PhD, Associate Professor, Laboratory of Rehabilitation Research, Doctoral Program of Rehabilitation Sciences, University Pitágoras Unopar (UNOPAR), Rua Marselha, 491 Jardim Piza, Londrina 86041-140, Parana, Brazil. karenparron@gmail.com

Abstract**BACKGROUND**

Low vitamin D levels are associated with a more severe case of knee osteoarthritis (OA). However, there are few published reports concerning an association between vitamin D deficiency and functional status of individuals with OA and no reports about postural balance in this population.

AIM

To analyze the relationship between vitamin D deficiency and severity, functional status, and balance in elderly patients with OA.

(No. 070/09).

Informed consent statement: All study participants provided written informed consent prior to study enrollment.

Conflict-of-interest statement: The authors declare having no conflicts of interest.

Data sharing statement: No additional data are available.

STROBE statement: The authors have read the STROBE Statement-checklist and the manuscript was prepared and revised according to the STROBE Statement.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Specialty type: Orthopedics

Country/Territory of origin: Brazil

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

Received: February 4, 2021

Peer-review started: February 4, 2021

First decision: March 31, 2021

Revised: April 15, 2021

Accepted: July 9, 2021

Article in press: July 9, 2021

Published online: November 6, 2021

METHODS

In this cross-sectional study, 105 elderly patients with hip and knee OA were included. The severity was assessed by the Kellgren-Lawrence criteria. The functional status was assessed with the Lequesne index. Postural balance was assessed using a force platform, and center-of-pressure parameters (velocity at anteroposterior and mediolateral axis) were used as the balance outcomes. Serum 25(OH) vitamin D levels were measured using a chemiluminescence method.

RESULTS

Most of the patients (mean age: 70.6 ± 6.5 years) were female ($n = 78$, 74.3%). In the group with vitamin D deficiency, 43 patients (56.6%) had severe OA, while 33 patients (43.4%) had mild or moderate OA (χ^2 test, $P = 0.04$). Patients with vitamin D deficiency showed a higher Lequesne index score (Mann-Whitney test, $P = 0.04$), indicating a worse functional impairment when compared to individuals with normal vitamin D levels. Additionally, patients with vitamin D deficiency had worse postural balance according to the Mann-Whitney test ($P = 0.03$).

CONCLUSION

Vitamin D deficiency is associated with worse severity, functional status, and postural balance in patients with OA.

Key Words: Osteoarthritis; Elderly; Vitamin D; Disability; Severity; Balance

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This study investigated the association between vitamin D deficiency and disease severity, functional status, and postural balance in elderly patients with osteoarthritis. The severity of osteoarthritis was assessed by the Kellgren-Lawrence criteria, and the functional status was evaluated with the Lequesne index. Additionally, postural balance was evaluated using a force platform and center-of-pressure parameters. We found that vitamin D deficiency was associated with worse severity and functional status as well as worse postural balance in elderly patients with osteoarthritis.

Citation: Montemor CN, Fernandes MTP, Marquez AS, Poli-Frederico RC, da Silva RA, Fernandes KBP. Vitamin D deficiency, functional status, and balance in older adults with osteoarthritis. *World J Clin Cases* 2021; 9(31): 9491-9499

URL: <https://www.wjgnet.com/2307-8960/full/v9/i31/9491.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v9.i31.9491>

INTRODUCTION

Osteoarthritis (OA) is the most common degenerative disease of the osteoarticular system. Progression can affect functional status and health-related quality of life, especially in elderly patients[1,2]. It is estimated to affect about 40% of people over 70 years of age, representing a major cause of morbidity and disability worldwide[2]. It is characterized by synovial inflammation, followed by subchondral abnormalities resulting in bone reshaping[3,4]. Genetic factors as well as biomechanical stresses or changes in the articular joint and synovial membrane are important factors in the etiology of OA. Risk factors for OA are obesity, trauma, periarticular injury, and occupational factors[5-8].

Vitamin D plays an important role in calcium homeostasis and mineralization because it controls calcium and parathormone homeostasis with direct effects on bone remodeling[9]. Additionally, there is clear evidence of the benefits over different cellular functions, including muscle mass and strength. Vitamin D receptors have been identified in skeletal muscles. Vitamin D deficiency decreases receptor expression in skeletal muscles and loss of muscle mass in elderly patients[10]. Vitamin D supplementation improves quality of life and physical performance in individuals with OA [11]. On the other hand, vitamin D deficiency is a predictor of reduced muscle strength

P-Reviewer: Patel J
S-Editor: Liu M
L-Editor: Wang TQ
P-Editor: Wang LYT



and mass as well as worse physical performance among men and women within 3 years[12].

There is evidence that vitamin D deficiency is associated with severe knee OA[13, 14]. However, there are contrasting results about this topic in different populations, and the mechanisms between vitamin D and OA are not so clear[15]. Furthermore, there are only a few published reports suggesting a possible association between vitamin D deficiency and functional status of individuals with OA[16,17]. There is also a variety of methods to assess functional status. The Western Ontario and McMaster Universities (WOMAC) index and Lequesne index are the gold standards, but they are not always used. Of concern, there are no studies evaluating the association between vitamin D deficiency in patients with OA and postural balance. Therefore, this study analyzed the association between vitamin D deficiency and functional impairment and balance in elderly patients with OA.

MATERIALS AND METHODS

Ethical procedures

This study was approved by the Research Ethics Committee of the University Pitagoras Unopar. Before any procedure was undertaken, patients received instruction on the nature of the work and signed a free and informed consent form in which they agreed to participate in the study.

Study design and population

One hundred and five elderly patients with OA from a subsample of a project on aging and longevity were included in this cross-sectional study. The project was developed in a mid-sized city in the south region of Brazil (Londrina, Paraná, Brazil), which aimed to examine the social and demographic conditions and health indicators of elderly individuals of this city. The total sample was 508 elderly patients, who were selected in a random and stratified manner from the records of the Basic Health Units from this city.

The criteria for inclusion were patients: (1) of both genders; (2) aged 60 years or older; (3) with a diagnosis of hip and/or knee OA; and (4) who had signed a free and informed consent form. The criteria for exclusion were individuals: (1) with full or partial prosthetics in any of the joints evaluated; (2) with a concomitant diagnosis of other musculoskeletal disorder, such as rheumatoid arthritis, fibromyalgia, systemic erythematosus lupus, and other rheumatic diseases; (3) with post-traumatic or postseptic arthritis; (4) with skeletal dysplasia; (5) who were human immunodeficiency virus-positive; (6) who use corticosteroids or immunosuppressive medications; and (7) who were submitted to arthroplasty (due to the impossibility of functional status assessment). Additionally, those who did not agree to sign the free and informed consent form were also excluded from the study.

General characteristics of the population study

A structured interview guide with questions about gender, age, weight, height, and race was used to characterize the profile of the elderly patients with OA. The formula weight (kg)/height² (m²) was used to calculate the body mass index. Body mass index is classified as normal range (18-24.9 kg/m²), overweight (25-29.9 kg/m²), and obesity (≥ 30 kg/m²).

Evaluation of functional status and severity related to OA

The Lequesne index was used to assess the functional status of these patients. This questionnaire was translated to and validated in Portuguese and is recommended by the European League of Rheumatology to evaluate OA functional status. It is comprised of 11 questions involving pain, discomfort, and functional limitation, specially related to functional status[8,18].

The diagnosis and severity of OA were based on the Kellgren-Lawrence criteria[19] by an experienced radiologist. The severity of OA was determined by the following radiographic findings scale: Grade 0, no radiographic changes; grade I, doubtful space narrowing and possible marginal osteophytes; grade II, possible joint space narrowing and defined osteophytes observed; grade III, defined joint space narrowing, moderate multiple osteophytes, subchondral sclerosis, and any possible deformity of bone contour; and grade IV, remarkable joint space narrowing, severe subchondral sclerosis, definite deformity of bone contour, and presence of large osteophytes.

Postural balance evaluation

Balance assessments were performed by the same trained examiner. It was quantified using a force platform (BIOMECH 400; EMG system do Brasil, SP, Ltda). The participants performed a one-legged stance on the preferred leg, barefoot with their arms at their sides parallel to their trunk and with eyes open, as described elsewhere [20]. The 95% confidence ellipse area of center of pressure (A-CoP in cm²) represents an ellipse area that includes 95% of all CoP positions during a trial. The mean CoP sway velocity (cm/s) in the anteroposterior direction was the parameter used to assess postural balance in this population[20].

Serological quantification of vitamin D

Serum 25 (OH) vitamin D levels were measured using a chemiluminescence kit (Architect iSR2000; Abbott, Chicago, IL, United States). The serum 25 (OH) D level represents the storage form of vitamin D. The Brazilian Society of Endocrinology and Metabolism recommends the classification of vitamin D deficiency as serum 25 (OH)₂D₃ levels of 20 ng/mL or less, vitamin D insufficiency between 20 ng/mL and 29 ng/mL, and vitamin D sufficiency as more than 30 ng/mL. The patients were divided into two subgroups of vitamin D levels: Normal and insufficiency/deficiency[21].

Statistical analysis

The Statistical Package for Social Sciences version 20.0 (IBM Corp., Armonk, NY, United States) was used for statistical data analyses. A CI of 95% and a significance level of 5% ($P < 0.05$) were used. The Mann-Whitney test was used to compare functional status and balance parameters of individuals with OA and insufficiency/deficiency or sufficiency of vitamin D. The χ^2 test was used to assess vitamin D levels and OA severity. The Spearman correlation was used to evaluate vitamin D serum levels and functional status of elderly patients with OA.

RESULTS

In this cross-sectional study, 105 elderly patients with OA were recruited. The study participants were mostly women (74.3%) with a mean age of 70.6 ± 6.5 years. The mean body mass index was 28.0 ± 4.7 kg/m². Thirty-six individuals (34.3%) were smokers (Table 1).

We compared vitamin D levels and severity of OA based on the Kellgren-Lawrence criteria. The data were reorganized into mild to moderate radiographic changes (grades I and II) and severe radiographic changes (grades III and IV). There were significant differences in the prevalence of radiographic OA and vitamin D deficiency. There were more severe cases of OA (56.6%) in elderly patients with vitamin D insufficiency/deficiency ($P = 0.04$; Table 2).

Individuals with vitamin D deficiency or insufficiency had a higher score on the Lequesne index (Mann-Whitney test, $P = 0.04$) when compared to patients with sufficient vitamin D levels (Figure 1). This indicated a worse functional impairment in patients with vitamin D deficiency or insufficiency.

Patients with vitamin D deficiency or insufficiency had a significantly higher velocity value at the anteroposterior axis (median: 3.44 ± 1.79) when compared to patients with sufficient vitamin D levels (median: 2.18 ± 0.1; Mann-Whitney test, $P = 0.03$; Figure 2). Similar results were observed regarding the velocity at the mediolateral axis for patients with vitamin D deficiency or insufficiency (median: 3.86 ± 1.64) when compared to patients with sufficient vitamin D levels (median: 2.71 ± 0.10; Mann-Whitney test, $P = 0.04$; Figure 2).

DISCUSSION

The capacity of humans to produce vitamin D decreases in old age[16]. Furthermore, other risk factors, like malabsorptive diseases, medications that interfere with absorption, and less exposure to sunlight, increase the likelihood of vitamin D deficiency/insufficiency in the elderly population[22]. In this study, vitamin D deficiency/insufficiency impacted the severity of hip and/or knee OA. There were more severe cases in elderly patients with vitamin D deficiency/insufficiency. This data agrees with a previous report where vitamin D deficiency was associated with an advanced stage of knee OA[16]. Additionally, a case-control study found a correlation

Table 1 Characteristics of the population

Variable	Mean	Standard deviation
Age; BMI	70.6; 28.0	6.5; 4.7
Gender; Male; Female; Total	Absolute frequency (n) ; 27; 78; 105	Relative frequency (%); 25.7; 74.3; 100.0
Tobacco use; Yes; No	Absolute frequency (n); 36; 69	Relative frequency (%); 34.3; 65.7

BMI: Body mass index.

Table 2 Association between vitamin D levels and osteoarthritis severity, n (%)

Vitamin D classification	Osteoarthritis severity		
	Mild or moderate	Severe	Total
Deficiency/insufficiency	33 (43.4)	43 (56.6)	76 (100.0)
Normal	19 (65.6)	10 (34.5)	29 (100.0)
Total	52 (49.5)	53 (50.5)	105 (100.0)

Statistically significant, χ^2 test, $P = 0.04$.




Figure 1 Comparison of functional status (analyzed with Lequesne index) of individuals with osteoarthritis in relation to vitamin D classification (deficiency/insufficiency and normal). The boxplots represent the median with interquartile range. The error bars represent minimum and maximum values for each group. ^a Statistically different, Mann-Whitney test, $P < 0.05$.

between the presence of knee osteophytes (indicating increased severity of OA) and a decrease of vitamin D[23].

A possible explanation for this observation is that vitamin D reduces the production of inflammatory cytokines[24,25]. Vitamin D influences cytokines in the synovial fluid that induce hyperalgesia and are related to the breakdown of cartilage. The major mediators are tumor necrosis factor-alpha and interleukin-1 beta, which are proinflammatory cytokines. They are produced by chondrocytes and stimulate the production of proteolytic enzymes, such as metalloproteinases. They also induce the production of other mediators, such as interleukin-6[26].

There are conflicting reports on the association between vitamin D deficiency and OA. A study of 805 Finnish patients with follow-up for 22 years found that vitamin D levels were not associated with the incidence of either knee or hip OA[24]. On the other hand, it was reported that vitamin D deficiency appeared to be associated with an increased risk of knee OA progression[27]. Additionally, the Osteoporosis Fractures in Men Study in the United States found a high prevalence of vitamin D deficiency in hip OA patients and found that these individuals were twice as likely to have hip OA [28]. The literature shows that vitamin D sufficiency can prevent the occurrence of osteoporosis of the femoral neck but not OA of the knee or lumbar spondylolysis[29].

Vitamin D deficiency or insufficiency was related to worse functional status when compared to normal levels in our patient population. Pain is the most prevalent symptom of OA that leads to functional disability. Studies comparing vitamin D levels




Figure 2 Comparison of velocity (cm/s) of individuals with osteoarthritis in relation to vitamin D classification (deficiency/insufficiency and normal). A: Antero-posterior velocity; B: Medio-lateral velocity. The boxplots represent the median with interquartile range. The error bars represent minimum and maximum values for each group. ^a Statistically different, Mann-Whitney test, $P < 0.05$. A/P: Anteroposterior; M/L: Mediolateral.

and pain also have contrasting results[30]. In a trial of 103 knee OA patients that received oral vitamin D supplements (60000 IU per day for 10 d followed by 60000 IU once a month for 12 mo) had slightly higher pain and functional scores compared to those receiving placebos after a follow-up period of 1 year[31]. Joint function in OA is a good predictor of patient quality of life, and decreased joint function has a negative socioeconomic impact. In a study of patients with OA undergoing arthroplasty surgery, vitamin D deficiency was associated with decreased functional status scores. After the surgery, lower functional status scores were observed in patients with vitamin D deficiency, but the difference did not reach statistical significance[22].

The most novel finding of this study is that vitamin D deficiency was associated with balance impairment. CoP parameters from a force platform can directly analyze balance deficits related to proprioception and postural adjustments (feedback and feedforward) of the neuromuscular system[32]. Although the relationship between vitamin D deficiency and muscle weakness is well documented, there is almost no evidence regarding the effects of vitamin D on balance. Pfeifer *et al*[33] found that treatment with 800 IU of vitamin D3 per day reduced body sway in 242 elderly patients when compared with patients who received the placebo. This evidence agrees with our data. However, future studies regarding the optimal dose of vitamin D supplementation for the maximum effect with minimal side effects should be performed.

This study had some limitations. The small sample size and lack of longitudinal assessment were the main limitations of this study. This can increase the difficulty to generalize these findings for other populations and scenarios. Further multicenter studies are necessary to confirm our results as well as to elucidate the association between vitamin D deficiency and joint function and balance. More research to understand the mechanisms of vitamin D deficiency in disease development and progression will further our understanding of using vitamin D as a preventative or treatment.

CONCLUSION

Elderly patients with vitamin D deficiency and OA have worse disease severity, poor functional status, and balance deficit. Future studies with larger sample sizes are necessary to confirm these observations.

ARTICLE HIGHLIGHTS

Research background

Vitamin D deficiency may be related to osteoarthritis (OA) severity and functional status. No previous studies have reported an association between vitamin D deficiency in patients with OA and postural balance.

Research motivation

Our study aimed to explore the severity, functional status, and postural balance in elderly patients with OA and their association with vitamin D levels. The differences might be helpful for proper evaluation and rehabilitation of these patients.

Research objectives

This study aimed to analyze if vitamin D deficiency/insufficiency is correlated to OA severity, functional status, and postural balance in elderly patients.

Research methods

Elderly patients with hip and/or knee OA were included. Serum 25 (OH) vitamin D levels were measured using a chemiluminescence method. The severity was assessed by the Kellgren-Lawrence criteria. The functional status was evaluated with the Lequesne index. Postural balance was assessed using a force platform, and center-of-pressure parameters (velocity at anteroposterior and mediolateral axis) were used as the balance outcomes.

Research results

In this OA population, the patients with vitamin D deficiency had lower functional status than patients with sufficient vitamin D. Additionally, patients with vitamin D deficiency had worse postural balance.

Research conclusions

Vitamin D deficiency is associated with worse severity and functional status. We found for the first time that individuals with OA and vitamin D deficiency have balance impairment.

Research perspectives

We preliminarily explored the relationship between vitamin D levels and functional status and postural balance in elderly patients with OA. Further studies should expand the sample size and adopt a prospective study design to understand the role of vitamin D deficiency and joint function and balance.

REFERENCES

- 1 **Krasnokutsky S**, Attur M, Palmer G, Samuels J, Abramson SB. Current concepts in the pathogenesis of osteoarthritis. *Osteoarthritis Cartilage* 2008; **16** Suppl 3: S1-S3 [PMID: [18723377](#) DOI: [10.1016/j.joca.2008.06.025](#)]
- 2 **Sharma L**, Kapoor D, Issa S. Epidemiology of osteoarthritis: an update. *Curr Opin Rheumatol* 2006; **18**: 147-156 [PMID: [16462520](#) DOI: [10.1097/01.bor.0000209426.84775.f8](#)]
- 3 **Johnson VL**, Hunter DJ. The epidemiology of osteoarthritis. *Best Pract Res Clin Rheumatol* 2014; **28**: 5-15 [PMID: [24792942](#) DOI: [10.1016/j.berh.2014.01.004](#)]
- 4 **Goldring MB**, Otero M, Tsuchimochi K, Ijiri K, Li Y. Defining the roles of inflammatory and anabolic cytokines in cartilage metabolism. *Ann Rheum Dis* 2008; **67** Suppl 3: iii75-iii82 [PMID: [19022820](#) DOI: [10.1136/ard.2008.098764](#)]
- 5 **Harvey WF**, Hunter DJ. Pharmacologic intervention for osteoarthritis in older adults. *Clin Geriatr Med* 2010; **26**: 503-515 [PMID: [20699168](#) DOI: [10.1016/j.cger.2010.03.008](#)]
- 6 **Vina ER**, Kwok CK. Epidemiology of osteoarthritis: literature update. *Curr Opin Rheumatol* 2018; **30**: 160-167 [PMID: [29227353](#) DOI: [10.1097/BOR.0000000000000479](#)]
- 7 **Martel-Pelletier J**, Barr AJ, Cicuttini FM, Conaghan PG, Cooper C, Goldring MB, Goldring SR, Jones G, Teichtahl AJ, Pelletier JP. Osteoarthritis. *Nat Rev Dis Primers* 2016; **2**: 16072 [PMID: [27734845](#) DOI: [10.1038/nrdp.2016.72](#)]
- 8 **Marx FC**, Oliveira LM, Bellini CG, Ribeiro MCC. Tradução e validação cultural do questionário algofuncional de Lequesne para osteoartrite de joelhos e quadris para a língua portuguesa. *Rev Bras Reumatol* 2006; **46**: 253-260
- 9 **Rizzoli R**, Stevenson JC, Bauer JM, van Loon LJ, Walrand S, Kanis JA, Cooper C, Brandi ML, Diez-Perez A, Reginster JY; ESCEO Task Force. The role of dietary protein and vitamin D in maintaining musculoskeletal health in postmenopausal women: a consensus statement from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Maturitas* 2014; **79**: 122-132 [PMID: [25082206](#) DOI: [10.1016/j.maturitas.2014.07.005](#)]
- 10 **Bischoff-Ferrari HA**, Borchers M, Gudat F, Dürmüller U, Stähelin HB, Dick W. Vitamin D receptor expression in human muscle tissue decreases with age. *J Bone Miner Res* 2004; **19**: 265-269 [PMID: [14969396](#) DOI: [10.1359/jbmr.2004.19.2.265](#)]
- 11 **Manoy P**, Yuktanandana P, Tanavalee A, Anomasiri W, Ngarmukos S, Tanpowpong T, Honsawek S.

Vitamin D Supplementation Improves Quality of Life and Physical Performance in Osteoarthritis Patients. *Nutrients* 2017; **9** [PMID: 28933742 DOI: 10.3390/nu9080799]

- 12 **Visser M**, Deeg DJ, Lips P; Longitudinal Aging Study Amsterdam. Low vitamin D and high parathyroid hormone levels as determinants of loss of muscle strength and muscle mass (sarcopenia): the Longitudinal Aging Study Amsterdam. *J Clin Endocrinol Metab* 2003; **88**: 5766-5772 [PMID: 14671166 DOI: 10.1210/jc.2003-030604]
- 13 **Houston DK**, Tooze JA, Neiberg RH, Hausman DB, Johnson MA, Cauley JA, Bauer DC, Cawthon PM, Shea MK, Schwartz GG, Williamson JD, Tylavsky FA, Visser M, Simonsick EM, Harris TB, Kritchevsky SB; Health ABC Study. 25-hydroxyvitamin D status and change in physical performance and strength in older adults: the Health, Aging, and Body Composition Study. *Am J Epidemiol* 2012; **176**: 1025-1034 [PMID: 23118104 DOI: 10.1093/aje/kws147]
- 14 **Boland RL**. VDR activation of intracellular signaling pathways in skeletal muscle. *Mol Cell Endocrinol* 2011; **347**: 11-16 [PMID: 21664245 DOI: 10.1016/j.mce.2011.05.021]
- 15 **Hussain S**, Singh A, Akhtar M, Najmi AK. Vitamin D supplementation for the management of knee osteoarthritis: a systematic review of randomized controlled trials. *Rheumatol Int* 2017; **37**: 1489-1498 [PMID: 28421358 DOI: 10.1007/s00296-017-3719-0]
- 16 **Jansen JA**, Haddad FS. High prevalence of vitamin D deficiency in elderly patients with advanced osteoarthritis scheduled for total knee replacement associated with poorer preoperative functional state. *Ann R Coll Surg Engl* 2013; **95**: 569-572 [PMID: 24165338 DOI: 10.1308/003588413x13781990150374]
- 17 **Mermerci Başkan B**, Yurdakul FG, Aydın E, Sivas F, Bodur H. Effect of vitamin D levels on radiographic knee osteoarthritis and functional status. *Turk J Phys Med Rehabil* 2018; **64**: 1-7 [PMID: 31453483 DOI: 10.5606/tftrd.2018.986]
- 18 **Lequesne MG**. The algofunctional indices for hip and knee osteoarthritis. *J Rheumatol* 1997; **24**: 779-781 [PMID: 9101517]
- 19 **Kellgren JH**, Lawrence JS. Radiological assessment of osteo-arthritis. *Ann Rheum Dis* 1957; **16**: 494-502 [PMID: 13498604 DOI: 10.1136/ard.16.4.494]
- 20 **Oliveira MR**, Vieira ER, Gil AWO, Fernandes KBP, Teixeira DC, Amorim CF, da Silva RA. One-legged stance sway of older adults with and without falls. *PLoS One* 2018; **13**: e0203887 [PMID: 30222769 DOI: 10.1371/journal.pone.0203887]
- 21 **Maeda SS**, Saraiva GL, Kunii IS, Hayashi LF, Cendoroglo MS, Ramos LR, Lazaretti-Castro M. Factors affecting vitamin D status in different populations in the city of São Paulo, Brazil: the São Paulo vitamin D Evaluation Study (SPADES). *BMC Endocr Disord* 2013; **13**: 14 [PMID: 23627369 DOI: 10.1186/1472-6823-13-14]
- 22 **Kupisz-Urbańska M**, Pludowski P, Marciniowska-Suchowierska E. Vitamin D Deficiency in Older Patients-Problems of Sarcopenia, Drug Interactions, Management in Deficiency. *Nutrients* 2021; **13** [PMID: 33920130 DOI: 10.3390/nu13041247]
- 23 **Hunter DJ**, Hart D, Snieder H, Bettica P, Swaminathan R, Spector TD. Evidence of altered bone turnover, vitamin D and calcium regulation with knee osteoarthritis in female twins. *Rheumatology (Oxford)* 2003; **42**: 1311-1316 [PMID: 12867590 DOI: 10.1093/rheumatology/keg373]
- 24 **Barker T**, Martins TB, Hill HR, Kjeldsberg CR, Dixon BM, Schneider ED, Henriksen VT, Weaver LK. Vitamin D sufficiency associates with an increase in anti-inflammatory cytokines after intense exercise in humans. *Cytokine* 2014; **65**: 134-137 [PMID: 24388225 DOI: 10.1016/j.cyto.2013.12.004]
- 25 **Konstari S**, Kaila-Kangas L, Jääskeläinen T, Heliövaara M, Rissanen H, Marniemi J, Knekt P, Arokoski J, Karppinen J. Serum 25-hydroxyvitamin D and the risk of knee and hip osteoarthritis leading to hospitalization: a cohort study of 5274 Finns. *Rheumatology (Oxford)* 2014; **53**: 1778-1782 [PMID: 24817700 DOI: 10.1093/rheumatology/keu178]
- 26 **Goldring MB**, Otero M. Inflammation in osteoarthritis. *Curr Opin Rheumatol* 2011; **23**: 471-478 [PMID: 21788902 DOI: 10.1097/BOR.0b013e328349c2b1]
- 27 **McAlindon TE**, Felson DT, Zhang Y, Hannan MT, Aliabadi P, Weissman B, Rush D, Wilson PW, Jacques P. Relation of dietary intake and serum levels of vitamin D to progression of osteoarthritis of the knee among participants in the Framingham Study. *Ann Intern Med* 1996; **125**: 353-359 [PMID: 8702085 DOI: 10.7326/0003-4819-125-5-199609010-00001]
- 28 **Chaganti RK**, Parimi N, Cawthon P, Dam TL, Nevitt MC, Lane NE. Association of 25-hydroxyvitamin D with prevalent osteoarthritis of the hip in elderly men: the osteoporotic fractures in men study. *Arthritis Rheum* 2010; **62**: 511-514 [PMID: 20112402 DOI: 10.1002/art.27241]
- 29 **Yoshimura N**, Muraki S, Oka H, Nakamura K, Kawaguchi H, Tanaka S, Akune T. Serum levels of 25-hydroxyvitamin D and the occurrence of musculoskeletal diseases: a 3-year follow-up to the road study. *Osteoporos Int* 2015; **26**: 151-161 [PMID: 25138262 DOI: 10.1007/s00198-014-2844-9]
- 30 **Mabey T**, Honsawek S. Role of Vitamin D in Osteoarthritis: Molecular, Cellular, and Clinical Perspectives. *Int J Endocrinol* 2015; **2015**: 383918 [PMID: 26229532 DOI: 10.1155/2015/383918]
- 31 **Sanghi D**, Mishra A, Sharma AC, Singh A, Natu SM, Agarwal S, Srivastava RN. Does vitamin D improve osteoarthritis of the knee: a randomized controlled pilot trial. *Clin Orthop Relat Res* 2013; **471**: 3556-3562 [PMID: 23904246 DOI: 10.1007/s11999-013-3201-6]
- 32 Winter DA. Human balance and posture control during standing and walking. *Gait Posture* 1995; **3**: 193-214 [DOI: 10.1016/0966-6362(96)82849-9]
- 33 **Pfeifer M**, Begerow B, Minne HW, Suppan K, Fahrleitner-Pammer A, Dobnig H. Effects of a long-term vitamin D and calcium supplementation on falls and parameters of muscle function in

community-dwelling older individuals. *Osteoporos Int* 2009; **20**: 315-322 [PMID: 18629569 DOI: 10.1007/s00198-008-0662-7]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

