

Microsoft Bing

国内版 国际版

Chat with Bing

Neoadjuvant chemotherapy for patients with resectable colorectal

ALL IMAGES VIDEOS

85,200 Results Any time

The use of **neoadjuvant chemotherapy** in **resectable colorectal cancer** with **liver metastases** is currently under debate. It is argued that **chemotherapy** in **patients with resectable** disease could eradicate any undetectable **metastases** at the time of diagnosis and prevent further progression.

[Neoadjuvant Treatment for Liver Metastases of Colorectal ...](https://touchoncology.com/colorectal-cancer/journal-articles/neoadjuvant-treatment-for-live...)  
touchoncology.com/colorectal-cancer/journal-articles/neoadjuvant-treatment-for-live...

Was this helpful?

therapeutic age  
One example is  
therapy prior to  
adenocarcinoma  
Neoadjuvant the  
size or extent of  
radical treatment  
making procedure  
succeed and re

### Neoadjuvant chemotherapy for resectable colorectal liver ...

<https://www.sciencedirect.com/science/article/pii/S1479666X14000997>

Apr 01, 2015 · A systematic literature search was performed to identify all studies published from January 2003 up to and including January 2014 regarding patients with initially resectable colorectal liver metastases. Data were examined for information about indications, operation, neoadjuvant and adjuvant therapies, perioperative results, and survival.

Cited by: 20 Author: Giuseppe Nigri, Nicolò Petrucciari, Fabio ...  
Publish Year: 2015

### IPDF! The role of neoadjuvant chemotherapy for resectable ...

<https://www.researchgate.net/profile/Jian-Guo-Zhou...>

Neoadjuvant chemotherapy is being increasingly accepted as an effective treatment of resectable colorectal liver metastases (CRLM), but it may also damage the hepatic parenchyma.

#### PEOPLE ALSO ASK

- What is hepatic resection for metastatic colorectal cancer?
- Is neoadjuvant chemotherapy safe?
- What is the treatment for liver metastases?
- What kind of chemotherapy is used for liver cancer?

Feedback

### (PDF) The role of neoadjuvant chemotherapy for resectable ...

[https://www.researchgate.net/publication/300407748\\_The\\_role\\_of\\_neoadjuvant...](https://www.researchgate.net/publication/300407748_The_role_of_neoadjuvant...)

PDF | Neoadjuvant chemotherapy is being increasingly accepted as an effective treatment of resectable colorectal liver metastases (CRLM), but it may... | Find, read and cite all the research you ...

### Neoadjuvant Treatment for Surgically Resectable Metastatic ...

<https://www.cancernetwork.com/view/neoadjuvant...>

Bosberger F, Albrecht H, Komuruk PC, et al. Neoadjuvant treatment with weekly high-dose 5-fluorouracil as a 24h-infusion, folinic acid and biweekly oxaliplatin in patients with primary resectable liver metastases of colorectal cancer: long-term results of a phase II ...

### Role of neoadjuvant chemotherapy in resectable synchronous ...

<https://onlinelibrary.wiley.com/doi/abs/10.1002/jso.23899>

Apr 09, 2015 · The use of neo-adjuvant chemotherapy in resectable synchronous liver metastasis is ill defined. The aim of this study was to evaluate neo-adjuvant chemotherapy on outcomes following liver resection for synchronous CLM.

Cited by: 25 Author: Glenn K Borney, Chris Coldham, Rene Ada...  
Publish Year: 2015

### Use of Neoadjuvant Chemotherapy Plus Molecular Targeted ...

<https://pubmed.ncbi.nlm.nih.gov/27174607>

Background: Surgery remains the standard of care for patients with colorectal liver metastases (CLMs), with a 5-year survival rate approaching 35%. Perioperative chemotherapy confers a survival benefit in selected patients with CLMs. The use of molecular targeted therapy combined with neoadjuvant chemotherapy for CLMs, however, remains controversial.

Cited by: 14 Author: Dhanusha Sabanathan, Guy D. Eslick, Jenn...  
Publish Year: 2016

### Systematic review of systemic adjuvant, neoadjuvant and ...

<https://europepmc.org/articles/PMC4913134>

Apr 20, 2016 · For patients receiving neoadjuvant chemotherapy, chemotherapy-associated liver injury remains a concern and is associated with a poorer short-term prognosis. 16 Although 20–25% of patients receiving neoadjuvant chemotherapy experience liver-related complications post-surgery, this has not been shown to be significantly different to those receiving surgery-alone. 17, 18 Other factors, such as obesity, may have a greater influence on liver-related ...

Cited by: 62 Author: Emily Khoo, Stephen O'Neill, Ewan Brown, S...  
Publish Year: 2016

### The Role of Neoadjuvant Chemotherapy in Repeat Local ...

<https://www.mdpi.com/2072-6694/13/3/378/htm>

This systematic review and meta-analysis assessed the potential additive value of neoadjuvant chemotherapy before repeat local treatment. The addition of neoadjuvant chemotherapy prior to repeat local treatment was suggested by merely all authors, though supporting evidence is lacking.

Author: Madelon Dijkstra, Sanne Nieuwenhuizen... Publish Year: 2021

### UpToDate

<https://www.uptodate.com/contents/neoadjuvant-chemotherapy-for-resectable-colorectal-liver-metastases>

### See results for

Make a difference for a nonprofit, simply by searching on Bing

MAYBE LATER YES

#### Neoadjuvant therapy

Neoadjuvant therapy is the administration of therapeutic agents before a main treatment. One example is neoadjuvant hormone therapy prior to rs...

#### Search Tools

Turn off Hover Translation (关闭翻译)

Name of Journal: World Journal of Clinical Cases

Manuscript NO: 64374

Manuscript Type: META-ANALYSIS

**Neoadjuvant chemotherapy for patients with resectable colorectal cancer liver metastases: A systematic review and meta-analysis**

Zhang Y *et al.* Meta-analysis of NAC for CRLM patients

Yue Zhang, Long Ge, Jun Weng, Wen-Yu Tuo, Bin Liu, Shi-Xun Ma, Ke-Hu Yang, Hui Cai

**Abstract**

**BACKGROUND**

In recent years, neoadjuvant chemotherapy (NAC) has been increasingly used in patients with resectable colorectal liver metastases. However, the efficacy

Match Overview

1	Crossref 264 words Bin Liu, Long Ge, Jing Wang, Ya-Qiang Chen, Shi-Xun Ma, Pei-Lan Ma, Yun-Qiang Zhang, Ke-Hu Yang, Hui Cai.	5%
2	Internet 73 words crawled on 24-Apr-2021 www.wjnet.com	1%
3	Crossref 67 words "Abstracts", HPB, 2008	1%
4	Internet 54 words crawled on 30-Apr-2020 www.ancotarget.com	1%
5	Internet 50 words crawled on 09-Nov-2020 ro-journal.biomedcentral.com	1%
6	Internet 47 words crawled on 24-Mar-2020 link.springer.com	1%
7	Internet 43 words crawled on 31-Oct-2020 journals.lww.com	1%
8	Internet 43 words crawled on 23-Dec-2020 test.dovepress.com	1%



### Neoadjuvant chemotherapy for resectable colorectal liver ...

<https://www.sciencedirect.com/science/article/pii/S1479666X14000997>

Apr 01, 2015 · A systematic literature search was performed to identify all studies published from January 2003 up to and including January 2014 regarding patients with initially resectable colorectal liver...

Cited by: 21 Author: Giuseppe Nigri, Niccolò Petrucciani, Fabi...  
Publish Year: 2015

### Is neoadjuvant chemotherapy necessary for patients with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4582468>

Dec 15, 2011 · INTRODUCTION. Hepatic resection has only guaranteed long-term survival in patients with colorectal liver metastases (CRLM). 1,2,3 Recent improvements in perioperative care and in...

Cited by: 1 Author: Sang-Yong Son, Nam-Joon Yi, Geun Hon...  
Publish Year: 2011

### [PDF] The role of neoadjuvant chemotherapy for resectable ...

<https://www.researchgate.net/profile/Jian-Guo-Zhou...>

Neoadjuvant chemotherapy is being increasingly accepted as an effective treatment of resectable colorectal liver metastases (CRLM), but it may also damage the hepatic parenchyma.

### Neoadjuvant Treatment for Surgically Resectable Metastatic ...

<https://www.cancernetwork.com/view/neoadjuvant...>

Overview Preoperative Treatment Appr... Discussion References

In the United States, colorectal cancer was predicted to have the fourth highest rate of incidence of new diagnoses among all cancers in 2015, with over 130,000 new cases expected.[1,2] Twenty percent of patients with newly diagnosed colorectal cancers present with metastatic disease. The average 5-year survival for metastatic colorectal cancer is 12.9%. Surgical resection is the only potentially curative option for patients with this disease. However, there are limitations to its use, since less than 15% of pa...

See more on [cancernetwork.com](http://cancernetwork.com)