

**[Guidance on the special care of liver or kidney ...**<https://pubmed.ncbi.nlm.nih.gov/32750019>

The efficacy and tolerability of hidroxychloroquin remains to be also questionable; enroll patients into clinical trial with remdesivir or favipiravir if available. COVID-19 is characterized by virus-induced endothelial dysfunction, procoagulant state and renin-angiotensin-aldosteron system imbalance.

Cited by: 1

Author: Ádám Rempert, Zsuzsanna Gerlei, Orsolya ...

Publish Year: 2020

## Search Tools

Turn on Hover Translation (开启悬停)

**Transplants | COVID-19 Treatment Guidelines**<https://www.covid19treatmentguidelines.nih.gov/special-populations/transplant> ▾

## Introduction

## Transplant Recipients with Covid...

## Treatment of Covid-19 in Transp...



Treating COVID-19 in solid organ transplant (SOT), hematopoietic cell transplant (HCT), and cellular immunotherapy recipients can be challenging due to the presence of coexisting medical conditions, transplant-related cytopenias, and the need for chronic immunosuppressive therapy to prevent graft rejection and graft-versus-host disease.

Transplant recipients may also potentially have increased exposure to severe acute resp...

[See more on covid19treatmentguidelines.nih.gov](https://www.covid19treatmentguidelines.nih.gov)

PRACTICE ADVISE FOR HEPATOLOGY AND

ALL IMAGES VIDEOS

577,000 Results Any time ▾

[Transplants](#) | [COVID-19 Treatment Guidelines](#)<https://www.covid19treatmentguidelines.nih.gov/special-populations/transplant> ▾ See more

## Introduction

## Transplant Recipients with Co...

## Treatment of Covid-19 in Trans...




Treating COVID-19 in solid organ transplant (SOT), hematopoietic cell transplant (HCT), and cellular immunotherapy recipients can be challenging due to the presence of coexisting medical conditions, transplant-related cytopenias, and the need for chronic immunosuppressive therapy to prevent graft rejection and graft-versus-host disease. Transplant recipients may also potentially have increased exposure to SARS-CoV-2 giv...

[See more on covid19treatmentguidelines.nih.gov](#)

20-Aug-2021 10:43PM

6062 words • 59 matches • 5 sources

FINQ

iThenticate

64621\_Auto\_Edited.docx

Quoted Excluded  
Bibliography Excluded  
7%  
0.00000

**Name of Journal:** *World Journal of Hepatology*

**Manuscript NO:** 64621

**Manuscript Type:** MINIREVIEWS

**COVID-19 in liver transplant patients. Clinical and therapeutic aspects**

Carmelo Loinaz-Seguro, Alberto Marcacuzco-Quinto, Mario Fernandez-Ruiz

**Abstract**

The coronavirus disease 2019 (COVID-19) pandemic has profoundly impacted liver transplant (LT) activity across the world, with notable decreases in the number of donations and procedures in most Western countries, in particular throughout the first

Match Overview

1	Internet: 151 words created on 20-May-2021 covid19dataportal.es	2%
2	Internet: 159 words created on 21-Jan-2021 www.aasld.org	2%
3	Internet: 52 words created on 02-Nov-2020 online.library.elsevier.com	1%
4	Internet: 85 words www.nctz.slm.nih.gov	1%
5	Crossref: 43 words Magdalena Menezes, Lucy Mounier, David Morquin, Kaida Khouche et al: "Abnormal liver tests in patients hospitalized ...	1%
6	Crossref: 41 words Shetty S, Reddy, Abdul R, Hameem, Taruqeet Khat, Frances sca Marcon et al: "Tritational Study Exploring the Early Ris	1%
7	Crossref: 58 words Alexandra T. Strauss, Brian J. Boyarsky, Jacqueline M. Gar seuk-Vuong, William Werbel et al: "Liver transplantation in t	1%
8	Crossref: 55 words Liane Rabinowich, Avelot Gruber, Roni Dersuch, Meray De	1%

PAGE: 1 OF 26

Test Only Report

国内版

国际版

COVID-19 in liver transplant patients: Clinical and therapeutic aspe



ALL

IMAGES

VIDEOS

577,000 Results

Any time ▾

## Transplants | COVID-19 Treatment Guidelines

<https://www.covid19treatmentguidelines.nih.gov/special-populations/transplant> ▾

Introduction

Transplant Recipients with C...

Treatment of Covid-19 in Tran...



Treating COVID-19 in solid organ transplant (SOT), hematopoietic cell transplant (HCT), and cellular immunotherapy recipients can be challenging due to the presence of coexisting medical conditions, transplant-related cytopenias, and the need for chronic **immunosuppressive therapy** to prevent graft rejection and **graft-versus-host** disease.

Transplant recipients may also potentially have increased exposure to SARS-CoV-2 giv...

See more on [covid19treatmentguidelines.nih.gov](https://www.covid19treatmentguidelines.nih.gov)

## Liver transplantation in the era of COVID-19

<https://pubmed.ncbi.nlm.nih.gov/32439237>

Liver transplantation in the era of COVID-19. Liver transplantation is considered the ultimate solution for patients with end-stage chronic liver disease or acute liver failure. Patients with liver transplant need special care starting from preoperative preparation, surgical intervention ending with postoperative care.

Cited by: 12

Author: Mohamed El Kassas, Mohamed Alboraie,...

Publish Year: 2020

## Fatal outcome in a liver transplant recipient with COVID-19

<https://pubmed.ncbi.nlm.nih.gov/32277591>

Liver injury is common in patients with COVID-19, but little is known about its clinical presentation and severity in the context of liver transplant. We describe a case of COVID-19 in a patient who underwent transplant 3 years ago for hepatocellular carcinoma. The patient came to clinic with sympto ...

Cited by: 78

Author: Jiao Feng Huang, Kenneth I. Zheng, Jaco...

Publish Year: 2020

## Clinical Characteristics and Outcomes of Liver ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8214214>

Because no effective drugs or therapies were available, countries have adopted different preventive and therapeutic strategies against COVID-19. The Turkish Health Ministry has recommended commencing hydroxychloroquine and/or favipiravir + enoxacin sodium to all patients with the diagnosis of COVID-19