

World Journal of *Gastrointestinal Oncology*

World J Gastrointest Oncol 2022 February 15; 14(2): 369-546



**EDITORIAL**

- 369 Anal human papilloma viral infection and squamous cell carcinoma: Need objective biomarkers for risk assessment and surveillance guidelines
Shenoy S

REVIEW

- 375 Microbiome and colorectal carcinogenesis: Linked mechanisms and racial differences
Tortora SC, Bodiwala VM, Quinn A, Martello LA, Vignesh S
- 396 Gastric epithelial histology and precancerous conditions
Yang H, Yang WJ, Hu B

MINIREVIEWS

- 413 Small bowel adenocarcinoma: An overview
Khosla D, Dey T, Madan R, Gupta R, Goyal S, Kumar N, Kapoor R
- 423 Relation between skeletal muscle volume and prognosis in rectal cancer patients undergoing neoadjuvant therapy
De Nardi P, Giani A, Maggi G, Braga M
- 434 Multimodal treatment in oligometastatic gastric cancer
Chevallay M, Wassmer CH, Iranmanesh P, Jung MK, Mönig SP

ORIGINAL ARTICLE**Basic Study**

- 450 Frankincense myrrh attenuates hepatocellular carcinoma by regulating tumor blood vessel development through multiple epidermal growth factor receptor-mediated signaling pathways
Zheng P, Huang Z, Tong DC, Zhou Q, Tian S, Chen BW, Ning DM, Guo YM, Zhu WH, Long Y, Xiao W, Deng Z, Lei YC, Tian XF
- 478 Comprehensive molecular characterization and identification of prognostic signature in stomach adenocarcinoma on the basis of energy-metabolism-related genes
Chang JJ, Wang XY, Zhang W, Tan C, Sheng WQ, Xu MD

Clinical and Translational Research

- 498 Association and prognostic significance of alpha-L-fucosidase-1 and matrix metalloproteinase 9 expression in esophageal squamous cell carcinoma
Yu XY, Lin SC, Zhang MQ, Guo XT, Ma K, Wang LX, Huang WT, Wang Z, Yu X, Wang CG, Zhang LJ, Yu ZT

Retrospective Study

- 511 Chemotherapy predictors and a time-dependent chemotherapy effect in metastatic esophageal cancer
Midthun L, Kim S, Hendifar A, Osipov A, Klempner SJ, Chao J, Cho M, Guan M, Placencio-Hickok VR, Gangi A, Burch M, Lin DC, Waters K, Atkins K, Kamrava M, Gong J
- 525 Predictive value of serum alpha-fetoprotein for tumor regression after preoperative chemotherapy for rectal cancer
Zhang DK, Qiao J, Chen SX, Hou ZY, Jie JZ

SYSTEMATIC REVIEWS

- 533 Endoscopic ultrasound-guided ablation of solid pancreatic lesions: A systematic review of early outcomes with pooled analysis
Spadaccini M, Di Leo M, Iannone A, von den Hoff D, Fugazza A, Galtieri PA, Pellegatta G, Maselli R, Anderloni A, Colombo M, Siersema PD, Carrara S, Repici A

LETTER TO THE EDITOR

- 543 Prevention of late complications of endoscopic resection of colorectal lesions with a coverage agent: Current status of gastrointestinal endoscopy
Miao YD, Tang XL, Wang JT, Mi DH

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Oncology*, Filippo Lococo, MD, PhD, Assistant Professor, Department of Thoracic Surgery, Catholic University, Rome 00168, Italy. filippo_lococo@yahoo.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Oncology* (WJGO, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, *etc.*

INDEXING/ABSTRACTING

The WJGO is now indexed in Science Citation Index Expanded (also known as SciSearch®), PubMed, PubMed Central, and Scopus. The 2021 edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJGO as 3.393; IF without journal self cites: 3.333; 5-year IF: 3.519; Journal Citation Indicator: 0.5; Ranking: 163 among 242 journals in oncology; Quartile category: Q3; Ranking: 60 among 92 journals in gastroenterology and hepatology; and Quartile category: Q3. The WJGO's CiteScore for 2020 is 3.3 and Scopus CiteScore rank 2020: Gastroenterology is 70/136.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*, Production Department Director: *Xiang Li*, Editorial Office Director: *Ya-Juan Ma*.

NAME OF JOURNAL

World Journal of Gastrointestinal Oncology

ISSN

ISSN 1948-5204 (online)

LAUNCH DATE

February 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Monjur Ahmed, Florin Burada

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

PUBLICATION DATE

February 15, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Endoscopic ultrasound-guided ablation of solid pancreatic lesions: A systematic review of early outcomes with pooled analysis

Marco Spadaccini, Milena Di Leo, Andrea Iannone, Daan von den Hoff, Alessandro Fugazza, Piera Alessia Galtieri, Gaia Pellegatta, Roberta Maselli, Andrea Anderloni, Matteo Colombo, Peter D Siersema, Silvia Carrara, Alessandro Repici

ORCID number: Marco Spadaccini 0000-0003-3909-9012; Milena Di Leo 0000-0003-3524-282X; Andrea Iannone 0000-0002-5468-9515; Daan von den Hoff 0000-0003-3621-226X; Alessandro Fugazza 0000-0003-0485-4903; Piera Alessia Galtieri 0000-0002-3253-6972; Gaia Pellegatta 0000-0003-0235-4905; Roberta Maselli 0000-0001-7291-9110; Andrea Anderloni 0000-0002-1021-0031; Matteo Colombo 0000-0003-0715-8233; Peter D Siersema 0000-0002-6940-8499; Silvia Carrara 0000-0003-4206-9463; Alessandro Repici 0000-0002-1621-6450.

Author contributions: Spadaccini M drafted the manuscript; Spadaccini M, Di Leo M, Iannone A, Fugazza A, Galtieri PA, Pellegatta G, Maselli R, Anderloni A, Colombo M, Siersema PD, Carrara S, and Repici A conducted the analyses and interpretation of the data; Di Leo M, Iannone A, von den Hoff D, Fugazza A, Galtieri PA, Pellegatta G, Maselli R, Anderloni A, Colombo M, Siersema PD, Carrara S, and Repici A made critical revisions related to the important intellectual content of the manuscript; All authors made substantial contributions to the conception and design of the study, acquisition of data, and final approval of the version of the article to be published.

Marco Spadaccini, Milena Di Leo, Alessandro Fugazza, Piera Alessia Galtieri, Gaia Pellegatta, Roberta Maselli, Andrea Anderloni, Matteo Colombo, Silvia Carrara, Alessandro Repici, Endoscopic Unit, Department of Gastroenterology, IRCCS Humanitas Research Hospital, Rozzano 20089, Milan, Italy

Marco Spadaccini, Roberta Maselli, Alessandro Repici, Department of Biomedical Sciences, Humanitas University, Pieve Emanuele 20090, Milan, Italy

Andrea Iannone, Endoscopy Unit, Azienda Ospedale Policlinico, Bari 70124, Italy

Daan von den Hoff, Peter D Siersema, Endoscopy Unit, Radboud University Medical Center, Nijmegen 6525 GA, Netherlands

Corresponding author: Silvia Carrara, MD, Doctor, Professor, Endoscopic Unit, Department of Gastroenterology, IRCCS Humanitas Research Hospital, Via Manzoni 56, Rozzano 20089, Milan, Italy. silvia.carrara@hunimed.eu

Abstract

BACKGROUND

Endoscopic ultrasound-guided radiofrequency ablation (EUS-RFA) is emerging as a complementary therapeutic approach for pancreatic solid masses. However, results of published data are difficult to interpret because of a retrospective design and small sample size.

AIM

To systematically review data on EUS-RFA for solid lesions and to pool the results of the different experiences in order to provide more consistent evidence in terms of safety and efficacy.

METHODS

A comprehensive systematic literature search on the main databases was performed to identify articles in which patients with pancreatic solid lesions underwent EUS-RFA. The primary outcomes were procedure-related adverse events (AEs) and mortality. Secondary outcomes were the technical success rate and the effects on primary tumor growth. Statistical analyses were performed using Stata version 14.0.

Conflict-of-interest statement: All the authors declare that they have no competing interests related to the topic.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Country/Territory of origin: Italy

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Received: February 22, 2021

Peer-review started: February 22, 2021

First decision: April 19, 2021

Revised: April 27, 2021

Accepted: December 31, 2021

Article in press: December 31, 2021

Published online: February 15, 2022

P-Reviewer: Ryoza S

S-Editor: Wu YXJ

L-Editor: Filipodia

P-Editor: Wu YXJ

RESULTS

In total, 14 studies were included, with 120 patients undergoing 153 ablations of 129 solid pancreatic lesions. The STARMED technology was used in seven studies, the Habib system in six studies, and the HybridTherm probe in one study. The pooled technical success rate was 99.0% (P : 25.82%). The pooled overall AE rate was 8.0% (P : 11.46%). Excluding mild AEs, the pooled rates of serious AEs was 1.0% (P : 0%). No mortality related to the procedure was reported.

CONCLUSION

The present pooled analysis confirms the safety and feasibility of EUS-RFA.

Key Words: Pancreatic cancer; Neuroendocrine tumors; Habib; Cryotechnology; Pancreas; Endoscopic ultrasound

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Endoscopic ultrasound-guided radiofrequency ablation (EUS-RFA) is emerging as a complementary therapeutic approach for pancreatic solid masses. However, results of published data are difficult to interpret because of a retrospective design and small sample size. Pooling results across 14 published experiences, the technical success rate was 99.0% among over 150 ablations, with an overall adverse event (AE) rate of 8.0%. Excluding mild AEs, the pooled rate of serious AEs was 1.0%, with no mortality. The present pooled analysis confirms the feasibility and safety of EUS-RFA.

Citation: Spadaccini M, Di Leo M, Iannone A, von den Hoff D, Fugazza A, Galtieri PA, Pellegatta G, Maselli R, Anderloni A, Colombo M, Siersema PD, Carrara S, Repici A. Endoscopic ultrasound-guided ablation of solid pancreatic lesions: A systematic review of early outcomes with pooled analysis. *World J Gastrointest Oncol* 2022; 14(2): 533-542

URL: <https://www.wjgnet.com/1948-5204/full/v14/i2/533.htm>

DOI: <https://dx.doi.org/10.4251/wjgo.v14.i2.533>

INTRODUCTION

Endoscopic ultrasound-guided radiofrequency ablation (EUS-RFA) is emerging as a complementary therapeutic approach for the multidisciplinary treatment of pancreatic solid masses or as a single treatment in case of patients unfit for any other therapy. It has been proposed for pancreatic lesions such as pancreatic ductal adenocarcinoma, pancreatic neuroendocrine tumors (PNETs), or pancreatic cystic neoplasms[1].

The minimally invasive technique and real-time imaging guidance of a probe inside the tumor make EUS the preferred modality for the minimally invasive treatment of lesions in the pancreas. The mechanism of action of RFA is high-frequency alternating current, generating high local temperatures that induce coagulative necrosis of the tissues[2]. Beside thermal damage (direct effect), RFA has also been suggested to also act with an indirect immune modulatory effect[3,4].

Three ablation devices specifically designed for EUS are currently available. The EUS RFA System (STARMED, Koyang, South Korea) consists of an 18 G needle with a monopolar RFA electrode and a VIVA RF generator (STARMED). The device is perfused internally with circulating chilled saline solution that cools the system during the ablation. The Habib EUS monopolar RFA probe (EMcision Ltd., London, United Kingdom, recently purchased by Boston Scientific) is a 1 Fr wire (0.33 mm) that is passed through a 19 G standard EUS needle and is connected to the RITA electrosurgical RF generator (RITA Medical Systems Inc., Fremont, CA, United States). The HybridTherm bipolar flexible probe (ERBE Elektromedizin, Tübingen, Germany) is a 14 G probe that combines bipolar RF ablation with Cryotechnology. It is used in conjunction with the VIO 300 D electrosurgical unit and the ERBECRYO 3 cryosurgical unit.

Current literature[5] suggests that EUS-RFA is feasible and safe; however, the significance of the majority of the studies is hampered by a small sample size and lack



of adequate follow-up. Thus, the aim of this paper was to systematically review data on EUS-RFA for solid lesions and to pool the results of the different experiences in order to have a more reliable point of view.

MATERIALS AND METHODS

The methodology used was in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (commonly known as PRISMA) recommendations[6]. Our systematic review protocol was registered in the International Prospective Register of Systematic Reviews (PROSPERO, www.crd.york.ac.uk/prospero/) on July 2019 (registration No. CRD42020151668).

Data sources and search strategy

A comprehensive systematic literature search was performed to identify full-text articles and abstracts in which patients with pancreatic solid lesions underwent EUS-guided RFA. We performed the literature search in PubMed, EMBASE, and SCOPUS (up to June 11, 2019) electronic databases. PROSPERO was searched for ongoing or recently completed systematic reviews. Electronic searches were supplemented by manual searches of references of included studies and review articles. We identified studies using the following medical subject headings (MeSH) and the keywords “pancreas,” “ablation,” and “EUS.” The search was restricted to the English language. The Medline search strategy was: “((Ultrasound-guided[tiab] OR US guided[tiab]) AND (Endoscop*[tiab] OR Endoscopy[Mesh]) OR (EUS[tiab] OR EUS-guided[tiab] OR Endosonography[Mesh] OR Endoscopic ultrasound[tiab])) AND (Pancreat*[tiab] OR Pancreas[tiab] OR Pancreas[Mesh]) AND (Ablati*[tiab] OR Ablation Techniques[Mesh]).”

Study selection

For the purpose of this systematic review, we included all clinical studies enrolling patients with solid lesions treated by EUS-RFA techniques and reporting the rate of adverse events (AEs) based on the number of procedures. Prospective and retrospective studies, published as full text or presented as abstract at major international meetings (*i.e.* DDW, UEGW) were considered for inclusion. Relevant studies were also searched from reference lists of all identified studies. To be included, studies had to be written in the English language. Studies enrolling patients treated for either cystic or solid pancreatic lesions were excluded if outcomes of interest were not subgrouped according to lesion type. Case reports were excluded.

Two review authors (Spadaccini M; Von den Hoff D) independently screened the titles and abstracts yielded by the search against the inclusion criteria. Full reports were obtained for all titles that appeared to meet the inclusion criteria or where there was any uncertainty. Then review author pairs screened the full text and abstract reports and decided whether these met the inclusion criteria. Disagreements were resolved through discussion among all authors. The reasons for excluding trials were recorded. The review authors were not blinded to the journal titles or to the study authors or institutions. When there were multiple articles for a single study, we used the latest publication and supplemented it, if necessary, with data from the more complete version.

Data extraction

Using standardized forms, two reviewers (Spadaccini M; Von den Hoff D) extracted data independently and in duplicate from each eligible study. Reviewers resolved disagreements by discussion. Unresolved disagreements were resolved by two arbitrators (Carrara S; Repici A). The following data were extracted from each study including the publication status, study design and location, the number of centers involved, the enrollment period, the number of patients, the number of all lesions ablated, the number of procedures, patient characteristics (mean age, gender), mean lesion size, lesion location (head, body or tail), RF technology and needle used, technical success rate, AEs (bleeding, pancreatitis, portal vein thrombosis, ductal stenosis, abdominal pain, deaths), histology of the ablated lesions (*i.e.* adenocarcinoma, neuroendocrine tumor, metastasis, solid pseudopapillary tumor), number of patients with follow-up, mean follow-up period, number of patients with symptomatic success, and radiologic response (complete response [CR], partial response [PR], no response).

Quality assessment

Quality was assessed by the modified Newcastle-Ottawa Scale for non-randomized studies, ranging from 0 (low quality) to 5 (high quality). Two reviewers (Spadaccini M; Von den Hoff D) assessed quality measures for included studies and discrepancies were adjudicated by collegial discussion.

Outcomes measures

The primary outcomes were procedure-related AEs and mortality. Secondary outcomes were technical success rate (feasibility), and the effects on primary tumor growth. In patients with hormone-producing neuroendocrine tumors (NETs), the clinical success rate was also assessed. The outcomes definitions were reported below.

AEs: Pancreatitis, bleeding, portal vein thrombosis, burn of the gastric or duodenal wall, and perforations were regarded as AEs. Any intra- or post-procedural AEs were recorded. We considered mild abdominal pain and asymptomatic ascites as “mild” AEs, while the remaining complication were classified as “serious AEs (SAEs).”

Technical success: The procedure was defined as feasible if the placement of the probe inside the target was successful and it was possible to apply the RF energy for a sufficient time.

Effects on primary tumor growth: The assessment of dimensional response was obtained by contrast-enhanced computed tomography, and/or magnetic resonance imaging, and/or contrast-enhanced EUS. Results were classified as CR (disappearance or complete necrosis), PR (significant decrease or partial necrosis), or failure (minimal decrease or no effect).

Clinical success: In patients undergoing EUS-RFA for hormone-producing NET, the procedure was defined as clinically successful if symptoms were completely resolved.

Statistical analysis

We expressed dichotomous variables as proportion with 95% confidence interval (95% CI) and continuous variables as mean with standard deviation. We calculated pooled estimates with 95% CIs in individual studies using the Freeman-Tukey Double Arcsine Transformation, which stabilizes the variances and allows the inclusion of proportions close to the margins[7]. We selected a random-effects model to summarize data from individual studies. We estimated heterogeneity between studies using the χ^2 test (Cochran Q) and I^2 statistics[8]. I^2 values of 25%, 50%, and 75% were considered low, moderate, and high heterogeneity, respectively. We performed subgroup analyses for the outcomes of success rate, overall AEs and SAEs by publication type and technology used. We estimated differences among subgroups by the Mantel-Haenszel test and heterogeneity in subgroups by the χ^2 test (Cochran Q) and I^2 statistics. We also planned metaregression analyses to estimate the impact of mean pancreatic lesion size, number of lesions localized in the pancreatic head, and number of adenocarcinoma lesions on the outcomes of SAEs and overall AEs. All statistical analyses and graphics were performed using Stata version 14.0 (College Station, TX, United States) using the Metaprop[9] and Metareg[10] commands. For all calculations, a two-tailed P value < 0.05 was considered statistically significant.

RESULTS

Study and patient characteristics


The literature search resulted in 437 articles (Figure 1). After preliminary screening of titles, 22 publications were selected to be reviewed as abstract and full text. Of these, 14 studies[11-24], published between 2012 and 2019, matched the selection criteria and were included in the final analyses (Figure 1). Nine studies (64%) were published as full text, while the remaining five (36%) were presented as abstract at international conferences (UEGW, DDW). Five studies were performed in Europe (54 patients) and the United States (29 patients), three in Asia (19 patients), and the remaining study was from Israel (22 patients). Five studies had a prospective design. All studies but four were single-center experiences. The average Newcastle Ottawa score was 4.7 (range: 3-6). Studies characteristics are summarized in Table 1.

The 14 studies reported outcomes of 120 patients who underwent 153 ablations to treat 129 solid pancreatic lesions. Of all patients, 54.4% were males (56 of 103 patients,

Table 1 Study characteristics

Author	Publication	Publication year	Country	Design	Centers involved	NOS	Patients, n	Lesions, n	Procedures, n	RF technology
Barthet	Full article	2019	France	Prospective	Multi	6	12	14	14	Starmed
Song	Full article	2016	Korea	Retrospective	Multi	4	6	6	6	Starmed
Oleinikov	Full article	2019	Israel	Retrospective	Multi	6	18	25	25	Starmed
Choi	Full article	2018	Korea	Prospective	Mono	6	10	10	16	Starmed
Scopelliti	Full article	2018	Italy	Prospective	Mono	6	10	10	10	Starmed
Crinò	Full article	2018	Italy	Retrospective	Mono	5	8	8	8	Starmed
Lakhtakia	Full article	2016	India	Retrospective	Mono	5	3	3	3	Starmed
Pai	Full article	2015	United Kingdom	Prospective	Multi	5	2	2	3	Habib
Arcidiacono	Full article	2012	Italy	Prospective	Mono	6	22	22	22	Hybrid-terminal
Wang	Abstract	2013	United States	Retrospective	Mono	3	3	3	5	Habib
Sharma	Abstract	2017	United States	Retrospective	Mono	3	2	2	2	Habib
Ramireddy	Abstract	2019	United States	Retrospective	Mono	4	10	10	18	Habib
Malikowski	Abstract	2017	United States	Retrospective	Mono	3	4	4	6	Habib
Guider	Abstract	2018	United States	Retrospective	Mono	4	10	10	15	Habib

NOS: Newcastle Ottawa Scale; RFA: Radiofrequency ablation; RF: Radiofrequency.

**Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow chart.**

from 11 studies) and the mean age was 58.3 ± 9.7 years (provided by 10 studies). The mean lesion size was 24.9 ± 13.3 mm (from 11 studies). Twelve studies reported the location of the treated mass, with the pancreatic head being the most frequent location (66 of 116 lesions, 56.9%), followed by the pancreatic body (39 of 116, 33.6%), and tail (11 of 116, 9.5%). In terms of histology, 68 of 129 (52.7%) lesions were locally advanced adenocarcinomas and 55 of 129 (42.6%) were NETs, of which 15 were

functional tumors, 4 (3.1%) were metastatic lesions, and 2 (1.6%) were solid pseudo-papillary pancreatic neoplasia. The STARmed technology was used in seven studies (76 lesions), the Habib system in six studies (31 lesions), and the HybridTherm probe in one study (22 lesions). Pre-procedural patients and lesions characteristics of each study are shown in [Supplementary Table 1](#).

Technical success

All included studies (14 studies, 120 patients) reported the technical success rate. The procedure was technically successful in 144 of 153 cases. The pooled technical success rate was 99.0% (95%CI: 94.0%-100.0%, I^2 : 25.82%) ([Figure 2](#)). Eleven of fourteen studies (78.6%, 100 procedures) reported a technical success rate of 100%. Arcidiacono *et al*[11] reported a technical success rate of 72.7% (16/22 procedures) with 6 patients in whom the procedure was not feasible because of stiffness of both the gastrointestinal wall and the tumor[11]. In the study by Oleinikov *et al*[18], a technically successful RFA ablation was achieved in 96% of lesions (24/25), based on the response rate visualized by EUS immediately after the procedure. A third study by Malikowski *et al*[17] reported a technically successful procedure in four of six RFAs. We also performed subgroup analysis by publication type and technology as shown in [Supplementary Figures 1 and 2](#), respectively. There was no statistically significant difference between subgroups divided according to publication type ($P = 0.91$) or type of RFA system ($P = 0.54$).

AEs

All of the included studies (14 studies, 120 patients) reported AEs related to the procedure. In 19 of 153 procedures, AEs were seen, yielding an overall pooled AE rate of 8.0% (95%CI: 3.0%-15.0%, I^2 : 11.46) ([Figure 3](#)). However, excluding mild AEs such as mild abdominal pain (10 cases) and asymptomatic ascites (2 cases), in 7 of 153 procedures, SAEs were reported, with a pooled SAEs rate of 1.0% (95%CI: 0.0-5.0; I^2 : 0%). No mortality related to the procedure was reported. Post procedural pancreatitis occurred in 1.0% of cases (5 cases, 95%CI: 0.0-4.0; I^2 : 0%), and one case of post procedural bleeding and one of pancreatic duct stenosis were reported. Safety outcomes of each study are shown in [Supplementary Table 2](#).

We also performed subgroup analysis by publication type and technology. There was no statistically significant difference between subgroups divided according to publication type ($P = 0.39$). In the sensitivity analysis for this outcome, considering only studies published as full text articles, 14% (95%CI: 7.0%-23.0%, I^2 : 0%) of procedures reported AEs ([Supplementary Figure 3](#)). However, excluding mild AEs, a pooled rate of 4.0% (95%CI: 0.0-10.0, I^2 : 0%) was reported (details in [Supplementary Figure 4](#)). No differences were reported among the different RFA systems in terms of safety ($P = 0.65$), considering only SAEs ($P = 0.92$) (details in [Supplementary Figures 5 and 6](#)). On univariate metaregression analysis, lesion size, location, or histology did not affect the risk of developing AEs, as shown in [Supplementary Figures 7-9](#).

Efficacy outcomes

The effect on primary tumor size was reported in nine studies. Across the five studies providing data on ablation of pancreatic NETs, 39 of 45 lesions had a CR, 4 had a PR, and 2 lesions did not respond. In all 15 patients with functional NETs, symptoms were completely resolved after a mean follow-up time ranging from 3 to 13 mo. Considering the four studies reporting outcomes on locally advanced adenocarcinoma or metastasis, 25 of 32 lesions partially responded to ablation and the remaining 7 did not significantly respond. None of these lesions were completely ablated.

DISCUSSION

To the best of our knowledge, this is the first systematic review and pooled analysis evaluating the impact of EUS-guided tissue ablation for the treatment of solid pancreatic lesions. Here, we analyzed 14 different studies that included 120 patients with a total number of 153 EUS-RFA procedures.

Our study confirmed that EUS-RFA had a high safety profile, with an overall pooled rate of 8.0% AEs. However, the pooled rate decreased to 1% when only SAEs were considered.

Furthermore, the present analysis confirmed the high technical success of EUS-RFA with a pooled technical success rate of 99.0%. Nonetheless, the study also demonstrated the lack of robust clinical data regarding the efficacy of the technique




Figure 2 Pooled success rate.




Figure 3 Pooled adverse event rate.

with limited information on long-term outcome. Only nine studies reported data with regard to tumor response, and there was no clear agreement on which imaging technique should be used to confirm the efficacy of the ablation. Moreover, in case of locally advanced adenocarcinoma or metastasis, none of these lesions were completely ablated. Indeed, currently, the absence of SAEs has favored the use of ablation




Figure 4 Endoscopic ultrasound view. Endoscopic ultrasound-guided radiofrequency ablation.

techniques in special clinical settings, such as the local treatment of functioning PNET in patients unfit for surgery. While ablation may be a therapeutic option in these patients to control the overproduction of hormones and resolution of symptoms, its use in small asymptomatic PNETs is questionable. Due to a lack of long-term clinical outcomes, European Neuroendocrine Tumor Society guidelines suggest the use of endoscopic or percutaneous ablative therapy in patients with resectable insulinoma non-fit for surgery (Figure 4), although surgical approach remains the standard cure [25].

Although our systematic literature search is a complete review of all manuscripts published to date, some limitations are present. First, we included different ablation techniques: the STARmed technology in seven studies, the Habib system in six studies and HybridTherm probe in one study. Despite having considered different techniques and different devices, the heterogeneity among the studies was mild to moderate for all outcomes.

Second, all of the included studies are observational studies, which may have inevitably introduced a bias. Thus, the lack of controlled trials shows that the clinical value of local ablation is still insufficiently known. A multicenter randomized controlled trial named Pancreatic Locally Advanced Irresectable Cancer Ablation (PELICAN) (NTR5517) is currently ongoing and will compare chemotherapy and surgical RFA with chemotherapy alone (U.S. National Library of Medicine. Clinical Trials. PELICAN. available online: <https://clinicaltrials.gov/ct2/show/NCT03690323>). The results are expected in 2022 and will probably provide evidence on the additive role of RFA as part of a multimodality approach to treat patients with locally advanced pancreatic cancer.

Finally, none of the studies published to date have evaluated the indirect effects of RFA. In fact, previous results have indicated that besides thermal damage (debulking cytoreductive effect), RFA also seems to have an indirect immune modulatory effect [26,27]. A peripheral zone with sub-lethal injury is often seen around the area of necrosis created by local injury. In this area, oxidative stress and inflammation have been suggested to lead to an indirect antitumoral systemic effect, generally mediating the immune response. Moreover, the local damage causes the release of tumor antigens that can trigger an immune response and stimulate the inflammatory response, acting on distant circulating neoplastic cells. Thus, the combination of ablation techniques with immunotherapy could potentially increase the efficacy of these kind of treatments, although currently only limited experimental evidence has been published[28].

CONCLUSION

In conclusion, we conducted the first pooled analysis on EUS-RFA, confirming its emerging role as a complementary therapeutic approach for pancreatic solid masses with limited risk of serious AEs. Further studies with longer follow-up are necessary to properly evaluate the clinical efficacy as a stand-alone approach or as a part of multimodality management of pancreatic neoplasia.

ARTICLE HIGHLIGHTS

Research background

Endoscopic ultrasound-guided radiofrequency ablation (EUS-RFA) is emerging as a complementary therapeutic approach for pancreatic solid masses.

Research motivation

Results of published data are difficult to interpret because of a retrospective design and small sample size.

Research objectives

The aim of this paper was to systematically review data on EUS-RFA for solid lesions and to pool the results of the different experiences in order to have a more reliable point of view.

Research methods

A comprehensive systematic literature search on the main databases was performed to identify articles in which patients with pancreatic solid lesions underwent EUS-RFA.

Research results

The pooled technical success rate was 99.0% (I^2 : 25.82%). The pooled overall adverse event (AE) rate was 8.0% (I^2 : 11.46%). Excluding mild AEs, the serious AE pooled rate was 1.0% (I^2 : 0%). No mortality related to the procedure was reported.

Research conclusions

The present pooled analysis confirms the safety and feasibility of the EUS-guided RFA.

Research perspectives

Further studies with longer follow-up are necessary to properly evaluate the clinical efficacy as a stand-alone approach or as a part of multimodality management of pancreatic neoplasia.

REFERENCES

- 1 Lakhtakia S, Seo DW. Endoscopic ultrasonography-guided tumor ablation. *Dig Endosc* 2017; **29**: 486-494 [PMID: 28171697 DOI: 10.1111/den.12833]
- 2 Paiella S, Salvia R, Ramera M, Girelli R, Frigerio I, Giardino A, Allegrini V, Bassi C. Local Ablative Strategies for Ductal Pancreatic Cancer (Radiofrequency Ablation, Irreversible Electroporation): A Review. *Gastroenterol Res Pract* 2016; **2016**: 4508376 [PMID: 26981115 DOI: 10.1155/2016/4508376]
- 3 Haen SP, Pereira PL, Salih HR, Rammensee HG, Gouttefangeas C. More than just tumor destruction: immunomodulation by thermal ablation of cancer. *Clin Dev Immunol* 2011; **2011**: 160250 [PMID: 22242035 DOI: 10.1155/2011/160250]
- 4 Chu KF, Dupuy DE. Thermal ablation of tumours: biological mechanisms and advances in therapy. *Nat Rev Cancer* 2014; **14**: 199-208 [PMID: 24561446 DOI: 10.1038/nrc3672]
- 5 Signoretti M, Valente R, Repici A, Delle Fave G, Capurso G, Carrara S. Endoscopy-guided ablation of pancreatic lesions: Technical possibilities and clinical outlook. *World J Gastrointest Endosc* 2017; **9**: 41-54 [PMID: 28250896 DOI: 10.4253/wjge.v9.i2.41]
- 6 Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA; PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* 2015; **350**: g7647 [PMID: 25555855 DOI: 10.1136/bmj.g7647]
- 7 Freeman MF, Tukey JW. Transformations Related to the Angular and the Square Root. *Ann Math Statist* 1950; **21**: 607-611 [DOI: 10.1214/aoms/1177729756]
- 8 Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003; **327**: 557-560 [PMID: 12958120 DOI: 10.1136/bmj.327.7414.557]
- 9 Nyaga VN, Arbyn M, Aerts M. Metaprop: a Stata command to perform meta-analysis of binomial data. *Arch Public Health* 2014; **72**: 39 [PMID: 25810908 DOI: 10.1186/2049-3258-72-39]
- 10 Harbord R, Higgins JPT. Meta-regression in Stata. *Stata J* 2008; **8**: 493-519 [DOI: 10.1177/1536867X0800800403]
- 11 Arcidiacono PG, Carrara S, Reni M, Petrone MC, Cappio S, Balzano G, Boemo C, Cereda S, Nicoletti R, Enderle MD, Neugebauer A, von Renteln D, Eickhoff A, Testoni PA. Feasibility and safety of EUS-guided cryothermal ablation in patients with locally advanced pancreatic cancer. *Gastrointest Endosc* 2012; **76**: 1142-1151 [PMID: 23021160 DOI: 10.1016/j.gie.2012.08.006]

- 12 **Barthet M**, Giovannini M, Lesavre N, Boustiere C, Napoleon B, Koch S, Gasmi M, Vanbiervliet G, Gonzalez JM. Endoscopic ultrasound-guided radiofrequency ablation for pancreatic neuroendocrine tumors and pancreatic cystic neoplasms: a prospective multicenter study. *Endoscopy* 2019; **51**: 836-842 [PMID: [30669161](#) DOI: [10.1055/a-0824-7067](#)]
- 13 **Choi JH**, Seo DW, Song TJ, Park DH, Lee SS, Lee SK, Kim MH. Endoscopic ultrasound-guided radiofrequency ablation for management of benign solid pancreatic tumors. *Endoscopy* 2018; **50**: 1099-1104 [PMID: [29727904](#) DOI: [10.1055/a-0583-8387](#)]
- 14 **Crinò SF**, D'Onofrio M, Bernardoni L, Frulloni L, Iannelli M, Malleo G, Paiella S, Larghi A, Gabbriellini A. EUS-guided Radiofrequency Ablation (EUS-RFA) of Solid Pancreatic Neoplasm Using an 18-gauge Needle Electrode: Feasibility, Safety, and Technical Success. *J Gastrointest Liver Dis* 2018; **27**: 67-72 [PMID: [29557417](#) DOI: [10.15403/jgld.2014.1121.271.eus](#)]
- 15 **Guider JC**, Kannadath BS, Cen P, Rowe J, Wray CJ, Bynon JS, Rahimi EF, DaVee T, Thosani N. Tu1410 Feasibility, safety, and efficacy of endoscopic ultrasound guided radiofrequency ablation (EUS-RFA) of pancreatic ductal adenocarcinoma: A single center us experience. *Gastrointestinal Endoscopy* 2018; **87**: AB585 [DOI: [10.1016/j.gie.2018.04.2263](#)]
- 16 **Lakhtakia S**, Ramchandani M, Galasso D, Gupta R, Venugopal S, Kalpala R, Reddy DN. EUS-guided radiofrequency ablation for management of pancreatic insulinoma by using a novel needle electrode (with videos). *Gastrointest Endosc* 2016; **83**: 234-239 [PMID: [26394384](#) DOI: [10.1016/j.gie.2015.08.085](#)]
- 17 **Malikowski T**, Gleeson FC, Block M, Chari ST, Abu Dayyeh BK, Kendrick ML, Pearson RK, Petersen BT, Smoot R, Truty MJ, Topazian M, Vege SS, Levyet MJ. Mo1268 Endoscopic Ultrasound Guided Radiofrequency Ablation (EUS-RFA): Initial Assessment of Safety and Efficacy. *Gastrointestinal Endoscopy* 2017; **85**: AB484 [DOI: [10.1016/j.gie.2017.03.1117](#)]
- 18 **Oleinikov K**, Dancour A, Epshtein J, Benson A, Mazeh H, Tal I, Matalon S, Benbassat CA, Livovsky DM, Goldin E, Gross DJ, Jacob H, Grozinsky-Glasberg S. Endoscopic Ultrasound-Guided Radiofrequency Ablation: A New Therapeutic Approach for Pancreatic Neuroendocrine Tumors. *J Clin Endocrinol Metab* 2019; **104**: 2637-2647 [PMID: [31102458](#) DOI: [10.1210/je.2019-00282](#)]
- 19 **Pai M**, Habib N, Senturk H, Lakhtakia S, Reddy N, Cicinnati VR, Kaba I, Beckebaum S, Drymoussis P, Kahaleh M, Brugge W. Endoscopic ultrasound guided radiofrequency ablation, for pancreatic cystic neoplasms and neuroendocrine tumors. *World J Gastrointest Surg* 2015; **7**: 52-59 [PMID: [25914783](#) DOI: [10.4240/wjgs.v7.i4.52](#)]
- 20 **Ramireddy S**, Cen P, Rowe J, Wray CJ, Bynon JS, Surabhi V, Patil P, Thosani N. Tu1353 EUS RFA likely offers survival benefits in patients with unresectable pancreatic cancer: results from a single center study with long term follow-up. *Gastrointestinal Endoscopy* 2019; **89**: AB585-AB586 [DOI: [10.1016/j.gie.2019.03.1010](#)]
- 21 **Scopelliti F**, Pea A, Conigliaro R, Butturini G, Frigerio I, Regi P, Giardino A, Bertani H, Painsi M, Pederzoli P, Girelli R. Technique, safety, and feasibility of EUS-guided radiofrequency ablation in unresectable pancreatic cancer. *Surg Endosc* 2018; **32**: 4022-4028 [PMID: [29766302](#) DOI: [10.1007/s00464-018-6217-x](#)]
- 22 **Song TJ**, Seo DW, Lakhtakia S, Reddy N, Oh DW, Park DH, Lee SS, Lee SK, Kim MH. Initial experience of EUS-guided radiofrequency ablation of unresectable pancreatic cancer. *Gastrointest Endosc* 2016; **83**: 440-443 [PMID: [26344883](#) DOI: [10.1016/j.gie.2015.08.048](#)]
- 23 **Wang D**, Jin Z, Lei W, Leung JW, Li Z. Mo1524 Endoscopic Ultrasound Guided Radiofrequency Ablation for the Treatment of Advanced Pancreatic Carcinoma. *Gastrointestinal Endoscopy* 2013; **77**: AB414 [DOI: [10.1016/j.gie.2013.03.272](#)]
- 24 **Sharma NR**, Sharma A, Perelman A, Zelt C. Spare the Whipple: Novel Endoscopic Radiofrequency Ablation Treatment for Insulin Secreting Tumors: 2017 Presidential Poster Award: 2060. *American J Gastroenter* 2017; **112**: S1133-S4 [DOI: [10.14309/0000434-201710001-02061](#)]
- 25 **Falconi M**, Eriksson B, Kaltsas G, Bartsch DK, Capdevila J, Caplin M, Kos-Kudla B, Kwekkeboom D, Rindi G, Klöppel G, Reed N, Kianmanesh R, Jensen RT; Vienna Consensus Conference participants. ENETS Consensus Guidelines Update for the Management of Patients with Functional Pancreatic Neuroendocrine Tumors and Non-Functional Pancreatic Neuroendocrine Tumors. *Neuroendocrinology* 2016; **103**: 153-171 [PMID: [26742109](#) DOI: [10.1159/000443171](#)]
- 26 **Napoletano C**, Taurino F, Biffoni M, De Majo A, Coscarella G, Bellati F, Rahimi H, Pauselli S, Pellicciotta I, Burchell JM, Gaspari LA, Ercoli L, Rossi P, Rugghetti A. RFA strongly modulates the immune system and anti-tumor immune responses in metastatic liver patients. *Int J Oncol* 2008; **32**: 481-490 [PMID: [18202772](#)]
- 27 **Zhao J**, Wen X, Tian L, Li T, Xu C, Melancon MP, Gupta S, Shen B, Peng W, Li C. Irreversible electroporation reverses resistance to immune checkpoint blockade in pancreatic cancer. *Nat Commun* 2019; **10**: 899 [PMID: [30796212](#) DOI: [10.1038/s41467-019-08782-1](#)]
- 28 **Lin M**, Alnaggar M, Liang S, Wang X, Liang Y, Zhang M, Chen J, Niu L, Xu K. An important discovery on combination of irreversible electroporation and allogeneic natural killer cell immunotherapy for unresectable pancreatic cancer. *Oncotarget* 2017; **8**: 101795-101807 [PMID: [29254205](#) DOI: [10.18632/oncotarget.21974](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

