

World Journal of *Psychiatry*

World J Psychiatry 2022 February 19; 12(2): 204-378



Contents

Monthly Volume 12 Number 2 February 19, 2022

EDITORIAL

- 204 Screening dementia and predicting high dementia risk groups using machine learning
Byeon H

REVIEW

- 212 Prenatal nicotine alters development of the laterodorsal tegmentum: Possible role for attention-deficit/hyperactivity disorder and drug dependence
Polli FS, Kohlmeier KA
- 236 Drug-induced stuttering: A comprehensive literature review
Nikvarz N, Sabouri S
- 264 Insights into myelin dysfunction in schizophrenia and bipolar disorder
Valdés-Tovar M, Rodríguez-Ramírez AM, Rodríguez-Cárdenas L, Sotelo-Ramírez CE, Camarena B, Sanabrais-Jiménez MA, Solís-Chagoyán H, Argueta J, López-Riquelme GO

MINIREVIEWS

- 286 Common outcome, different pathways: Social information-processing deficits in autism spectrum disorder and attention-deficit/hyperactivity disorder
Chan JKY, Leung PWL

ORIGINAL ARTICLE

Retrospective Cohort Study

- 298 Associated mortality risk of atypical antipsychotic medication in individuals with dementia
Phiri P, Engelthaler T, Carr H, Delanerolle G, Holmes C, Rathod S

Observational Study

- 308 Reduced paraoxonase 1 activities may explain the comorbidities between temporal lobe epilepsy and depression, anxiety and psychosis
Michelin AP, Maes MHJ, Supasitthumrong T, Limotai C, Matsumoto AK, de Oliveira Semeão L, de Lima Pedrão JV, Moreira EG, Kanchanatawan B, Barbosa DS
- 323 Importance of communication in medical practice and medical education: An emphasis on empathy and attitudes and their possible influences
Steinmair D, Zervos K, Wong G, Löffler-Stastka H
- 338 Cross-sectional study of traumatic stress disorder in frontline nurses 6 mo after the outbreak of the COVID-19 in Wuhan
Zhou ZQ, Yuan T, Tao XB, Huang L, Zhan YX, Gui LL, Li M, Liu H, Li XD

SYSTEMATIC REVIEWS

- 348** Catatonia in older adults: A systematic review

Jaimes-Albornoz W, Ruiz de Pellon-Santamaria A, Nizama-Via A, Isetta M, Albajar I, Serra-Mestres J

SCIENTOMETRICS

- 368** Burnout amongst radiologists: A bibliometric study from 1993 to 2020

Qureshi MFH, Mohammad D, Shah SMA, Lakhani M, Shah M, Ayub MH, Sadiq S

ABOUT COVER

Peer Reviewer of *World Journal of Psychiatry*, Ali Gorji, MD, Professor, Epilepsy Research Center, Münster University, Münster 48149, Germany. gorjial@uni-muenster.de

AIMS AND SCOPE

The primary aim of *World Journal of Psychiatry* (WJP, *World J Psychiatry*) is to provide scholars and readers from various fields of psychiatry with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJP mainly publishes articles reporting research results and findings obtained in the field of psychiatry and covering a wide range of topics including adolescent psychiatry, biological psychiatry, child psychiatry, community psychiatry, ethnopsychology, psychoanalysis, psychosomatic medicine, etc.

INDEXING/ABSTRACTING

The WJP is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, and PubMed Central. The 2021 edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJP as 4.571; IF without journal self cites: 4.429; 5-year IF: 7.697; Journal Citation Indicator: 0.73; Ranking: 46 among 156 journals in psychiatry; and Quartile category: Q2.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yu; Production Department Director: Xu Guo; Editorial Office Director: Jia-Ping Yan.

NAME OF JOURNAL

World Journal of Psychiatry

ISSN

ISSN 2220-3206 (online)

LAUNCH DATE

December 31, 2011

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Rajesh R Tampi, Ting-Shao Zhu, Panteleimon Giannakopoulos

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2220-3206/editorialboard.htm>

PUBLICATION DATE

February 19, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/gerinfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/gerinfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/gerinfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Drug-induced stuttering: A comprehensive literature review

Naemeh Nikvarz, Salehe Sabouri

Specialty type: Psychiatry

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Ahmad DS, Tran MT

Received: February 26, 2021

Peer-review started: February 26, 2021

First decision: July 15, 2021

Revised: July 29, 2021

Accepted: November 25, 2021

Article in press: December 25, 2021

Published online: February 19, 2022



Naemeh Nikvarz, Department of Clinical Pharmacy, Faculty of Pharmacy, Kerman University of Medical Sciences, Kerman 7616911319, Iran

Salehe Sabouri, Neuroscience Research Center, Institute of Neuropharmacology, Kerman 7616911319, Iran

Salehe Sabouri, Department of Pharmaceutical Biotechnology, Kerman University of Medical Sciences, Kerman 7616911319, Iran

Corresponding author: Salehe Sabouri, PharmD, PhD, Assistant Professor, Neuroscience Research Center, Institute of Neuropharmacology, Somayyeh Cross Road, Shariati Street, Kerman 7616911319, Iran. ssabouri@kmu.ac.ir

Abstract

Drug-induced stuttering (DIS) is a type of neurogenic stuttering (NS). Although DIS has not been reported as frequently as other cases of NS in the literature, it is not a negligible adverse drug reaction (ADR) which can significantly affect the quality of life if not treated. This literature review aims to evaluate the epidemiological and clinical characteristics of DIS and suggests some pathophysiological mechanisms for this ADR. Relevant English-language reports in Google Scholar, PubMed, Web of Science, and Scopus were identified and assessed without time restriction. Finally, a total of 62 reports were included. Twenty-seven drugs caused 86 episodes of stuttering in 82 cases. The most episodes of DIS were related to antipsychotic drugs (57%), mostly including clozapine, followed by central nervous system agents (11.6%) and anticonvulsant drugs (9.3%). The majority of the cases were male and between the ages of 31 and 40 years. Repetitions were the most frequent core manifestations of DIS. In 55.8% of the episodes of DIS, the offending drug was withdrawn to manage stuttering, which resulted in significant improvement or complete relief of stuttering in all cases. Based on the suggested pathophysiological mechanisms for developmental stuttering and neurotransmitters dysfunctions involved in speech dysfluency, it seems that the abnormalities of several neurotransmitters, especially dopamine and glutamate, in different circuits and areas of the brain, including cortico-basal ganglia-thalamocortical loop and white matter fiber tracts, may be engaged in the pathogenesis of DIS.

Key Words: Drug; Dysfluency; Dysphemia; Psychotropics; Speech; Stammering

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Stuttering has two main types, developmental and acquired stuttering. Acquired stuttering is a manifestation of psychogenic or neurogenic disorders. Neurogenic stuttering is caused by brain injury, stroke, drugs, *etc.* Because most drugs inducing stuttering are used in the management of psychiatric and/or neurologic disorders, clinicians may merely attribute a new-onset stuttering to the worsening of the underlying disorder and neglect drugs as the causes of stuttering. Therefore, in this review, reports of drug-induced stuttering (DIS) are collected to provide information about epidemiological and clinical characteristics of DIS. Moreover, some pathophysiological changes are proposed as the underlying mechanisms of DIS.

Citation: Nikvarz N, Sabouri S. Drug-induced stuttering: A comprehensive literature review. *World J Psychiatry* 2022; 12(2): 236-263

URL: <https://www.wjgnet.com/2220-3206/full/v12/i2/236.htm>

DOI: <https://dx.doi.org/10.5498/wjp.v12.i2.236>

INTRODUCTION

Speech production is a complex process involving various areas of the brain. Stuttering is a fluency disorder classified as developmental or acquired stuttering. Developmental stuttering, which is mentioned as childhood-onset fluency disorder in the diagnostic and statistical manual of mental disorders, fifth edition[1], often manifests between the ages of 2 and 6 years and spontaneously remits in most cases[2,3]. Acquired stuttering which has a secondary cause can occur in both children and adults. There are two types of acquired stuttering, psychogenic and neurogenic stuttering (NS). NS is caused by the traumatic brain injury, stroke, neurodegenerative disorders like Parkinson's disease (PD) and multiple sclerosis, seizure disorders, drugs, *etc.*[4]. In contrast to the other cases of NS, in which injuries to the brain areas involved in the speech production result in neuroanatomical and neurochemical abnormalities leading to stuttering, in the cases of drug-induced stuttering (DIS), short intervals between the initiation of culprit drug and the occurrence of stuttering and also between the dose reduction or discontinuation of the drug and the relief of stuttering suggest that DIS may be caused merely by neurochemical changes in the brain[5].

Although DIS has not been reported as frequently as other cases of NS in the literature, it is not a rare and negligible adverse drug reaction as Trenque *et al*[6] have reported that 724 individual case safety reports (ICSRs) containing the lowest level terms "stuttering" or "stutter" have been registered in Vigibase, the world health organization international pharmacovigilance database, up to May 31, 2020. The aim of this review is to describe the reported cases of DIS, including their demographic characteristics, medical history, predominant manifestations of stuttering, and the interventions done to manage stuttering and propose some probable pathophysiological mechanisms of this type of NS.

SEARCH STRATEGY

The electronic databases Google Scholar, PubMed, Web of Science, and Scopus were searched by two reviewers without time limitation to find the relevant data. The keywords "stutter", "stuttering", "speech dysfluency", "drug-induced stuttering", "medication-induced stuttering", "psychotropics", "antipsychotics", "antiepileptics", "antiseizure drug", "anticonvulsants", "antidepressants", "clozapine", and "mood stabilizers" were used as search terms. The references of published articles were also examined to find any additional relevant reports. Case reports and case series were included. The reports whose full texts were not available and those being written in non-English language were excluded. According to the above-mentioned inclusion and exclusion criteria, 63 articles reporting DIS in 82 cases were considered in this review.

Twenty-seven drugs caused 86 episodes of stuttering in 82 cases. In four cases, two drugs caused stuttering[8-11]. Of 86 episodes of DIS, 49 (57%) were caused by antipsychotic drugs, followed by 10 (11.6%) by central nervous system (CNS) agents and 8 (9.3%) by anticonvulsant drugs (Table 1). As mentioned above, Trenque *et al*[6] have done a disproportionality analysis using reports registered in Vigibase to estimate the association between exposure to a drug and occurrence of stuttering. Of 22632669 reports registered in this database, 724 ICSRs contained the lowest level term "stuttering" or "stutter". Consistent with our findings, the most reports of stuttering were related to clozapine ($n = 40$), pregabalin ($n = 33$) methylphenidate ($n = 27$), adalimumab ($n = 26$), and olanzapine ($n = 25$)[6]. The results of the disproportionality analysis done by Trenque *et al*[6] showed that the following drugs had the highest reported odds ratio: Methylphenidate (19.57; 95%CI: 13.3-28.7); topiramate (12.48; 95%CI:

Table 1 Number of episodes of drug-induced stuttering reported in the online literature and included in the present review

Therapeutic category	Number of episodes of stuttering, <i>n</i> = 86 (%)
Antipsychotics	
Clozapine	30 (34.9)
Olanzapine	8 (9.3)
Risperidone	4 (4.6)
Aripiprazole	3 (3.5)
Trifluoperazine	1 (1.2)
Chlorpromazine	1 (1.2)
Fluphenazine	1 (1.2)
Levomepromazine	1 (1.2)
Anticonvulsants	
Phenytoin	2 (2.3)
Divalproex	2 (2.3)
Pregabalin	2 (2.3)
Gabapentin	1 (1.2)
Lamotrigine	1 (1.2)
Central nervous system agents	
Methylphenidate	3 (3.5)
Memantine	2 (2.3)
Levodopa	4 (4.6)
Dextroamphetamine and amphetamine salts (Adderall®)	1 (1.2)
Atomoxetine	1 (1.2)
Pemoline	1 (1.2)
Antidepressants	
Sertraline	3 (3.5)
Bupropion	3 (3.5)
Desipramine	1 (1.2)
Bipolar agents	
Lithium	3 (3.5)
Respiratory tract agents	
Theophylline	4 (4.6)
Anxiolytics	
Alprazolam	1 (1.2)
Antineoplastics	
Methotrexate	1 (1.2)
Pyrethrin	1 (1.2)
Total	86 (100)

7.1-22.1); olanzapine (11.98; 95% CI: 8-17.9); golimumab (10.25; 95% CI: 5.5-19.1); clozapine (8.44; 95% CI: 6.1-11.6); and pregabalin (8.36; 95% CI: 5.9-11.9).

DEMOGRAPHIC CHARACTERISTICS OF THE PATIENTS

Twenty-eight (34.6%) cases were female. Most patients were in the age range of 31 to 40 years. Fifteen cases were less than 12 years old. Gender and age of a patient were not reported in one case report[12] (Table 2).

MANIFESTATIONS OF STUTTERING

The primary or core behaviors of stuttering include sound, syllables, and monosyllabic whole-word repetitions, sound prolongations, and speech blocks[4]. The core behaviors were described only in 40 cases. Repetitions ($n = 42$) were the most frequent primary behavior, followed by blockages ($n = 21$) and sound prolongations ($n = 16$). Twenty-one cases had more than one type of primary behavior (Table 3).

Individuals with developmental stuttering and persistent developmental stuttering (PDS) have anxiety related to stuttering[4]. However, patients with acquired stuttering may be annoyed by stuttering but do not experience anxiety. None of the reports included in this review described the psychological presentations of patients when producing a dysfluent speech.

Individuals with developmental stuttering[4,13] as well as patients with NS[13-15] often develop secondary behaviors including eye blinking, facial grimacing, interjections, word or phrase substitution, *etc.* The secondary behaviors are acts that are learnt in long term to cope with stuttering[13]. In the cases included in the present review, the secondary behaviors related to the stuttering were not reported. It may not be unusual because in comparison to the developmental stuttering, secondary behaviors are less prominent in the acquired stuttering[13]. Furthermore, we believe that since the development of the secondary behaviors requires a sufficient period, it is unlikely that patients with DIS have enough time to present these behaviors because interventions such as dose reduction or discontinuation of the culprit drug are carried out as soon as possible, which result in the complete relief or significant improvement of stuttering.

MANAGEMENT

Therapeutic measures that resulted in significant improvement or complete relief of stuttering were drug withdrawal in 48 (55.8%) episodes, dose reduction in 18 (21%), addition of a new drug in three (3.6%), addition of a new drug besides dose reduction of the culprit drug in three (3.5%), and slow dose titration in two (2.3%). Moreover, in three (3.5%) episodes, the stuttering spontaneously remitted despite continuation of the offending drug with no dose reduction[16-18]. However, three (3.5%) episodes did not respond to the dose reduction[19-21]. For four (4.6%) episodes, the authors did not describe the actions taken to manage stuttering[22-25]. In two (2.3%) cases with PDS and PD, stuttering was exacerbated during levodopa-on periods, and levodopa was not discontinued[26,27] (Table 3).

PATHOPHYSIOLOGY

In this section, the abnormalities suggested in the pathogenesis of developmental stuttering and PDS are described. Then, we propose some mechanisms for DIS based on the underlying impairments involved in the pathogenesis of PDS and developmental stuttering.

For producing a fluent speech, orofacial as well as respiratory muscles must work properly. Different areas of the brain including several parts of the cerebral cortex and subcortical structures such as the basal ganglia (BG) and cerebellum are involved in speech production. Functional abnormalities of each part of these areas may cause speech dysfluency. It has been suggested that dysfunction in the different parts and networks of cerebral cortex engaged in speech production[28], the impairments in the neural connections between the cerebral cortex, the BG, and the thalamus, which are called cortico-BG-thalamocortical circuit (CBTC)[28,29], and/or the dysfunction of the BG are involved in the pathogenesis of developmental stuttering[28,29].

CBTC AND STUTTERING

Several neural circuits are engaged in the process of speech production. One of these circuits that has received much attention in the pathogenesis of stuttering is CBTC[28,29]. The BG consist of input, intrinsic, and output nuclei. The input nuclei receive information from different parts of the brain, especially the cerebral cortex, the thalamus, and the substantia nigra, and send signals to the intrinsic nuclei for further processing and then to the output nuclei. The output nuclei relay signals to the thalamus which sends signals back to the part of the cerebral cortex from which the primary signal was

Table 2 Demographic characteristics and clinical history of patients

Characteristic/history ¹	n (%)
Gender	
Female	28 (34.6)
Male	53 (65.4)
Age (yr)	
< 12	15 (18.5)
12-20	3 (3.7)
21-30	15 (18.5)
31-40	16 (19.7)
41-50	13 (16)
51-60	10 (12.3)
> 60	9 (11.1)
History of speech dysfluency	11 (13.4)

¹The total number of the patients is 82, but the age and gender of one patient was not reported.

originated. The input nuclei consist of the putamen and the caudate nucleus, collectively named the striatum. The intrinsic nuclei consist of the external segment of the globus pallidus (GPe), the subthalamic nucleus (STN), and the substantia nigra pars compacta (SNpc). The internal segment of the globus pallidus (GPi) and the substantia nigra pars reticulata (SNpr) make the output nuclei[30].

The striatum has two types of neurons including gamma-aminobutyric acid-ergic (GABAergic) medium-sized spiny neurons (MSNs) representing 90%-95% of the striatal neurons and GABA-ergic/cholinergic interneurons. The striatum receives inputs from different parts of the brain including excitatory glutamatergic inputs from different parts of the cerebral cortex, motor cortex, supplementary motor area (SMA), pre-SMA, auditory cortex, somatosensory cortex, and the thalamus and dopaminergic inputs from SNpc[30]. BG match motor, sensory, and cognitive context received from different parts of the cerebral cortex[31] and send signals to the thalamus to stimulate the cerebral cortex to initiate the desired movement, *e.g.*, terminating production of the previous phoneme and initiating the next phoneme in the case of speaking, and inhibit competing movements[32]. The cerebral cortex sends signals through corticobulbar tracts to the orofacial and respiratory muscles including the muscles of larynx, pharynx, lips, and tongue[33], activates the proper group of muscles, and inhibits competing muscles whose activation interferes with the production of the desired movement.

The signals that are received by the input nuclei of BG are transmitted *via* two pathways: Direct and indirect. The activation of the direct pathway stimulates the cerebral cortex and therefore activates the right motor program while the activation of the indirect pathway inhibits the cerebral cortex and therefore, all other competing motor programs[2,28].

In the direct pathway, MSNs, which have N-methyl-D-aspartate (NMDA) glutamate receptors, D1 dopamine receptors, A1 adenosine receptors, and M4 muscarinic receptors[34] and release gamma-aminobutyric acid (GABA), substance P, and dynorphin[35], project to the SNpr and GPi, the output nucleus of BG. MSNs are stimulated by glutamatergic projections from the cerebral cortex. The activated MSNs release GABA, which inhibits the output nucleus. GABAergic projections from the output nucleus inhibit the glutamatergic neurons of the thalamus. Therefore, the inhibition of the output nucleus by MSNs disinhibits thalamic neurons which ultimately activate the cerebral cortex and increase locomotor activity[35] (Figure 1).

MSNs in the indirect pathway, which have NMDA glutamate receptors, D2 dopamine receptors, A2A adenosine receptors, and M1 muscarinic receptors[34] and release GABA and enkephalin, project to the output nucleus through the GPe and STN. The GPe has GABAergic neurons that inhibit STN neurons which are glutamatergic cells. The release of glutamate because of the activation of the cerebral cortex stimulates MSNs in the indirect pathway to release GABA. GABA inhibits the GPe and therefore disinhibits the STN. Release of glutamate from the STN stimulates the inhibitory GABAergic neurons of the output nucleus, which inhibits thalamic neurons and therefore cerebral cortex, resulting in decreased locomotor activity (Figure 1). It has been proposed that the direct and indirect pathways have interactions with each other, and their integration and balanced activation during movement selection is required for the proper execution of motor programs[36,37].

Table 3 Summary of the case reports of drug-induced stuttering

Offending drug (dosage)	Patients' gender/age (yr)	Main indication of drug administration	Concomitant medications (dosage)	Onset/aggravation of stuttering	Primary behaviors	Concomitant symptoms	Management, response	Recurrence of stuttering after medication resumption	Concomitant disorders	Ref.
Adderall XR® (20 mg/d)	Male/10	ADHA	No other drugs	Within two weeks after the initiation of Adderall XR®	Single word and syllable repetitions and audible/silent sound prolongations	Increased tic behaviors, increased levels of social anxiety and communication related frustration	DC of Adderall XR® and start of atomoxetine (10 mg/d), significant reduction of stuttering	NR	Developmental stuttering, Tourette Syndrome, allergies, chronic ear infections, frequent phonic and motor tics	Donaher <i>et al</i> [82]
Alprazolam (1 mg)	Female/22	Anxiety and depression	No other drugs	Shortly after increasing the dose	Not restricted to initial syllables, occurred on small grammatical words and substantive words, persisted during singing, not associated with secondary symptomatology such as facial grimacing or fist clenching	A right carotid bruit and a grade II/IV systolic murmur without a click, Minimal late systolic mitral valve prolapse and mild stenosis of both internal carotid arteries	DC of alprazolam, complete relief after two days	Within one hour after a single morning dose of 0.5 mg alprazolam, stuttering started, then 10 to 12 h later it was stopped. Stuttering did not happen with placebo	No history of speech dysfluency	Elliott <i>et al</i> [83]
Aripiprazole (2 mg/d)	Male/8	ADHD combined-type	Atomoxetine (25 mg/d)	After 10 d of starting aripiprazole	NR	NR	DC of Aripiprazole, complete relief	NR	Developmental stuttering	Ünay <i>et al</i> [84]
Aripiprazole (10 mg/d)	Male/11	Mild intellectual disability	No other drugs	4 wk after increasing the dose to 10 mg/d	NR	Addition of clonazepam 0.75 mg/d, no improvement. Reduction of aripiprazole dose to 5 mg, complete relief over 10 d	NR	Increasing the dose to 10 mg resulted in re-emergence of stuttering which responded to DC of aripiprazole	No history of speech dysfluency	Naguy <i>et al</i> [85]
Atomoxetine (started at 25 mg/d and gradually increased to 40 mg/d)	Male/14	ADHD	No other drugs	Three weeks after the initiation of atomoxetine	NR	NR	Dose reduction to 25 mg/d, no improvement. DC of atomoxetine and initiation of methylphenidate, complete relief of atomoxetine-induced stuttering and	NR	Developmental stuttering since the age of 7 yr. ADHD predominantly inattentive	Cicek <i>et al</i> [86]

							considerable reduction of developmental stuttering				
Bupropion (SR) 150 mg BID	Female/59	Major depressive disorder	No other drugs	Four days after starting the drug	Sound prolongations, silent blocking, word production with excess physical tension, and monosyllabic whole-word repetitions. The stuttering was anxiogenic and restricted to initial syllables	Slight finger dysdiadochokinesia	DC of bupropion, complete relief of stuttering after 2 d	NR	No history of speech dysfluency	Fetterolf <i>et al</i> [78]	
Bupropion SR (300 mg/d)	Male/38	Major depressive disorder	No other drugs	Two days after increasing the dosage from 150 to 300 mg/d	Involuntary silent pauses or blocks, repetitions, prolongations of sounds, syllables, and words, affected rhythm of speech	NR	DC of bupropion, complete relief of stuttering	Re-administration of bupropion 150 mg after 1 wk caused stammering, and the drug was stopped immediately	A history of occasional smoking, no history of speech dysfluency	Bhatia <i>et al</i> [79]	
Bupropion XL (300 mg/d)	Male/53	Depression	No other drugs	After increasing the dosage of Bupropion	Difficulty starting words and repetition of syllables	NR	Administration of 5 mg oral haloperidol, stuttering was improved after 3 h and completely relieved after 7 h	Medication was continued	No history of speech dysfluency	McAllister <i>et al</i> [80]	
Clozapine (up to 400 mg/d)	Female/32	Paranoid schizophrenia	No other drugs	4 wk after the initiation of clozapine	NR	Pharyngeal dystonia and buccolingual and facial dyskinesia associated with laryngeal dystonia	DC of clozapine, complete relief after 5 d	Clozapine was reintroduced at 100 mg/d. All symptoms reoccurred and relieved by clozapine cessation	History of neuroleptic-induced parkinsonism but not concomitant with dysarthria, no history of speech dysfluency	Thomas <i>et al</i> [63]	
Clozapine (was initiated at 400 mg/d and gradually increased to 900 mg/d)	Female/28	schizoaffective disorder	No other drugs	Shortly after the initiation of clozapine at 400 mg/d and not relieved during the gradual increase in the dosage to 900 mg/d	NR	NR	Dosage reduction to ≤ 700 mg/d, complete relief	The dose was not increased again	No history of speech dysfluency	Ebeling <i>et al</i> [87]	
Clozapine (450-750 mg/d)	Female/49	Psychosis	No other drugs	Stuttering was initiated when the clozapine dosage was	NR	Generalized seizure followed by myoclonic jerks of	The addition of phenytoin and then sodium valproate and	Clozapine was continued at 600 mg/d in addition to sodium	History of neuroleptic-induced acute	Supprian <i>et al</i> [59]	

				increased to 700 mg/d		her arms at the clozapine dosage of 750 mg/d	the reduction of clozapine dosage to 600 mg/d, complete relief	valproate 900 mg/d with no recurrence of stuttering	dystonia, no history of speech dysfluency	
Clozapine (300 mg/d)	Male/28	Paranoid schizophrenia	No other drugs	Stuttering was initiated when the dosage of clozapine was increased from 150 mg to 300 mg/d and worsened with further increases in the clozapine dosage	NR	Generalized tonic colonic seizure at 425 mg/d along with the increased severity of stuttering	The reduction in the dosage of clozapine to 200 mg/d and addition of sodium valproate, significant improvement	The clozapine dosage was increased to 300 mg/d, but stuttering was not reoccurred albeit in the presence of sodium valproate 800 mg/d	No history of speech dysfluency	Duggal <i>et al</i> [64]
Clozapine (300 mg/d)	Male/57	Schizoaffective disorder	Lithium (900 mg/d), sodium valproate (600 mg/d)	Four days after the initiation of clozapine	NR	NR	Dose reduction and DC of clozapine, complete relief after 7 d	NR	History of alcohol dependency, diabetes mellitus, no history of speech dysfluency	Bar <i>et al</i> [15]
Clozapine (up to 500 mg/d)	Not mentioned	Schizophrenia	No other drugs	A few days after the initiation of clozapine at 300 mg/d	NR	Myoclonic jerks at night and facial tics	Addition of sodium valproate, significant improvement, reducing the dosage of clozapine from 500 to 300 mg/d, complete relief	Clozapine was not discontinued	No history of speech dysfluency	Begum <i>et al</i> [11]
Clozapine (700 mg/d)	Female/33	Schizophrenia	No other drugs	After reaching the daily dose to 700 mg (interval was not reported)	NR	Facial tics, seizure (seizure was initiated after the occurrence of stuttering)	Reduction in the dosage of clozapine to 600 mg/d, remarkable improvement, addition of sodium valproate to control seizure, no effect on stuttering	Clozapine was not discontinued	No history of speech dysfluency	Hallahan <i>et al</i> [58]
Clozapine (300 mg/d)	Female/34	Schizophrenia	No other drugs	2 wk after the initiation of clozapine	NR	Orofacial dyskinesia	Clozapine dosage reduction to 50 mg/d, complete relief	Clozapine was not discontinued	No history of speech dysfluency	Hallahan <i>et al</i> [58]
Clozapine (50-125 mg)	Male/62	Delusional disorder	No other drugs	NR	Unsustained phonation, hesitation, irregular articulatory break down, sound repetition (not related to any specific sound, occurred at irregular word positions)	Orofacial dyskinesia, laryngeal and pharyngeal tardive dystonia, harsh and strangled voice	Addition of tetrabenazine, patient could not tolerate the clozapine dosages more than 100 mg/d, DC of clozapine, complete relief	Clozapine was not restarted	No history of speech dysfluency	Lyall <i>et al</i> [9]

Risperidone and then clozapine (450 mg/d and 75 mg/d)	Male/55	Schizophrenia	No other drugs	NR	Occasional blocking, prolongation on word-initial sounds and repetitions of speech elements including one-syllable words at the beginning of his speech utterances	Stammering and unusual limb and trunk movements related to risperidone, belching, persistent hiccupping, worsening of the facial tic, and the orofacial dyskinesia involving the lips and tongue related to clozapine	Risperidone-induced stuttering; NR, the first episode of clozapine-induced stuttering, dose reduction to 125 mg/d and cessation of clozapine; significant improvement and complete relief of stuttering; the second time of clozapine-induced stuttering: addition of sodium valproate, considerable improvement	Clozapine was restarted at 75 mg/d, recurrence of stuttering, the addition of sodium valproate, 600 mg/d, significant improvement in the stuttering	History of head injury resulting in problems with executive functioning and a significant discrepancy, between the patient's verbal IQ and performance IQ, making various clicking noises and blowing sounds when speaking before the initiation of antipsychotic drugs	Lyall <i>et al</i> [9]
Clozapine (up to 600 mg/d)	Male/35	Schizotypal personality disorder	No other drugs	At clozapine dosage of 250 mg/d and progressive worsening with dose escalation	NR	NR	Reducing the dosage of clozapine to 200 mg/d, complete relief	Clozapine was continued at 200 mg/d without causing stuttering	History of trifluoperazine-induced truncal dystonia, no history of speech dysfluency	Krishnakanth <i>et al</i> [88]
Clozapine (200 mg/d)	Male/24	Paranoid schizophrenia	No other drugs	After increase in the dosage of clozapine to 200 mg/d	NR	NR	DC of clozapine, complete relief	Clozapine was not restarted, amisulpiride was started and did not cause stuttering	No history of speech dysfluency	Krishnakanth <i>et al</i> [88]
Clozapine (250 mg/d)	Male/23	Paranoid schizophrenia	No other drugs	At clozapine dosage of 250 mg/d (interval was not reported)	NR	NR	Clozapine dosage reduction to 150 mg/d, complete relief	Clozapine was not discontinued	History of neuroleptic-induced tardive dyskinesia, no history of speech dysfluency	Krishnakanth <i>et al</i> [88]
Clozapine (350 mg/d)	Male/15	Undifferentiated schizophrenia	Clomipramine (225 mg/d)	Three years after the initiation of clozapine and clomipramine	Repetitions of syllables and transient accelerations of speech rate	Involuntary paroxysmal perioral movements, facial tic-like movements, myoclonic jerks of the upper limbs, GTC seizure	Addition of valproic acid at 500 mg/d, complete relief of stuttering within days	Clozapine was continued with valproic acid without reoccurrence of seizure and speech dysfluency during 2 yrs of follow-up	Symptoms of obsessive-compulsive disorder, no history of epilepsy or speech dysfluency	Horga <i>et al</i> [66]
Clozapine (up to 250 mg/d)	Male/29	Undifferentiated schizophrenia	No other drugs	After the clozapine dosage titration from 137.5 mg/d to 150 mg/d	Frequent repetition and prolongation of syllables or words with frequent hesitations, blocking and	No focal dystonia or any evidence of seizure-like activity	Reducing and splitting the dose of clozapine to 50 mg in morning and 75 mg at night, improvement of stuttering	Reoccurrence of stuttering at clozapine dosage of 250 mg/d, improvement of stuttering after dose reduction to 225 mg/d, a later increase in the	History of antipsychotic-induced extrapyramidal symptoms, no history of speech dysfluency	Grover <i>et al</i> [61]

				pauses				dosage to 300 mg/d did not cause recurrence of stuttering		
Clozapine (400 mg/d)	Female/33	Severe MDD with psychotic features	No other drugs	Stuttering was started after increasing the dosage of clozapine to 400 mg/d and worsened when the dosage was increase to 450 mg/d	Excessive prolongation of syllables or words	Sialorrhea	Addition of benztropine, no improvement. Reduction of the dosage of clozapine to 350 mg/d, complete relief	Stuttering recurred 16 d after increasing the clozapine dosage to 400 mg/d, but completely relieved after dosage reduction to 300 mg/d	None	Kumar <i>et al</i> [89]
Clozapine (up to 650 mg/d)	Male/32	Paranoid schizophrenia	Sertraline (300 mg/d), lamotrigine (500 mg/d), haloperidol (4 mg/d), clonazepam (1 mg/d)	Noticeable stuttering at clozapine dosages of ≥ 600 mg/d	Expressive speech dysfluency with hesitancy and frequent pauses	Involuntary twitching of muscles of jaw	Clozapine dose reduction by 25 mg, improvement of stuttering	Clozapine was not discontinued	No history of speech dysfluency	Murphy <i>et al</i> [20]
Clozapine (400 mg/d)	Male/43	Schizoaffective disorder	Paroxetine (20 mg/d)	Stuttering became noticeable when the clozapine daily dose was increased to more than 350 mg	Expressive speech dysfluency	NR	Clozapine dose reduction by 50 mg, improvement of stuttering	Clozapine was not discontinued	No history of speech dysfluency	Murphy <i>et al</i> [20]
Clozapine (450 mg/d)	Male/33	Paranoid schizophrenia	No other drugs	Stuttering was developed during the initiation and dose titration of clozapine	Intermittent stuttering of speech	NR	Reducing the rate of dose titration, improvement of stuttering	Clozapine was not discontinued	No history of speech dysfluency	Murphy <i>et al</i> [20]
Clozapine (up to 300 mg/d)	Female/46	Delusional disorder	No other drugs	Stuttering was developed during the initiation and dose titration of clozapine	Hesitancy with specific syllables	Orofacial dyskinesia	Clozapine dose reduction to 50 mg, improvement of stuttering	Clozapine was not discontinued	No history of speech dysfluency	Murphy <i>et al</i> [20]
Clozapine (325 mg/d)	Male/67	Schizoaffective disorder	Duloxetine (60 mg/d), hyoscine (30 mg/d), aripiprazole (10 mg/d)	Stuttering was developed during the initiation and dose titration of clozapine	Expressive speech dysfluency	Orofacial twitching, upper limb jerking, hypersalivation	Reducing the rate of clozapine dose titration, improvement of stuttering	The clozapine dose was increased again to control psychotic symptoms, but nothing about the recurrence of stuttering was reported	Hearing impairment, hypertension	Murphy <i>et al</i> [20]
Clozapine (650 mg)	Female/63	Paranoid schizophrenia	Amisulpride 200 mg/d, amitriptyline 25 mg/d, paroxetine 20 mg/d, zopiclone 3.75 mg/d	Stuttering was developed on a stable dose of clozapine	Expressive speech dysfluency with hesitancy	NR	Reducing the dose of clozapine by 50 mg, no improvement	Clozapine at 650 mg/d was recommenced, but authors did not report its effects on the recurrence of stuttering	No history of speech dysfluency	Murphy <i>et al</i> [20]
Clozapine (100 mg), aripiprazole (7.5	Female/21	Schizophrenia	No other drugs	At clozapine dosage of 100 mg/d and	NR	NR	Reduction of the dose of clozapine and	The drugs were not discontinued	Turner syndrome, no history of	Ertekin <i>et al</i> [8]

mg/d)				aripiprazole dosage of 7.5 mg/d			addition of aripiprazole (5 mg/d), complete relief. Reduction of the dose of aripiprazole from 5 to 7.5 mg/d, complete relief		speech dysfluency	
Clozapine (gradually increased to 450 mg/d)	Male/16	Schizoaffective disorder	Citalopram (NR), clonazepam (NR), atenolol (NR), lithium (NR)	Approximately 22 d after increasing the clozapine dosage to 400 mg/d	Persistent stuttering (difficulties with the pronunciation of letters "L," "D," and "T")	Orofacial dyskinesia with perioral twitching (started at clozapine dosage of 350 mg/d), microseizure according to EEG (at clozapine dosage of 400 mg/d)	Substituting lithium with divalproex sodium, improvement in stuttering 4 wk after receiving divalproex sodium at 500 mg BID	Clozapine was not discontinued because of its considerable therapeutic effects	History of type 1 DM, DKA with episodic hallucinations, GERD, cerebral contusion, occasional cocaine use, anxiety- induced intermittent stuttering, family history of stuttering	Rachamalla <i>et al</i> [62]
Clozapine (up to 600 mg/d)	Female/22	Schizophrenia	Fluoxetine (60 mg/d)	Stuttering was developed after the clozapine dose escalation to 300 mg/d	NR	NR	Reduction in the clozapine dose and initiation of ECT, minimal improvement	Clozapine was not discontinued	NR	Das <i>et al</i> [19]
Clozapine (450 mg/d)	Man/in early 40s	NR	No other drugs	After increasing the clozapine daily dose from 400 mg to 450 mg	NR	Marked increase in seizure activity	DC of clozapine, nothing was clearly reported by the authors	NR	NR	Kranidiotis <i>et al</i> [24]
Clozapine (200 mg/d)	Male/38	Schizophrenia	No other drugs	Stuttering was evident at 200 mg/d and became so disabling at 350 mg/d	NR	NR	Dose reduction of clozapine and addition of amisulpiride and BDZ, reduction of stuttering, DC of clozapine, complete relief	Clozapine was not restarted	NR	Kranidiotis <i>et al</i> [24]
Clozapine (300 mg BID)	Male/57	Paranoid schizophrenia	Risperidone, IM injection (37.5 mg every 2 wk), Risperidone, oral (1.5 mg/d which increased to 2 mg BID on admission)	Two days after admission (the dosage of clozapine, 300 mg BID, was not changed on admission)	NR	Orofacial and extremities myoclonic jerks, drop attacks	Clozapine dosage reduction to 100 mg BID, resolution of stuttering within two days	The patient was discharged on clozapine 150 mg BID, but author reported nothing about stuttering at follow-up	History of COPD, hypertension, DM, and chronic back pain, cigarette smoking	Chochol <i>et al</i> [60]
Clozapine (125 mg/d)	Male/29	Schizophrenia	No other drugs	A few days after titrating the clozapine dosage to 125 mg/d	Frequent repetitions of words that included broken words	NR	Reducing the clozapine dosage to 100 mg/d, significant improvement	Clozapine dosage was not re-escalated	No history of speech dysfluency	Nagendrappa <i>et al</i> [90]

Clozapine (up to 200 mg/d)	female/25	Schizophrenia	No other drugs	At clozapine dosage dose of 150 mg/d (interval was not mentioned)	NR	Tonic-clonic epileptic seizure	DC of clozapine and start of amisulpiride and biperiden, complete relief of stuttering and seizure	Clozapine was not rechallenged	No history of speech dysfluency	Gica <i>et al</i> [65]
Divalproex sodium (600 mg/d)	Male/45	Affective instability and irritability	Citalopram (30 mg/d), promazine (100 mg/d)	Four days after initiation of divalproex sodium	Sound repetitions and prolongations (not restricted to the initial syllable and caused pronounced difficulty in starting and completing his sentences)	NR	DC of divalproex, complete relief after 3 d	Divalproex sodium was not restarted	A 10-yr history of post-traumatic stress disorder and alcoholism, no history of speech dysfluency	Aukst-Margetić <i>et al</i> [91]
Divalproex sodium (1500 mg/d in divided dose)	Male/56	Bipolar affective disorder	Olanzapine (10 mg/d), lorazepam (4 mg/d, gradually stopped along with increase in the dose of divalproex)	Two weeks after increasing the dosage of divalproex from 1000 to 1500 mg/d	A moderately pressured speech, articulation of speech, alterations in intensity and timings of utterance segments, Involuntary repetitions and prolongations of sounds, syllables, words or phrases, involuntary silent pauses or blocks	NR	DC of divalproex, instant amelioration of the stuttering	Re-initiation of the drug after one week caused resurgence of symptoms, so the drug was stopped	No history of speech dysfluency	Mukherj <i>et al</i> [92]
Desipramin (300 mg/d)	Male/28	Dystimia, primary type, major depression	Doxepin (50 mg at bed time)	Two months after increasing the dosage of desipramine	Stuttering with difficulty in articulation	Minimal dryness of mouth before stuttering, myoclonic jerking (twitching movements around his jaw) concomitant with stuttering	DC of both drugs, complete relief after 48 h	Twenty-four hours after restarting both drugs stuttering happened again, the desipramin dosage was decreased to 250 mg/d, but stuttering was persisted occasionally, on 4 different occasions, desipramin was discontinued and stuttering was solved within 24-48 h; an increase in the doxepin dosage to 200 mg at bed time had not resulted in stuttering	Opiate and alcohol dependence in remission, retinal detachment and ruptured disc and chronic back pain in the past, no history of speech dysfluency	Masand <i>et al</i> [93]
Fluphenazine (up to 50 mg/d)	Male/35	Schizophrenia	Benztropine mesylate (4 mg/d)	12 d after increasing fluphenazine dosage to 50 mg/d	NR	EPS	Dosage reduction to 30 mg/d, complete relief	Increasing the dosage of fluphenazine to 40 mg/d caused stuttering	No history of speech dysfluency	Nurnberg <i>et al</i> [10]

										recurrence
Gabapentin (NR)	Female/58	Intractable seizure	Phenytoin (NR)	NR	NR	NR	DC of gabapentin, relief after 4 d	NR	No history of speech dysfluency	Nissani <i>et al</i> [94]
Lamotrigine (up to 5 mg/kg/d)	Female/5	BECTS	Valproic acid (30 mg/kg/d)	Stuttering was initiated after increasing the dosage of lamotrigine to 5 mg/kg/d	NR	Frequent diurnal absence seizures, poor concentration and forgetfulness, clumsiness and poor coordination, emotional lability, dysarthria, and slurred speech	DC of lamotrigine, speech improvement in a couple of days	Lamotrigine was not rechallenged	NR	Catania <i>et al</i> [95]
Levodopa/carbidopa (100/25 mg TID)	Male/44	PD	NR	Patient had a history of PDS, and stuttering was exacerbated during on periods, 1 h after levodopa/carbidopa intake	NR	Dyskinesia during drug-on phases and akinesia, bradykinesia, resting tremors, and rigidity in drug-off phases	The severity of stuttering return to baseline during levodopa-off periods	Levodopa was not discontinued	PDS	Anderson <i>et al</i> [25]
Levodopa (200 mg/d)	Male/72	PD	None	Nearly one month after increasing the dosage to 200 mg/d	NR	Palilalia, speech freezing	DC of levodopa and initiation of pramipexole, return to the baseline level of dysfluency	Reinitiating levodopa caused stuttering	Speech dysfluency due to PD	Louis <i>et al</i> [22]
Levodopa (up to 1000 mg/d)	Male/42	PD	Pergolide (1.5 mg/d), quetiapine (50 mg at bed time)	After increasing the levodopa dosage to 300 mg/d	Pressured speech and sound repetition	Palilalia, speech freezing	NR	NR	None	Louis <i>et al</i> [22]
Levodopa 600 mg/d	Male/57	PD	Cabergoline (4 mg/d), selegiline (10 mg/d), amantadine (300 mg/d)	Patient had a history of PDS, and stuttering was exacerbated during on phases after levodopa consumption	Speech repetitions and speech blocks	Speech problems associated with PD including hypokinetic dysarthria and hypophonia occurred during levodopa-off phases	Severity of stuttering return to baseline during levodopa-off periods	Levodopa was not discontinued	PDS	Burghause <i>et al</i> [26]
Levomepromazine (50 mg at bed time)	Male/65	Bipolar disorder	Quetiapine (NR), valproate semisodium (NR), zolpidem, moxonidin (NR), propafenone (NR), insulin (NR)	Five days after the initiation of levomepromazin	NR	NR	DC of Levomepromazin, complete relief three days later	Levomepromazin was not recommended	History of drug induced EPS, supraventricular tachycardia, type 2 DM, HTN, and mild cognitive impairment	Margetic <i>et al</i> [96,97]
Lithium (1200 mg at bed time)	Male/48	Bipolar affective disorder	Fluoxetine (20 mg/d)	One month after the initiation of lithium	Worsening his developmental stuttering, a repetitive word	Lightheadedness, hand tremor	Tapering off lithium, stuttering returned to baseline within a few weeks	Valproic acid (2750 mg/d) was started instead of lithium	PDS, depression	Netski <i>et al</i> [98]

					stutter that severely limited his verbal communication ability					
Lithium (900 mg twice daily)	Male/10	Bipolar disorder	Risperidone (4 mg bed time), clonidine (0.1 mg 3 times daily), melatonin (3 mg at bed time), famotidine (20 mg BID)	Two days after increasing the dose of lithium, stuttering was worsened	Syllable repetitions, occurred only at the beginning of sentences	NR	Dose adjustment of lithium to 600 mg in the morning and 900 mg at night, stuttering returned to baseline after 2 d	Lithium was not discontinued	History of developmental stuttering, bipolar disorder not otherwise specified, ADHD, and conduct disorder	Gulack <i>et al</i> [99]
Lithium (the dose was not mentioned, but lithium was used for a long time)	Female/86	Bipolar disorder	Donepezil (NR), primidone (NR), risperidone (NR)	After a chronic use of lithium, stuttering was started and stayed for 3 more mo. The lithium level was elevated (2.0 mmol/L)	Starting a few words fluently, then repeating syllables and words and terminating the sentence abruptly	NR	DC of lithium, complete relief of stuttering after two weeks	Lithium was not restarted	Past medical history of dementia and epilepsy, no history of speech dysfluency	Sabillo <i>et al</i> [100]
Memantine (10 mg/d)	Male/9	Autistic disorder	No other drugs	After increasing the dose	Deterioration of primary behaviors of developmental stuttering including sound repetition, and sound prolongation on first and middle vowels, and difficulty for starting to speak. His parents explained that the child could only start to speak after a deep and audible breath	NR	Reduction of memantine dosage to 7.5 mg/d, improvement of acquired stuttering after several days. DC of memantine, stuttering was reduced to baseline after 3 wk	Risperidone was used instead	Developmental stuttering	Alaghband-Rad <i>et al</i> [17]
Memantine (5 mg/d)	Male/4	Autism	No other drugs	After increasing the dose	The difficulty was with the start of the speech and the child could only start to speak after a deep and audible breath	NR	The drug was continued at the same dose as the difficulty was tolerable, and gradually was increased to 7.5 mg/d, relief of speech difficulty	Medication was continued, and its dose was gradually increased	No history of speech dysfluency	Alaghband-Rad <i>et al</i> [17]
Methotrexate (cumulative dose of 62.5 mg, IT)	Female/22	Pre-B acute lymphoblastic leukemia	NR	After achieving cumulative dose of 62.5 mg (26 d after	NR	Dysphasia progressed to aphasia, mild	Three months after initiation of symptoms (no intervention was	NR	No history of speech dysfluency	Shuster <i>et al</i> [21]

				initiating IT MTX)		headache, low-grade fever, behavioral problems	described)			
Methylphenidate (10 mg/d)	Male/7	ADHD	No other drugs	10 d after the initiation of the drug	Sound prolongations, silent blocking, word production with excess physical tension, monosyllabic whole-word repetitions	NR	DC of methylphenidate, speech returned to normal after 1 wk	Atomoxetine was used instead	No history of speech dysfluency	Alpaslan <i>et al</i> [101]
Methylphenidate (5 mg in the morning and 5 mg at noon)	Male/7	ADHD	No other drugs	One day after drug initiation	Troubles during the pronouncing the first syllables and repetitions of some syllables	NR	DC of Methylphenidate, improvement after 10 d	Methylphenidate was restarted at 10 mg in the morning and 5 mg at noon. After 10 d, stuttering was returned	NR	Copur <i>et al</i> [102]
Methylphenidate (2.5 mg BID) and pemoline (9.375 mg/d) after DC of methylphenidate	Girl/3	Pervasive hyperactivity	None	Three days after starting methylphenidate, four days after starting pemoline	Repetition of the first syllable of word which gradually worsened	NR	DC of methylphenidate, relief of stuttering, DC of pemoline, relief of stuttering	Methylphenidate and pemoline were not restarted	NR	Burd <i>et al</i> [7]
Olanzapine (15 mg/d)	Male/56	Depression	Intrathecal morphine (7.5 mg/d), clomipramine (225 mg/d)	Four days after the initiation of clozapine	Constant word repetition (acquired)	NR	DC of olanzapine, complete relief after two days	NR	Chronic pain syndrome, no history of speech dysfluency	Bar <i>et al</i> [15]
Olanzapine (7.5-10 mg/d)	Male/72	Psychotic depression	Clomipramine (50-150 mg/d)	3 wk after the initiation of olanzapine	Repetition and retention of first syllables and prolongation of phonemes	NR	DC of olanzapine, complete relief after 5 d	NR	Brain cortical atrophy, no history of speech dysfluency	Bar <i>et al</i> [15]
Olanzapine (5 mg/d)	Female/36	Manic episode	Sodium valproate (300 mg/d), prednisolone (75 mg/d)	7 d after the initiation of olanzapine	Repetition of syllables and words	NR	DC of olanzapine, complete relief after 4 d	NR	Ulcerative colitis and celiac disease, no history of speech dysfluency	Bar <i>et al</i> [15]
Olanzapine (10 mg/d)	Female/43	Schizophrenia	No other drugs	Approximately 21 d after the initiation of olanzapine	Repetition of first syllables and word prolongation	NR	DC of olanzapine, complete relief after 3-5 d	NR	Mild cluttering at the age of 19	Bar <i>et al</i> [15]
Olanzapine (2.5 mg/d)	Female/51	Depression	Sertraline (100 mg/d), promethazine (50 mg at night); both was started 14 wk before initiation of	14 d after the initiation of olanzapine	Blocking of speech and prolongation of phonemes	NR	Increase in olanzapine dose to 5 mg/d, relief of stuttering during the next weeks	Olanzapine was not discontinued	Symmetrical cerebellar hypoplasia and generalized cortical atrophy, no history of	Bar <i>et al</i> [15]

			olanzapine						speech dysfluency	
Olanzapine (10 mg/d)	Male/42	Schizophrenia	Zopiclone (7.5 mg/d)	Two days after the initiation of olanzapine	Difficulty in articulating words properly	NR	DC of olanzapine, complete relief after two days	NR (patient was not followed-up)	A fall without loss of consciousness 2 d before initiation of stuttering, no history of speech dysfluency	Bar <i>et al</i> [15]
Olanzapine (10 mg/d)	Male/42	Paranoid ideation	Venlafaxine (150 mg/d), promazine (200 mg/d)	Four days after the initiation of olanzapine	Repetition and retention of first syllables and prolongation of phonemes	NR	DC of olanzapine, complete relief after two days		PTSD, adjustment disorders, no history of speech dysfluency	Lasic <i>et al</i> [103]
Olanzapine (10 mg/d)	Male/21	Psychotic disorder	No other drugs	Three days after the initiation of olanzapine	disturbance in the fluency and time patterning of speech, repetition of sounds and syllables, blocking between words	NR	DC of olanzapine and start of quetiapine, complete relief after three days	Olanzapine was not restarted	No history of speech dysfluency	Asan <i>et al</i> [104]
Phenytoin (200 mg/d)	Male/42	Seizure due to head injury	No other drugs	Shortly after initiation of phenytoin	Predominantly part-word repetitions and prolongation	Abnormality of speech muscle fine motor control	Addition of CBZ and gradual DC of phenytoin, sustain decrease in the frequency of dysfluencies and improved motor performance	Phenytoin was not restarted	No history of speech dysfluency	Mcclean <i>et al</i> [105]
Phenytoin (20 mg/kg LD and 5 mg/kg/d MD)	Male/3	GTC seizure due to head trauma	No other drugs	10 d after the initiation of phenytoin	NR	NR	DC of phenytoin and initiation of sodium valproate, complete relief 10 d after DC of phenytoin	Phenytoin was not rechallenged	No history of speech dysfluency	Ekici <i>et al</i> [106]
Pregabalin (75 mg twice daily)	Female/31	Complex regional pain syndrome	No other drugs	After taking the second dose of pregabalin on the first day	A slurred speech	NR	DC of pregabalin, complete relief after one week	Pregabalin was not restarted	No history of speech dysfluency	Giray <i>et al</i> [107]
Pregabalin (75 mg twice daily)	Female/68	Herpes zoster	Acyclovir (800 mg five times daily)	Three days after the initiation of pregabalin	NR	Frequent blepharospasm	DC of pregabalin; alleviated of symptoms after four days and complete relief after one week	A 75 mg pregabalin capsule consumption after 4 wk resulted in stuttering and frequent blepharospasm	No history of speech dysfluency	Ge <i>et al</i> [108]
Pyrethrin product containing 0.33% pyrethrum extract and 4% piperonyl butoxide (3 times	Female/2 (the child's mother, who was breastfeeding	Repeated episodes of head lice	No other drugs	Two days after the last period of mother's treatment	An acute onset of stuttering especially at the initiation of the speech	An increase in clumsiness, slight erythematous rash of approximately 3 cm × 2 cm on the	Six weeks postexposure	Pyrethrin was not repeated	No history of speech dysfluency	Hammond <i>et al</i> [81]

over a period of 12 d left on the scalp for 10 min)	her at least one time per day, were receiving this topical product)						occiput of the scalp			
Risperidone (4 mg/d, then 8 mg/d)	Male/32	Aggravated psychotic disorder	Lorazepam (1 mg/d)	Stuttering was initiated after the dose increase to 4 mg/d, and worsened 16 d after the dose increase to 8 mg/d	Severe sound repetitions and interjections in a way that it was difficult to understand his words	Slight akathisia-like symptoms such as anxiety and restlessness (not prominent)	No action, stuttering diminished 23 d later	He continued taking risperidone at 8 mg/d with only a slight stuttering	A 10-yr history of Schizophrenia. His friend during junior high school was a stutter, and the patient used to mimic his stuttering. He began stuttering at that time for 1 yr	Lee <i>et al</i> [16]
Risperidone (4 mg)	Female/48	Psychosis	Lorazepam (1 mg PRN), procyclidine (5 mg BID for treatment of EPS)	11 d after taking risperidone	Repetitions in the speech, pausing within a word and her speech, an excess of physical tension in the speech	NR	A little decrease in risperidone dose, a bit reduction in stuttering	Risperidone was not discontinued	No history of speech dysfluency	Yadav <i>et al</i> [18]
Risperidone (at a dose of 1 mg/d for 2 yr)	Male/21	Behavioral disorder	No other drugs	After chronic treatment with low-dose of risperidone	Prolongation of sounds, hearable blocks, repetitions of single-syllable words	NR	No action, stuttering was decreased to a minimal level after 17 d	Risperidone was not discontinued	Moderate mental retardation because of perinatal asphyxia, no history of speech dysfluency	İnci <i>et al</i> [23]
Sertraline (100 mg daily)	Male/36	Major depression	Alprazolam (0.25 mg 3 TID)	Two weeks after increasing the dosage from 50 to 100 mg/d	Normal vocabulary, decreased rate of speech, normal tone, interrupted words	NR	DC of sertraline, speech problem resolved after one day	Medication was not restarted. Later, administering phenelzine, imipramine, and fluoxetine caused milder speech hesitancy	No history of speech dysfluency	Makela <i>et al</i> [109]
Sertraline (50 mg daily)	Female/32	Recurrent depression	No other drugs	During the third week of starting the drug, stuttering occurred and worsened over a 3-d period	Difficulty in starting and completing the sentences	Feeling nervous, increased restlessness, and insomnia two days before the onset of stuttering	DC of sertraline, complete relief of stuttering after 3 d	Previously, patient has received sertraline and experienced stuttering, so discontinued the medication. Medication was not restarted. Desipramine was started and did not cause stuttering	No history of speech dysfluency	Christensen <i>et al</i> [76]

Sertraline (150 mg daily)	Female/22	Bulimia nervosa, anorexia nervosa, posttraumatic stress disorder, recurrent depression, panic disorder	Clonazepam (0.5 mg QID), trimethoprim-sulfamethoxazole (BID)	One week after increasing the dosage of sertraline	NR	Hyperreflexia and mild tremulousness, generalized muscle twitching (myoclonus), restlessness, and mild confusion	DC of sertraline and Antibiotic, gradual normalization of speech over two to three days	Seven days after restarting sertraline at 50 mg/d, stuttering and other symptoms returned, then the drug was discontinued	No history of speech dysfluency	Brewerton <i>et al</i> [77]
Theophylline (200 mg BID to 100 mg QID)	Male/the age of the onset of theophylline-induced stuttering was not reported, but it surely occurred when he was between 1.5 and 4 yr old	Asthma	Nothing was clearly mentioned	The patient only experienced stuttering during the autumn when he was receiving theophylline for the management of asthma attacks	Repeating whole words, six or seven times usually at the beginnings of the sentences, no dysfluency while singing	Being tense, having insomnia, and be frustrated by his speech problem	DC of theophylline at the end of autumn before age 4 yr, complete relief of stuttering. Changing the dosage from 100 mg QID to 200 mg BID at age 4 yr, complete relief after 7 d with no recurrence of stuttering	The patients had stuttered each time that he was on Theophylline regimen 200 mg BID	No history of speech dysfluency	Rosenfield <i>et al</i> [110]
Theophylline (130 mg TID and sometimes QID)	Female/6.5	Asthma	Metaproterenol sulphate (PRN)	Within a few days after increasing the theophylline dosage to 130 mg TID	Multiple repetitions of the word "I", especially at the beginning of sentences, she could speak better when speak more slowly. Stuttering was worse when she was excited	NR	DC of theophylline, complete relief within two days	Resumption of theophylline resulted in the recurrence of stuttering which responded to drug withdrawal. Several months after the discontinuation of theophylline, the drug was resumed without causing any dysfluency	No history of speech dysfluency	Rosenfield <i>et al</i> [110]
Theophylline 200 mg BID to 200 mg TID	Male/4 yr and 3 mo	Asthma	Beclomethasone dipropionate and Theo-Dur sprinkle (200 mg BID) (at age 4 yr and 4 mo). Addition of metaproterenol sulphate, isoetharine HCL and atropine (at age 4 yr and 10 mo, DC of all drugs and initiation of cromolyn capsules (20 mg TID) (at age 5 yr)	Nine months after the initiation of theophylline	Repeating "ah, ah, ah" in the middle of sentences, stuttering was worse when he was excited	Anxiety, sleep problems	Withdrawal of theophylline at age 5 yr, complete relief within two weeks	After complete relief of stuttering, the patient only received theophylline during asthma attacks and experienced no stuttering	No history of speech dysfluency	Rosenfield <i>et al</i> [110]
Theophylline (400 mg BID)	Male/73	A long-standing chronic obstructive lung	Steroids and ranitidine as well as being on oxygen	One month after the introduction of theophylline	An intense tonic-clonic stuttering without any	NR	DC of theophylline, stuttering was diminished within 48	Theophylline was readministered 2 wk later at the same	No history of speech dysfluency	Gerard <i>et al</i> [111]

		disease secondary to pneumoconiosis			extrapyramidal components			h	dosage, and the same speech disorder recurred within a few days and persisted until treatment was stopped		
1. Trifluoperazine (30 mg/d) 2. Chlorpromazine (up to 800 mg/d)	Male/40	Schizophrenia	Trihexyphenidyl (5 mg/d)	1. Four days after increasing the trifluoperazine dosage to 30 mg/d 2. After increasing chlorpromazine dosage to 800 mg/d	NR	NR		1. Increasing dosage of trihexyphenidyl, no improvement. DC of trifluoperazine, complete relief 2. Addition of benztropine, no improvement. Reduction of chlorpromazine dose to 400 mg/d, complete relief	Increasing the dosage of chlorpromazine to 700 mg/d caused the return of stuttering; reducing the dosage of chlorpromazine to 400 mg/d caused cessation of stuttering	No history of speech dysfluency	Nurnberg <i>et al</i> [10]

ADHD: Attention-deficit hyperactivity disorder; BECTS: Benign focal epilepsy of childhood with centrottemporal spikes; BID: Two times per day; BDZ: Benzodiazepine; CBZ: Carbamazepine; COPD: Chronic obstructive pulmonary disease; DC: Discontinue; DM: Diabetes mellitus; EPS: Extrapyramidal side effect; GERD: Gastroesophageal reflux disease; GTC: Generalized tonic-colonic seizure; HTN: Hypertension; IM: Intramuscular; IT: Intrathecal; IV: Intravenous; LD: Loading dose; MD: Maintenance dose; MDD: Major Depressive Disorder; MTX: Methotrexate; NR: Not reported; PD: Parkinson disease; PDS: Persistent developmental stuttering; PRN: Pro re nata (as needed); QID: Four times a day; TID: Three times a day; VPA: Valproic acid.

ROLE OF DOPAMINE IN THE BG MOTOR CIRCUIT

As mentioned above, D1, A1, and M4 receptors are colocalized on MSNs in the direct pathway[34]. The stimulation of D1 receptors activates MSNs in the direct pathway and stimulates this pathway[28]. Blocking A1 and M4 receptors facilitates the dopamine neurotransmission. Therefore, A1 and M4 antagonists also stimulate the direct pathway[34]. Neurons of the indirect pathway have D2, A2A, and M1 receptors[34]. The activation of D2 receptors inhibits the indirect pathway[28]. A2A and M1 receptors antagonists also increase the dopamine signaling and therefore suppress the indirect pathway. Since these two pathways have a cross talk and coordinated activity, disturbing their coordination can cause movement disorders such as dystonia, dyskinesia, and stuttering. Therefore, both increased and decreased dopamine neurotransmission in the striatum may cause stuttering by impairing the balance between the direct and indirect pathways.

WHITE MATTER FIBER TRACTS AND STUTTERING

Other changes that can affect connectivity between different areas of the brain involved in the speech motor control are the white matter abnormalities. The white matter tracts, bundles of myelinated axons, relay signals between different areas of the brain and therefore coordinate their communication and function[38]. Several *in vitro* and animal studies and a small number of human studies have found multiple neurotransmitters, including glutamate, adenosine, GABA, glycine, dopamine, serotonin,




Figure 1 Schematic graph of direct and indirect pathways of basal ganglia. A: Direct pathway; B: Indirect pathway. A1: A1 adenosine receptor; A2A: A2A adenosine receptor; D1: D1 dopamine receptor; D2: D2 dopamine receptor; GABA: Gamma-aminobutyric acid; Glu: Glutamate; Gpe: External segment of the globus pallidus; Gpi: Internal segment of the globus pallidus; M1: M1 muscarinic receptor; M4: M4 muscarinic receptor; MSN: Medium-sized spiny neurons; NMDA: N-methyl-D-aspartate receptor; SNpr: Substantia nigra pars reticulata; STN: Subthalamic nucleus.

acetyl choline, histamine, norepinephrine, and substance P, and their receptors in the white matter. It has been proposed that glutamate and purine signaling have the most prominent effects on the white matter functioning; however, it seems that the white matter requires a coordinated action of all of these neurotransmitters for conduction of action potentials and maintaining signal integrity through very long signal transmission axonal pathways[38]. Different studies in adults[33,39] and adolescents and young people[40] with developmental stuttering demonstrated the reduced integrity of the white matter fiber tracts. It may be proposed that an agent that impairs the normal activity of one or some of the neurotransmitters in the white matter may impair signal transmission between different areas of the brain engaged in speech motor control and result in stuttering. Moreover, in some psychiatric disorders such as schizophrenia and bipolar disorder, one of the pathological findings in the CNS is myelin loss or disruption[41]. Of cases included in the present review, 25, 5, and 6 had schizophrenia, schizoaffective disorder, and bipolar disorder, respectively. As a result, it can be suggested that drugs disrupting normal neurotransmitter balance in the white matter, which already has an underlying impairment in these patients, may exacerbate white matter dysfunction.

STUTTERING INDUCED BY DRUGS AFFECTING DOPAMINE NEUROTRANSMISSION

Increased dopamine neurotransmission in BG and stuttering

Evidence that supports the role of the dopamine excess in the pathophysiology of stuttering includes the reduction of stuttering by antipsychotic drugs, which are dopamine blockers, such as haloperidol, risperidone, olanzapine, aripiprazole, and asenapine[42], the finding of Wu *et al*[43] that showed the excessive striatal dopamine activity and increased uptake of fluoro-L-3, 4-dihydroxy-phenylalanine, a precursor of dopamine, in several parts of the brain in persons who stutter in comparison to healthy controls, and computational modeling of stuttering by Civier *et al*[44]. Furthermore, it has been shown that children aged 2.5-3 years, the age of onset of developmental stuttering in most children, have more density of D2 than D1 receptors and therefore low D1/D2 density in the striatum in comparison to older children. Therefore, drugs like haloperidol, which is a highly selective D2 antagonist, decrease this D1/D2 imbalance and the severity of stuttering. Besides the above-mentioned studies conducted in persons with developmental stuttering, cases of exacerbation of stuttering by levodopa in patients with PD[23,26,27] also propose a role for dopamine excess in the pathogenesis of NS. Chang *et al*[28] have suggested that the inhibition of the indirect pathway in the states of dopamine excess decreases the suppressing effect of this pathway on the competing motor programs. Therefore, choosing correct motor

program over incorrect ones becomes difficult, which could ultimately delay the initiation of the right motor program. This delay may cause speech blockage or sound prolongation. Furthermore, in this situation, the proper signal that originated from the direct pathway and stimulates the right motor program may be initiated but suffers premature termination which may lead to the repetitions[28].

In conclusion, drugs such as levodopa, a precursor of dopamine, methylphenidate that increases the extracellular level of dopamine in the striatum[45], amphetamines that increase the release of catecholamines mainly dopamine and nerve-end particles (NEP) from presynaptic nerves and inhibit the reuptake of dopamine and NEP into presynaptic neurons[46], and phenytoin that has been proposed as a dopamine enhancer in the BG pathways may cause stuttering by increasing the dopamine neurotransmission in BG[47,48].

Decreased dopamine neurotransmission in BG and stuttering

Reduced dopamine neurotransmission in the striatum also can cause stuttering. For example, some patients with PD, the disorder that is mainly characterized by the dopamine depletion from BG, experience new-onset NS[49], exacerbation of PDS[26,27], or re-emergence of developmental stuttering [50]. Chang *et al*[28] proposed that in this state, the decreased excitation of the direct pathway results in reduced stimulation of the correct motor program and its ability to compete other motor programs. This also may result in unstable or delayed production of signals initiating the right speech motor program.

ANTIPSYCHOTICS-INDUCED STUTTERING

Although some studies have shown the relative efficacy of antipsychotic drugs in the treatment of stuttering[42], there are case reports of antipsychotics-induced stuttering. All studies that reported efficacy of antipsychotic drugs in reducing stuttering were conducted in patients with PDS. However, all cases of antipsychotics-induced stuttering had a psychotic disorder, which was schizophrenia in the majority of them. Elevated dopamine levels and excessive dopamine activity in the striatum are present in both developmental stuttering[42,43] and schizophrenia. Therefore, the opposite effects of dopamine blockers in these disorders, improving stuttering in some cases of developmental stuttering but causing stuttering in some patients with schizophrenia, indicate that effects of these drugs on the dopamine activity in other parts of the brain and on other neurotransmitters may be responsible for their different effects on speech motor control. Furthermore, abnormalities in the brain of patients with psychotic disorders are extensive and are not comparable to persons with developmental stuttering. These differences also may justify why a dopamine blocker can be a therapeutic option in PDS but a causative agent of stuttering in the psychotic disorders.

However, it should be noted that studies by Fish *et al*[51] and Langova *et al*[52] demonstrated that not all persons with developmental stuttering respond to the dopamine blockers. In the study conducted by Fish *et al*[51], 14 out of 28 persons who stuttered Prader-Willi syndrome (PWS) and received amphetamine experienced improvement in stuttering while two got worse. Of 12 PWS who did not improve by amphetamine, eight got better on trifluoperazine, a D2 blocker. Four participants did not show any improvement neither by each medication nor by their combination. Similarly, in the study conducted by Langova *et al*[52], 88% of PWS got better on phenmetrazine, a stimulant, while 67% deteriorated using chlorpromazine. These findings have led to the suggestion of the hypothesis that persons with developmental stuttering may be classified as the dopamine blocker-responsive or stimulant-responsive[29].

In the present review and study conducted by Trenque *et al*[6], the majority of cases of DIS were caused by clozapine. Regarding the effects of antipsychotics on dopamine receptors, all antipsychotics except than clozapine, olanzapine, ziprasidone, and asenapine have a higher affinity for D2 receptors than D1 receptors. Clozapine, olanzapine, ziprasidone, and asenapine equally block both D1 and D2 receptors. Furthermore, clozapine and asenapine have a lower affinity for dopamine receptors in comparison to olanzapine and ziprasidone[53]. Clozapine is one of the most effective antipsychotic drugs[54]; however, because of its serious side effects such as agranulocytosis, seizure, and cardiovascular adverse effects[55], clozapine is considered as one of the last options in the treatment of schizophrenia and other psychotic disorders. Therefore, many patients had received several antipsychotics prior to the initiation of clozapine. Long-term blockage of D2 receptors while sparing D1 receptors caused by other antipsychotics results in the supersensitivity of D2 receptors[56]. The affinity of clozapine for blocking D2 receptors is lower than that of many other antipsychotics[53]. Therefore, after the initiation of clozapine, decreased D2 blocking combined with the supersensitivity of D2 receptors creates a state of increased D2 stimulation which finally inhibits the indirect pathway. On the other hand, antagonizing the D1 receptors inhibits the direct pathway. The inhibition of the indirect pathway impairs the suppression of the competing motor programs. Besides, the inhibition of the direct pathway decreases cerebral cortex stimulation and locomotor activity, which causes difficulties in initiating next segment in a movement sequence like speaking. Clozapine also is a 5HT_{2a} and 5HT_{2c} receptor blocker[55]. Reducing the serotonin neurotransmission increases the dopamine transmission in the prefrontal cortex, which may also cause stuttering. Although clozapine is a potent M1 antagonist

[55], the effect that facilitates the dopamine neurotransmission in the direct pathway, the final result of antagonizing all above-mentioned receptors is the inhibition of both direct and indirect pathways and therefore impaired speech motor control. Moreover, changing the normal function of dopamine, serotonin, acetylcholine, and norepinephrine, by blocking $\alpha 1$ receptors, and histamine, by blocking H1 receptors[55], can disturb normal functions of the white matter and cause stuttering.

Other antipsychotics that are included in the present review are olanzapine, risperidone, aripiprazole, chlorpromazine, fluphenazine, and trifluoperazine. Olanzapine is very similar to clozapine regarding the affinity for different receptors including D1, D2, 5HT_{2A}, 5HT_{2C}, M1, $\alpha 1$, and H1 receptors[57]. Other antipsychotics such as risperidone, chlorpromazine, and fluphenazine are potent inhibitors of D2 receptors without any considerable effect on D1 receptors. Chlorpromazine also is a potent M1 antagonist while risperidone and fluphenazine have no considerable effect on M1 receptor. The efficacy of risperidone in antagonizing 5HT receptors is comparable to that of olanzapine and more than the efficacy of chlorpromazine, fluphenazine, and trifluoperazine. Aripiprazole is a partial agonist of D2 and 5HT_{1A} receptors and antagonist of 5HT_{2A} receptor[57]. We suggest that these antipsychotics can cause extrapyramidal side effects (EPS) which may manifest as stuttering as well as other movement disorders by impairing the balanced and coordinated activity of the direct and indirect pathways. Furthermore, increasing the dopamine neurotransmission in the prefrontal cortex by blocking the serotonin effects as well as disturbing neurotransmitters' functioning in the white matter can be the other underlying mechanisms of the antipsychotics-induced stuttering.

The following section is focused on clozapine as the most prevalent cause of DIS.

CLOZAPINE-INDUCED STUTTERING

Clozapine is the drug with most reports of inducing a new episode of stuttering or worsening pre-existing stuttering both in the present review and analysis carried out by Trenque *et al*[6]. In the cases included in this review, clozapine induced stuttering in a wide variety of dosages ranging from 50 mg/d[10] to 700 mg/d[58,59]. However, in most cases, clozapine caused stuttering at the daily doses of 250 mg to 450 mg. It seems that stuttering is a dose-dependent adverse effect of clozapine as in 13 (43.3%) cases stuttering was significantly improved or completely vanished following dose reduction (Table 3).

We have suggested likely mechanisms of clozapine-induced stuttering in the previous section. Furthermore, based on the concomitant signs and symptoms that patients experienced with stuttering, it has been proposed that clozapine-induced stuttering may be a manifestation of the movement disorders such as focal segmental dystonia in orofacial muscles, akathisia, or dyskinesia[10,21,58,60-63] or a seizure disorder.

Although clozapine is an antipsychotic with a low potential for causing EPS[55], of 30 cases, seven experienced a type of movement disorder concomitant with stuttering[10,21,58]. Grover *et al*[61] reported a case who experienced clozapine-induced stuttering and had a history of EPS associated with other antipsychotics. Although clozapine did not cause other manifestations of EPS, they proposed that stuttering might be a symptom of movement disorders induced by clozapine. Concerning the management of clozapine-induced stuttering the dose reduction or withdrawal of clozapine resulted in significant improvement or complete relief of both stuttering and the movement disorders in two cases [58]. In contrast, in one case reported by Lyall *et al*[10], substitution of clozapine with zuclopenthixol decanoate relieved stuttering but not dyskinetic movements, and restarting clozapine resulted in reoccurrence of stuttering which responded to sodium valproate despite no electroencephalogram (EEG) abnormality. For four cases, the authors did not report whether their intervention improved the movement disorders in addition to stuttering or not[10,21].

Regarding the other likely mechanisms of clozapine-induced stuttering, it has been suggested that stuttering may be a manifestation of seizure. Clozapine-induced stuttering was associated with seizure or EEG abnormalities without typical symptoms of seizure in nine out of 30 case reports[12,25,59,60,62-66], and stuttering was significantly improved or completely relieved by addition of sodium valproate and the dose reduction of clozapine in five cases[12,59,62,64,66]. In the other four cases[25,60,63,65], anticonvulsant drugs were not tried, but discontinuation or dose reduction of clozapine resulted in complete relief of stuttering in three cases[60,63,65]. In contrast, one of the cases reported by Hallahan *et al*[58] experienced seizure after the development of stuttering. After the addition of sodium valproate, the patient had no seizure, but stuttering did not improve. It is worth mentioning that three cases had stuttering with both movement disorders and EEG abnormalities[60,62,63], which may demonstrate that some movement disorders induced by clozapine may be due to the epileptiform activity in the brain rather than EPS.

Collectively, the mechanism of clozapine-induced stuttering is multifactorial. Any of the following impairments or a combination of them can be a cause of clozapine-induced stuttering: Imbalance between direct and indirect pathways in the BG which may cause just stuttering or other movement disorders in addition to stuttering, abnormalities in the white matter neurotransmitters, and seizure or EEG abnormalities.

Finally, we suggest that in the cases of clozapine-induced stuttering at first, clinicians must do the electroencephalography to rule out any abnormal electrical activity of the brain which may progress to convulsion. If the patient has seizure or any abnormalities in EEG, the addition of sodium valproate is recommended. Even Varma *et al*[67] have recommended that one of the indications of initiating sodium valproate in patients receiving clozapine is stuttering with or without any types of seizure or EEG abnormalities. After ruling out the abnormal electrical brain activities, considering stuttering as an EPS and typical management of these side effects or dose reduction or discontinuation of clozapine may be considered if stuttering is annoying the patient.

STUTTERING INDUCED BY DRUGS AFFECTING GLUTAMATE NEUROTRANSMISSION

As described above, as the input nucleus of the BG, the striatum receives glutamatergic projection from different parts of the cerebral cortex and thalamus. Furthermore, the STN stimulates inhibitory GABAergic neurons of the output nuclei of the BG by releasing glutamate. All ionotropic NMDA, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid, and kainite, and metabotropic glutamate receptors, mGlu1-8, are expressed in the BG, and glutamate is one of the important neurotransmitters that mediate signal transmission in the BG motor circuit. In addition, the computational modeling of stuttering[44] and the data of diffusion tensor imaging obtained from children who stutter[44] indicate that there is an impaired connectivity between different parts of the cerebral cortex and striatum in developmental stuttering. This impaired connection causes that BG cannot optimally detect cognitive and sensory motor context to terminate the previous phoneme and initiate the next phoneme[29] in a timely manner to produce a fluent speech. Therefore, drugs like memantine by inhibiting NMDA receptors[68] and pregabalin, gabapentin, and lamotrigine by inhibiting the release of glutamate[69,70] change the normal function of glutamate in corticostriatal pathway or in the different parts of the BG motor circuit can impair balance between the direct and indirect pathways and cause motor abnormalities like stuttering. Moreover, glutamate is the main neurotransmitter in the white matter and antagonizing its effects also disturbs signal transmission through white matter fiber tracts.

Theophylline

Theophylline increases dopamine release and transmission by inhibiting GABA receptors on the SNpc and adenosine receptors on MSNs of both direct and indirect pathways. Theophylline also increases glutamate release. Therefore, it can disturb the normal balance between the mentioned neurotransmitters in the BG and cause stuttering. Another mechanism may be impairing the normal function of neurotransmitters in the white matter by increasing glutamate, the main neurotransmitter of the white matter[5].

STUTTERING INDUCED BY OTHER DRUGS

Divalproex

Although the main mechanism of action of valproate is the blockage of voltage-dependent sodium channels[71], it has several other mechanisms of action that justify its broad anticonvulsant activity, effects in the prophylaxis of migraine headache, and mood stabilizing properties. Animal studies in rats have demonstrated that valproate increases GABA concentration in both the striatum and substantia nigra, but its effect is more pronounced in the substantia nigra[72]. Valproate also increases the firing pattern and frequency of neurons of the SNpr[73]. We propose that an increased level of GABA in the substantia nigra increases its inhibitory effects on the thalamus through the direct pathway, and therefore, reduces the brain cortex stimulation to execute the desired movement which may cause stuttering.

Atomoxetine

Atomoxetine is a selective NEP reuptake inhibitor. It increases the extracellular concentrations of NEP and dopamine in the prefrontal cortex[45]. However, studies that examined the effects of atomoxetine on the concentration of dopamine in the striatum obtained opposite results[45,74]. Because of uncertainty about the effect of atomoxetine on the dopamine levels in the striatum, we do not focus on dopamine as a mediator of likely effects of atomoxetine on speech motor control and stuttering. A study measured the blood oxygenation level dependent response using pharmacological magnetic resonance imaging in different regions of the rat brain following acute administration of atomoxetine. That study showed that atomoxetine increased SNpr and STN activity in the BG[75]. These increased activities decrease the stimulatory activity of the thalamocortical pathway. Therefore, we suggest that because of the decreased stimulation of the cerebral cortex, the favorable motor program is not executed, which can result in the inappropriate activation of the orofacial muscles and cause stuttering.

Sertraline

It is proposed that sertraline-induced stuttering may be related to the serotonergic inhibition of the dopaminergic neurons. The cell bodies of these neurons are located in the ventral tegmental area. Therefore, inhibition of the dopamine pathways in the nigrostriatum can be considered as a mechanism of promoting stuttering by sertraline or selective serotonin reuptake inhibitors drugs in general[76,77].

Bupropion

Bupropion is able to increase dopamine levels in the prefrontal cortex, which may cause stuttering[78-80].

Pyrethrin

It seems that the stuttering induced by a topical pyrethrin product in a child is related to its neurotoxicity since the metabolism of pyrethrin in children is slow. The product also had contained piperonyl butoxide, which can inhibit the hepatic metabolism of the compound and potentiate the toxicity[81].

CONCLUSION

In this review, 82 cases of DIS were collected. Most cases were related to antipsychotic drugs. Similar to the developmental stuttering, the majority of persons who experienced an episode of DIS were male. The repetitions followed by speech blocks were the most frequent core manifestations of stuttering. In 55.8% of cases, drug withdrawal was the therapeutic measure that was used to manage the stuttering.

Although we tried to provide a complete feature of the epidemiological and clinical characteristics of DIS, much information such as the core behaviors of stuttering, the interval between the initiation or increase in the dose of offending medications and the occurrence of stuttering and between the drug withdrawal or dose reduction and the improvement of stuttering, and concurrent psychological symptoms with stuttering was not reported in several cases. As a result, future cases of DIS must be reported with more detailed information since these data give others a comprehensive feature of this type of the NS.

By focusing on the cortico-BG-thalamocortical loop and the white matter fiber tracts and their neurotransmitters such as dopamine and glutamate, we suggest some likely mechanisms for DIS. However, dysfunctions in other areas of the brain like the cerebral cortex and cerebellum and other neurotransmitters are not addressed in this review. In addition, we consider stuttering as a speech motor disorder, but cognitive and sensory disorders may also play roles in the pathogenesis of DIS. Therefore, it is suggested that these subjects should be considered in the future papers discussing the underlying mechanisms of DIS. In spite of many hypotheses that can be proposed for the pathogenesis of DIS, experimental studies will provide the most robust evidence in this field. Since advanced brain imaging facilities may not be available in every setting where clinicians encounter a case of DIS to find the areas of the brain that act abnormally, animal studies evaluating the changes in the functions of the brain and different neurotransmitters are required to shed a light on the underlying mechanisms of DIS.

FOOTNOTES

Author contributions: Both authors substantially contributed to conception and design of the review, searched and interpreted the relevant data, drafted the manuscript, and approved the final version of the article to be published.

Conflict-of-interest statement: The authors declare no conflict of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Iran

ORCID number: Naemeh Nikvarz 0000-0002-0144-3415; Salehe Sabouri 0000-0002-9479-212X.

S-Editor: Wang JL

L-Editor: Wang TQ

P-Editor: Wang JL

REFERENCES

- 1 **Black DW**, Grant JE. DSM-5® guidebook: the essential companion to the diagnostic and statistical manual of mental disorders: American Psychiatric Pub, 2014
- 2 **Craig-McQuaide A**, Akram H, Zrinzo L, Tripoliti E. A review of brain circuitries involved in stuttering. *Front Hum Neurosci* 2014; **8**: 884 [PMID: [25452719](#) DOI: [10.3389/fnhum.2014.00884](#)]
- 3 **Neef NE**, Anwander A, Friederici AD. The Neurobiological Grounding of Persistent Stuttering: from Structure to Function. *Curr Neurol Neurosci Rep* 2015; **15**: 63 [PMID: [26228377](#) DOI: [10.1007/s11910-015-0579-4](#)]
- 4 **Ashurst JV**, Wasson MN. Developmental and persistent developmental stuttering: an overview for primary care physicians. *J Am Osteopath Assoc* 2011; **111**: 576-580 [PMID: [22065298](#)]
- 5 **Movsessian P**. Neuropharmacology of theophylline induced stuttering: the role of dopamine, adenosine and GABA. *Med Hypotheses* 2005; **64**: 290-297 [PMID: [15607558](#) DOI: [10.1016/j.mehy.2004.07.026](#)]
- 6 **Trenque T**, Morel A, Trenque A, Azzouz B. Drug induced stuttering: pharmacovigilance data. *Expert Opin Drug Saf* 2021; **20**: 373-378 [PMID: [33337944](#) DOI: [10.1080/14740338.2021.1867101](#)]
- 7 **Burd L**, Kerbeshian J. Stuttering and stimulants. *J Clin Psychopharmacol* 1991; **11**: 72-73 [PMID: [2040720](#) DOI: [10.1097/00004714-199102000-00020](#)]
- 8 **Ertekin H**, Ertekin YH, Sahin B, Yayla S, Turkyilmaz E, Kara M. Clozapine and Aripiprazole-Induced Stuttering: A Case Report of Turner Syndrome with Schizophrenia. *Klinik Psikofarmakoloji Bülteni-Bulletin Clin Psychopharmacol* 2016; **26**: 422-425 [DOI: [10.5455/bcp.20151204115654](#)]
- 9 **Lyall M**, Pryor A, Murray K. Clozapine and speech dysfluency: two case reports. *Psychiatr Bull* 2007; **31**: 16-18 [DOI: [10.1192/pb.31.1.16](#)]
- 10 **Nurnberg HG**, Greenwald B. Stuttering: an unusual side effect of phenothiazines. *Am J Psychiatry* 1981; **138**: 386-387 [PMID: [6110346](#) DOI: [10.1176/ajp.138.3.386](#)]
- 11 **Begum M**. Clozapine-induced stuttering, facial tics and myoclonic seizures: a case report. *Aust N Z J Psychiatry* 2005; **39**: 202 [PMID: [15701074](#) DOI: [10.1080/j.1440-1614.2005.01549.x](#)]
- 12 **Prasse JE**, Kikano GE. Stuttering: an overview. *Am Fam Physician* 2008; **77**: 1271-1276 [PMID: [18540491](#)]
- 13 **Tani T**, Sakai Y. Stuttering after right cerebellar infarction: a case study. *J Fluency Disord* 2010; **35**: 141-145 [PMID: [20609334](#) DOI: [10.1016/j.jfludis.2010.03.001](#)]
- 14 **Theys C**, Van Wieringen A, Tuyls L, De Nil L. Acquired stuttering in a 16-year-old boy. *J Neurolinguist* 2009; **22**: 427-435 [DOI: [10.1016/j.jneuroling.2009.02.001](#)]
- 15 **Bär KJ**, Häger F, Sauer H. Olanzapine- and clozapine-induced stuttering. A case series. *Pharmacopsychiatry* 2004; **37**: 131-134 [PMID: [15179972](#) DOI: [10.1055/s-2004-818992](#)]
- 16 **Lee HJ**, Lee HS, Kim L, Lee MS, Suh KY, Kwak DI. A case of risperidone-induced stuttering. *J Clin Psychopharmacol* 2001; **21**: 115-116 [PMID: [11199937](#) DOI: [10.1097/00004714-200102000-00024](#)]
- 17 **Alaghband-Rad J**, Nikvarz N, Tehrani-Doost M, Ghaeli P. Memantine-induced speech problems in two patients with autistic disorder. *Daru* 2013; **21**: 54 [PMID: [23819879](#) DOI: [10.1186/2008-2231-21-54](#)]
- 18 **Yadav DS**. Risperidone induced stuttering. *Gen Hosp Psychiatry* 2010; **32**: 559.e9-559.10 [PMID: [20851282](#) DOI: [10.1016/j.genhosppsych.2010.01.004](#)]
- 19 **Das S**, Manjunatha N, Thirthali J. Clozapine-induced Weight Loss and Stuttering in a Patient with Schizophrenia. *Indian J Psychol Med* 2018; **40**: 385-387 [PMID: [30093754](#) DOI: [10.4103/IJPSYM.IJPSYM_523_17](#)]
- 20 **Murphy R**, Gallagher A, Sharma K, Ali T, Lewis E, Murray I, Hallahan B. Clozapine-induced stuttering: an estimate of prevalence in the west of Ireland. *Ther Adv Psychopharmacol* 2015; **5**: 232-236 [PMID: [26301079](#) DOI: [10.1177/2045125315590060](#)]
- 21 **Shuster J**. Methotrexate Neurotoxicity Causes Speech Problem; Severe, Irreversible Sensory Neuropathy Due to Long-Term Use of Linezolid; Hypomania with Topiramate; Bortezomib-Induced Hepatitis; Steroid Dementia—An Overlooked Diagnosis? *Hosp Pharm* 2005; **40**: 383-386 [DOI: [10.1177/001857870504000503](#)]
- 22 **Louis ED**, Winfield L, Fahn S, Ford B. Speech dysfluency exacerbated by levodopa in Parkinson's disease. *Mov Disord* 2001; **16**: 562-565 [PMID: [11391759](#) DOI: [10.1002/mds.1081](#)]
- 23 **Atay İM**, Tanritanir B, Akpınar A, Demirdağ A. A Case of Risperidone Induced Stuttering as a Paradox. *Noro Psikiyatr Ars* 2014; **51**: 403-404 [PMID: [28360662](#) DOI: [10.5152/npa.2014.6946](#)]
- 24 **Kranidiotis L**, Thomas S. Clozapine-induced speech dysfluency: further cases. *Psychiatr Bull* 2007; **31**: 191-191 [DOI: [10.1192/pb.31.5.191b](#)]
- 25 **Anderson JM**, Hughes JD, Rothi LJ, Crucian GP, Heilman KM. Developmental stuttering and Parkinson's disease: the effects of levodopa treatment. *J Neurol Neurosurg Psychiatry* 1999; **66**: 776-778 [PMID: [10329754](#) DOI: [10.1136/jnnp.66.6.776](#)]
- 26 **Burghaus L**, Hilker R, Thiel A, Galldiks N, Lehnhardt FG, Zaro-Weber O, Sturm V, Heiss WD. Deep brain stimulation of the subthalamic nucleus reversibly deteriorates stuttering in advanced Parkinson's disease. *J Neural Transm (Vienna)* 2006; **113**: 625-631 [PMID: [16075183](#) DOI: [10.1007/s00702-005-0341-1](#)]
- 27 **Chang SE**, Guenther FH. Involvement of the Cortico-Basal Ganglia-Thalamocortical Loop in Developmental Stuttering. *Front Psychol* 2019; **10**: 3088 [PMID: [32047456](#) DOI: [10.3389/fpsyg.2019.03088](#)]
- 28 **Alm PA**. Stuttering and the basal ganglia circuits: a critical review of possible relations. *J Commun Disord* 2004; **37**: 325-369 [PMID: [15159193](#) DOI: [10.1016/j.jcomdis.2004.03.001](#)]
- 29 **Lanciego JL**, Luquin N, Obeso JA. Functional neuroanatomy of the basal ganglia. *Cold Spring Harb Perspect Med* 2012; **2**: a009621 [PMID: [23071379](#) DOI: [10.1101/cshperspect.a009621](#)]
- 30 **Leisman G**, Braun-Benjamin O, Melillo R. Cognitive-motor interactions of the basal ganglia in development. *Front Syst Neurosci* 2014; **8**: 16 [PMID: [24592214](#) DOI: [10.3389/fnsys.2014.00016](#)]
- 31 **Mink JW**. The Basal Ganglia and involuntary movements: impaired inhibition of competing motor patterns. *Arch Neurol* 2003; **60**: 1365-1368 [PMID: [14568805](#) DOI: [10.1001/archneur.60.10.1365](#)]

- 32 Connally EL, Ward D, Howell P, Watkins KE. Disrupted white matter in language and motor tracts in developmental stuttering. *Brain Lang* 2014; **131**: 25-35 [PMID: 23819900 DOI: 10.1016/j.bandl.2013.05.013]
- 33 Silkis I. The cortico-basal ganglia-thalamocortical circuit with synaptic plasticity. II. Mechanism of synergistic modulation of thalamic activity via the direct and indirect pathways through the basal ganglia. *Biosystems* 2001; **59**: 7-14 [PMID: 11226622 DOI: 10.1016/S0303-2647(00)00135-0]
- 34 Galvan A, Kuwajima M, Smith Y. Glutamate and GABA receptors and transporters in the basal ganglia: what does their subsynaptic localization reveal about their function? *Neuroscience* 2006; **143**: 351-375 [PMID: 17059868 DOI: 10.1016/j.neuroscience.2006.09.019]
- 35 Calabresi P, Picconi B, Tozzi A, Ghiglieri V, Di Filippo M. Direct and indirect pathways of basal ganglia: a critical reappraisal. *Nat Neurosci* 2014; **17**: 1022-1030 [PMID: 25065439 DOI: 10.1038/nn.3743]
- 36 Conn PJ, Battaglia G, Marino MJ, Nicoletti F. Metabotropic glutamate receptors in the basal ganglia motor circuit. *Nat Rev Neurosci* 2005; **6**: 787-798 [PMID: 16276355 DOI: 10.1038/nrn1763]
- 37 Butt AM, Fern RF, Matute C. Neurotransmitter signaling in white matter. *Glia* 2014; **62**: 1762-1779 [PMID: 24753049 DOI: 10.1002/glia.22674]
- 38 Kronfeld-Duenias V, Amir O, Ezrati-Vinacour R, Civier O, Ben-Shachar M. The frontal aslant tract underlies speech fluency in persistent developmental stuttering. *Brain Struct Funct* 2016; **221**: 365-381 [PMID: 25344925 DOI: 10.1007/s00429-014-0912-8]
- 39 Watkins KE, Smith SM, Davis S, Howell P. Structural and functional abnormalities of the motor system in developmental stuttering. *Brain* 2008; **131**: 50-59 [PMID: 17928317 DOI: 10.1093/brain/awm241]
- 40 Haroutunian V, Katsel P, Roussos P, Davis KL, Altschuler LL, Bartzokis G. Myelination, oligodendrocytes, and serious mental illness. *Glia* 2014; **62**: 1856-1877 [PMID: 25056210 DOI: 10.1002/glia.22716]
- 41 Maguire GA, Yeh CY, Ito BS. Overview of the diagnosis and treatment of stuttering. *J Exp Clin Med* 2012; **4**: 92-97 [DOI: 10.1016/j.jecm.2012.02.001]
- 42 Wu JC, Maguire G, Riley G, Lee A, Keator D, Tang C, Fallon J, Najafi A. Increased dopamine activity associated with stuttering. *Neuroreport* 1997; **8**: 767-770 [PMID: 9106763 DOI: 10.1097/00001756-199702100-00037]
- 43 Civier O, Bullock D, Max L, Guenther FH. Computational modeling of stuttering caused by impairments in a basal ganglia thalamo-cortical circuit involved in syllable selection and initiation. *Brain Lang* 2013; **126**: 263-278 [PMID: 23872286 DOI: 10.1016/j.bandl.2013.05.016]
- 44 Bymaster FP, Katner JS, Nelson DL, Hemrick-Luecke SK, Threlkeld PG, Heiligenstein JH, Morin SM, Gehlert DR, Perry KW. Atomoxetine increases extracellular levels of norepinephrine and dopamine in prefrontal cortex of rat: a potential mechanism for efficacy in attention deficit/hyperactivity disorder. *Neuropsychopharmacology* 2002; **27**: 699-711 [PMID: 12431845 DOI: 10.1016/S0893-133X(02)00346-9]
- 45 Fleckenstein AE, Volz TJ, Riddle EL, Gibb JW, Hanson GR. New insights into the mechanism of action of amphetamines. *Annu Rev Pharmacol Toxicol* 2007; **47**: 681-698 [PMID: 17209801 DOI: 10.1146/annurev.pharmtox.47.120505.105140]
- 46 García-Ramos R, Moreno Ramos T, Villarejo Galende A, Porta Etessam J. Phenytoin-induced acute orofacial dyskinesia. *Neurologia* 2013; **28**: 193-194 [PMID: 22595500 DOI: 10.1016/j.nrl.2012.02.005]
- 47 Zaatreh M, Tennison M, D'Cruz O, Beach RL. Anticonvulsants-induced chorea: a role for pharmacodynamic drug interaction? *Seizure* 2001; **10**: 596-599 [PMID: 11792164 DOI: 10.1053/seiz.2001.0555]
- 48 Toft M, Dietrichs E. Aggravated stuttering following subthalamic deep brain stimulation in Parkinson's disease-two cases. *BMC Neurol* 2011; **11**: 44 [PMID: 21477305 DOI: 10.1186/1471-2377-11-44]
- 49 Shahed J, Jankovic J. Re-emergence of childhood stuttering in Parkinson's disease: a hypothesis. *Mov Disord* 2001; **16**: 114-118 [PMID: 11215569 DOI: 10.1002/1531-8257(200101)16:1<114::aid-mds1004>3.0.co;2-2]
- 50 Fish CH, Bowling E. Stuttering. The effect of treatment with D-amphetamine and a tranquilizing agent, trifluoperazine. A preliminary report on an uncontrolled study. *Calif Med* 1965; **103**: 337-339 [PMID: 5836893]
- 51 Langova J, Moravek M. Some Results of Experimental Examinations among Stutterers and Clutterers. *Folia Phoniatr (Basel)* 1964; **16**: 290-296 [PMID: 14188546]
- 52 Perez-Costas E, Melendez-Ferro M, Roberts RC. Basal ganglia pathology in schizophrenia: dopamine connections and anomalies. *J Neurochem* 2010; **113**: 287-302 [PMID: 20089137 DOI: 10.1111/j.1471-4159.2010.06604.x]
- 53 Rampino A, Marakhovskaia A, Soares-Silva T, Torretta S, Veneziani F, Beaulieu JM. Antipsychotic Drug Responsiveness and Dopamine Receptor Signaling: Old Players and New Prospects. *Front Psychiatry* 2018; **9**: 702 [PMID: 30687136 DOI: 10.3389/fpsy.2018.00702]
- 54 Essali A, Al-Haj Haasan N, Li C, Rathbone J. Clozapine versus typical neuroleptic medication for schizophrenia. *Cochrane Database Syst Rev* 2009; **2009**: CD000059 [PMID: 19160174 DOI: 10.1002/14651858.CD000059.pub2]
- 55 Gardner DM, Baldessarini RJ, Warch P. Modern antipsychotic drugs: a critical overview. *CMAJ* 2005; **172**: 1703-1711 [PMID: 15967975 DOI: 10.1503/cmaj.1041064]
- 56 Yin J, Barr AM, Ramos-Miguel A, Procyshyn RM. Antipsychotic Induced Dopamine Supersensitivity Psychosis: A Comprehensive Review. *Curr Neuropsychopharmacol* 2017; **15**: 174-183 [PMID: 27264948 DOI: 10.2174/1570159x14666160606093602]
- 57 Alldredge BK, Corelli RL, Ernst ME, Guglielmo BJ, Jacobson PA, Kradjan WA, Williams BR. Koda-kimble and Young's applied therapeutics: the clinical use of drugs. 10th ed. Philadelphia, USA: Wolters Kluwer Health Adis (ESP), 2013: 1931-1932
- 58 Hallahan B, Murray I, Doyle PG. Clozapine induced stuttering. *Ir J Psychol Med* 2007; **24**: 121 [PMID: 30290494 DOI: 10.1017/S079096670001048X]
- 59 Supprian T, Retz W, Deckert J. Clozapine-induced stuttering: epileptic brain activity? *Am J Psychiatry* 1999; **156**: 1663-1664 [PMID: 10518185 DOI: 10.1176/ajp.156.10.1663]
- 60 Chochol MD, Kataria L, O'Rourke MC, Lamotte G. Clozapine-Associated Myoclonus and Stuttering Secondary to Smoking Cessation and Drug Interaction: A Case Report. *J Clin Psychopharmacol* 2019; **39**: 275-277 [PMID: 30925500 DOI: 10.1097/JCP.0000000000001032]

- 61 Grover S, Verma AK, Nebhinani N. Clozapine-induced stuttering: a case report and analysis of similar case reports in the literature. *Gen Hosp Psychiatry* 2012; **34**: 703.e1-703.e3 [PMID: 22516217 DOI: 10.1016/j.genhosppsych.2012.02.010]
- 62 Rachamalla V, Haq A, Song MM, Aligeti M. Clozapine-Induced Microseizures, Orofacial Dyskinesia, and Speech Dysfluency in an Adolescent with Treatment Resistant Early Onset Schizophrenia on Concurrent Lithium Therapy. *Case Rep Psychiatry* 2017; **2017**: 7359095 [PMID: 28835863 DOI: 10.1155/2017/7359095]
- 63 Thomas P, Lalaux N, Vaiva G, Goudemand M. Dose-dependent stuttering and dystonia in a patient taking clozapine. *Am J Psychiatry* 1994; **151**: 1096 [PMID: 8010372 DOI: 10.1176/ajp.151.7.1096a]
- 64 Duggal HS, Jagadheesan K, Nizamie SH. Clozapine-induced stuttering and seizures. *Am J Psychiatry* 2002; **159**: 315 [PMID: 11823281 DOI: 10.1176/appi.ajp.159.2.315]
- 65 Gica S, Kiliç C, Karamustafalioglu N. Clozapine-Associated Stuttering: A Case Report. *Am J Ther* 2020; **27**: e624-e627 [PMID: 31219808 DOI: 10.1097/MJT.0000000000001016]
- 66 Horga G, Horga A, Baeza I, Castro-Fornieles J, Lázaro L, Pons A. Drug-induced speech dysfluency and myoclonus preceding generalized tonic-clonic seizures in an adolescent male with schizophrenia. *J Child Adolesc Psychopharmacol* 2010; **20**: 233-234 [PMID: 20578939 DOI: 10.1089/cap.2009.0010]
- 67 Varma S, Bishara D, Besag FM, Taylor D. Clozapine-related EEG changes and seizures: dose and plasma-level relationships. *Ther Adv Psychopharmacol* 2011; **1**: 47-66 [PMID: 23983927 DOI: 10.1177/2045125311405566]
- 68 Johnson JW, Kotermanski SE. Mechanism of action of memantine. *Curr Opin Pharmacol* 2006; **6**: 61-67 [PMID: 16368266 DOI: 10.1016/j.coph.2005.09.007]
- 69 Lapidus KA, Soleimani L, Murrough JW. Novel glutamatergic drugs for the treatment of mood disorders. *Neuropsychiatr Dis Treat* 2013; **9**: 1101-1112 [PMID: 23976856 DOI: 10.2147/NDT.S36689]
- 70 Sills GJ. The mechanisms of action of gabapentin and pregabalin. *Curr Opin Pharmacol* 2006; **6**: 108-113 [PMID: 16376147 DOI: 10.1016/j.coph.2005.11.003]
- 71 Johannessen CU. Mechanisms of action of valproate: a commentary. *Neurochem Int* 2000; **37**: 103-110 [PMID: 10812195 DOI: 10.1016/S0197-0186(00)00013-9]
- 72 Löscher W. Valproate enhances GABA turnover in the substantia nigra. *Brain Res* 1989; **501**: 198-203 [PMID: 2508993 DOI: 10.1016/0006-8993(89)91044-5]
- 73 Töllner K, Wolf S, Löscher W, Gernert M. The anticonvulsant response to valproate in kindled rats is correlated with its effect on neuronal firing in the substantia nigra pars reticulata: a new mechanism of pharmacoresistance. *J Neurosci* 2011; **31**: 16423-16434 [PMID: 22072692 DOI: 10.1523/JNEUROSCI.2506-11.2011]
- 74 Easton N, Shah YB, Marshall FH, Fone KC, Marsden CA. Guanfacine produces differential effects in frontal cortex compared with striatum: assessed by pHMRI BOLD contrast. *Psychopharmacology (Berl)* 2006; **189**: 369-385 [PMID: 17016709 DOI: 10.1007/s00213-006-0558-1]
- 75 Easton N, Marshall F, Fone K, Marsden C. Atomoxetine produces changes in cortico-basal thalamic loop circuits: assessed by pHMRI BOLD contrast. *Neuropharmacology* 2007; **52**: 812-826 [PMID: 17140608 DOI: 10.1016/j.neuropharm.2006.09.024]
- 76 Christensen RC, Byerly MJ, McElroy RA. A case of sertraline-induced stuttering. *J Clin Psychopharmacol* 1996; **16**: 92-93 [PMID: 8834434 DOI: 10.1097/00004714-199602000-00025]
- 77 Brewerton TD, Markowitz JS, Keller SG, Cochrane CE. Stuttering with sertraline. *J Clin Psychiatry* 1996; **57**: 90-91 [PMID: 8591976]
- 78 Fetterolf F, Marceau M. A case of bupropion-induced stuttering. *Gen Hosp Psychiatry* 2013; **35**: 574.e7-574.e8 [PMID: 22959418 DOI: 10.1016/j.genhosppsych.2012.07.003]
- 79 Bhatia MS. Bupropion-Induced Stuttering. *Prim Care Companion CNS Disord* 2015; **17** [PMID: 26693041 DOI: 10.4088/PCC.15101777]
- 80 McAllister MW, Woodhall DM. Bupropion-induced stuttering treated with haloperidol. *Clin Toxicol (Phila)* 2016; **54**: 603 [PMID: 27159769 DOI: 10.1080/15563650.2016.1179749]
- 81 Hammond K, Leikin JB. Topical pyrethrin toxicity leading to acute-onset stuttering in a toddler. *Am J Ther* 2008; **15**: 323-324 [PMID: 18645333 DOI: 10.1097/MJT.0b013e318160c2d7]
- 82 Donaher J, Healey EC, Zobel A. The effects of ADHD medication changes on a child who stutters. *Perspect Fluency Fluency Disord* 2009; **19**: 95-98 [DOI: 10.1044/ffd19.3.95]
- 83 Elliott RL, Thomas BJ. A case report of alprazolam-induced stuttering. *J Clin Psychopharmacol* 1985; **5**: 159-160 [PMID: 2860135]
- 84 Ünay M, Adanir AS, Özatalay E. Aripiprazole-induced stuttering in an 8 year-old boy with ADHD. *Klin Psikofarmakol Bul* 2018; **28**: 230-231
- 85 Naguy A, Moodliar S, Elson DH, Alamiri B. Dose-Dependent Aripiprazole-Induced Stuttering in a Child With Mild Intellectual Disability. *Am J Ther* 2020 [PMID: 32167942 DOI: 10.1097/MJT.0000000000001158]
- 86 Cicek AU. Aggravating influence of atomoxetine on the severity of stuttering and its successful treatment with methylphenidate: a case report. *Dusunen Adam* 2020; **33**: 210-213 [DOI: 10.14744/DAJPN.2020.00081]
- 87 Ebeling TA, Compton AD, Albright DW. Clozapine-induced stuttering. *Am J Psychiatry* 1997; **154**: 1473 [PMID: 9326837 DOI: 10.1176/ajp.154.10.1473a]
- 88 Krishnakanth M, Haridas Phutane V, Muralidharan K. Clozapine-induced stuttering: a case series. *Prim Care Companion J Clin Psychiatry* 2008; **10**: 333-334 [PMID: 18787667 DOI: 10.4088/pcc.v10n0411e]
- 89 Kumar T, Kathpal A, Longshore CT. Dose dependent stuttering with clozapine: a case report. *Asian J Psychiatr* 2013; **6**: 178-179 [PMID: 23466117 DOI: 10.1016/j.ajp.2012.08.004]
- 90 Nagendrappa S, Sreeraj VS, Venkatasubramanian G. "I Stopped Hearing Voices, Started to Stutter" - A Case of Clozapine-Induced Stuttering. *Indian J Psychol Med* 2019; **41**: 97-98 [PMID: 30783319 DOI: 10.4103/IJPSYM.IJPSYM_157_18]
- 91 Aukst-Margetić B, Margetić B. Stuttering as a side-effect of divalproex sodium. *Psychiatry Clin Neurosci* 2008; **62**: 748 [PMID: 19068016 DOI: 10.1111/j.1440-1819.2008.01878.x]

- 92 **Mukherjee S**, Sen S, Chatterjee SS, Biswas A, Tripathi SK. Divalproex-induced stuttering: A rare case report. *Eur J Psychol Educat Studies* 2015; **2**: 25 [DOI: [10.4103/2395-2555.161419](https://doi.org/10.4103/2395-2555.161419)]
- 93 **Masand P**. Desipramine-induced oral-pharyngeal disturbances: stuttering and jaw myoclonus. *J Clin Psychopharmacol* 1992; **12**: 444-445 [PMID: [1474184](https://pubmed.ncbi.nlm.nih.gov/1474184/) DOI: [10.1097/00004714-199212000-00014](https://doi.org/10.1097/00004714-199212000-00014)]
- 94 **Nissani M**, Sanchez EA. Stuttering caused by gabapentin. *Ann Intern Med* 1997; **126**: 410 [PMID: [9054293](https://pubmed.ncbi.nlm.nih.gov/9054293/) DOI: [10.7326/0003-4819-126-5-199703010-00018](https://doi.org/10.7326/0003-4819-126-5-199703010-00018)]
- 95 **Catania S**, Cross H, de Sousa C, Boyd S. Paradoxical reaction to lamotrigine in a child with benign focal epilepsy of childhood with centrottemporal spikes. *Epilepsia* 1999; **40**: 1657-1660 [PMID: [10565596](https://pubmed.ncbi.nlm.nih.gov/10565596/) DOI: [10.1111/j.1528-1157.1999.tb02053.x](https://doi.org/10.1111/j.1528-1157.1999.tb02053.x)]
- 96 **Margetic B**. Stuttering (first report) in an elderly patient: case report. *Reactions* 2009; **1257**: 27-28
- 97 **Margetic B**, Aukst-Margetic B, Krajinovic B. A case of stuttering during treatment with levomepromazine. *Psychopharmacol Bull* 2009; **42**: 8-10 [PMID: [19204648](https://pubmed.ncbi.nlm.nih.gov/19204648/)]
- 98 **Netski AL**, Piasecki M. Lithium-induced exacerbation of stutter. *Ann Pharmacother* 2001; **35**: 961 [PMID: [11485152](https://pubmed.ncbi.nlm.nih.gov/11485152/) DOI: [10.1345/aph.10202](https://doi.org/10.1345/aph.10202)]
- 99 **Gulack BC**, Puri NV, Kim WJ. Stutter exacerbated by lithium in a pediatric patient with bipolar disorder. *Ann Pharmacother* 2011; **45**: e57 [PMID: [21917558](https://pubmed.ncbi.nlm.nih.gov/21917558/) DOI: [10.1345/aph.1Q140](https://doi.org/10.1345/aph.1Q140)]
- 100 **Sabillo S**, Samala RV, Ciocon JO. A stuttering discovery of lithium toxicity. *J Am Med Dir Assoc* 2012; **13**: 660-661 [PMID: [22749636](https://pubmed.ncbi.nlm.nih.gov/22749636/) DOI: [10.1016/j.jamda.2012.05.014](https://doi.org/10.1016/j.jamda.2012.05.014)]
- 101 **Alpaslan AH**, Coşkun KŞ, Kocak U, Görtüçü Y. Stuttering Associated With the Use of Short-Acting Oral Methylphenidate. *J Clin Psychopharmacol* 2015; **35**: 739-741 [PMID: [26436866](https://pubmed.ncbi.nlm.nih.gov/26436866/) DOI: [10.1097/JCP.0000000000000403](https://doi.org/10.1097/JCP.0000000000000403)]
- 102 **Copur M**, Copur S. Emergence of stuttering in an attention deficit hyperactivity disorder patient treated with methylphenidate. *Dusunen Adam* 2018; **31**: 222-224 [DOI: [10.5350/DAJPN2018310212](https://doi.org/10.5350/DAJPN2018310212)]
- 103 **Lasić D**, Cvitanović MŽ, Krnić S, Uglešić B. Olanzapine induced stuttering: a case report. *Psychiatr Danub* 2016; **28**: 299-300 [PMID: [27658840](https://pubmed.ncbi.nlm.nih.gov/27658840/)]
- 104 **Asan O**, Yaylaci ET, Okay IT, Goka E. A case of stuttering due to olanzapine treatment. *Dusunen Adam J Psychiatry Neurolog Sci* 2018; **31**: 405 [DOI: [10.5350/DAJPN2018310410](https://doi.org/10.5350/DAJPN2018310410)]
- 105 **McClean MD**, McLean Jr A. Case report of stuttering acquired in association with phenytoin use for post-head-injury seizures. *J Fluency Disord* 1985; **10**: 241-255 [DOI: [10.1016/0094-730X\(85\)90023-3](https://doi.org/10.1016/0094-730X(85)90023-3)]
- 106 **Ekici MA**, Ekici A, Ozdemir O. Phenytoin-induced stuttering: an extremely rare association. *Pediatr Neurol* 2013; **49**: e5 [PMID: [23859866](https://pubmed.ncbi.nlm.nih.gov/23859866/) DOI: [10.1016/j.pediatrneurol.2013.03.011](https://doi.org/10.1016/j.pediatrneurol.2013.03.011)]
- 107 **Giray E**, Şanal Toprak C, Saçaklıdır R, Gündüz OH. Pregabalin-Associated Stuttering in a Patient With Complex Regional Pain Syndrome: A Case Report. *J Clin Psychopharmacol* 2016; **36**: 740-742 [PMID: [27755156](https://pubmed.ncbi.nlm.nih.gov/27755156/) DOI: [10.1097/JCP.0000000000000609](https://doi.org/10.1097/JCP.0000000000000609)]
- 108 **Ge L**, Li A, Wang N, Li P, Xin H, Li W. Pregabalin-associated stuttering and frequent blepharospasm: case report and review. *Daru* 2020; **28**: 815-818 [PMID: [32632575](https://pubmed.ncbi.nlm.nih.gov/32632575/) DOI: [10.1007/s40199-020-00354-9](https://doi.org/10.1007/s40199-020-00354-9)]
- 109 **Makela EH**, Sullivan P, Taylor M. Sertraline and speech blockage. *J Clin Psychopharmacol* 1994; **14**: 432-433 [PMID: [7884029](https://pubmed.ncbi.nlm.nih.gov/7884029/) DOI: [10.1097/00004714-199412000-00015](https://doi.org/10.1097/00004714-199412000-00015)]
- 110 **Rosenfield DB**, McCarthy M, McKinney K, Viswanath NS, Nudelman HB. Stuttering induced by theophylline. *Ear Nose Throat J* 1994; **73**: 914, 918-920 [PMID: [7882883](https://pubmed.ncbi.nlm.nih.gov/7882883/) DOI: [10.1177/014556139407301208](https://doi.org/10.1177/014556139407301208)]
- 111 **Gérard JM**, Delecluse F, Robience Y. Theophylline-induced stuttering. *Mov Disord* 1998; **13**: 847-848 [PMID: [9756159](https://pubmed.ncbi.nlm.nih.gov/9756159/) DOI: [10.1002/mds.870130518](https://doi.org/10.1002/mds.870130518)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

