

Reviewer's Comment	Our response	Changes made to the manuscript
This is a guideline that outlines how to screen, treat, and monitor ROP under pandemic situations caused by the Covid-19 virus.	We thank the Reviewer for the kind comments in recognizing this as a special guideline for ROP screening under the given COVID-19 pandemic conditions	None suggested
In this situation, it would be a good idea to think about writing once.	We agree. This has not been submitted to any other journal	None suggested
In the case of a solution in the text, such as alcohol or water, it would be good to clearly state the correct concentration and what kind of water it is.	We agree We have replaced all proprietary names with an explanation of liquid mixture of 1- Propanol and 2- Propanol with total alcohol concentration of 75% to replace Sterillium	Replaced Sterillium / Bacillocid with "liquid mixture of 1- Propanol and 2- Propanol with total alcohol concentration of 75%"
Check the spelling in "13. Between each patient, hands must be washed and Strerlium be used and allowed to dry before handling the patient".	Thank you for pointing out the typo	Replaced with the correct spelling
Please refer to the most important tables, Tables 2 and 3. It would be good if you could attach a background reference to each of the reasons for this follow-up or treatment.	Thank you for the suggestion Whereas table 2 are the committee's suggestions and hence there is no background reference, we have added the references to Table 3 in the appropriate rows	Table 3 has been edited to include 2 background references – ETROP (ref 14) and APROP (ref 15) in row 1 and 2 respectively