

Match Overview

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 65826

Manuscript Type: CASE REPORT

Malignant peripheral nerve sheath tumor in an elderly patient with superficial spreading melanoma: A case report

Chong-Miao Yang, Jia-Min Li, Rui Wang, Li-Gong Lu

Abstract

BACKGROUND

Malignant peripheral nerve sheath tumor (MPNST) is a type of spindle cell sarcoma originating from the peripheral nerve, which usually results in the corresponding nerve



Malignant peripheral nerve sheath tumor of the left knee in an elder



ALL IMAGES VIDEOS

26,800 Results Any time ▾ Open links in new tab ☒

Malignant melanoma with neural differentiation: an ... europepmc.org/articles/PMC4079032 ▾

Jan 01, 2014 · The differential diagnostic considerations associated with neurotropic **melanoma** may include a variety of other spindle cell lesions, including benign and **malignant peripheral nerve shea...**

Metastatic Malignant Melanoma Resembling Malignant ... <https://www.researchgate.net/publication/12716017>...

We report 16 cases of metastatic **malignant melanoma** presenting clinically as lymphadenopathy or a soft-tissue mass and histologically resembling **malignant peripheral nerve sheath tumor** (MPNST).

Ewing Sarcoma Treatment (PDQ®)—Health Professional ... <https://www.cancer.gov/types/bone/hp/ewing-treatment-pdq> ▾

Patients with Ewing sarcoma as a second **malignant** neoplasm were older (secondary Ewing sarcoma, mean age of 47.8 years; primary Ewing sarcoma, mean age of 22.5 years), more likely to have a primar...

Soft tissue sarcoma - Symptoms and causes - Mayo Clinic <https://www.mayoclinic.org/diseases-conditions/>...

Overview Symptoms Causes Risk Factors

Soft tissue sarcoma is a rare type of cancer that begins in the tissues that connect, support and surround other body structures. This includes muscle, fat, blood vessels, nerves, tendons and the lining of your joints. More than 50 subtypes of **soft tissue sarcoma** exist. Some types are more likely to affect children, while others affect mostly adults. These tumors can be difficult to diagnose because they may be mistaken for many other types of growths. **Soft tissue sarcoma** can occur anywhere in...

[See more on mayoclinic.org](https://www.mayoclinic.org)

Published: Dec 02, 2020

Malignant peripheral nerve sheath tumor arising from the ... <https://www.researchgate.net/publication/7296215>...

A **malignant peripheral nerve sheath tumor** (MPNST) is a rare neoplasm arising from **peripheral nerve sheath**. Here, we report the first case of MPNST arising in the colon and also the youngest ca...



Malignant peripheral nerve sheath tumor in an elderly patient with



ALL

IMAGES

VIDEOS

21,900 Results

Any time ▾

[Rare primary malignant peripheral nerve sheath tumor of ...](https://www.spandidos-publications.com/10.3892/mco.2021.2306)

<https://www.spandidos-publications.com/10.3892/mco.2021.2306> ▾

Malignant peripheral nerve sheath tumor (MPNST) is a rare disease. The present study discusses the case of a 30-year-old male **patient** who presented with complaints of pain in the left testis and groin ov...

[Malignant Peripheral Nerve Sheath Tumor - an overview ...](https://www.sciencedirect.com/.../malignant-peripheral-nerve-sheath-tumor)

<https://www.sciencedirect.com/.../malignant-peripheral-nerve-sheath-tumor>

Melanoma tends to be a dermal-based **tumor** arising in sun-damaged skin of the head and neck in **elderly patients**. **Malignant peripheral nerve sheath tumor** tends to be a deep soft tissue-based...

[Malignant Peripheral Nerve Sheath Tumor of the Orbit: Case ...](https://www.researchgate.net/publication/7615023...)

<https://www.researchgate.net/publication/7615023...>

Duane M. Ilstrup. A review was done of 120 cases of **malignant peripheral nerve sheath tumor** (MPNST) seen during a 71-year period. Of the 120 **patients**, 52 were males and 68 were females with ...

Malignant peripheral nerve sheath tumor

Symptoms

Diagnosis



A malignant peripheral nerve sheath tumor is a form of cancer of the connective tissue surrounding nerves. Given its origin and behavior it is classified as a sarcoma. About half the cases are diagnosed in people with neurofibromatosis; the lifetime risk for an MPNST in patients with neurofibromatosis type 1 is 8–13%. MPNST with rhabdomyoblastomatous component are

ALL IMAGES VIDEOS MAPS NEWS SHOPPING

Also try: malignant peripheral nerve sheath tumor pathology outlines · triton tumor pathology

31,800 Results Any time ▾

Malignant Peripheral Nerve Sheath Tumor - an overview ...

<https://www.sciencedirect.com/.../malignant-peripheral-nerve-sheath-tumor>

Melanoma tends to be a dermal-based tumor arising in sun-damaged skin of the head and neck in elderly patients. Malignant peripheral nerve sheath tumor tends to be a deep soft tissue-based tumor nodule...

What Is a Soft Tissue Sarcoma? - American Cancer Society

<https://www.cancer.org/cancer/soft-tissue-sarcoma/about/soft-tissue-sarcoma> ▾

Malignant peripheral nerve sheath tumors include neurofibrosarcomas, malignant schwannomas, and neurogenic sarcomas. These are sarcomas that start in the cells that surround a nerve....

Types of Soft Tissue Sarcomas

There are more than 50 different types of soft tissue sarcomas. Some are quite rare, and not all are listed here: 1. Adult fibrosarcoma usually aff...

Intermediate Soft Tissue Tumors

These may grow and invade nearby tissues and organs, but they tend to not spread to other parts of the body. 1. Dermatofibrosarcoma protuberans is...

Benign Soft Tissue Tumors

Many benign tumors, or tumors that are not cancer, can start in soft tissues. These include: 1. Elastofibromas: benign tumors of fibrous tissue 2....

Recurrent epithelioid malignant peripheral nerve sheath ...

<https://www.spandidos-publications.com/10.3892/ol.2019.10676> ▾

< Introduction Case Report Discussion Acknowledgements >

Neurofibromatosis type-1 (NF-1) is a common,autosomal dominant neurocutaneous disorder with an incidence of1:3,000 worldwide, which primarily involves the skin, bone andnervous system (1). The standardclinical diagnostic criteria for NF-1 have existed since 1987, with≥2 criteria required for a positive diagnosis. The criteria aresummarized as follows: ≥6 'café-au-lait' spots, size >5 mm inprepubertal children or >15 mm in post pubertal individuals,axillary or inguinal freckling, ≥2 cutaneous neurofibroma...

See more on spandidos-publications.com

Cited by: 1 Author: Peng Du, Jia Zhu, Zhen-Dong Zhang, Chon...

Malignant peripheral nerve sheath tumor

Symptoms Diagnosis Causes Treatments >

A malignant peripheral nerve sheath tumor is a form of cancer of the connective tissue surrounding nerves. Given its origin and behavior it is classified as a sarcoma. About half the cases are diagnosed in people with neurofibromatosis; the lifetime risk for an MPNST in patients with neurofibromatosis type 1 is 8–13%. MPNST with rhabdomyoblastomatous component are called malignant triton tumors.

Wikipedia

People also search for

- Schwannoma
- Neurofibroma
- Neurofibromatosis
- Neuroblastoma
- Ganglioneuroma
- See more ▾

Data from: Wikipedia
Suggest an edit