

[\(PDF\) Massive haemoperitoneum due to rupture of splenic ...](#)<https://www.researchgate.net/publication/13523600...>

There are no previous reports of rupture of an abdominal organ during chronic peritoneal dialysis. However massive intraperitoneal hemorrhage in CAPD has been described secondary to abdominal ...

[\(PDF\) Hemoperitoneum: bloody peritoneal dialysate in ESRD ...](#)<https://www.academia.edu/23466092> ▾

Hemoperitoneum after catheter placement can result in catheter failure (60%) and adhesion formation, especially if the abdomen is dry. When the abdomen was irrigated until the PD fluid was clear, followed...

[Images of Abdominal Hemorrhage After Peritoneal Dialysis C...](#)<bing.com/images>[See all images >](#)[\(PDF\) Ruptured Abdominal Aortic Aneurysm As a Cause of ...](#)<https://www.researchgate.net/publication/51696538...>

Bleeding into the peritoneal cavity (hemoperitoneum) is one of the complications of chronic peritoneal dialysis (PD). Its occurrence varies from 6.1 to 8.4% of patients receiving this treatment.

[Hemoperitoneum: Bloody Peritoneal Dialysate in Esrd ...](#)<https://journals.sagepub.com/doi/full/10.1177/089686080702700303>

Hemoperitoneum can develop shortly after peritoneal catheter insertion. Bleeding diathesis due to

国内版 国际版

Abdominal hemorrhage after peritoneal dialysis catheter insertion -



ALL IMAGES VIDEOS

50,800 Results

Any time ▾

(PDF) Massive haemoperitoneum due to rupture of splenic ...

<https://www.researchgate.net/publication/13523600...>

There are no previous reports of rupture of an abdominal organ during chronic peritoneal dialysis. However massive intraperitoneal hemorrhage in CAPD has been described secondary to abdominal ...

Hemoperitoneum: Bloody Peritoneal Dialysate in Esrd ...

<https://journals.sagepub.com/doi/full/10.1177/089686080702700303>

May 01, 2007 · Hemoperitoneum after catheter placement can result in catheter failure (60%) and adhesion formation, especially if the abdomen is dry. When the abdomen was irrigated until the PD fluid was clear, followed by low volume (500 – 1000 mL) short dwells, there was no catheter failure .

Cited by: 35

Author: Susie Q. Lew

Publish Year: 2007

PEOPLE ALSO ASK

Can hemoperitoneum cause peritoneal dialysis? ▾

What is peritoneal bleeding? ▾

How to manage hemoperitoneum? ▾

When does hemoperitoneum occur with retrograde menstruation? ▾

Feedback

(PDF) Hemoperitoneum: bloody peritoneal dialysate in ESRD ...

<https://www.academia.edu/23466092> ▾

Hemoperitoneum after catheter placement can result in catheter failure (60%) and adhesion formation, especially if the abdomen is dry. When the abdomen was irrigated until the PD fluid was clear, followed by low volume (500 -1000 mL) short dwells, there was no catheter failure (57).

(PDF) Ruptured Abdominal Aortic Aneurysm As a Cause of ...

<https://www.researchgate.net/publication/51696538...>

Bleeding into the peritoneal cavity (hemoperitoneum) is one of the complications of chronic peritoneal ...

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 66648

Manuscript Type: CASE REPORT

**Abdominal hemorrhage after peritoneal dialysis catheter insertion – a rare reason for
luteal rupture: A case report**

Gan LW *et al.* Postoperative luteal rupture

Lin-Wang Gan, Qian-Cheng Li, Zhao-Lan Yu, Li-Ling Zhang, Qi Liu, Ying Li, San-Tao
Ou

Match Overview

1	Internet 31 words created on 16-Oct-2019 www.xipnut.com	2%
2	Internet 13 words created on 16-Jan-2019 casereports.bmj.com	1%

国内版 国际版

Abdominal hemorrhage after peritoneal dialysis catheter insertion -



ALL IMAGES VIDEOS

50,200 Results

Any time ▾

(PDF) Massive haemoperitoneum due to rupture of splenic ...

<https://www.researchgate.net/publication/13523600...>

There are no previous reports of rupture of an abdominal organ during chronic peritoneal dialysis. However massive intraperitoneal hemorrhage in CAPD has been described secondary to abdominal ...

Estimated Reading Time: 4 mins

Hemoperitoneum: Bloody Peritoneal Dialysate in Esrd ...

<https://journals.sagepub.com/doi/full/10.1177/089686080702700303>

May 01, 2007 · Hemoperitoneum after catheter placement can result in catheter failure (60%) and adhesion formation, especially if the abdomen is dry. When the abdomen was irrigated until the PD fluid was clear, followed by low volume (500 – 1000 mL) short dwells, there was no catheter failure .

Cited by: 35

Author: Susie Q. Lew

Publish Year: 2007

PEOPLE ALSO ASK

Can hemoperitoneum cause peritoneal dialysis? ▾

What is peritoneal bleeding? ▾

How to manage hemoperitoneum? ▾

When does hemoperitoneum occur with retrograde menstruation? ▾

Feedback

(PDF) Hemoperitoneum: bloody peritoneal dialysate in ESRD ...

<https://www.academia.edu/23466092> ▾

Hemoperitoneum after catheter placement can result in catheter failure (60%) and adhesion formation, especially if the abdomen is dry. When the abdomen was irrigated until the PD fluid was clear, followed by low volume (500 -1000 mL) short dwells, there was no catheter failure (57).

Estimated Reading Time: 10 mins

(PDF) Ruptured Abdominal Aortic Aneurysm As a Cause of ...

<https://www.researchgate.net/publication/51606538>