

Including results for successful outcome of retrograde **pancreaticojejunostomy** for chronic pancreatitis and infected pancreatic cysts a case report.

Do you want results only for Successful outcome of retrograde pancreatojejunostomy for chronic pancreatitis and infected pancreatic cysts: a case report?

### Outcome of Pancreaticojejunostomy After Previous ...

<https://www.researchgate.net/publication/11027213...>

During a 10 year period, 69 patients with pancreatic duct dilation of 7 mm or more and intractable pain from chronic pancreatitis underwent Roux-Y drainage either as a lateral pancreatojejunostomy ...

### Pancreaticojejunostomy For Severe Symptomatic Chronic ...

<https://www.researchgate.net/publication/21544457...>

The role of operation, particularly pancreaticojejunostomy, in the treatment of abdominal pain from chronic pancreatitis is controversial, but relief of pancreatic duct obstruction may decrease ...

#### PEOPLE ALSO ASK

How does pain decrease and duration of chronic pancreatitis? ▾

How big can a pseudocyst in the pancreas get? ▾

What causes fluid in the pseudocyst of the pancreas? ▾

How long does it take for a pseudocyst to resolve? ▾

Feedback

### Successful surgical internal drainage of postoperative ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4560123>

Background. Pancreatic pseudocyst (PPC) is associated with acute pancreatitis and chronic pancreatitis, and develops as a postoperative complication []. PPC is a localized collection of amylase-rich fluid located within or adjacent to the pancreas and is devoid of an epithelial wall []. Treatment is only required for persisting PPC symptoms such as abdominal pain, infection, or compression of ...

Author: Kazuya Sakata, Daisuke Hashimoto, ... Publish Year: 2015

**Successful surgical internal drainage of postoperative ...**

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4560123>

Jun 26, 2015 · Background. Pancreatic pseudocyst (PPC) is associated with acute pancreatitis and chronic pancreatitis, and develops as a postoperative complication []. PPC is a localized collection of amylase-rich fluid located within or adjacent to the pancreas and is devoid of an epithelial wall []. Treatment is only required for persisting PPC symptoms such as abdominal pain, infection, or ...

Author: Kazuya Sakata, Daisuke Hashimoto, ... Publish Year: 2015

**The Management of Acute and Chronic Pancreatitis**

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2886461>

Approximately 30% of patients with chronic pancreatitis develop an inflammatory enlargement in the pancreatic head with a concomitant obstruction of the pancreatic duct causing diffuse dilatation of the duct. 4 In these cases, there are several possible surgical procedures. One is called the "Frey procedure," in which the head of the pancreas is cored out and drained with the same loop of the defunctionalized jejunum that is utilized to decompress the dilated main pancreatic ...

Cited by: 107 Author: Peter A. Banks, Darwin L. Conwell, Philli... Publish Year: 2010

PEOPLE ALSO ASK

- What is the mortality rate from pancreatic resection? ▾
- What is the rate of recurrent pancreatitis in men? ▾
- How often does pancreatitis relapse after 10 years? ▾
- What is the goal of pancreatic enzyme replacement therapy? ▾

Feedback

**Chronic obstructive pancreatitis due to a pancreatic cyst ...**

<https://gut.bmj.com/content/42/1/131>

Jan 01, 1998 · Patient —The first case of painful chronic obstructive pancreatitis due to a true pancreatic cyst in a patient with autosomal dominant polycystic kidney disease is reported. Abdominal transperitoneal and endoscopic ultrasonography, computed tomography, and endoscopic retrograde

**Name of Journal:** *World Journal of Clinical Cases*  
**Manuscript NO:** 66786  
**Manuscript Type:** CASE REPORT

**Successful outcome of retrograde pancreatojejunostomy for chronic pancreatitis and infected pancreatic cysts: A case report**

Kimura K *et al.* Retrograde pancreatojejunostomy for chronic pancreatitis

**Abstract**

**BACKGROUND**

Chronic pancreatitis occasionally requires surgical treatment that can be performed with various techniques. Often, this type of surgery presents with postoperative complications. We report a case of a successful retrograde pancreatojejunostomy for chronic pancreatitis and infected pancreatic cysts.

Match Overview

Match	Source	Words	Created	Match %
1	Internet	35 words	Created on 27-Mar-2021 bmc.org, bmcjournal.com	2%
2	Internet	15 words	Created on 26-Jun-2019 publishing.bwh.com, windows.net	1%

国内版

国际版

Successful outcome of retrograde pancreatojejunostomy for chron



ALL

IMAGES

VIDEOS

12,200 Results

Any time ▾

### Successful surgical internal drainage of postoperative ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4560123>

Jun 26, 2015 · Background. Pancreatic pseudocyst (PPC) is associated with acute pancreatitis and chronic pancreatitis, and develops as a postoperative complication [].PPC is a localized collection of amylase-rich fluid located within or adjacent to the pancreas and is devoid of an epithelial wall [].Treatment is only required for persisting PPC symptoms such as abdominal pain, infection, or ...

Author: Kazuya Sakata, Daisuke Hashimoto, ... Publish Year: 2015

### Diagnostic and therapeutic recommendations for chronic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6173070>

Pancreatic pseudocysts are formed in the course of CP in 20–40% of patients, and spontaneously regress in 40% of cases, more often in case of lesions measuring less than 3 cm. Symptoms (pain, severe obstruction, jaundice, weight loss) and infection of the lesion are indications for the treatment of a pseudocyst . It was found that large cysts (> 6 cm) are much more often symptomatic and therefore ...

Cited by: 5 Author: Roland Kadaj-Lipka, Michał Lipiński, Krys...  
Publish Year: 2018

### Chronic obstructive pancreatitis due to a pancreatic cyst ...

<https://gut.bmj.com/content/42/1/131> ▾

Jan 01, 1998 · chronic obstructive pancreatitis; pancreatic cyst; autosomal dominant polycystic kidney disease; distal pancreatectomy; Autosomal dominant polycystic kidney disease (ADPKD) is the most frequent inherited polycystic disease, affecting 1 in 400 to 1 in 1000 people.1 It is a systemic disorder. Numerous and bilateral renal cysts are usually observed in the second decade of life and their ...

Cited by: 19 Author: D. Malka, P. Hammel, V. Vilgrain, J.-F. Fl...  
Publish Year: 1998 Estimated Reading Time: 11 mins

### Chronic pancreatitis with ductal stones in the pancreatic ...

<https://academic.oup.com/jscr/article/2020/10/rjaa352/5918937> ▾

Oct 07, 2020 · INTRODUCTION. Chronic pancreatitis (CP) involves progressive inflammatory and fibrotic changes of the exocrine pancreas owing to repeated episodes of acute inflammation over a long period [].Main risk factors for CP include alcohol abuse, smoking, gene mutations, autoimmune syndromes, metabolic disturbances, environmental conditions and anatomical abnormalities, which in ...

Author: Alberto Mayer, Andrea Johann Males Publish Year: 2020