



66947\_Auto\_Edited.docx

Quotes Excluded  
Bibliography Excluded11%  
SIMILAR**Name of Journal:** *World Journal of Clinical Cases***Manuscript NO:** 66947**Manuscript Type:** CASE REPORT**Prenatal Ultrasonographic Findings in Klippel-Trenaunay Syndrome: A Case Report**

The ultrasonographic findings of a case of KTS in utero

Houqing Pang, Qianqian Gao

## Match Overview

- | Rank | Source  | Words    | Similarity |
|------|---|----------|------------|
| 1    | <b>Crossref</b> Dongmei Yu, Lingyu Sun, Taotao Chen. "Prenatal ultrasound diagnosis of syndrome associated with umbilical cord hemang             | 67 words | 4%         |
| 2    | <b>Crossref</b> Olga Ivanitskaya, Elena Andreeva, Natalia Odegova. "Prenatal diagnosis of Klippel-Trenaunay syndrome: Series of four case         | 41 words | 2%         |
| 3    | <b>Internet</b> crawled on 24-Feb-2021<br><a href="http://www.cureus.com">www.cureus.com</a>  | 35 words | 2%         |
| 4    | <b>Crossref</b> Lu Wang, Yi Feng, Lu-Yang Jiang. "Anesthetic management of bilateral pheochromocytoma resection in Von Hippel-Lindau sy           | 16 words | 1%         |
| 5    | <b>Crossref</b> P. R. Coombs. "Sonographic identification of lower limb venous hypoplasia in the prenatal diagnosis of Klippel-Trenaunay syndrome | 13 words | 1%         |

Prenatal Ultrasonographic Findings in Klippel-Trenaunay Syndrome:



ALL

IMAGES

VIDEOS

93,000 Results

Any time ▾

## Early prenatal ultrasonic findings in Klippel-Trenaunay ...

<https://www.ncbi.nlm.nih.gov/pubmed/6087308>

Early **prenatal** ultrasonic **findings in Klippel-Trenaunay-Weber syndrome**. Seoud M, Santos-Ramos R, Friedman JM. Routine sonography prior to genetic amniocentesis revealed the presence of a large complex mass with pulsating channels over the anterior fetal chest wall.

**Cited by:** 9**Author:** Muhieddine Seoud, Rigoberto Santo...**Publish Year:** 1984

## Sonographic Identification of Klippel-Trenaunay-Weber ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3866887>

Dec 03, 2013 · We report the **prenatal sonographic findings** in a case of **Klippel-Trenaunay-Weber (KTW) syndrome** including fetal limb hypertrophy and large subcutaneous cystic lesions. **Prenatal diagnosis** is possible by **ultrasound** examination and recognition important for prevention of complications and future management.

**Cited by:** 4**Author:** Yigit Cakiroglu, Emek Doğer, Sule Yil...**Publish Year:** 2013

## Klippel-Trenaunay Syndrome: A Case Report

<https://pubmed.ncbi.nlm.nih.gov/27885298>

**Klippel-Trenaunaysyndrome** (KTS) is a rare congenital condition usually presenting with port wine stains, excessive growth of bones and soft tissue and varicose veins which most commonly occurs in the legs, but it also may affect the arms, face, head, or internal organs. We **report** a **case** of term male neonate with clinical findings of Port-wine stain, multiple cystic swellings with **ultrasonographic findings** ...

**Cited by:** 1**Author:** R K Sah, S Sharma, S Ghimire, B B B...**Publish Year:** 2016

## Prenatal sonographic findings of Klippel-Trénaunay-Weber ...

<https://onlinelibrary.wiley.com/doi/10.1002/jcu.20323>

Abstract We describe the **prenatal sonographic findings in a case** of Klippel-Trénaunay-Weber **syndrome** including fetal ascites and subcutaneous cystic lesions associated with ...

**Author:** Chih-Ping Chen, Shuan-Pei Lin, Tun...

Prenatal Ultrasonographic Findings in Klippel-Trenaunay Syndrome:



Sign in



ALL

IMAGES

VIDEOS

Add the Give with Bing extension &gt;

21,700 Results

Any time ▾

### [Sonographic Identification of Klippel-Trenaunay-Weber ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3866887)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3866887>

Dec 03, 2013 · We report the **prenatal sonographic findings** in a case of **Klippel-Trenaunay-Weber (KTW) syndrome** including fetal limb hypertrophy and large subcutaneous cystic lesions. **Prenatal diagnosis** is possible by **ultrasound** examination and recognition important for prevention of complications and future management.

**Cited by:** 4**Author:** Yigit Cakiroglu, Emek Doğer, Sule Yildiri...**Publish Year:** 2013

### [Klippel-Trenaunay Syndrome: A Case Report](https://pubmed.ncbi.nlm.nih.gov/27885298)

<https://pubmed.ncbi.nlm.nih.gov/27885298>

**Klippel-Trenaunaysyndrome** (KTS) is a rare congenital condition usually presenting with port wine stains, excessive growth of bones and soft tissue and varicose veins which most commonly occurs in the legs, but it also may affect the arms, face, head, or internal organs. We **report a case** of term male neonate with clinical findings of Port-wine stain, multiple cystic swellings with **ultrasonographic findings**

...

**Cited by:** 2**Author:** R K Sah, S Sharma, S Ghimire, B B Bag...**Publish Year:** 2016

### [Prenatal diagnosis of Klippel-Trenaunay syndrome: Series ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7254946)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7254946>

Oct 17, 2019 · Klippel-Trenaunay syndrome (KTS, OMIM 149000) is a rare disease, characterised by a classic triad of port wine stains, varicose veins, and bony and soft tissue hypertrophy of an extremity. Other anomalies reported include osseous haemangiomas, which may involve any region of

## Klippel-Trenaunay Syndrome

Medical condition

Symptoms

Diagnosis

Causes

Treatments



Klippel-Trenaunay (klih-PEL tray-no-NAY) syndrome is a rare disorder found at birth (congenital) involving abnormal development of blood vessels, soft tissues, bones and the lymphatic system.

[MayoClinic](#)

### Symptoms

Symptoms of Klippel-Trenaunay syndrome include:

- Many port wine stains or other blood vessel problems, including dark spots on the skin
- Varicose veins (may be seen in early infancy, but are more likely to be seen later in childhood or adolescence)...

[Read more on MedlinePlus](#)

### Causes

Most people with Klippel-Trenaunay syndrome are born with a port-wine stain. This type of birthmark is caused by swelling of small blood vessels near the surface of the skin. Port-wine stains are typically flat and can vary

Sonographic Identification of Klippel-Trenaunay-Weber Syndrome

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3866887

Dec 03, 2013 · We report the prenatal sonographic findings in a case of Klippel-Trenaunay-Weber (KTW) syndrome including fetal limb hypertrophy and large subcutaneous cystic lesions. Prenatal diagnosis is possible by ultrasound examination and recognition important ...

Cited by: 4

Author: Yigit Cakiroglu, Emek Doğer, Sule Yildirim K...

Publish Year: 2013

Klippel-Trenaunay Syndrome: A Case Report

https://pubmed.ncbi.nlm.nih.gov/27885298

Klippel-Trenaunaysyndrome (KTS) is a rare congenital condition usually presenting with port wine stains, excessive growth of bones and soft tissue and varicose veins which most commonly occurs in the legs, but it also may affect the arms, face, head, or internal organs. We report a case of term male neonate with clinical findings of Port-wine stain, multiple cystic swellings with ultrasonographic findings ...

Cited by: 2

Author: R K Sah, S Sharma, S Ghimire, B B Bagale, ...

Publish Year: 2016

Prenatal diagnosis of Klippel–Trenaunay syndrome: Series ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7254946

Oct 17, 2019 · Klippel–Trenaunay syndrome (KTS, OMIM 149000) is a rare disease, characterised by a classic triad of port wine stains, varicose veins, and bony and soft tissue hypertrophy of an extremity.10ther anomalies reported include cavernous haemangiomas, which may involve any region of the body (particularly limbs and trunk) and any organ (especially the

Author: Olga Ivanitskaya, Elena Andreeva, Natal...

Publish Year: 2020

Prenatal sonographic findings of Klippel-Trénaunay-Weber ...

https://onlinelibrary.wiley.com/doi/abs/10.1002/jcu.20323

The full text of this article hosted at iucr.org is unavailable due to technical difficulties.

Cited by: 13

4.5/5 ★★★★★ (3)

Author: Chih-Ping Chen, Shuan-Pei Lin, Tung-Ya...

Publish Year: 2007

Prenatal sonographic findings of Klippel-Trénaunay-Weber ...

https://onlinelibrary.wiley.com/doi/10.1002/jcu.20323

Klippel-Trenaunay Syndrome

Medical condition

Symptoms

Diagnosis

Causes

Treatments

>

Klippel-Trenaunay (klih-PEL tray-no-NAY) syndrome is a rare disorder found at birth (congenital) involving abnormal development of blood vessels, soft tissues, bones and the lymphatic system.

MayoClinic

Symptoms

Symptoms of Klippel-Trenaunay syndrome include:

- Many port wine stains or other blood vessel problems, including dark spots on the skin
- Varicose veins (may be seen in early infancy, but are more likely to be seen later in childhood or adolescence)...

Read more on MedlinePlus

Causes

Most people with Klippel-Trenaunay syndrome are born with a port-wine stain. This type of birthmark is caused by swelling of small blood vessels near the surface of the skin. Port-wine stains are typically flat and can vary from pale pink to deep...

Read more on NIH

Treatments

You and your doctor can work together to determine which of the following treatments are most appropriate for you:

Compression therapy. Bandages or elastic garments are wrapped around affected limbs to help prevent swelling, varicose veins and...

Read more on MayoClinic