

Removal of Ingested Magnetic Bodies via Laparoscopic ...

<https://europepmc.org/article/MED/33101833> ▾

Jan 01, 2020 · Search worldwide, life-sciences literature Search. Advanced Search Coronavirus articles and preprints Search examples: "breast cancer" Smith J

Single-incision assisted laparoscopic surgery (SILS) in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4457091>

Jun 02, 2015 · In this case report, we describe the successful treatment of a small-bowel intussusception, which was caused by a 3cm solitary hamartomatous polyp, with single-incision...

Cited by: 8

Author: Michael Pitiakoudis, Konstantinos Roma...

Publish Year: 2015

Search Tools

Turn off Hover Translation (关闭悬停)

PEOPLE ALSO ASK

How is a laparoscopic small bowel resection performed? ▾

What kind of surgery is used to remove the large intestine? ▾

How are the cut ends of the small bowel sewn together? ▾

激活 Windows
转到“设置”以激活 Windows。

ALL

IMAGES

VIDEOS

12,000 Results

Any time ▾

Single-incision assisted laparoscopic surgery (SILS) in ...<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4457091>

Jun 02, 2015 · In this **case report**, we describe the **successful** treatment of a **small-bowel** intussusception, which was caused by a 3cm solitary hamartomatous polyp, with single-incision laparoscopic surgery. Single-incision laparoscopic surgery is a minimally invasive surgical procedure with important advantages that allows the reduction of the intussusception and the resection of the polyp.

Cited by: 8

Author: Michael Pittakoudis, Konstantinos Romani...

Publish Year: 2015

Colonoscopic perforation: Incidence, risk factors ...<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2811793>

Jan 28, 2010 · A **small** perforation site caused by therapeutic **colonoscopy** has been shown to have a better **success** rate with medical treatment. Colonic stricture following conservative treatment of a colonoscopic perforation has been reported in the literature[57], but this can be safely managed by either endoscopic dilatation or surgery.

Cited by: 217

Author: Varut Lohsirirwat

Publish Year: 2010

Search Tools

Turn off Hover Translation (关闭取词)

People also ask

How does a surgeon remove the large intestine? ▾

What was the name of the surgery to remove the colon? ▾

国内版

国际版

Successful removal of two magnets in the small intestine by laparo



ALL

IMAGES

VIDEOS

12,000 Results

Any time ▾

Single-incision assisted laparoscopic surgery (SILS) in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4457091>

Jun 02, 2015 · In this **case report**, we describe the **successful** treatment of a **small-bowel** intussusception, which was caused by a 3cm solitary hamartomatous polyp, with single-incision laparoscopic surgery. Single-incision laparoscopic surgery is a minimally invasive surgical procedure with important advantages that allows the reduction of the intussusception and the resection of the polyp.

Cited by: 8

Author: Michael Pitiakoudis, Konstantinos Roma...

Publish Year: 2015

Colonoscopic perforation: Incidence, risk factors ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2811793>

Jan 28, 2010 · A **small** perforation site caused by therapeutic **colonoscopy** has been shown to have a better **success** rate with medical treatment. Colonic stricture following conservative treatment of a colonoscopic perforation has been reported in the literature[57], but this can be safely managed by either endoscopic dilatation or surgery.

Cited by: 217

Author: Varut Lohsiriwat

Publish Year: 2010

PEOPLE ALSO ASK

How is the small intestine removed in laparoscopic surgery? ▾

What is the name of the procedure to remove the right side of the colon? ▾

What's the name of the procedure to cut the small bowel? ▾

How are polyps removed from the large intestine? ▾

Feedback

[PDF] A Case Report of Complete Intraoperative Double- ...

www.scielo.org.co/pdf/rcg/v29n1/en_v29n1a11.pdf

A Case Report of Complete Intraoperative Double- ... Peutz-Jeghers syndrome is characterized by hamartomatous polyps primarily located in the **small intestine**. Most of these patients undergo several