

Natural history of untreatable hepatocellular carcinoma: A ...

<https://pubmed.ncbi.nlm.nih.gov/23060970>

Natural history of untreatable hepatocellular carcinoma: A retrospective cohort study. This study confirms the heterogeneous behavior of untreated HCC. BCLC staging remains an important prognostic guide and may be important in decision-making for palliative treatment.

Cited by: 81

Author: Giuseppe Cabibbo, Marcello Maida, Chiara ...

Publish Year: 2012

Combination therapy with sorafenib and radiofrequency ...

<https://pubmed.ncbi.nlm.nih.gov/25403366>

Methods: One hundred and twenty-eight patients with HCC from 12 centers were enrolled in this retrospective study; 64 patients who received Sorafenib plus RFA (Sorafenib-RFA) were compared with a control group treated with RFA alone. The two patient groups were selected with a predefined criterion and matched in terms of their clinical and tumor characteristics at baseline.

Clinical characteristics of non-B non-C hepatocellular ...

<https://pubmed.ncbi.nlm.nih.gov/22156485>

Non-ALD-HCC tended to be diagnosed at a more advanced stage, whereas liver function was worse, and tumor recurrence rate was higher in ALD-HCC patient ... Clinical characteristics of non-B non-C hepatocellular carcinoma: a single-center retrospective study

Early versus late recurrence of centrally located ...

<https://www.ncbi.nlm.nih.gov/pubmed/31169667>

Early versus late recurrence of centrally located hepatocellular carcinoma after mesohepatectomy: A cohort study based on the STROBE guidelines. Zhao J(1), Li W(2), Mao J(1). Author information:



Study on the clinical stages of recurrent hepatocellular carcinoma:



ALL IMAGES VIDEOS

172,000 Results Any time ▾

Predictors of Recurrence in Hepatitis C Virus Related ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4261437>

Jul 22, 2013 · This is a **retrospective cohort study** of 208 cases of HCC in hepatitis C virus positive patients with cirrhotic livers who underwent first-time liver resection, in Gastroenterology Surgical Centre, Mansoura University, Egypt during the period from January 2002 to December 2011. Shapiro-Wilk test was used to assess normality of data.

Cited by: 6 Author: Wahab Ma, Shehta A, Hamed H, El Nake...

Publish Year: 2014



Natural history of untreatable hepatocellular carcinoma: A ...


<https://pubmed.ncbi.nlm.nih.gov/23060970>

Aim: To investigate the clinical course of untreatable hepatocellular carcinoma (HCC) identified at any

30-Jul-2021 05:00PM

1710 words • 7 matches • 3 sources

FAQ

iThenticate

67641_Auto_Edited.docx

Quotient Excluded
Bibliography Excluded
8%
0.0000

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 67641

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Study on the clinical stages of recurrent hepatocellular carcinoma : a retrospective cohort study

Si-Yang Yao, Bin Liang, Yuan-Yuan Chen, Yun-Tian Tang, Xiao-Feng Dong, Tian-Qi Liu

Abstract

BACKGROUND

Match Overview

1	Internet 70 words created on 05-May-2015 www.tdfr	4%
2	Internet 65 words created on 04-Jun-2021 proquest.broadus.org	3%
3	Internet 21 words created on 20-Jul-2021 www.springer.com	1%

Page: 1 of 2

Full-Only Report

国内版

国际版

Study on the clinical stages of recurrent hepatocellular carcinoma:



ALL

IMAGES

VIDEOS

171,000 Results

Any time ▾

Predictors of Recurrence in Hepatitis C Virus Related ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4261437>

Jul 22, 2013 · This is a **retrospective cohort study** of 208 cases of HCC in **hepatitis C virus** positive patients with cirrhotic livers who underwent first-time liver resection, in Gastroenterology Surgical Centre, Mansoura University, Egypt during the period from January 2002 to December 2011. Shapiro-Wilk test was used to assess normality of data.

Cited by: 6

Author: Wahab Ma, Shehta A, Hamed H, El Nake...

Publish Year: 2014

Natural history of untreatable hepatocellular carcinoma: A ...

<https://pubmed.ncbi.nlm.nih.gov/23060970>

Aim: To investigate the **clinical** course of untreatable **hepatocellular carcinoma** (HCC) identified at any **stage** and to identify factors associated with mortality. Methods: From January 1999 to December 2010, 320 out of 825 consecutive patients with a diagnosis of HCC and not appropriate for curative or palliative treatments were followed and managed with supportive therapy.

Cited by: 91

Author: Giuseppe Cabibbo, Marcello Maida, Chiar...

Publish Year: 2012

Radiomic Features at Contrast-enhanced CT Predict ...

<https://pubmed.ncbi.nlm.nih.gov/31934830>

Background Early **stage hepatocellular carcinoma** (HCC) is the ideal candidate for resection in patients with preserved liver function; however, **cancer** will recur in half of these patients and no reliable prognostic tool has been established. Purpose To investigate the effectiveness of radiomic featur ...

Cited by: 22

Author: Gu-Wei Ji, Fei-Peng Zhu, Qing Xu, Ke W...

Publish Year: 2020

Comparisons of recurrence-free survival and overall ...

<https://www.ncbi.nlm.nih.gov/pubmed/31918433>

Jan 09, 2020 · Comparisons of **recurrence-free survival** and overall survival between microwave versus radiofrequency ablation treatment for **hepatocellular carcinoma**: A multiple centers **retrospective cohort study** with propensity score matching. Du S(1), Yang JZ(2), Chen J(3), Zhou WG(2), Sun YY(1).

Cited by: 6

Author: Shibin Du, Jian-Zhi Yang, Jing Chen, Wei...

Publish Year: 2020