

[A new modified Gant-Miwa-Thiersch combined with submucosal ...](#)<https://pubmed.ncbi.nlm.nih.gov/34090388>

Background: **Full-thickness rectal prolapse (FTRP)** frequently occurs in **elderly women**, and more than 100 surgical **procedures** have been proposed to restore **FTRP**. The **Gant-Miwa-Thiersch (GMT) procedure** is the most used treatment in China. However, the recurrence rate of **FTRP post-GMT**, which is as high as 23.8%, is concerning.

[A new modified Gant-Miwa-Thiersch combined with submucosal ...](#)<https://europepmc.org/article/MED/34090388> ▾

Jun 05, 2021 · **Full-thickness rectal prolapse (FTRP)** frequently occurs in **elderly women**, and more than 100 surgical **procedures** have been proposed to restore **FTRP**. The **Gant-Miwa-Thiersch (GMT) procedure** is the most used treatment in China. However, the recurrence rate of **FTRP post-GMT**, which is as high as 23.8%, is concerning.

PEOPLE ALSO ASK

Which is the best surgical treatment for rectal prolapse? ▾

Why are rectal prolapses so common in men? ▾

What's the difference between incomplete and incomplete rectal prolapse? ▾

Modified Gant-Miwa-Thiersch procedure (mucosal plication ...<https://pubmed.ncbi.nlm.nih.gov/30673147>

Aim: Rectal prolapse (RP) is usually associated with elderly women and is well recognized as having a detrimental effect on quality of life. A number of surgical procedures for RP are available, but morbidity and mortality are substantial. The Gant-Miwa-Thiersch procedure ...

Cited by: 2

Author: Y. Iida, K. Honda, H. Saitou, Y. Munemot...

Publish Year: 2019

Modified Gant-Miwa-Thiersch procedure (mucosal plication ...<https://www.ncbi.nlm.nih.gov/pubmed/30673147>

AIM: Rectal prolapse (RP) is usually associated with elderly women and is well recognized as having a detrimental effect on quality of life. A number of surgical procedures for RP are available, but morbidity and mortality are substantial. The Gant-Miwa-Thiersch procedure ...

Cited by: 2

Author: Y. Iida, K. Honda, H. Saitou, Y. Munemot...

Publish Year: 2019

PEOPLE ALSO ASK

Which is a useful operation for complicated rectal prolapse? ▾

How does full thickness rectal prolapse affect the elderly? ▾

Can a perineal approach be used for pelvic surgery? ▾

[Feedback](#)**A new modified Gant-Miwa-Thiersch combined with ...**<https://pubmed.ncbi.nlm.nih.gov/34090388>

Background: Full-thickness rectal prolapse (FTRP) frequently occurs in elderly women, and more than 100 surgical procedures have been proposed to restore FTRP. The Gant-Miwa-Thiersch (GMT) procedure is the most used treatment in China. However, the recurrence rate of FTRP post-GMT, which is as high as 23.8%, is concerning.

Author: Jinxi Wang, Huiyu Li, Xiaoming Ma, G...

Publish Year: 2021

国内版

国际版

Modified Gant procedure for the treatment of internal rectal prolap



ALL

IMAGES

VIDEOS

21,300 Results

Any time ▾

A new modified Gant-Miwa-Thiersch combined with ...

<https://pubmed.ncbi.nlm.nih.gov/34090388>

Background: Full-thickness rectal prolapse (FTRP) frequently occurs in elderly women, and more than 100 surgical procedures have been proposed to restore FTRP. The Gant-Miwa-Thiersch (GMT) procedure is the most used treatment in China. However, the recurrence rate of FTRP post-GMT, which is as high as 23.8%, is concerning.

Author: Jinxi Wang, Huiyu Li, Xiaoming Ma, G... Publish Year: 2021

Modified Gant-Miwa-Thiersch procedure (mucosal plication ...

<https://www.ncbi.nlm.nih.gov/pubmed/30673147>

Aim: Rectal prolapse (RP) is usually associated with elderly women and is well recognized as having a detrimental effect on quality of life. A number of surgical procedures for RP are available, but morbidity and mortality are substantial. The Gant-Miwa-Thiersch procedure ...

Cited by: 2 Author: Y. Iida, K. Honda, H. Saitou, Y. Munemot...
Publish Year: 2019

PEOPLE ALSO ASK

Which is the best surgical treatment for rectal prolapse?



How did Hippocrates treat a rectal prolapse?



What does a complete rectal prolapse look like?



Can a rectal prolapse be caused by incontinence?



Feedback

Modified Gant-Miwa-Thiersch procedure (mucosal plication ...

<https://pubmed.ncbi.nlm.nih.gov/30673147>

Aim: Rectal prolapse (RP) is usually associated with elderly women and is well recognized as having a detrimental effect on quality of life. A number of surgical procedures for RP are available, but morbidity and mortality are substantial. The Gant-Miwa-Thiersch procedure ...

Cited by: 2 Author: Y. Iida, K. Honda, H. Saitou, Y. Munemot...