

Safety of upper endoscopy in patients with active cocaine use



ALL

IMAGES

VIDEOS

27,500,000 Results

Any time ▾

[PDF] Appropriate use of GI endoscopy - ASGE

<https://www.asge.org/docs/default-source/education/...>

E. GI bleeding or melena with a demonstrated **upper** GI source. Colonoscopy is generally contraindicated in: A. Fulminant colitis. B. Documented acute diverticulitis. FS FS is generally indicated for: A. Screening of asymptomatic, average-risk **patients** at risk of colonic neoplasia. Appropriate **use of GI endoscopy**

UpToDate

<https://www.uptodate.com/contents/approach-to-acute...>

Mar 09, 2021 · Limited data suggest that proceeding with **upper endoscopy in patients** with thrombocytopenia is generally **safe** , though whether there is a lower limit below which **endoscopy** should be delayed is unclear . Our approach is to perform an **upper endoscopy** if the platelet count is >20,000/microL, though if the **patient** is suspected to have **active** bleeding, we attempt to raise the platelet count to >50,000/microL prior to **endoscopy**.

PEOPLE ALSO ASK

Can a woman be sedated during an upper endoscopy? ▾

Is it safe to use benzodiazepines for upper endoscopy? ▾

What kind of drugs are used for an upper endoscopy? ▾

Do you need anesthesia for upper GI endoscopy? ▾

Match Overview

1	Crossref 17 words Tiffany S. Moon, Michael X. Gonzales, Joshua J. Sun, Agnes Kim et al. "Recent cocaine use and the incidence of hem..."	1%
2	Internet 13 words en.wikipedia.org	1%
3	Internet 12 words crawled on 21-Feb-2021 bmcpublichealth.biomedcentral.com	1%
4	Internet 12 words crawled on 30-Apr-2020 eventscribe.com	1%

Name of Journal: *World Journal of Gastrointestinal Endoscopy*

Manuscript NO: 68133

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Safety of upper endoscopy in patients with active cocaine use

Liyen Cartelle A *et al.* Upper endoscopy in active cocaine users

Anabel Liyen Cartelle, Alexander Nguyen, Parth M Desai, Vikram Kotwal, Jinal Makhija,

Jie Yu, John Erikson L Yap

Safety of upper endoscopy in patients with active cocaine use



ALL

IMAGES

VIDEOS

7,450,000 Results

Any time ▾

[PDF] [Guidelines for sedation and anesthesia in GI endoscopy](#)

https://www.asge.org/docs/default-source/education/practice_guidelines/piis...

Sedation is a **drug**-induced depression in the level of consciousness. The clinical objectives of administering sedation for GI **endoscopy** are to relieve **patient** anxiety and discomfort, improve the outcome of the examination, and diminish the **patient's** memory of the event. A number of different sedatives and analgesics can be used to achieve

EXPLORE FURTHER

[Practice Guidelines for Moderate Procedural Sedation and ...](#)

[pubs.asahq.org](#)

[Sedation for Endoscopy - American College of Gastroenterology](#)

[gi.org](#)

[Sedation and anesthesia in GI endoscopy](#)

[c8g3e5x4.rocketcdn.me](#)

[Sedation and anesthesia in GI endoscopy](#)

[www.sages.org](#)

[Colonoscopy Anesthesia: Types, Levels, and Possible Side ...](#)

[gastrocenternj.com](#)

Recommended to you based on what's popular • Feedback

[Indications and Contraindications for Upper ...](#)

<https://www.sciencedirect.com/science/article/pii/S1052515718304896>

Jul 01, 1994 · All reports, however, that have evaluated the yield of **upper endoscopy** in this population have noted positive findings in a significant number of **patients**, particularly with nonsteroidal anti-inflammatory **drug use** or **upper** gastrointestinal symptoms.^{8,14, 28} Importantly, two studies that were limited to **patients** with iron deficiency anemia⁸ ...

Search

Turn on