



WJG 20th Anniversary Special Issues (14): Pancreatic cancer

Novel therapeutic targets for pancreatic cancer

Shing-Chun Tang, Yang-Chao Chen

Shing-Chun Tang, Yang-Chao Chen, School of Biomedical Sciences, Faculty of Medicine, the Chinese University of Hong Kong, Hong Kong, China

Yang-Chao Chen, Shenzhen Research Institute, The Chinese University of Hong Kong, Hong Kong, China

Author contributions: Chen YC and Tang SC wrote the paper. Correspondence to: Yang-Chao Chen, Professor, School of Biomedical Sciences, Faculty of Medicine, The Chinese University of Hong Kong, Shatin, NT, Hong Kong, China. frankch@cuhk.edu.hk

Telephone: +852-39435728 Fax: +852-26035123

Received: October 28, 2013 Revised: February 13, 2014

Accepted: April 5, 2014

Published online: August 21, 2014

Abstract

Pancreatic cancer has become the fourth leading cause of cancer death in the last two decades. Only 3%-15% of patients diagnosed with pancreatic cancer had 5 year survival rate. Drug resistance, high metastasis, poor prognosis and tumour relapse contributed to the malignancies and difficulties in treating pancreatic cancer. The current standard chemotherapy for pancreatic cancer is gemcitabine, however its efficacy is far from satisfactory, one of the reasons is due to the complex tumour microenvironment which decreases effective drug delivery to target cancer cell. Studies of the molecular pathology of pancreatic cancer have revealed that activation of KRAS, overexpression of cyclooxygenase-2, inactivation of p16^{INK4A} and loss of p53 activities occurred in pancreatic cancer. Co-administration of gemcitabine and targeting the molecular pathological events happened in pancreatic cancer has brought an enhanced therapeutic effectiveness of gemcitabine. Therefore, studies looking for novel targets in hindering pancreatic tumour growth are emerging rapidly. In order to give a better understanding of the current findings and to seek the direction in future pancreatic cancer research; in this review we will focus on targets suppressing tumour metastasis and progression, KRAS

activated downstream effectors, the relationship of Notch signaling and Nodal/Activin signaling with pancreatic cancer cells, the current findings of non-coding RNAs in inhibiting pancreatic cancer cell proliferation, brief discussion in transcription remodeling by epigenetic modifiers (*e.g.*, HDAC, BMI1, EZH2) and the plausible therapeutic applications of cancer stem cell and hyaluronan in tumour environment.

© 2014 Baishideng Publishing Group Inc. All rights reserved.

Key words: Pancreatic cancer; CTHRC1; RAC1; RalGEF-RAI; Notch Signaling; Nodal/Activin Signaling; NDRG1; Hypoxic condition; DR5; PAR2; HER3; IAP; Non-coding RNA; HDAC; BMI1; EZH2; Pancreatic cancer stem cell; Tumour microenvironment

Core tip: Some of the targets discussed here have been discovered to enhance the effectiveness of gemcitabine upon co-administration of the corresponding agents, for instance, hyaluronidase can deplete hyaluronan in stromal region to enhance gemcitabine delivery. Besides, some signaling molecules, *e.g.*, RalGEF-RAI, Rac1, and PAR2 are being targeted to suppress metastasis. Tumour proliferation is limited upon DR5 activated apoptosis and others promising therapeutic areas like epigenetic modifiers; IAP, miR, lncRNA, and cancer stem cells-tumour microenvironment will also be discussed.

Tang SC, Chen YC. Novel therapeutic targets for pancreatic cancer. *World J Gastroenterol* 2014; 20(31): 10825-10844 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v20/i31/10825.htm> DOI: <http://dx.doi.org/10.3748/wjg.v20.i31.10825>

INTRODUCTION

Pancreatic cancer is the fourth leading cause of cancer death in the last two decades because of various obsta-

cles in its treatment^[1]. Late and poor prognosis are two of the causes of high fatality rate^[2]. Patients diagnosed having pancreatic cancer are usually at their very late stage and spreading of the highly metastatic pancreatic cancer cell into the lymphatic system and vicinal organs limited the choices of effective treatments^[3].

Gemcitabine is the current standard chemotherapy for pancreatic cancer^[4], however, due to the complex tumour microenvironment^[5] and high metastatic property of pancreatic cancer. The effectiveness of gemcitabine in treating pancreatic cancer is unsatisfactory. Studies of targeting the molecular pathology have been carried out to quest for more potential targets; for instance, activation of oncoprotein V-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog (KRAS)^[6], overexpression of cyclooxygenase-2^[1], inactivation of p16^{INK4A}^[6] and loss of p53 activities^[6] mark the onset of pancreatic cancer. Because, the treatments of targeting these molecules can enhance the efficacy of gemcitabine in pancreatic cancer^[3], these imply that combinatorial therapies may be the future direction in treating pancreatic cancer^[7]. Therefore, in the following context of this review we are going to briefly evaluate plausible therapeutic targets, in terms of the molecular and cellular level which covers the roles of several signal transducers, signaling pathways, surface proteins, receptor proteins, non-coding RNA, epigenetic modifiers and tumour microenvironment in driving pancreatic cancer and to explore any possibilities of combinatorial therapy among them.

SIGNAL TRANSDUCERS

CTHRC1

Collagen Triple Helix Repeat Containing-1 (CTHRC1) is a secretory protein^[8], which participates in vascular remodeling through limiting collagen matrix deposition^[9], and also morphogenesis but most importantly enhancing cell migratory ability and adhesiveness in tumour cells^[8]. CTHRC1 is found expressed in a wide spectrum of human cancer cells, and is in particular found highly expressed in pancreatic cells^[10]. The CTHRC1 protein is found highly expressed in invasive melanoma but weakly expressed or absent in benign nevi or non-invasive melanoma^[8]. Over expression of CTHRC1 in pancreatic cancer has enhanced the tumour cells migration and metastatic properties; studies of using induced hyper CTHRC1 expressed pancreatic cancer cell, MiaPaCa-2-CTHRC1 and shRNA-CTHRC1 suppressed pancreatic cancer cells, BxPC3 and Panc1, are used to evaluate CTHRC1 on pancreatic cancer cells metastatic in *in vivo* mice model^[10]. The result has revealed a wider metastatic spread of hyper CTHRC1 expressed pancreatic cancer cell to secondary organs while the hypo CTHRC1 expressed pancreatic tumour cells has reduced tumour cells spreading to neighboring organs when compared with the tumour cells transfected with control shRNA^[10]. The phosphorylation of Focal adhesion kinase (FAK)-steroid receptor coactivator (Src) cascade and extracel-

lular signal-regulated kinases (Erk) are the causes of the enhanced metastasis^[10], it is found that the binding of CTHRC1 onto the wntless-type MMTV integration site family protein, member 5A (Wnt5a) can stabilize the Wnt receptor complex^[11] and the facilitated binding of the Wnt5a into its Wnt receptor complex will activate paxillin which leads to phosphorylation of Src-FAK signaling cascade and Erk^[10], as both Src and Erk signaling pathways could lead to tumour progression and enhanced motility^[12], overexpression of CTHRC1 has increased the phosphorylation of Src and Erk, and vice versa^[10], these indicating the CTHRC1 plays a critical role in controlling pancreatic tumour cell adhesiveness and metastasis. Besides, activating the fore mentioned kinases, CTHRC1 is reported to repress the production of collagen I into the stromal environment of pancreatic cancer^[8], supporting of its role as a cancer metastasis enhancing gene.

As suppressing CTHRC1 can reduce the metastatic and motility of pancreatic cancer cell, future studies can investigate on the feasibility of combining CTHRC1 targeted therapy with current anti pancreatic cancer drugs. CTHRC1 appears as a promising target in sequestering pancreatic cancer from spreading to neighboring organs, however, whether it could sensitize the tumour cells to current anti-cancer treatments in pancreatic cancer is not yet published. CTHRC1 would be a more promising target if it is proved to sequester pancreatic cancer during chemotherapy, providing a higher chance in elimination of tumour cells in the patient.

RAC1

RAS-related C3 botulinum toxin substrate 1 (Rac1)^[13] is found to be an important factor in regulating pancreatic islet morphogenesis^[14], failure of cell spreading has been reported on gelatin-coated culture by blocking Rac1 in isolated islet cells^[14]. Apart from its vital role in directing organogenesis, Rac1 is one of the Rat sarcoma (Ras) effectors^[15] and is being overexpressed in pancreatic cancer^[16]. It has been found diminishing the formation of acinar-ductal metaplasia (ADM), pancreatic intraepithelial neoplasia (PanIN) and tumours when its expression is ablated in K-Ras^{G12D} induced pancreatic ductal adenocarcinoma (PDAC) mice model^[15]. In cancer biology, Rac1 is found to promote tumour migration and metastasis through lamellipodia production^[17]. Studies of targeting Rac1 may be beneficial in slowing down the spreading of pancreatic cancer cells.

Two guanine nucleotide exchange factors (GEFs) have been reported activating Rac1, dynamin 2 (Dyn2) has been reported regulating Rac1 in an undefined mechanism^[15]; Dyn2 is found associated with vav 1 guanine nucleotide exchange factor (Vav1) in coimmunoprecipitation, an onco-protein acts as a guanine nucleotide exchange factor (GEF) in Rac1 activation, and Vav1 is stabilized by the degradation of lysozyme and heat shock cognate 70 upon binding with Dyn2^[15]. Truncated form of Dyn2 has found unable to associate with Vav1 and

leading to reduced activation of Rac1 by 50%^[15]. However, cell lines deficit in Vav1 expression (*e.g.*, Panc1) would undermine this therapeutic direction^[15]. Another GEF, T lymphoma invasion and metastasis 1 (Tiam1), which is reported as an oncogene and associated with various cancers, Tiam1 directs Rac1 to enhance tumour proliferation and metastasis through the Wnt signaling pathway, however, suppressions of Tiam1 and Rac1 will lead to the activation of another oncoprotein, RhoA, which also promotes pancreatic tumour cells aggressiveness and metastasis^[17]. The tumour growth and long term survival are significantly suppressed and enhanced respectively, upon simultaneous inhibition of Rac1 and RhoA^[17]. These revealed the invasiveness and tumour migration of pancreatic tumour cells are under complex controls, balanced Rac1 and RhoA expression level is suggested to be one of those^[17], and the possibility of the participation of Dyn2 in between Rac1 and RhoA, as the effect of reduction of activated Rac1 in truncated Dyn2 experiment on RhoA is unknown.

However, when shifting to microRNA research, microRNA-124 (miR-124) is found able to suppress Rac1 mRNA and protein levels in pancreatic cancer cells, through the binding onto 3'-UTR of the Rac1 mRNA^[16]. Although the suppression of RhoA by miR-124 is yet to be determined; miR-143 is found able to suppress Rac1 and RhoA at the same time, producing a decreased tumour migration result in a pancreatic tumour cell xenograft model^[18].

From the recent findings, Rac1 is difficult to target and obtain therapeutic value, owing to switching on another oncoprotein RhoA, however, the discovery of miR-143 is exemplifying; microRNA could be the way out in tackling target that is similar to Rac1 which has an antagonist carries the similar tumour proliferative and metastasis function. It is worth to investigate on how miR-143 suppressing this "double fused" system in pancreatic tumour metastasis enhancement and its effect on long term survival.

RalGEF-Ral effector signaling network

Ras-like guanine nucleotide exchange factors (RalGEFs) and Ras-like (Ral) protein (which is also named as Ral small GTPase) have drawn increasing attention in cancers mediated by Ras, because RalGEFs are one of the direct effectors of activated Ras^[19] and the discoveries of the important roles of Ral proteins in tumorigenesis and metastasis^[20], however, the exact mechanism of the signaling network requires further studies to complete. There are more than four kinds of RalGEFs (*e.g.*, RalGDS, Rgl1, Rgl2, and Rgl3) and two homologues of Ral are found, Ral-A and Ral-B, in which they share same nucleotide sequence but differ in 82% of amino acid sequence^[19-21]. It is known that activated Ras will activate RalGEFs and in turn the activated RalGEF will convert the GDP bound Ral into GTP bound Ral, the activated Ral GTPase will then activate its downstream targets, for instance RalBP1, filamin, PLC δ 1, PLD1, *etc.*, bringing

out the corresponding biological responses^[19]. Although the general mechanism is elucidated nowadays, the exact RalGEFs activating RalA and RalB are remain unknown, so do the identity of the exact Ras proteins in activating a particular RalGEFs^[19].

There are two homologues of Ral small GTPase which are named RalA and RalB, their roles are distinct in tumorigenesis^[19], but are seemingly overlapped in metastasis and invasiveness^[21], ubiquitinated form of RalA and RalB have been found and it is in a non-degradative manner for selective localization modulation and functional regulations of Ral^[22]. Studies have shown that mutated RalA in a constitutively active state can cause transformation of human cells but not in the same mutant of RalB^[23], stable suppression of RalA in pancreatic cancer cells has brought a significant inhibition in the anchorage-independent growth^[19], and inhibition of RalA can delay the tumorigenesis K-Ras mutants PDAC in mouse model^[19], and the binding of RalA onto RalBP1 or Sec5 is found crucial in Ras - mediated transformation^[23]. Aurora A kinase (AAK) is a kind of RalA inhibitors which prevents RalA phosphorylation, in fact an AAK, MLN8237 has been entered phase III clinical trials, and such targeting is not effective in suppressing RalA signaling^[19].

On the other hand, suppression of RalB alone does not reduce tumorigenesis and transformation but bringing a more pronounced effect in metastatic tumour growth suppression when compared to RalA inhibited alone pancreatic cell lines. In addition, enhanced apoptosis in RalB suppressed cells in suspension state^[24]; these are suggesting RalB has a more significant role in the control of metastatic growth of cancer cells than RalA^[21]. However, when abrogating the expression of either RalA or RalB in pancreatic cell lines, reduced invasiveness is observed in some pancreatic cancer cells with RalA or RalB suppression but not in all kinds of pancreatic cancer cells, *e.g.*, reduced invasiveness is observed in RalA and RalB suppressed Capan-1 cell line, while in Panc-1 cell line suppressed RalA boosted the cancer cell invasiveness and RalB can bring a reduced invasiveness, and in T3M4 cell line RalB suppression cannot bring down the cancer cell invasiveness but RalA suppression can bring a reduced invasiveness^[21]. Thus, RalA and RalB may participate in the control of the invasiveness of pancreatic cancer cells, but there should be some other signaling pathways in control to this tumour malignancy, as the invasiveness reduction cannot be observed in all types of pancreatic cancer cell lines^[21]. In regard to the observations, RalB has been suggested in maintaining the viability of the cancer cells in the circulatory system and ensuring tumour cells invasiveness to other organs^[21].

Nevertheless, the localization of Ral proteins may also have their roles in the control of the cancer malignancies and is in relation to their ubiquitination and phosphorylation status, as de - ubiquitination of RalA in lipid raft microdomains is reported at the loss of cell-matrix interactions, and ubiquitination of RalA promotes lipid raft microdomains exposure on the cell membrane

when the cell got re-adhered^[22]. Since lipid raft microdomains served as the platform for various signaling pathways, when cancer cell is in detached state, its growth is inhibited due to the loss of related signaling cascade in the lipid raft microdomains^[24,25]. Therefore, the prevention of the re-exposure of the lipid raft microdomains onto the membrane can be a direction in the RalGEF - Ral signaling cascade for inhibiting the cancer metastasis by targeting the ubiquitination and activation of RalA.

All in all, the Ral proteins in the RalGEF - Ral signaling cascade play important roles in the control of cancer malignancies, targeting the RalGEF would seem to be efficient in shutting down the transduction of the signaling cascade, however, the question of the availability of inhibitors to RalGEF is concerning, as it is a Ras like signaling molecules, the design of an effective inhibitor to RalGEF may not be easy, and the effectors downstream of this signaling cascade should be closely investigated to aid the discovery of inhibitors that can block the signal transduction downstream of this pathway.

SIGNALING PATHWAYS

Notch signaling pathway

Notch signaling is found to be an important pathway in pancreas development, however the exact mechanism of how Notch regulates pancreatic development and the effectors it recruits are not fully understood^[26]. Notch signaling pathway has been reported to maintain a pool of pancreatic progenitor cells at the early stage of pancreatic development, and governs pancreatic ductal cell differentiation which found to be triggered by the intensity of the Notch activation^[26]. Implying Notch signaling mediates different effectors depends on cell type, and the stage of organogenesis.

In the pancreatic cancer, Notch signaling molecules are over-expressed^[26] and could produce oncogenic, anti-tumour, and drug resistance^[27] activities basing on the cellular context^[26]. In an ADM study using mouse PDAC model has shown that subject carrying mutant KRAS^[28], Notch is constitutively activated and up-regulated in the absence of EGFR^[28], while in wild type KRAS carrier, Notch activation requires EGFR activation to induce ADM^[29]. Implying the mutation of Ras could alter the activation pathway of Notch. Moreover, the anti-tumour activity of Notch signaling is brought out by Notch2 receptor deletion in mutant KRAS carrier^[29], which showed PanIN development is inhibited and subject survival is risen^[29]. For the same model, deletion of Notch1 resulted an opposite effect, PanIN development is accelerated and subject median survival is decreased^[28,30]. As these two Notch receptors are localized in different compartment of a pancreatic cell, and the exact location of them is not yet concluded^[26]. Thus, studying the distribution of Notch1 and Notch2 in pancreatic cell may help to understand their roles in pancreatic cancer and the effectors downstream of this signaling pathway.

As the functions of the Notch1 and Notch2 receptors

appeared to be distinct and the involvement of EGFR for activation, the roles of Notch receptors may act as the decision maker in deciding how the cell behave according to the external environment. Due to the complex environment during cancer development, figuring out the roles of Notch at each stage of the pancreatic cancer development will definitely help sorting out targets this signaling pathway that can compromise pancreatic cancer.

Nodal/Activin signaling pathway

Nodal and Activin are morphogens which are being secreted into extracellular region^[31,32] to mediate gene expression in target cell through phosphorylating the transcription factor mothers against decapentaplegic homolog 2, 3 and 4 (Smad2, Smad3, and Smad4)^[33], and the signal intensity is found to be able to determine the cell fate decision that the target cell would execute^[33]. Thus it is an important switch in deciding cell differentiation, self-renewal and pluripotency maintenance^[33], the decision of the cell fate control is found to be related to the signal intensity and signal gradient generated by this pathway^[33].

It is found that these two morphogens are over-expressed in pancreatic stem cells and pancreatic stellate cells, their expression levels are barely detectable in highly differentiated pancreatic cancer cell and normal pancreas or other developed tissues^[34]. Moreover, it has been suggested that a small population of cancer stem cell is encompassed in pancreatic carcinomas^[34], therefore, taking these two characteristics together this signaling pathway can be a specific therapeutic target for pancreatic cancer.

The common receptors of Nodal and Activin which are named Activin-like type I receptor 4 and 7 (Alk4 and Alk7, also written as Alk4/7), are being targeted by the inhibitor SB431542^[34]. Targeting Alk4/7 is to abrogate the signal transduction from Nodal/Activin receptors to the transcription factors Smad 2, Smad 3, and Smad 4; and preventing the downstream genes transcription which favor tumour malignancies expression^[34]. Under *in vivo* condition, pancreatic cancer cell L3.6pl pre-treated with co-administration of SB431542 and gemcitabine before implanting onto immunocompromised mice, have resulted a significant increase in apoptosis of cell carrying CD133⁺ surface marker, implying such regimen can deplete the population of cancer stem cell in pancreatic cancer, and prevented the tumorigenicity of the cancer cell in this xenograft model; while such observations cannot be obtained in either single treatment alone^[34].

However, such regimen is challenged by the abundant stroma in the xenograft model employing primary pancreatic cancer tissue, co-administration of SB431542 and gemcitabine cannot inhibit the tumour growth in such model, overcoming the sheltering effect of stroma to the pancreatic cancer cell is vital for efficient drug delivery to the tumour cell^[35]. The triple-administration of SB431542, gemcitabine and CUR199691 resulted in an enhanced depletion of cancer stem cell population, as CUR199691 is an inhibitor targets hedgehog signaling pathway and ultimately deplets the stroma^[34].

Besides, pancreatic tumour cells with certain mutations on Smad 4 gene have showed to be less responsive towards the regimen^[34]. As Smad 4 is one of the factors for the signal transduction in the Notch/Activin signaling cascade^[34], thus it is essential for the future studies to identify others up-stream targets controlling mutated Smad 4, so as to provide regimen for pancreatic cancer patients with mutations in Smad 4.

Nodal/Activin signaling is a promising target in elimination of pancreatic cancer stem cell from the studies presented here, despite its limitation in pancreatic cancer patients with mutations in Smad 4 gene, its effectiveness in wild type Smad 4 still makes it an attractive target in primary pancreatic cancer tissue model with the use of hedgehog inhibitor.

Metastatic suppressor-N-myc downstream-regulated gene-1

The N-myc downstream-regulated gene-1 (NDRG1) has recently been identified as a metastasis suppressor in several human cancer types^[36], including human pancreatic cancer^[37]. NDRG1 is found to increase the expression of tumour suppressor gene Smad4, which further inhibits the phosphatidylinositol-3 kinase (PI3K)/phosphorylated protein kinase B (AKT) signaling and extracellular signal-regulated kinase (ERK) pathway^[36], besides, NDRG1 inhibits broad signaling molecules in nuclear factor - kappa B (NF- κ B) signaling pathway, which resulted in reduced cancer metastasis^[37]. As these three signaling pathways contribute to cancer cell proliferation and metastasis promotion, and cross-talk activities among them^[36], therefore, NDRG1 is playing a role of modulator in orchestrating the signals in this triad networks.

The regulation of NDRG1 is debatable; numerous of studies have found out that hypoxia condition^[39], epigenetic regulation^[39] and iron depletion^[40] can up-regulate NDRG1 expression level and such up-regulation seems to correlate with the differentiation status of the cancer cell.

It is worth to note that in a human PDAC model, under 2% of oxygen supply the NDRG1 mRNA and protein levels are elevated in differentiated pancreatic cancer cells but there are no change in the mRNA and protein levels in poorly differentiated cell lines^[38]. Suggesting NDRG1 expression depends on both hypoxic condition and differentiation status of the tumour cell. The main focus would be on the rationale behind this phenomenon, as undifferentiated pancreatic cancer cell is comparatively more invasive and metastatic than highly differentiated counterpart^[38]. In light of this, poorly differentiated pancreatic cancer cell (*e.g.*, Panc1) would have its NDRG1 level being suppressed in order to maintain high CXCL chemokines^[37] and high pro-angiogenic factor vascular endothelial growth factor (VEGF) expression^[38] to direct cancer cell proliferation and angiogenesis. While NDRG1 over-expression has found down-regulating of these two signaling molecules and leading to suppression of tumour growth and angiogenesis^[37].

The low NDRG1 expression in undifferentiated pancreatic cancer cells is related to the epigenetic regulation, as treating the undifferentiated pancreatic cancer cells with methyltransferase inhibitor 5-aza-2'-deoxycytidine has enhanced NDRG1 protein expression level, however, the epigenetic control on NDRG1 is not directly acting on the NDRG1 promoter, as there is no significant DNA methylation in the NDRG1 promoter region; suggesting other genes being silenced are essential for the NDRG1 expression^[39].

As NDRG1 expression is affected by numerous factors, studies of targeting the molecular events downstream of NDRG1 are carried out, for instance an novel synthetic derivative of curcumin (CDF) has shown its inhibitory effect of the expression of VEGF, hypoxia inducible factor-1 α (HIF-1 α), miR-210 and cancer stem cell self-renewal properties under hypoxia condition and are crucial for pancreatic cancer cell to promote tumour angiogenesis^[41].

Although the exact mechanism of controlling the NDRG1 remains unclear, the current findings have suggested maintaining a high NDRG1 expression level in undifferentiated cell is able to suppress the tumour malignancies. Therefore, studies in finding enhancing NDRG1 expression genes is important in suppressing pancreatic cancer growth and metastasis.

Energy metabolism

As mentioned in the previous sections, the low vascularity structure of pancreatic tumour leading to a hypoxic environment and the adaptation of pancreatic cancer cells in hypoxic conditions through enhanced proliferation, angiogenesis and metastasis have been described. However, the primary element for cell survival is energy source which normally generated in glycolysis and Krebs Cycle, as pancreatic cancer cells have an oxygen scarcity issue^[42]; metabolic changes in pancreatic cancer cells allow them to cope with hypoxia.

First, the utilization of glucose would rely heavily on TCA-independent pathways, for example, there is up-regulation of pentose-phosphate pathway, anaerobic respiration for ATP production in hypoxic environment^[43]. Secondly, glutamine metabolism is also elevated in hexosamine biosynthetic pathway which is crucial for the production of UDP-N-acetylglucosamine, and it is used for glycosylating proteins in proteins modification^[44]. Glutaminolysis is also employed by hypoxia pancreatic cancer cell which metabolizing glutamine to generate glutamate and can be further metabolized in TCA cycle to produce pyruvate and lactate for further ATP production^[44]. Lactate production is important for tumour cell invasiveness and neighboring cell proliferation, as inhibition of the enzyme glutamine fructose-6-phosphate amidotransferase by azaserine can cause significant reduction in hypoxic pancreatic cell proliferation^[42]. Apart from targeting glutaminolysis, cannabinoids are found to suppress TCA cycle and induce the reactive oxygen species which leads to AMP-activated protein kinase

level increase to mediate autophagy in pancreatic cancer cells^[45-47]. The ROS signaling activation could also abrogate the electron transport chain in mitochondria with unclear mechanism^[48], leading to depletion of ATP in the cell and the AMPK dependent autophagy would be mediated^[45].

By considering the founding in the energy metabolism of PDAC, targeting glutamine, glucose metabolism and increase the ROS production in hypoxic region in pancreatic tumour can elicit autophagy in PDAC. It is important to evaluate the effects of targeting them in *in vivo* model, as blocking major metabolic pathways is very likely to damage normal tissues, specific targeting the metabolic pathways in PDAC would minimize such drawback and enhancing the therapeutic value of targeting the energy metabolism pathway.

RECEPTOR PROTEINS

DR5

The death receptor 5 (DR5), is found to be frequently expressed in pancreatic cancer stem cell^[49] and mediates cancer cells apoptosis *via* caspase 8 recruitment upon interacting with another receptor, Tumour necrosis factor-related apoptosis-inducing ligand (Apo2L/TRAIL)^[50,51], forming the death-inducing signal complex (DISC) to induce apoptosis, thus this enables the elimination of pancreatic cancer stem cell specifically, and reduces the occurrence of tumour relapse and overcoming the chemoresistance of pancreatic cancer stem cell^[52].

Several studies on targeting activation of Apo2L/TRAIL induced apoptosis have been carried out, by combing with chemotherapies to obtain a synergistic effect in shrinking cancer stem cell population in pancreatic cancer^[53]. Co-administration of DR5 agonist Tigatuzumab and gemcitabine has recorded more tumour regression on PDA xenografts than administrating either agent alone^[49]. Moreover the up-regulation of several signaling molecules, for instance, cell surface death receptor Fas, Fas-associated death domain, and tumour necrosis factor receptor 1-associated death domain in the apoptotic pathway are also recorded^[49]. Indicating the co-administration of Tigatuzumab and gemcitabine can result in cell growth inhibition and apoptosis for cell expressing DR5^[49].

Another study has showed that dihydroartemisinin can increase intracellular ROS concentration and would lead to DR5 expression elevation and in turn mediate apoptosis *via* Apo2L/topoisomerase, TNF- α -related apoptosis-inducing ligand (TRAIL)^[52]. Revealing the apoptotic pathway activation through DR5 requires high intracellular ROS^[52]. Therefore, eliciting apoptosis in DR5 over-expressed cancer cell is a promising therapy for pancreatic cancer^[50].

PAR2

The Protease-activated Receptor-2 (PAR2) is a member of the G-protein coupled receptor family and is acti-

vated by trypsin^[54]. PAR2 is able to promote angiogenesis through two distinct pathways. The first one is *via* the activation of the mitogen-activated protein kinase (MAPK) to mediate VEGF release^[55], another pathway involves the tissue factor to bind with integrin-linked kinase to up-regulate HIF-1 α expression *via* AKT phosphorylation and eventually enhanced VEGF expression^[56]. Hence, PAR2 is essential for tumour survival under hypoxic condition in the micro environment, as PAR2 maintains a constitutive high level of HIF-1 α for angiogenesis promotion and this also explains the high metastatic property of pancreatic cancer cell in hypoxia region.

Besides, the role of PAR2 in pancreatic cancer cell migration is also being reported, PAR2 is found to mediate MAPK-epidermal growth factor receptor 1/2 (EGF1/2) signaling pathway with the utilization of extracellular ATP, blocking the cross talk between PAR2 and extracellular ATP can be a target in reducing pancreatic cancer metastasis^[57].

HER3

The Human Epidermal Growth Factor Receptor (HER) family consists of four members in which they are all type 1 transmembrane receptor with tyrosine kinase properties^[58], except HER3^[59], a member of HER which is found overexpressed 41% in pancreatic cancer^[60]. Because of lacking tyrosine kinase activity in HER3, it requires phosphorylation by another HER receptor to activate PI3K/AKT signaling pathway to mediate cell angiogenesis and metastasis^[61]. The expression level of HER3 has been correlated with tumour progression^[59].

Therefore, HER3 has been an important target for suppressing tumour angiogenesis and metastasis by using humanized monoclonal antibodies, *e.g.* U3-1287 which has gone through phase 1 clinical trials with well tolerance in solid tumour patients, MM-121 and tyrosine kinase inhibitors^[62,63]. The anti-HER3 agents block the activation site on the HER3 receptor, preventing the activation of HER3 during heterodimerization with HER2 receptor^[59] and promoting receptor internalization upon binding onto its extracellular domain^[58]. This reduces the activation of PI3K/AKT signaling pathway and its downstream effectors activation, resulting tumour growth suppression^[59].

In view of this, because of HER3 over-expression in PDAC and its crucial role in activating the signaling pathway essential for cell growth, it is a valuable and specific therapeutic target upon co-administration of anti-HER3 agent and gemcitabine has resulted an enhanced anti-tumour effect^[64], confirming the therapeutic value of anti-HER3 agents in PDAC and is worth investing in more clinical studies.

All in all, we have described three receptor proteins which carry out apoptosis, tumour proliferation and metastasis. Current studies are focusing on how to trigger the signaling molecule that could induce apoptosis, inhibit the receptors that favor cancer proliferation and metastasis, so as to reduce tumour progression. How-

ever, the possibility of combining these two approaches in the same model is not yet published, in which the total effect on tumour clearance is expecting to be more efficient.

CELL SURFACE PROTEIN

E-Cadherin

E-Cadherin is a transmembrane protein^[65] and is a member of cadherins family in which its expression in epithelial cells is controlled by intracellular signaling molecules^[66]. E-Cadherin directs the positioning of the cell during morphogenesis, controlling cell migration and tissue structure maintenance^[67]. It is reported that during epithelial-mesenchymal transition (EMT) the E-Cadherin level in neoplastic epithelial cells is down-regulated, suggesting triggering the dedifferentiation of neoplastic epithelial cells into mesenchymal cell with higher motility^[65].

The fading E-cadherin expression is frequently reported in undifferentiated, noncohesive pancreatic cancers^[68], and it is found that the silencing of E-cadherin is mediated by Snail/ histone deacetylase 1 (HDAC1)/histone deacetylase 2 (HDAC2) complex^[65] and Enhancer of Zeste Homolog 2 (EZH2)^[69] through hypermethylation of its promoter region^[68]. Inhibition of Snail and HDAC2 are also carried out to validate the E-cadherin expression is under such complex governance^[70].

Since the absence of E-Cadherin marks the onset of metastasis and PDAC progression, a study targeting E - Cadherin restoration by using microRNA 101, has inhibited the EZH2 binding on E-Cadherin promoter region in PANC1 preventing E - Cadherin silencing and in turn inhibited its tumorigenicity xenograft^[69].

The key for targeting E-Cadherin to obtain therapeutic value in PDAC is to up - hold the E - Cadherin expression by down - regulating its inhibitor, as described above, inhibiting EZH2, and Snail/HDAC1/HDAC2 can reduce E-Cadherin depression and suppresses the tumorigenicity of pancreatic cancer, in the future studies, discovering targets that suppress E-Cadherin expression is important for therapy involving E - Cadherin.

Galectin-4

Galectin-4 (Gal-4) is a glycan binding proteins which belongs to the galectin family. Gal-4 is found over-expressed in cystic tumours of the human pancreas, PDAC and cancer stromal cell^[71]. Activated galectins carry out several functions; include cell-cell adhesion, cell proliferation^[72], mediation of intracellular signaling^[73] and tumour metastasis^[74], *etc.*, However the mechanisms behind are remain unknown.

A study has evaluated the inhibition effect of Gal-4 in a pancreatic cancer cell, PaTu-S cell, can lead to enhanced tumour migration^[74]. The exact reason is yet to be elucidated, but it is speculated that the reduction of Gal-4 on the cell membrane would destabilize cell-cell interaction, allowing tumour cells escape^[75]. Another important implication suggests Gal-4 expression may be

dependent on the tumour development stage, and is vital for tumorigenesis as it promotes cell-cell adhesion^[76].

Because of Gal-4 multi-roles in expressing tumour malignancies, and the little knowledge on how Gal-4 control cell migration and tumour metastasis, it is worth to investigate its related signaling pathways and identifying possible inhibitors so as to enable targeting Gal-4 in treating pancreatic cancer.

TMPRSS4

Transmembrane Protease, Serine 4 (TMPRSS4) is found highly expressed in several cancer cells, including pancreatic cancer cell^[77]. However its regulation mechanism is poorly known^[78], several studies have shown that TMPRSS4 can promote EMT, metastasis and invasiveness in human epithelial pancreatic^[79], lung and colon cancer^[77].

It is reported that EMT mediation is not solely rely on TMPRSS4 up-regulated integrin $\alpha 5$ to activate FAK/ERK signaling pathway and enhanced invasiveness^[79] but also count on the down-regulation of E-Cadherin in TMPRSS4 up-regulated cancer cells^[79].

Another downstream target of TMPRSS4 is the urokinase-type plasminogen activator (uPA) gene^[80]. TMPRSS4 would activate the transcription factors of μ PA *via* c-Jun N-terminal kinase mechanism before promoting μ PA transcription in a cell-type dependent manner^[80]. And the increased μ PA gene transcription marks the increased tumour cell aggressiveness.

The coupling effects of TMPRSS4 on tumour aggravated invasiveness and metastasis with other signaling molecules (*e.g.*, integrin $\alpha 5$, uPA), moreover, the inverse expression pattern of TMPRSS4 and E-cadherin suggests TMPRSS4 can be suppressed by targeting E-cadherin inhibitors as previously mentioned and should be investigated in future studies. TMPRSS4 expression is affected by various signaling molecules and by considering its important role in expressing tumours malignancies, it is a target with multiple approaches for suppression.

IAP

Inhibitor of apoptosis protein (IAP) is a group of proteins bind to caspases and inhibit caspases apoptotic effect resulting apoptosis abortion^[81]. The importance of apoptosis mediation in cancer therapies has an irreplaceable place, and therapies incapable to induce cell death would be meaningless. However, most therapies nowadays involve the elicitation of apoptosis at their end, and resistance of the corresponding therapies developed due to the presence of IAP^[82]. Therefore IAP is the obstacle to tackle with, so as to ameliorate the effectiveness of therapies targeting apoptosis induction.

Two of the IAP members would be discussed here which are X-linked IAP (XIAP) and survivin, because of the reports of their close interaction in triggering anti-apoptotic effect^[82].

Survivin's action has been controversial in anti-apoptotic activity^[82], it is reported that survivin carries out neurogenesis, angiogenesis, cell cycle progression in can-

cer cell^[83] and displays caspase inhibitory effect through associating with XIAP and stabilizes XIAP *via* their baculovirus-inhibitor of apoptosis repeat (BIR) domain^[84]. Most of the survivin inhibitors that are under clinical trials have improved the effectiveness of chemotherapies *e.g.*, topoisomerase, TRAIL^[82]. Revealing survivin's submissive and supporting role in IAP targeted treatment.

XIAP is the most studied IAP, it is found able to suppress caspase-3, caspase-7 and caspase-9 apoptotic activities^[85]. Inhibition of its caspase binding domains, which are named, BIR-2 and BIR-3, with the use of phenylurea-based chemical inhibitors of XIAP (XAntags) could make pancreatic cancer cells more vulnerable to apoptosis^[85].

Because of the anti-tumour effect in inhibiting XIAP and the supportive role of surviving in apoptotic inhibition, co - suppressing XIAP and survivin has also been performed in Panc-1 cell^[86], resulting cell proliferation hindrance, and enhanced gemcitabine effectiveness in XIAP and survivin suppressed model than sole suppression of either IAP^[86].

Nevertheless, there is no cell toxicity recorded in XIAP knocked out mouse model and *in vitro* cell model, possibly by the compensatory up-regulation of other cIAPs, and the masking effect of such up - regulation requires further studies^[81]. From the recent findings in IAP, inhibiting IAP is a promising therapeutic direction in promoting apoptosis progression in PDAC cells, thus enhancing the effectiveness of current chemotherapies upon co - administration in treating PDAC.

NON-CODING RNA

MicroRNA

MicroRNAs (miRNAs) consist of 18-24 base pair which are small and non-coding-sequence^[87]. They execute target gene expression control by binding miRNA 3'UTR on to the target gene mRNA^[87], and only when perfect binding of miRNA on to the target mRNA could mediate mRNA cleavage, otherwise, it would result into inhibited protein production^[88]. miRNAs which induce over-expression of oncogenes are termed the oncogenic miRs (onco-miRs), on the other hand, miRNAs which suppress cell transformation are named tumour suppressor miRs (TSG-miRs)^[89]. The abnormal expression levels of these two kinds of miRNAs are observed in pancreatic cancer^[90]. Current studies are either suppressing onco - miRs or reconstituting the TSG-miRs level^[91], therefore, in the following we will discuss some TSG-miRs which are promising therapeutic targets in pancreatic cancer.

miR-34: miR-34 is reported to be up-regulated by p53^[92], inducing cell cycle arrest in primary and tumour derived cell lines^[93]. A significant reduction of miR - 34 expression level in gastric cancer cells with p53 mutation has been observed and reconstituted miR - 34 expression by transfecting pancreatic cancer cells with letivirus carrying vector expressing miR - 34^[94], and resulted in decreased

Notch2 and Bcl-2 protein production, reduced tumour-sphere formation from cancer stem cell (CSC)^[94]. Although the relationship between miR-34 and p53 is still unclear^[94], the encouraging results generated by miR-4 in p53 deficient pancreatic cancer cells^[93] have make it a worthy therapeutic target.

miR-143: miR-143 has been studied for its anti-metastasis and anti-tumour proliferation in liver undergone metastasis and a pancreatic cancer xenograft in mouse model, respectively^[95]. miR-143 expression level in KRAS mutant pancreatic cancer cells is also being ablated^[96], re-expressing miR-143 in its deficit cell lines has performed, GEF, RAC1, matrix metalloproteases (MMPs) and KRAS are the inhibition targets for miR-143^[95], as described previously lessened RAC1 level can inhibit metastasis and tumorigenesis, while inhibiting KRAS is even more important, which implies a board spectrum of signaling pathways diminishing effect.

Another tumour growth promotion factor that miR-143 targets is the cyclooxygenase (COX-2)^[97], COX-2 is reported as an essential factor for prostaglandin synthesis to mediate inflammation and cancer cell growth and survival^[98]. In pancreatic cancer cell, miR - 143 was found to be repressed by prostaglandin^[99], and restoration of miR-143 level can decrease both mRNA and protein level of COX-2 and inhibited cell growth^[98].

miR-200: miR-200 is a family of miRNAs related to EMT^[100], reconstituted expression level of miR-200 has restored the phosphatase and tensin homolog (PTEN) expression level^[100], as PTEN is widely down regulated in various cancer cell lines and is a tumour suppressor gene in which reduced expression would lead to enhanced tumour aggressiveness^[101,102], while membrane type-1-matrix metalloproteinase (MT1-MMP) is up-regulated and lead to aggravated cancer invasion^[101-103]. Restoration of miR-200 by using CDF, which is a synthetic analog of curcumin, and a natural compound, BR - DIM are reported and are found able to enhance PTEN expression level and a decreased MT1-MPP promoted invasiveness^[100]. Therefore, agents which could enhance miR-200 expression would have promising therapeutic value in curbing pancreatic cancer aggressiveness for enhanced treatments efficiency.

The three TSG-miRs exemplified the diverse roles of miRNAs in anti-tumour activities, up-regulation of TSG-miRs can suppress tumour malignancies expression, however, suppressing onco-miRs that can up-regulate oncogenes also have tumour malignancies suppression effect, therefore screening and studying the agents that can up-regulate TSG-miRs and down-regulate onco-miRs are vital for PDAC therapy development.

Long non-coding RNA

Long non-coding RNAs (lncRNAs) are transcribed from intergenic and intronic regions in human genome^[104] by RNA polymerase II^[105], which lengths more than 200 bp^[106] and their biological functions have been reported,

for instance, epigenetic control, transcription regulation, pre and post-translational regulation^[107], cell cycle and differentiation control and even governing the apoptosis process^[108]. lncRNA is used as a diagnostic parameter and can be a therapeutic target in cancers^[104]. However, the definition and discovery of lncRNAs are expected to keep on changing as very little is known in this emerging area^[109]. In the following, two lncRNAs that are highly expressed in pancreatic cancer will be discussed.

HOTAIR: HOX transcript antisense RNA (HOTAIR) is a lncRNA which is highly expressed in a range of primary tumours and metastatic cell^[110], in which its expression pattern is variable but in general is also over-expressed pancreatic cancer^[111]. HOTAIR carries out tumour supporting effect by inhibiting anti - tumour genes activity, in which the interaction of HOTAIR and a Polycomb-group (PcG) family protein named, EZH2, would promote chromosomal histone protein histone 3 protein at lysine 27 (H3K27) trimethylation, which leads to repressed transcription of multiple gene targets^[112]. However, there are some genes inhibited in an EZH2-independent manner^[113].

Suppressed HOTAIR expression by using RNAi in pancreatic cancer cell has caused retarded cell growth, diminished tumour aggressiveness; altered cell-cycle progression and apoptosis induction^[114]. Thus, relieving the repressed transcription of the tumour suppressor genes by suppressing HOTAIR expression is therapeutically valuable in treating PDAC. As the studies of genes activation mechanism and the genes that are targeted by HOTAIR are still ongoing^[114], and the mechanism of genes being independently regulated by EZH2 but dependent on HOTAIR only, are currently under studies.

Studies of targeting HOTAIR in PDAC cell lines have achieved reduced tumour malignancies expression, indicating the relieved tumour suppressor genes expression by targeting HOTAIR has made HOTAIR an attractive target in pancreatic cancer therapies development. However, cautions should be taken on the over-expressed genes induced by HOTAIR, as HOTAIR induced and suppressed multiple genes at the same time and some of the over-expressed genes expression level do not reduce with HOTAIR suppression^[114], suggesting another mechanism may exist in down-regulating them. In conclusion, a more thorough understanding on the regulation and the functions of HOTAIR induced and suppressed genes, it could lead to a more rounded target in promoting tumour suppressors genes functions while inhibiting oncogenes activities.

MALAT1: Metastasis-Associated Lung Adenocarcinoma Transcript 1 (MALAT1), as known as the Nuclear-Enriched Abundant Transcript 2^[115] is found highly expressed in normal pancreatic and lung tissues with high abundance and highly conserved among mammalian^[116]. Intensive studies of MALAT1 in non-small-cell lung carcinoma revealed its metastasis and tumorigenicity promotion activities^[117]. Although inadequate studies of MALAT1 in

pancreatic cancer cell model, it has been reported for its promotion of tumour malignancies expression in various cancer types^[116]. For instance, in colorectal cancer, a Chinese herb extract Resveratrol is shown to down-regulate MALAT1 and cause suppression of Wnt/ β signaling *via* decreasing β -catenin nuclear localization and eventually inhibited the invasiveness and metastasis of colorectal cancer^[118].

Apart from promoting tumourigenesis, invasiveness and metastasis, MALAT1 also participates in the control of cell cycle progression, oncogenic transcription factor B-MYB is found up-regulated with the expression of MALAT1, silencing of B-MYB in fibroblast model resulted into cell cycle arrest in G1/S and S phase^[119], moreover, another transcription factor, E2F transcription factor 1 (E2F1), which is essential for cell cycle progression and is also under the modulation of MALAT1, however down - regulated MALAT1 brought down E2F1 expression has elicited p53 expression enhancement and lead to cell cycle arrest and hence reduced cell proliferation^[119]. This implies MALAT1 could induce DNA damage response *via* an unknown mechanism^[119].

With regard to the findings of MALAT1 in other cancer, MALAT1 expression in PDAC is also very likely to correlate to pancreatic cancer progression. Although MALAT1 expression level is high in normal pancreatic tissue, its expression level in pancreatic cancer is not yet reported and also the role of MALAT1 in pancreatic tumour activities. Thus, if MALAT1 has a similar tumour malignancies promotion role in pancreatic cancer as in other cancer types, it would be a promising therapeutic target for PDAC treatment development.

EPIGENETIC MODIFIER

HDAC

Histone Deacetylases (HDACs) are a group of four classes of deacetylases^[120], each class of the enzyme contributes to different tumour malignancies expression, for example, as mentioned previously, HDAC1 is responsible for the acceleration of EMT and metastasis in PDAC^[121], while HDAC2 would desensitize the PDAC towards DNA damage response and decreased prop - apoptotic proteins^[122], however, only the third class, which is named the human hist proteins (SIRT's) did not respond to HDAC inhibitors (HDACIs) under current clinical trials^[120], but a HDACI named Sirtinol is able to induce apoptosis with the administration of Sirtinol^[121], and its effect is further enhanced with the co - administration with gemcitabine^[123].

The exact mechanism of HDACIs in mediating anti-tumour activities remain further elucidation. However, studies have shown that it is not necessary for HDACIs to inhibit the expression of HDACs in mediating tumour suppression, for instance, a class I and II HDACI did not cause changes in the expression level of HDAC1, and other tumour suppressor genes but has shown reduced cell proliferation in cervical tumour cell^[124]. Moreover, a class I and II HDACI inhibitor is found able to cause

a changes in the expression profile of class III HDAC, SIRT6^[121]. These evidences suggest the working mechanism of HDACs involve complex molecular control. On the other hand, Rel/p65 (NF- κ B) is found relating to the expression level of HDAC, for instance, over-expression of class I HDAC in pancreatic cancer cells, their expression level of NF- κ B is also high, besides, a class I HDACI valproic acid (VPA) is studied and found to cause a decreased expression in pancreatic cancer cells which leads to enhanced PDAC apoptosis^[125], in which over-expression of NF- κ B has been reported for enhanced tumour growth, angiogenesis, chemo-resistance and metastasis^[125]. Blocking NF- κ B activity by VPA can obtain anti-tumour effect in such case^[126].

The action of HDAC on gene silencing is mediated by deacetylating the histone proteins in the chromatin leading to chromatin condensation, resulting silenced genes transcription^[121]. As the genes being silenced in cancer are mostly related to tumour suppressors, anti - apoptosis, and often resulted in drug resistance, therefore, targeting HDAC by using HDACI is believed to reduce the these tumour malignancies expression by suppressing the related signaling pathways of the PDAC and synergistically enhance the anti - tumour effect of current chemotherapy.

BMI1: B-Cell-specific Moloney murine leukemia virus Insertion site 1 (BMI1), belongs to the polycomb group (PcG) which represses transcriptional activity of various genes^[127]. Over-expression of BMI1 in a board spectrum of cancer cells is observed, it strengthens tumour growth by providing anti-apoptotic activities and participate in tumour metastasis by up-regulating PI3K/AKT signaling pathway^[127]. In *in vitro* experiment, PDAC with BMI1 suppressed using shRNA has shown enhanced cell death in response to gemcitabine treatment, a significant decrease for its cell surface markers CD44⁺CD24⁺ESA⁺, loss of self-renewal ability, reduced tumour sphere formation by CSCs and reduced tumour size in xenograft model^[127].

Because of the diversified anti-tumour effects of silencing BMI1 in pancreatic cancer cell, such as reduced invasiveness, tumourigenesis^[127], metastasis, CSC phenotypes, cell proliferation^[128] and also chemo-resistance^[127]. Besides, CSC is reported to be the causes for tumour relapse in pancreatic cancer^[129], thus, the diversified roles of BMI1 in pancreatic cancer have made it a very attractive target, future studies targeting BMI1 inhibition and its downstream effectors would benefit PDAC treatment development.

EZH2

The polycomb repressor complex 2 member, Enhancer of Zeste Homolog 2 (EZH2) is a histone methyltransferase which is highly expressed in pancreatic cancer cells^[130], EZH2 mediates tumour suppressor genes transcription inhibition through trimethylation of the H3K27^[131], such as suppressing Rap1GAP expression in squamous car-

cinoma^[132], E-Cadherin in pancreatic cancer^[131]. Besides, several reports have suggested EZH2 suppresses miRNAs in contributing to pancreatic cancer progression, *e.g.*, microRNA-218 (miR-218), microRNA-26a^[133,134], miR-218 is essential in suppressing tumour proliferation and metastasis in nude mouse model^[133], EZH2 is believed to interact with two polycomb repressive complexes (PRCs), PRC1 and PRC2, and promoting the methylation of the target miRNA promoter region to silence the miRNAs expression in pancreatic cancer^[133]. It is found that with the administration of EZH2 inhibitor, such as 3-deazaneplanocin A (DZNep), can reduce EZH2 expression of EZH2 and rescued the expression of miR-218 leading to tumour malignancies reduction^[133,135].

Apart from suppressing miRNAs in tumour progression, studies of the role of EZH2 as a tumourigenesis initiator have found that EZH2 also suppresses tumour suppressor gene p16^{INK4}, in which it suppresses tumour proliferation and regeneration, enhanced EZH2 expression has caused p16^{INK4} down - regulation, counteracting the suppression effects exerted by p16^{INK4}^[136]. Such control is crucial for the regeneration of the injured acinar pancreatic cell, in which the injured cell undergone de-differentiation into metaplastic epithelial intermediate, depleted p16^{INK4} allows the cell to re-differentiate into acinar cell from metaplastic epithelial intermediate^[136]. Thus in combination with the early appearance of PaIN lesion in pancreatic cell baring KRAS mutation and the loss of p16^{INK4} expression due to enhanced EZH2, invasive and metastatic tumour development are accelerated, demonstrating the linkage between regeneration and tumourigenesis under the influence of mutant KRAS^[137].

Further studies of the role of EZH2 in pancreatic CSC has found it is essential in maintaining the CSC population in pancreatic cancer, suppressing EZH2 has decreased the degree of H3K27 methylation, reduced CSC population in pancreatic cancer, enhanced genes expressions for cell differentiation and migration^[130]. Since the trimethylation of H3K27 and the expression is correlated with the CSC population, it is suggested that the H3K27 trimethylation by EZH2 can be used as a marker for the CSC population which allows rapid evaluation for the population of CSC when compared to conventional methods, hence, speeding up the studies of the effectiveness of compounds towards pancreatic CSC^[130].

From the current findings of suppressing EZH2 in pancreatic cancer, EZH2 has an important role in tumour development initiation and supporting cancer stemness, and co-administration of DZNep and gemcitabine has achieved promising anti-tumour effects. Nevertheless, EZH2 has also demonstrated its possibility to act as an indicator for CSC population estimation, and CSC elimination is an important factor for researchers to evaluate the efficacy of the compounds under studies, thus EZH2 is a versatile targets that possess both therapeutic and assay values and screening compounds suppressing EZH2 would definitely help speeding up therapies development

in PDAC.

PANCREATIC CANCER STEM CELL (PANCREATIC CSC)

The tumour cell population is reported to encompass a population of cancer stem cell (CSC)^[138], and it is reported to give rise to the cancer stemness in various cancers, by carrying out self-renewal, metastasis, invasiveness enhancement^[139], and drug resistance for pancreatic tumour^[140]. Studies in CSC have led to the discovery of distinguishable cell surface markers presented in various cancer types, and this allowed the isolation of cancer stem cell for various studies^[141]. In this section, we will briefly discuss how CSC contributes to enhanced cancer malignancies, and the plausible targets in CSC that have been reported to have therapeutic value.

Signaling pathways in CSC

There are three members of hedgehog proteins in the hedgehog signaling^[141], a member of the hedgehog family, sonic hedgehog is found over-expressed in both pancreatic cancer cell and CSC^[142]. The up-regulated sonic hedgehog signaling molecules facilitates the development of PanIN and enhanced accumulation of mutations in KRAS while inhibiting the hedgehog signaling pathway by the hedgehog signaling inhibitor cyclopamine has resulted decelerated tumour growth and on set of apoptosis^[142], another inhibitor GDC-0449 is reported to produce reduced cell viability, caspase-3 mediated apoptosis, reduced tumour sphere formation in pancreatic CSC^[143]. Sonic hedgehog has displayed its critical role in tumorigenesis initiation and tumour proliferation, targeting hedgehog signaling is therefore advantageous in the early development of pancreatic cancer.

In Notch signaling pathway, over-expressed Notch-1 promotes EMT and tumour sphere formation^[144], which is confirmed by the increase expression of CD44 and EpCAM cell surface markers on CSC^[144], suggesting Notch as a factor in pancreatic tumorigenesis in CSC, but the role of it in CSC self-renewal requires further studies^[141]. Notch mediates signaling by nuclear translocation and is modified by γ -secretase before entering the nucleus, thus inhibitors of γ -secretase have been used to study the role of Notch in pancreatic cancer and also pancreatic CSC^[145], a Notch inhibitor, PF-03084014 is found able to bring a reduction of CSC population, tumour re-growth and inhibited several cancer malignancies, *e.g.*, tumour growth, angiogenesis in pancreatic cancer xenograft model with the co-administration with gemcitabine^[146], therefore its effect on pancreatic cancer is expectable.

The CXCR4 signaling pathway which comprises the ligand, stromal cell - derived factor-1/CXCR chemokine ligand 12 (SDF-1/CXCL12) and the G-protein coupled receptor, CXCR4. This signaling pathway is up-regulated in pancreatic cancer cell due to enhanced expression of CXCR4, and resulting into tumour metastasis promo-

tion; enhanced migration and strengthened stromal adhesion^[141]. In pancreatic CSC, co-expression of CD133⁺ and CXCR4⁺ on the CSC signified a highly metastatic characteristic and contributes to tumour metastasis, therefore, disrupting the SDF-1 mediated CXCR4 signaling and depletion of the CD133⁺ CSC can abrogate the metastatic characteristic of pancreatic tumour^[146]. Although targeting the CXCR4 in stopping pancreatic tumour metastasis looks promising, CXCR4 inhibitors are found highly toxic and non-specific reaction are the drawbacks that must have to be overcome before translating into clinical trials or practices^[147].

Forkhead Box M1 (FoxM1) is a transcription factor found over-expressed in pancreatic cancer in which it promotes the expression of EMT characteristic^[141], which is deduced by the increased mesenchymal cell markers expression including, zinc-finger E-box binding homeobox 1 (ZEB1), ZEB2, E-cadherin, and vimentin, and also enhanced tumour sphere formation which marks the strengthened self-renewal ability for CSC^[148], as enhanced EMT is having close resemblance to CSC in giving rise to cancer stemness^[149]. A natural compound genistein can inhibit the FoxM1 signaling pathway by down-regulating the expression of FoxM1 and its downstream gene targets (*e.g.*, VEGF, MMP-9) leading to reduced EMT and reduced tumour sphere formation and have resulted into reduced tumour growth and enhanced apoptosis^[150]. The exact mechanism of the regulation of genistein on FoxM1 and its target genes is not clear yet, however, the application genistein can rescue the microRNA-200 (miR-200) expression by attenuated FoxM1 expression, and enhanced expression of miR-200 can inhibit the EMT progression^[147]. Because of the important role for FoxM1 plays in the EMT and CSC progression, and the availability of FoxM1 inhibitor has made FoxM1 an attractive target and should evaluate the anti-tumour effects under co-administration of genistein and gemcitabine.

Cell surface marker on CSC

There are several cell surface markers on CSC which are not only be used to isolate CSC, but also have important functions towards CSC. For instance, expression of CD44⁺/CD24⁺/ESA⁺ mark the pancreatic cancer cell that function as CSC, with several signaling pathways (*e.g.*, BMI1, sonic hedgehog) up-regulated and self-renewal and tumorigenesis enhancement are observed^[151], while ablated CD133 would lead to loss of CSC self-renewal ability^[146]. This demonstrates the markers presented on the CSC can provide some clue on the de-regulated signaling pathways in the tumour which can help deciding the targets of the sub-population of the pancreatic tumour. Nevertheless, a novel CSC marker, c-Met, is found to be essential for tumour growth, tumour sphere formation and metastasis, inhibiting the expression of c-Met have suppressed these tumour malignancies^[152], possibly *via* the downstream signaling pathways of c-Met, such as Ras-MAPK, and PI3K-AKT^[153]. With the emerging knowl-

edge of the cell surface markers and their downstream signaling, options for targeting signaling transduction in PDAC is ever growing, besides, the surface markers can also act as the reference reflecting the de-regulated signaling pathways, hence, facilitating the therapeutic direction formulation.

TUMOUR MICROENVIRONMENT

Matrix metalloproteinase

MMPs are a group of zinc-dependent endopeptidases which hypothetically can degrade almost all proteins in the extracellular matrix (ECM)^[154]. MMPs are over-expressed in pancreatic cancer and the biological roles of MMPs in cancer are to digest the proteins in basement membrane in the ECM which leads to enhanced migration of tumour cell^[155], an evidence of migration signal mediation by cleavage of laminin-5 in ECM has been reported^[156]. Moreover, cleavage of E-Cadherin by MMP-3, MMP-7^[157] and A disintegrin and metalloprotease 10 (ADAM10) are observed, the loss of E-Cadherin not only enhanced cell mobility but also enhanced the tumour invasiveness and migration^[158]. In addition, the release of pro-angiogenic inflammatory cytokine (TNF- α)^[157] and VEGF^[159] are correlated with the MMPs activity and all these confirm the role of MMPs in EMT and tumour metastasis promotion^[160].

MMP is found related to the pancreatic stellate cell, which will be described in the next section, the TGF- α up-regulation in pancreatic stellate cell correlates with up-regulated MMP-1, suggesting a possibility of the association of these two molecules overexpression in enhanced tumour cell invasion^[161].

As the importance of MMPs in tumour angiogenesis and metastasis is undeniable, studies of formulating MMP inhibitors (MMPi) are undergoing; a MMPi, SB-3CT is able to reconvert the MMP-2 into its pro-enzyme state and has brought down liver metastasis^[162]. However due to the usual late stage discovery of pancreatic cancer in real life^[163] the use of MMPi is limited and also MMPs activities have been observed to be stage dependent^[164], therefore MMPs can be targeted for PDAC patients with early detection and can be applied widely when early detection method for PDAC is developed.

Pancreatic stellate cell

Pancreatic stellate cell (PaSC) resides in the exocrine of the pancreas and has dual roles in normal pancreatic tissue^[165], first it acts as a storage of vitamin A^[166], secondly upon pancreatic injury, the PaSC would be activated to acquire a myo-fibroblast-like phenotype which is called activated PaSC^[167-169], activated PaSC will secrete proteins into the ECM^[170] resulting into pancreatic fibrosis^[165] and on setting chronic pancreatitis which could lead to high risk of PDAC development. As mentioned in 10.1, high TGF- α expression level correlates with the high MMP-1 expression level in inducing PaSC migration, inhibition of MMP-1 has showed such migration induction is curbed by using MMP-1 tissue inhibitor and siRNA of

MMP-1^[161], indicating PaSC activity can be modulated *via* MMPs.

Other studies by targeting PaSC proliferation and migration have been carried out, transgelin has been reported to be over-expressed in activated PaSC but not in normal acinar cell which could cause pancreatic fibrosis^[171]. Moreover, knocking down transgelin expression has reduced cell proliferation and migration abilities in *in vitro* experiment^[171], providing a biomarker for specific therapeutic target in knocking down PaSC population in the future.

Hedgehog signaling pathway

The Hedgehog (hh) signaling pathway, involves the secreted signaling molecules hedgehog proteins, which is classified into 3 subcategories, namely, Sonic hedgehog (Shh), Indian hedgehog (Ihh), and Desert hedgehog (Dhh)^[172]. Among these 3 hh, Shh is found over-expressed in 70% of primary PDAC^[173]. Hedgehog biological roles have been described as an essential factor in embryonic development and regulate cell proliferation^[172].

The mediation of hedgehog signaling is triggered upon the binding of Shh to the Patched 12-transmembrane domain receptor (Ptch) which further activates another transmembrane signal transducer, smoothed (Smo), that would lead to localization of transcription factors in the nucleus and initiate transcription of downstream effectors^[174], for instance, Cyclin D2, FoxM1, jagged 2 (JAG2), *etc*^[175].

It is reported that tumourigenesis and tumour proliferation requires constitutive activated hedgehog signaling, and in pancreatic stromal cell in PDAC, Smo is over-expressed and direct tumour cell growth in the vicinity of stromal cell, leading to a therapy targeting hedgehog signaling in tumour-stromal interaction^[172]. Besides, report of Shh activation in CSC is crucial for CSC proliferation^[176], and it has been discussed for the CSC in aggravating pancreatic cancer treatment, such as heightened drug resistance and tumour relapse. Therefore, studying hedgehog signaling inhibitors is beneficial for pancreatic cancer treatments. A hh signaling inhibitor, Sulforaphane has been found to inhibit self-renewal capacity in CSC *via* Shh signaling inhibition leading to downstream effectors *e.g.*, Nanog and Oct-4 suppression^[176]. Moreover, inhibition of hedgehog in pancreatic cancer cells and tissue are performed and it is found that a marked decrease in EMT with EMT related transcription factors (Snail and Slug) down-regulation and had suppressed PI3K/AKT signaling, which is downstream of hedgehog signaling with an association of decreased cell proliferation^[177].

Because of diversified roles of hedgehog signaling in tumour malignancies and CSC population maintenance, and the cross talk among other signaling pathways, *e.g.* FoxM1, Notch (*via* JAG2), targeting hedgehog may have a centralized effect in weakening the malignancies of pancreatic tumour.

Stromal environment-hyaluronan

The microenvironment of pancreatic cancer, has been

accused to be the major challenge in drug delivery because of its highly dense ECM, the penetration of even small drug molecules gemcitabine is prevented^[178]. Besides, stromal cells which are the activated fibroblasts and PaSC, inflammatory cells^[178]; and distorted vascular structure of blood and lymphatic vessel composed the stromal environment^[179]. And the production of stroma is mediated by various factors involved in numerous signaling pathways in an autocrine and paracrine action, TGF- β , insulin growth factor 1, and EGF are the examples^[180].

Among the molecules in ECM, hyaluronan or hyaluronic acid (HA) is secreted by PDAC^[181], and is a repeat of N-acetylglucosamin/glucuronic acid disaccharide^[170]. It is able to interact with a hyaladherin, CD44, to regulate tyrosine kinase receptor and to facilitate angiogenesis, EMT, and chemoresistance^[182]. It is also one of the main components that contributes to high intra-tumoral fluidic pressure (IFP) through solvating with water molecule, hence, impeded the diffusion of drug molecules into the target tumour cell^[183]. It is found that co-administration of hyaluronidase with gemcitabine or other drugs prolonged the localization of the accompanied drug in the tumour^[183]. Therefore, it would become a trend for future development of drug target, *e.g.*, well incorporated with hyaluronidase to facilitate drug delivery, or even HA can be a target to disintegrate the condensed ECM, enhancing the responsiveness of the tumour cell to the treatment.

CONCLUSION

A range of therapeutic targets in PDAC have been briefly described in this article, in which their anti - tumour and oncogenic activities are characterized through various experiments and can be taken as potential target for PDAC therapies development.

Nevertheless, numerous of the targets are found overlapped with each other in producing certain kinds of tumour malignancies, *e.g.*, over-expression of CXCR4, Rac1, BMI1, and *etc.*, in pancreatic tumour cell have observed a metastasis enhancement. In light of this, and hypothetically, in order to prevent metastasis, suppressing these targets should have a more pronounced effect in metastasis inhibition. Moreover, such outflanked approach may also prevent the tumour cell from switching into other signaling pathways producing the same tumour malignancies, and achieving elimination ultimately. Besides, the current knowledge of each of these targets is insufficient, categorizing these targets by the tumour malignancies produced and identify if there is any relationship between them and understand the mechanism behind, would allow the discovery of linkages among them in terms of proteins and mRNA expression levels and mechanistic studies.

Last but not least, screening of suitable inhibitors for these targets is crucial in putting these targets into practice. Toxicity of some of the inhibitors mentioned is

reported, while, traditional Chinese medicine (TCM) may be a good source for screening inhibitors that are less or non-toxic compounds, *e.g.*, an EZH2 inhibitor, davidian, is extracted from TCM *Polygonum capitatum* without toxicity observed in xenograft model^[184].

Single effort from one side is far from enough in pancreatic tumour elimination due to its high malignancy and complex tumour microenvironment, multiple targets have to be considered in developing PDAC therapies, therefore, the way of applying these targets and which targets should be applied require further effort.

REFERENCES

- 1 **Sarkar FH**, Banerjee S, Li Y. Pancreatic cancer: pathogenesis, prevention and treatment. *Toxicol Appl Pharmacol* 2007; **224**: 326-336 [PMID: 17174370 DOI: 10.1016/j.taap.2006.11.007]
- 2 **Silvestris N**, Gnoni A, Brunetti AE, Vincenti L, Santini D, Tonini G, Merchionne F, Maiello E, Lorusso V, Nardulli P, Azzariti A, Reni M. Target therapies in pancreatic carcinoma. *Curr Med Chem* 2014; **21**: 948-965 [PMID: 23992319]
- 3 **Iovanna J**, Mallmann MC, Gonçalves A, Turrini O, Dagorn JC. Current knowledge on pancreatic cancer. *Front Oncol* 2012; **2**: 6 [PMID: 22655256 DOI: 10.3389/fonc.2012.00006]
- 4 **Hill R**, Rabb M, Madureira PA, Clements D, Gujar SA, Waisman DM, Giacomantonio CA, Lee PW. Gemcitabine-mediated tumour regression and p53-dependent gene expression: implications for colon and pancreatic cancer therapy. *Cell Death Dis* 2013; **4**: e791 [PMID: 24008735 DOI: 10.1038/cddis.2013.307]
- 5 **Algül H**, Treiber M, Lesina M, Schmid RM. Mechanisms of disease: chronic inflammation and cancer in the pancreas - a potential role for pancreatic stellate cells? *Nat Clin Pract Gastroenterol Hepatol* 2007; **4**: 454-462 [PMID: 17667994 DOI: 10.1038/ncpgasthep0881]
- 6 **Cowley MJ**, Chang DK, Pajic M, Johns AL, Waddell N, Grimmond SM, Biankin AV. Understanding pancreatic cancer genomes. *J Hepatobiliary Pancreat Sci* 2013; Epub ahead of print [PMID: 23660961 DOI: 10.1007/s00534-013-0610-6]
- 7 **Danov SA**, Wong HH, Lemoine NR. Targeted therapies for pancreatic cancer. *Br Med Bull* 2008; **87**: 97-130 [PMID: 18753179 DOI: 10.1093/bmb/ldn027]
- 8 **Tang L**, Dai DL, Su M, Martinka M, Li G, Zhou Y. Aberrant expression of collagen triple helix repeat containing 1 in human solid cancers. *Clin Cancer Res* 2006; **12**: 3716-3722 [PMID: 16778098 DOI: 10.1158/1078-0432.CCR-06-0030]
- 9 **Pygay P**, Heroult M, Wang Q, Lehnert W, Belden J, Liaw L, Friesel RE, Lindner V. Collagen triple helix repeat containing 1, a novel secreted protein in injured and diseased arteries, inhibits collagen expression and promotes cell migration. *Circ Res* 2005; **96**: 261-268 [PMID: 15618538]
- 10 **Park EH**, Kim S, Jo JY, Kim SJ, Hwang Y, Kim JM, Song SY, Lee DK, Koh SS. Collagen triple helix repeat containing-1 promotes pancreatic cancer progression by regulating migration and adhesion of tumor cells. *Carcinogenesis* 2013; **34**: 694-702 [PMID: 23222813 DOI: 10.1093/carcin/bgs378]
- 11 **Wang Y**. Wnt/Planar cell polarity signaling: a new paradigm for cancer therapy. *Mol Cancer Ther* 2009; **8**: 2103-2109 [PMID: 19671746 DOI: 10.1158/1535-7163.MCT-09-0282]
- 12 **Avizienyte E**, Frame MC. Src and FAK signalling controls adhesion fate and the epithelial-to-mesenchymal transition. *Curr Opin Cell Biol* 2005; **17**: 542-547 [PMID: 16099634 DOI: 10.1016/j.ceb.2005.08.007]
- 13 **Didsbury J**, Weber RF, Bokoch GM, Evans T, Snyderman R. rac, a novel ras-related family of proteins that are botulinum toxin substrates. *J Biol Chem* 1989; **264**: 16378-16382 [PMID: 2674130]

- 14 **Heid I**, Lubeseder-Martellato C, Sipos B, Mazur PK, Lesina M, Schmid RM, Siveke JT. Early requirement of Rac1 in a mouse model of pancreatic cancer. *Gastroenterology* 2011; **141**: 719-730, 730.e1-7 [PMID: 21684285 DOI: 10.1053/j.gastro.2011.04.043]
- 15 **Razidlo GL**, Wang Y, Chen J, Krueger EW, Billadeau DD, McNiven MA. Dynamin 2 potentiates invasive migration of pancreatic tumor cells through stabilization of the Rac1 GEF Vav1. *Dev Cell* 2013; **24**: 573-585 [PMID: 23537630 DOI: 10.1016/j.devcel.2013.02.010]
- 16 **Wang P**, Chen L, Zhang J, Chen H, Fan J, Wang K, Luo J, Chen Z, Meng Z, Liu L. Methylation-mediated silencing of the miR-124 genes facilitates pancreatic cancer progression and metastasis by targeting Rac1. *Oncogene* 2014; **33**: 514-524 [PMID: 23334332 DOI: 10.1038/onc.2012.598]
- 17 **Guo X**, Wang M, Jiang J, Xie C, Peng F, Li X, Tian R, Qin R. Balanced Tiam1-rac1 and RhoA drives proliferation and invasion of pancreatic cancer cells. *Mol Cancer Res* 2013; **11**: 230-239 [PMID: 23322732 DOI: 10.1158/1541-7786.MCR-12-0632]
- 18 **Lazer G**, Katzav S. Guanine nucleotide exchange factors for RhoGTPases: good therapeutic targets for cancer therapy? *Cell Signal* 2011; **23**: 969-979 [PMID: 21044680 DOI: 10.1016/j.cellsig.2010.10.022]
- 19 **Neel NF**, Martin TD, Stratford JK, Zand TP, Reiner DJ, Der CJ. The RalGEF-Ral Effector Signaling Network: The Road Less Traveled for Anti-Ras Drug Discovery. *Genes Cancer* 2011; **2**: 275-287 [PMID: 21779498 DOI: 10.1177/1947601911407329]
- 20 **Kashatus DF**. Ral GTPases in tumorigenesis: emerging from the shadows. *Exp Cell Res* 2013; **319**: 2337-2342 [PMID: 23830877 DOI: 10.1016/j.jcyexr.2013.06.020]
- 21 **Lim KH**, O'Hayer K, Adam SJ, Kendall SD, Campbell PM, Der CJ, Counter CM. Divergent roles for RalA and RalB in malignant growth of human pancreatic carcinoma cells. *Curr Biol* 2006; **16**: 2385-2394 [PMID: 17174914 DOI: 10.1016/j.cub.2006.10.023]
- 22 **Neyraud V**, Aushev VN, Hatzoglou A, Meunier B, Cascone I, Camonis J. RalA and RalB proteins are ubiquitinated GTPases, and ubiquitinated RalA increases lipid raft exposure at the plasma membrane. *J Biol Chem* 2012; **287**: 29397-29405 [PMID: 22700969 DOI: 10.1074/jbc.M112.357764]
- 23 **Lim KH**, Baines AT, Fiordalisi JJ, Shipitsin M, Feig LA, Cox AD, Der CJ, Counter CM. Activation of RalA is critical for Ras-induced tumorigenesis of human cells. *Cancer Cell* 2005; **7**: 533-545 [PMID: 15950903 DOI: 10.1016/j.ccr.2005.04.030]
- 24 **Chien Y**, White MA. RAL GTPases are linchpin modulators of human tumour-cell proliferation and survival. *EMBO Rep* 2003; **4**: 800-806 [PMID: 12856001 DOI: 10.1038/sj.embor.embor899]
- 25 **Balasubramanian N**, Meier JA, Scott DW, Norambuena A, White MA, Schwartz MA. RalA-exocyst complex regulates integrin-dependent membrane raft exocytosis and growth signaling. *Curr Biol* 2010; **20**: 75-79 [PMID: 20005108 DOI: 10.1016/j.cub.2009.11.016]
- 26 **Avila JL**, Kissil JL. Notch signaling in pancreatic cancer: oncogene or tumor suppressor? *Trends Mol Med* 2013; **19**: 320-327 [PMID: 23545339 DOI: 10.1016/j.molmed.2013.03.003]
- 27 **Wang Z**, Li Y, Ahmad A, Azmi AS, Banerjee S, Kong D, Sarkar FH. Targeting Notch signaling pathway to overcome drug resistance for cancer therapy. *Biochim Biophys Acta* 2010; **1806**: 258-267 [PMID: 20600632 DOI: 10.1016/j.bbcan.2010.06.001]
- 28 **Miyamoto Y**, Maitra A, Ghosh B, Zechner U, Argani P, Iacobuzio-Donahue CA, Sriuranpong V, Iso T, Meszoely IM, Wolfe MS, Hruban RH, Ball DW, Schmid RM, Leach SD. Notch mediates TGF alpha-induced changes in epithelial differentiation during pancreatic tumorigenesis. *Cancer Cell* 2003; **3**: 565-576 [PMID: 12842085 DOI: 10.1016/S1535-6108(03)00140-5]
- 29 **Hanlon L**, Avila JL, Demarest RM, Troutman S, Allen M, Ratti F, Rustgi AK, Stanger BZ, Radtke F, Adsay V, Long F, Capobianco AJ, Kissil JL. Notch1 functions as a tumor suppressor in a model of K-ras-induced pancreatic ductal adenocarcinoma. *Cancer Res* 2010; **70**: 4280-4286 [PMID: 20484026 DOI: 10.1158/0008-5472.CAN-09-4645]
- 30 **Weijzen S**, Rizzo P, Braid M, Vaishnav R, Jonkheer SM, Zlobin A, Osborne BA, Gottipati S, Aster JC, Hahn WC, Rudolf M, Siziopikou K, Kast WM, Miele L. Activation of Notch-1 signaling maintains the neoplastic phenotype in human Ras-transformed cells. *Nat Med* 2002; **8**: 979-986 [PMID: 12185362 DOI: 10.1038/nm754]
- 31 **Gurdon JB**, Bourillot PY. Morphogen gradient interpretation. *Nature* 2001; **413**: 797-803 [PMID: 11677596 DOI: 10.1038/35101500]
- 32 **Gurdon JB**, Harger P, Mitchell A, Lemaire P. Activin signaling and response to a morphogen gradient. *Nature* 1994; **371**: 487-492 [PMID: 7935761 DOI: 10.1038/371487a0]
- 33 **Lee KL**, Lim SK, Orlov YL, Yit le Y, Yang H, Ang LT, Poellinger L, Lim B. Graded Nodal/Activin signaling titrates conversion of quantitative phospho-Smad2 levels into qualitative embryonic stem cell fate decisions. *PLoS Genet* 2011; **7**: e1002130 [PMID: 21731500 DOI: 10.1371/journal.pgen.1002130]
- 34 **Lonardo E**, Hermann PC, Mueller MT, Huber S, Balic A, Miranda-Lorenzo I, Zagorac S, Alcalá S, Rodriguez-Arabaolaza I, Ramirez JC, Torres-Ruiz R, Garcia E, Hidalgo M, Cebrián DÁ, Heuchel R, Löhr M, Berger F, Bartenstein P, Aicher A, Heeschen C. Nodal/Activin signaling drives self-renewal and tumorigenicity of pancreatic cancer stem cells and provides a target for combined drug therapy. *Cell Stem Cell* 2011; **9**: 433-446 [PMID: 22056140 DOI: 10.1016/j.stem.2011.10.001]
- 35 **Olive KP**, Jacobetz MA, Davidson JC, Gopinathan A, McIntyre D, Honess D, Madhu B, Goldgraben MA, Caldwell ME, Allard D, Frese KK, Denicola G, Feig C, Combs C, Winter SP, Ireland-Zecchini H, Reichelt S, Howat WJ, Chang A, Dhara M, Wang L, Rückert F, Grützmann R, Pilarsky C, Izeradjene K, Hingorani SR, Huang P, Davies SE, Plunkett W, Egorin M, Hruban RH, Whitebread N, McGovern K, Adams J, Iacobuzio-Donahue C, Griffiths J, Tuveson DA. Inhibition of Hedgehog signaling enhances delivery of chemotherapy in a mouse model of pancreatic cancer. *Science* 2009; **324**: 1457-1461 [PMID: 19460966 DOI: 10.1126/science.1171362]
- 36 **Sun J**, Zhang D, Bae DH, Sahni S, Jansson P, Zheng Y, Zhao Q, Yue F, Zheng M, Kovacevic Z, Richardson DR. Metastasis suppressor, NDRG1, mediates its activity through signaling pathways and molecular motors. *Carcinogenesis* 2013; **34**: 1943-1954 [PMID: 23671130 DOI: 10.1093/carcin/bgt163]
- 37 **Lv XH**, Chen JW, Zhao G, Feng ZZ, Yang DH, Sun WW, Fan JS, Zhu GH. N-myc downstream-regulated gene 1/Cap43 may function as tumor suppressor in endometrial cancer. *J Cancer Res Clin Oncol* 2012; **138**: 1703-1715 [PMID: 22678098 DOI: 10.1007/s00432-012-1249-4]
- 38 **Angst E**, Sibold S, Tiffon C, Weimann R, Gloor B, Candinas D, Stroka D. Cellular differentiation determines the expression of the hypoxia-inducible protein NDRG1 in pancreatic cancer. *Br J Cancer* 2006; **95**: 307-313 [PMID: 16832411 DOI: 10.1038/sj.bjc.6603256]
- 39 **Angst E**, Dawson DW, Nguyen A, Park J, Go VL, Reber HA, Hines OJ, Eibl G. Epigenetic regulation affects N-myc downstream-regulated gene 1 expression indirectly in pancreatic cancer cells. *Pancreas* 2010; **39**: 675-679 [PMID: 20173668 DOI: 10.1097/MPA.0b013e3181c8b476]
- 40 **Kovacevic Z**, Chikhani S, Lui GY, Sivagurunathan S, Richardson DR. The iron-regulated metastasis suppressor NDRG1 targets NEDD4L, PTEN, and SMAD4 and inhibits the PI3K and Ras signaling pathways. *Antioxid Redox Signal* 2013; **18**: 874-887 [PMID: 22462691 DOI: 10.1089/ars.2011.4273]
- 41 **Bao B**, Ali S, Ahmad A, Azmi AS, Li Y, Banerjee S, Kong D, Sethi S, Aboukameel A, Padhye SB, Sarkar FH. Hypoxia-induced aggressiveness of pancreatic cancer cells is due to increased expression of VEGF, IL-6 and miR-21, which can

- be attenuated by CDF treatment. *PLoS One* 2012; **7**: e50165 [PMID: 23272057 DOI: 10.1371/journal.pone.0050165]
- 42 **Guillaumond F**, Leca J, Olivares O, Lavaut MN, Vidal N, Berthezène P, Dusetti NJ, Loncle C, Calvo E, Turrini O, Iovanna JL, Tomasini R, Vasseur S. Strengthened glycolysis under hypoxia supports tumor symbiosis and hexosamine biosynthesis in pancreatic adenocarcinoma. *Proc Natl Acad Sci USA* 2013; **110**: 3919-3924 [PMID: 23407165 DOI: 10.1073/pnas.1219555110]
- 43 **Le A**, Lane AN, Hamaker M, Bose S, Gouw A, Barbi J, Tsukamoto T, Rojas CJ, Slusher BS, Zhang H, Zimmerman LJ, Liebler DC, Slebos RJ, Lorkiewicz PK, Higashi RM, Fan TW, Dang CV. Glucose-independent glutamine metabolism via TCA cycling for proliferation and survival in B cells. *Cell Metab* 2012; **15**: 110-121 [PMID: 22225880 DOI: 10.1016/j.cmet.2011.12.009]
- 44 **Ying H**, Kimmelman AC, Lyssiotis CA, Hua S, Chu GC, Fletcher-Sananikone E, Locasale JW, Son J, Zhang H, Coloff JL, Yan H, Wang W, Chen S, Viale A, Zheng H, Paik JH, Lim C, Guimaraes AR, Martin ES, Chang J, Hezel AF, Perry SR, Hu J, Gan B, Xiao Y, Asara JM, Weissleder R, Wang YA, Chin L, Cantley LC, DePinho RA. Oncogenic Kras maintains pancreatic tumors through regulation of anabolic glucose metabolism. *Cell* 2012; **149**: 656-670 [PMID: 22541435 DOI: 10.1016/j.cell.2012.01.058]
- 45 **Dando I**, Donadelli M, Costanzo C, Dalla Pozza E, D'Alessandro A, Zolla L, Palmieri M. Cannabinoids inhibit energetic metabolism and induce AMPK-dependent autophagy in pancreatic cancer cells. *Cell Death Dis* 2013; **4**: e664 [PMID: 23764845 DOI: 10.1038/cddis.2013.151]
- 46 **Wolpin BM**, Bao Y, Qian ZR, Wu C, Kraft P, Ogino S, Stampfer MJ, Sato K, Ma J, Buring JE, Sesso HD, Lee IM, Gaziano JM, McTiernan A, Phillips LS, Cochrane BB, Pollak MN, Manson JE, Giovannucci EL, Fuchs CS. Hyperglycemia, insulin resistance, impaired pancreatic β -cell function, and risk of pancreatic cancer. *J Natl Cancer Inst* 2013; **105**: 1027-1035 [PMID: 23847240 DOI: 10.1093/jnci/djt123]
- 47 **Donadelli M**, Dando I, Zaniboni T, Costanzo C, Dalla Pozza E, Scupoli MT, Scarpa A, Zappavigna S, Marra M, Abbruzzese A, Bifulco M, Caraglia M, Palmieri M. Gemcitabine/cannabinoid combination triggers autophagy in pancreatic cancer cells through a ROS-mediated mechanism. *Cell Death Dis* 2011; **2**: e152 [PMID: 21525939 DOI: 10.1038/cddis.2011.36]
- 48 **Murray J**, Taylor SW, Zhang B, Ghosh SS, Capaldi RA. Oxidative damage to mitochondrial complex I due to peroxytrite: identification of reactive tyrosines by mass spectrometry. *J Biol Chem* 2003; **278**: 37223-37230 [PMID: 12857734 DOI: 10.1074/jbc.M305694200]
- 49 **Rajeshkumar NV**, Rasheed ZA, García-García E, López-Ríos F, Fujiwara K, Matsui WH, Hidalgo M. A combination of DR5 agonistic monoclonal antibody with gemcitabine targets pancreatic cancer stem cells and results in long-term disease control in human pancreatic cancer model. *Mol Cancer Ther* 2010; **9**: 2582-2592 [PMID: 20660600 DOI: 10.1158/1535-7163.MCT-10-0370]
- 50 **Thorburn A**. Death receptor-induced cell killing. *Cell Signal* 2004; **16**: 139-144 [PMID: 14636884 DOI: 10.1016/j.cellsig.2003.08.007]
- 51 **Kischkel FC**, Lawrence DA, Chuntharapai A, Schow P, Kim KJ, Ashkenazi A. Apo2L/TRAIL-dependent recruitment of endogenous FADD and caspase-8 to death receptors 4 and 5. *Immunity* 2000; **12**: 611-620 [PMID: 10894161 DOI: 10.1016/S1074-7613(00)80212-5]
- 52 **Kong R**, Jia G, Cheng ZX, Wang YW, Mu M, Wang SJ, Pan SH, Gao Y, Jiang HC, Dong DL, Sun B. Dihydroartemisinin enhances Apo2L/TRAIL-mediated apoptosis in pancreatic cancer cells via ROS-mediated up-regulation of death receptor 5. *PLoS One* 2012; **7**: e37222 [PMID: 22666346 DOI: 10.1371/journal.pone.0037222]
- 53 **Ashkenazi A**. Targeting death and decoy receptors of the tumour-necrosis factor superfamily. *Nat Rev Cancer* 2002; **2**: 420-430 [PMID: 12189384 DOI: 10.1038/nrc821]
- 54 **Chang LH**, Pan SL, Lai CY, Tsai AC, Teng CM. Activated PAR-2 regulates pancreatic cancer progression through ILK/HIF- α -induced TGF- α expression and MEK/VEGF-A-mediated angiogenesis. *Am J Pathol* 2013; **183**: 566-575 [PMID: 23764046 DOI: 10.1016/j.ajpath.2013.04.022]
- 55 **Dutra-Oliveira A**, Monteiro RQ, Mariano-Oliveira A. Protease-activated receptor-2 (PAR2) mediates VEGF production through the ERK1/2 pathway in human glioblastoma cell lines. *Biochem Biophys Res Commun* 2012; **421**: 221-227 [PMID: 22497886 DOI: 10.1016/j.bbrc.2012.03.140]
- 56 **Tan C**, Cruet-Hennequart S, Troussard A, Fazli L, Costello P, Sutton K, Wheeler J, Gleave M, Sanghera J, Dedhar S. Regulation of tumor angiogenesis by integrin-linked kinase (ILK). *Cancer Cell* 2004; **5**: 79-90 [PMID: 14749128 DOI: 10.1016/S1535-6108(03)00281-2]
- 57 **Shi K**, Queiroz KC, Stap J, Richel DJ, Spek CA. Protease-activated receptor-2 induces migration of pancreatic cancer cells in an extracellular ATP-dependent manner. *J Thromb Haemost* 2013; **11**: 1892-1902 [PMID: 23899344 DOI: 10.1111/jth.12361]
- 58 **Desai MD**, Saroya BS, Lockhart AC. Investigational therapies targeting the ErbB (EGFR, HER2, HER3, HER4) family in GI cancers. *Expert Opin Investig Drugs* 2013; **22**: 341-356 [PMID: 23316969 DOI: 10.1517/13543784.2013.761972]
- 59 **Lazrek Y**, Dubreuil O, Garambois V, Gaborit N, Larboret C, Le Clorennec C, Thomas G, Leconet W, Jarlier M, Pugnère M, Vié N, Robert B, Monnet C, Bouayadi K, Kharrat H, Mondon P, Pèlerin A, Chardès T. Anti-HER3 domain 1 and 3 antibodies reduce tumor growth by hindering HER2/HER3 dimerization and AKT-induced MDM2, XIAP, and FoxO1 phosphorylation. *Neoplasia* 2013; **15**: 335-347 [PMID: 23479511 DOI: 10.1593/neo.121960]
- 60 **Hirakawa T**, Nakata B, Amano R, Kimura K, Shimizu S, Ohira G, Yamada N, Ohira M, Hirakawa K. HER3 overexpression as an independent indicator of poor prognosis for patients with curatively resected pancreatic cancer. *Oncology* 2011; **81**: 192-198 [PMID: 22067729 DOI: 10.1159/000333825]
- 61 **Hsieh AC**, Moasser MM. Targeting HER proteins in cancer therapy and the role of the non-target HER3. *Br J Cancer* 2007; **97**: 453-457 [PMID: 17667926 DOI: 10.1038/sj.bjc.6603910]
- 62 **Berlin J**, Keedy VL, Janne PA, Yee L, Rizvi NA, Jin X, Copigneaux C, Hettmann T, Beaupre DM, LoRusso P. A first-in-human phase I study of U3-1287 (AMG 888), a HER3 inhibitor, in patients (pts) with advanced solid tumors. 2011 ASCO Annual Meeting. Available from: URL: <http://meetinglibrary.asco.org/content/84026-102>
- 63 **Schoeberl B**, Faber AC, Li D, Liang MC, Crosby K, Onsum M, Burenkova O, Pace E, Walton Z, Nie L, Fulgham A, Song Y, Nielsen UB, Engelman JA, Wong KK. An ErbB3 antibody, MM-121, is active in cancers with ligand-dependent activation. *Cancer Res* 2010; **70**: 2485-2494 [PMID: 20215504 DOI: 10.1158/0008-5472.CAN-09-3145]
- 64 **Yotsumoto F**, Fukami T, Yagi H, Funakoshi A, Yoshizato T, Kuroki M, Miyamoto S. Amphiregulin regulates the activation of ERK and Akt through epidermal growth factor receptor and HER3 signals involved in the progression of pancreatic cancer. *Cancer Sci* 2010; **101**: 2351-2360 [PMID: 20726858 DOI: 10.1111/j.1349-7006.2010.01671.x]
- 65 **von Burstin J**, Eser S, Paul MC, Seidler B, Brandl M, Messer M, von Werder A, Schmidt A, Mages J, Pagel P, Schnieke A, Schmid RM, Schneider G, Saur D. E-cadherin regulates metastasis of pancreatic cancer in vivo and is suppressed by a SNAIL/HDAC1/HDAC2 repressor complex. *Gastroenterology* 2009; **137**: 361-371, 371.e1-5 [PMID: 19362090 DOI: 10.1053/j.gastro.2009.04.004]
- 66 **Teppass U**, Truong K, Godt D, Ikura M, Peifer M. Cadherins in embryonic and neural morphogenesis. *Nat Rev Mol Cell Biol* 2000; **1**: 91-100 [PMID: 11253370 DOI: 10.1038/35040042]
- 67 **Gumbiner BM**. Regulation of cadherin-mediated adhesion

- in morphogenesis. *Nat Rev Mol Cell Biol* 2005; **6**: 622-634 [PMID: 16025097 DOI: 10.1038/nrm1699]
- 68 **Carneiro P**, Figueiredo J, Bordeira-Carriço R, Fernandes MS, Carvalho J, Oliveira C, Seruca R. Therapeutic targets associated to E-cadherin dysfunction in gastric cancer. *Expert Opin Ther Targets* 2013; **17**: 1187-1201 [PMID: 23957294 DOI: 10.1517/14728222.2013.827174]
- 69 **Qazi AM**, Gruzdyn O, Semaan A, Seward S, Chamala S, Dhulipala V, Sethi S, Ali-Fehmi R, Philip PA, Bouwman DL, Weaver DW, Gruber SA, Batchu RB. Restoration of E-cadherin expression in pancreatic ductal adenocarcinoma treated with microRNA-101. *Surgery* 2012; **152**: 704-711; discussion 711-713 [PMID: 22943841 DOI: 10.1016/j.surg.2012.07.020]
- 70 **Winter JM**, Ting AH, Vilardeil F, Gallmeier E, Baylin SB, Hruban RH, Kern SE, Iacobuzio-Donahue CA. Absence of E-cadherin expression distinguishes noncohesive from cohesive pancreatic cancer. *Clin Cancer Res* 2008; **14**: 412-418 [PMID: 18223216 DOI: 10.1158/1078-0432.CCR-07-0487]
- 71 **Jung EJ**, Moon HG, Cho BI, Jeong CY, Joo YT, Lee YJ, Hong SC, Choi SK, Ha WS, Kim JW, Lee CW, Lee JS, Park ST. Galectin-1 expression in cancer-associated stromal cells correlates tumor invasiveness and tumor progression in breast cancer. *Int J Cancer* 2007; **120**: 2331-2338 [PMID: 17304502 DOI: 10.1002/ijc.22434]
- 72 **Horiguchi N**, Arimoto K, Mizutani A, Endo-Ichikawa Y, Nakada H, Taketani S. Galectin-1 induces cell adhesion to the extracellular matrix and apoptosis of non-adherent human colon cancer Colo201 cells. *J Biochem* 2003; **134**: 869-874 [PMID: 14769876 DOI: 10.1093/jb/mvg213]
- 73 **Paclik D**, Danese S, Berndt U, Wiedenmann B, Dignass A, Sturm A. Galectin-4 controls intestinal inflammation by selective regulation of peripheral and mucosal T cell apoptosis and cell cycle. *PLoS One* 2008; **3**: e2629 [PMID: 18612433 DOI: 10.1371/journal.pone.0002629]
- 74 **Belo AI**, van der Sar AM, Tefsen B, van Die I. Galectin-4 Reduces Migration and Metastasis Formation of Pancreatic Cancer Cells. *PLoS One* 2013; **8**: e65957 [PMID: 23824659 DOI: 10.1371/journal.pone.0065957]
- 75 **Boll M**, Fuchs G, Meier C, Trautwein A, El Kasmi A, Ragsdale SW, Buchanan G, Lowe DJ. Redox centers of 4-hydroxybenzoyl-CoA reductase, a member of the xanthine oxidase family of molybdenum-containing enzymes. *J Biol Chem* 2001; **276**: 47853-47862 [PMID: 11602591]
- 76 **Rumilla KM**, Erickson LA, Erickson AK, Lloyd RV. Galectin-4 expression in carcinoid tumors. *Endocr Pathol* 2006; **17**: 243-249 [PMID: 17308361]
- 77 **Jung H**, Lee KP, Park SJ, Park JH, Jang YS, Choi SY, Jung JG, Jo K, Park DY, Yoon JH, Park JH, Lim DS, Hong GR, Choi C, Park YK, Lee JW, Hong HJ, Kim S, Park YW. TMPRSS4 promotes invasion, migration and metastasis of human tumor cells by facilitating an epithelial-mesenchymal transition. *Oncogene* 2008; **27**: 2635-2647 [PMID: 17968309 DOI: 10.1038/sj.onc.1210914]
- 78 **Dawelbait G**, Winter C, Zhang Y, Pilarsky C, Grützmann R, Heinrich JC, Schroeder M. Structural templates predict novel protein interactions and targets from pancreas tumour gene expression data. *Bioinformatics* 2007; **23**: i115-i124 [PMID: 17646287 DOI: 10.1093/bioinformatics/btm188]
- 79 **Kim S**, Kang HY, Nam EH, Choi MS, Zhao XF, Hong CS, Lee JW, Lee JH, Park YK. TMPRSS4 induces invasion and epithelial-mesenchymal transition through upregulation of integrin alpha5 and its signaling pathways. *Carcinogenesis* 2010; **31**: 597-606 [PMID: 20118200 DOI: 10.1093/carcin/bgq024]
- 80 **Min HJ**, Lee Y, Zhao XF, Park YK, Lee MK, Lee JW, Kim S. TMPRSS4 upregulates uPA gene expression through JNK signaling activation to induce cancer cell invasion. *Cell Signal* 2014; **26**: 398-408 [PMID: 23978400 DOI: 10.1016/j.cellsig.2013.08.002]
- 81 **Schimmer AD**. Inhibitor of apoptosis proteins: translating basic knowledge into clinical practice. *Cancer Res* 2004; **64**: 7183-7190 [PMID: 15492230 DOI: 10.1158/0008-5472.CAN-04-1918]
- 82 **Mita AC**, Mita MM, Nawrocki ST, Giles FJ. Survivin: key regulator of mitosis and apoptosis and novel target for cancer therapeutics. *Clin Cancer Res* 2008; **14**: 5000-5005 [PMID: 18698017 DOI: 10.1158/1078-0432.CCR-08-0746]
- 83 **Delvaeye M**, De Vriese A, Zwerts F, Betz I, Moons M, Aurtiero M, Conway EM. Role of the 2 zebrafish survivin genes in vasculo-angiogenesis, neurogenesis, cardiogenesis and hematopoiesis. *BMC Dev Biol* 2009; **9**: 25 [PMID: 19323830 DOI: 10.1186/1471-213X-9-25]
- 84 **Dohi T**, Okada K, Xia F, Wilford CE, Samuel T, Welsh K, Marusawa H, Zou H, Armstrong R, Matsuzawa S, Salvesen GS, Reed JC, Altieri DC. An IAP-IAP complex inhibits apoptosis. *J Biol Chem* 2004; **279**: 34087-34090 [PMID: 15218035 DOI: 10.1074/jbc.C400236200]
- 85 **Karikari CA**, Roy I, Tryggestad E, Feldmann G, Pinilla C, Welsh K, Reed JC, Armour EP, Wong J, Herman J, Rakheja D, Maitra A. Targeting the apoptotic machinery in pancreatic cancers using small-molecule antagonists of the X-linked inhibitor of apoptosis protein. *Mol Cancer Ther* 2007; **6**: 957-966 [PMID: 17339366 DOI: 10.1158/1535-7163.MCT-06-0634]
- 86 **Zai HY**, Yi XP, Li YX, You XY, Cao LP, Liu H. [X-linked inhibitor of apoptosis protein (XIAP) and Survivin suppression on human pancreatic cancer cells Panc-1 proliferation and chemosensitivity]. *Beijing Daxue Xuebao* 2013; **45**: 242-249 [PMID: 23591345]
- 87 **Chang TC**, Mendell JT. microRNAs in vertebrate physiology and human disease. *Annu Rev Genomics Hum Genet* 2007; **8**: 215-239 [PMID: 17506656 DOI: 10.1146/annurev.genom.8.080706.092351]
- 88 **Tang S**, Bonaroti J, Unlu S, Liang X, Tang D, Zeh HJ, Lotze MT. Sweating the small stuff: microRNAs and genetic changes define pancreatic cancer. *Pancreas* 2013; **42**: 740-759 [PMID: 23774697 DOI: 10.1097/MPA.0b013e3182854ab0]
- 89 **Pramanik D**, Campbell NR, Karikari C, Chivukula R, Kent OA, Mendell JT, Maitra A. Restitution of tumor suppressor microRNAs using a systemic nanovector inhibits pancreatic cancer growth in mice. *Mol Cancer Ther* 2011; **10**: 1470-1480 [PMID: 21622730 DOI: 10.1158/1535-7163.MCT-11-0152]
- 90 **Kent OA**, Mullendore M, Wentzel EA, López-Romero P, Tan AC, Alvarez H, West K, Ochs MF, Hidalgo M, Arking DE, Maitra A, Mendell JT. A resource for analysis of microRNA expression and function in pancreatic ductal adenocarcinoma cells. *Cancer Biol Ther* 2009; **8**: 2013-2024 [PMID: 20037478 DOI: 10.4161/cbt.8.21.9685]
- 91 **Liu C**, Kelnar K, Liu B, Chen X, Calhoun-Davis T, Li H, Patravala L, Yan H, Jeter C, Honorio S, Wiggins JF, Bader AG, Fagin R, Brown D, Tang DG. The microRNA miR-34a inhibits prostate cancer stem cells and metastasis by directly repressing CD44. *Nat Med* 2011; **17**: 211-215 [PMID: 21240262 DOI: 10.1038/nm.2284]
- 92 **Chang TC**, Wentzel EA, Kent OA, Ramachandran K, Mullendore M, Lee KH, Feldmann G, Yamakuchi M, Ferlito M, Lowenstein CJ, Arking DE, Beer MA, Maitra A, Mendell JT. Transactivation of miR-34a by p53 broadly influences gene expression and promotes apoptosis. *Mol Cell* 2007; **26**: 745-752 [PMID: 17540599 DOI: 10.1016/j.molcel.2007.05.010]
- 93 **Ji Q**, Hao X, Meng Y, Zhang M, Desano J, Fan D, Xu L. Restoration of tumor suppressor miR-34 inhibits human p53-mutant gastric cancer tumorspheres. *BMC Cancer* 2008; **8**: 266 [PMID: 18803879 DOI: 10.1186/1471-2407-8-266]
- 94 **Ji Q**, Hao X, Zhang M, Tang W, Yang M, Li L, Xiang D, Desano J, Bommer GT, Fan D, Fearon ER, Lawrence TS, Xu L. MicroRNA miR-34 inhibits human pancreatic cancer tumor-initiating cells. *PLoS One* 2009; **4**: e6816 [PMID: 19714243 DOI: 10.1371/journal.pone.0006816]
- 95 **Hu Y**, Ou Y, Wu K, Chen Y, Sun W. miR-143 inhibits the metastasis of pancreatic cancer and an associated signaling

- pathway. *Tumour Biol* 2012; **33**: 1863-1870 [PMID: 23070684 DOI: 10.1007/s13277-012-0446-8]
- 96 **Kent OA**, Chivukula RR, Mullendore M, Wentzel EA, Feldmann G, Lee KH, Liu S, Leach SD, Maitra A, Mendell JT. Repression of the miR-143/145 cluster by oncogenic Ras initiates a tumor-promoting feed-forward pathway. *Genes Dev* 2010; **24**: 2754-2759 [PMID: 21159816 DOI: 10.1101/gad.1950610]
- 97 **Pham H**, Rodriguez CE, Donald GW, Hertzner KM, Jung XS, Chang HH, Moro A, Reber HA, Hines OJ, Eibl G. miR-143 decreases COX-2 mRNA stability and expression in pancreatic cancer cells. *Biochem Biophys Res Commun* 2013; **439**: 6-11 [PMID: 23973710 DOI: 10.1016/j.bbrc.2013.08.042]
- 98 **Greenhough A**, Smartt HJ, Moore AE, Roberts HR, Williams AC, Paraskeva C, Kaidi A. The COX-2/PGE2 pathway: key roles in the hallmarks of cancer and adaptation to the tumour microenvironment. *Carcinogenesis* 2009; **30**: 377-386 [PMID: 19136477 DOI: 10.1093/carcin/bgp014]
- 99 **Moore AE**, Young LE, Dixon DA. A common single-nucleotide polymorphism in cyclooxygenase-2 disrupts microRNA-mediated regulation. *Oncogene* 2012; **31**: 1592-1598 [PMID: 21822307 DOI: 10.1038/onc.2011.349]
- 100 **Soubani O**, Ali AS, Logna F, Ali S, Philip PA, Sarkar FH. Re-expression of miR-200 by novel approaches regulates the expression of PTEN and MT1-MMP in pancreatic cancer. *Carcinogenesis* 2012; **33**: 1563-1571 [PMID: 22637745 DOI: 10.1093/carcin/bgs189]
- 101 **Kim S**, Huang W, Mottillo EP, Sohail A, Ham YA, Conley-Lacomb MK, Kim CJ, Tzivion G, Kim HR, Wang S, Chen YQ, Fridman R. Posttranslational regulation of membrane type 1-matrix metalloproteinase (MT1-MMP) in mouse PTEN null prostate cancer cells: Enhanced surface expression and differential O-glycosylation of MT1-MMP. *Biochim Biophys Acta* 2010; **1803**: 1287-1297 [PMID: 20620173 DOI: 10.1016/j.bbamcr.2010.06.011]
- 102 **Zhang D**, Brodt P. Type 1 insulin-like growth factor regulates MT1-MMP synthesis and tumor invasion via PI 3-kinase/Akt signaling. *Oncogene* 2003; **22**: 974-982 [PMID: 12592384 DOI: 10.1038/sj.onc.1206197]
- 103 **Petrella BL**, Brinckerhoff CE. PTEN suppression of YY1 induces HIF-2 activity in von-Hippel-Lindau-null renal-cell carcinoma. *Cancer Biol Ther* 2009; **8**: 1389-1401 [PMID: 19483472 DOI: 10.4161/cbt.8.14.8880]
- 104 **Reis EM**, Verjovski-Almeida S. Perspectives of Long Non-Coding RNAs in Cancer Diagnostics. *Front Genet* 2012; **3**: 32 [PMID: 22408643 DOI: 10.3389/fgene.2012.00032]
- 105 **Mercer TR**, Dinger ME, Mattick JS. Long non-coding RNAs: insights into functions. *Nat Rev Genet* 2009; **10**: 155-159 [PMID: 19188922 DOI: 10.1038/nrg2521]
- 106 **Derrien T**, Johnson R, Bussotti G, Tanzer A, Djebali S, Tilgner H, Guernec G, Martin D, Merkel A, Knowles DG, Lagarde J, Veeravalli L, Ruan X, Ruan Y, Lassmann T, Carninci P, Brown JB, Lipovich L, Gonzalez JM, Thomas M, Davis CA, Shiekhattar R, Gingeras TR, Hubbard TJ, Notredame C, Harrow J, Guigó R. The GENCODE v7 catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and expression. *Genome Res* 2012; **22**: 1775-1789 [PMID: 22955988 DOI: 10.1101/gr.132159.111]
- 107 **Sana J**, Faltejskova P, Svoboda M, Slaby O. Novel classes of non-coding RNAs and cancer. *J Transl Med* 2012; **10**: 103 [PMID: 22613733 DOI: 10.1186/1479-5876-10-103]
- 108 **Li CH**, Chen Y. Targeting long non-coding RNAs in cancers: progress and prospects. *Int J Biochem Cell Biol* 2013; **45**: 1895-1910 [PMID: 23748105 DOI: 10.1016/j.biocel.2013.05.030]
- 109 **Kapranov P**, St Laurent G, Raz T, Ozsolak F, Reynolds CP, Sorensen PH, Reaman G, Milos P, Arcenci RJ, Thompson JF, Triche TJ. The majority of total nuclear-encoded non-ribosomal RNA in a human cell is 'dark matter' un-annotated RNA. *BMC Biol* 2010; **8**: 149 [PMID: 21176148 DOI: 10.1186/1741-7007-8-149]
- 110 **Tahira AC**, Kubrusly MS, Faria MF, Dazzani B, Fonseca RS, Maracaja-Coutinho V, Verjovski-Almeida S, Machado MC, Reis EM. Long noncoding intronic RNAs are differentially expressed in primary and metastatic pancreatic cancer. *Mol Cancer* 2011; **10**: 141 [PMID: 22078386 DOI: 10.1186/1476-4598-10-141]
- 111 **Stratford JK**, Bentrem DJ, Anderson JM, Fan C, Volmar KA, Marron JS, Routh ED, Caskey LS, Samuel JC, Der CJ, Thorne LB, Calvo BF, Kim HJ, Talamonti MS, Iacobuzio-Donahue CA, Hollingsworth MA, Perou CM, Yeh JJ. A six-gene signature predicts survival of patients with localized pancreatic ductal adenocarcinoma. *PLoS Med* 2010; **7**: e1000307 [PMID: 20644708 DOI: 10.1371/journal.pmed.1000307]
- 112 **Khalil AM**, Guttman M, Huarte M, Garber M, Raj A, Rivea Morales D, Thomas K, Presser A, Bernstein BE, van Oudenaarden A, Regev A, Lander ES, Rinn JL. Many human large intergenic noncoding RNAs associate with chromatin-modifying complexes and affect gene expression. *Proc Natl Acad Sci USA* 2009; **106**: 11667-11672 [PMID: 19571010 DOI: 10.1073/pnas.0904715106]
- 113 **Wang KC**, Chang HY. Molecular mechanisms of long non-coding RNAs. *Mol Cell* 2011; **43**: 904-914 [PMID: 21925379 DOI: 10.1016/j.molcel.2011.08.018]
- 114 **Kim K**, Jutooru I, Chadalapaka G, Johnson G, Frank J, Burghardt R, Kim S, Safe S. HOTAIR is a negative prognostic factor and exhibits pro-oncogenic activity in pancreatic cancer. *Oncogene* 2013; **32**: 1616-1625 [PMID: 22614017 DOI: 10.1038/onc.2012.193]
- 115 **Gutschner T**, Hämmerle M, Diederichs S. MALAT1 -- a paradigm for long noncoding RNA function in cancer. *J Mol Med (Berl)* 2013; **91**: 791-801 [PMID: 23529762 DOI: 10.1007/s00109-013-1028-y]
- 116 **Ji P**, Diederichs S, Wang W, Böing S, Metzger R, Schneider PM, Tidow N, Brandt B, Buerger H, Bulk E, Thomas M, Berdel WE, Serve H, Müller-Tidow C. MALAT-1, a novel noncoding RNA, and thymosin beta4 predict metastasis and survival in early-stage non-small cell lung cancer. *Oncogene* 2003; **22**: 8031-8041 [PMID: 12970751 DOI: 10.1038/sj.onc.1206928]
- 117 **Gutschner T**, Hämmerle M, Eissmann M, Hsu J, Kim Y, Hung G, Revenko A, Arun G, Stentrup M, Gross M, Zörnig M, MacLeod AR, Spector DL, Diederichs S. The noncoding RNA MALAT1 is a critical regulator of the metastasis phenotype of lung cancer cells. *Cancer Res* 2013; **73**: 1180-1189 [PMID: 23243023 DOI: 10.1158/0008-5472.CAN-12-2850]
- 118 **Ji Q**, Liu X, Fu X, Zhang L, Sui H, Zhou L, Sun J, Cai J, Qin J, Ren J, Li Q. Resveratrol inhibits invasion and metastasis of colorectal cancer cells via MALAT1 mediated Wnt/ β -catenin signal pathway. *PLoS One* 2013; **8**: e78700 [PMID: 24244343 DOI: 10.1371/journal.pone.0078700]
- 119 **Tripathi V**, Shen Z, Chakraborty A, Giri S, Freier SM, Wu X, Zhang Y, Gorospe M, Prasanth SG, Lal A, Prasanth KV. Long noncoding RNA MALAT1 controls cell cycle progression by regulating the expression of oncogenic transcription factor B-MYB. *PLoS Genet* 2013; **9**: e1003368 [PMID: 23555285 DOI: 10.1371/journal.pgen.1003368]
- 120 **Schneider G**, Krämer OH, Fritsche P, Schüler S, Schmid RM, Saur D. Targeting histone deacetylases in pancreatic ductal adenocarcinoma. *J Cell Mol Med* 2010; **14**: 1255-1263 [PMID: 19929947 DOI: 10.1111/j.1582-4934.2009.00974.x]
- 121 **Ouaïssi M**, Cabral S, Tavares J, da Silva AC, Mathieu Daude F, Mas E, Bernard J, Sastre B, Lombardo D, Ouaiissi A. Histone deacetylase (HDAC) encoding gene expression in pancreatic cancer cell lines and cell sensitivity to HDAC inhibitors. *Cancer Biol Ther* 2008; **7**: 523-531 [PMID: 18296916 DOI: 10.4161/cbt.7.4.5480]
- 122 **Fritsche P**, Seidler B, Schüler S, Schnieke A, Göttlicher M, Schmid RM, Saur D, Schneider G. HDAC2 mediates therapeutic resistance of pancreatic cancer cells via the BH3-only protein NOXA. *Gut* 2009; **58**: 1399-1409 [PMID: 19528037]

- DOI: 10.1136/gut.2009.180711]
- 123 **Gong DJ**, Zhang JM, Yu M, Zhuang B, Guo QQ. Inhibition of SIRT1 combined with gemcitabine therapy for pancreatic carcinoma. *Clin Interv Aging* 2013; **8**: 889-897 [PMID: 23898224 DOI: 10.2147/CIA.S45064]
 - 124 **Weichert W**, Boehm M, Gekeler V, Bahra M, Langrehr J, Neuhaus P, Denkert C, Imre G, Weller C, Hofmann HP, Niesporek S, Jacob J, Dietel M, Scheiderei C, Kristiansen G. High expression of RelA/p65 is associated with activation of nuclear factor-kappaB-dependent signaling in pancreatic cancer and marks a patient population with poor prognosis. *Br J Cancer* 2007; **97**: 523-530 [PMID: 17622249 DOI: 10.1038/sj.bjc.6603878]
 - 125 **Lehmann A**, Denkert C, Budczies J, Buckendahl AC, Darb-Esfahani S, Noske A, Müller BM, Bahra M, Neuhaus P, Dietel M, Kristiansen G, Weichert W. High class I HDAC activity and expression are associated with RelA/p65 activation in pancreatic cancer in vitro and in vivo. *BMC Cancer* 2009; **9**: 395 [PMID: 19912635 DOI: 10.1186/1471-2407-9-395]
 - 126 **Hu J**, Colburn NH. Histone deacetylase inhibition down-regulates cyclin D1 transcription by inhibiting nuclear factor-kappaB/p65 DNA binding. *Mol Cancer Res* 2005; **3**: 100-109 [PMID: 15755876 DOI: 10.1158/1541-7786.MCR-04-0070]
 - 127 **Song W**, Tao K, Li H, Jin C, Song Z, Li J, Shi H, Li X, Dang Z, Dou K. Bmi-1 is related to proliferation, survival and poor prognosis in pancreatic cancer. *Cancer Sci* 2010; **101**: 1754-1760 [PMID: 20426791 DOI: 10.1111/j.1349-7006.2010.01577.x]
 - 128 **Yin T**, Wei H, Leng Z, Yang Z, Gou S, Wu H, Zhao G, Hu X, Wang C. Bmi-1 promotes the chemoresistance, invasion and tumorigenesis of pancreatic cancer cells. *Chemotherapy* 2011; **57**: 488-496 [PMID: 22248802 DOI: 10.1159/000334103]
 - 129 **Proctor E**, Waghay M, Lee CJ, Heidt DG, Yalamanchili M, Li C, Bednar F, Simeone DM. Bmi1 enhances tumorigenicity and cancer stem cell function in pancreatic adenocarcinoma. *PLoS One* 2013; **8**: e55820 [PMID: 23437065 DOI: 10.1371/journal.pone.0055820]
 - 130 **van Vlerken LE**, Kiefer CM, Morehouse C, Li Y, Groves C, Wilson SD, Yao Y, Hollingsworth RE, Hurt EM. EZH2 is required for breast and pancreatic cancer stem cell maintenance and can be used as a functional cancer stem cell reporter. *Stem Cells Transl Med* 2013; **2**: 43-52 [PMID: 23283488 DOI: 10.5966/sctm.2012-0036]
 - 131 **Fujii S**, Fukamachi K, Tsuda H, Ito K, Ito Y, Ochiai A. RAS oncogenic signal upregulates EZH2 in pancreatic cancer. *Biochem Biophys Res Commun* 2012; **417**: 1074-1079 [PMID: 2222375 DOI: 10.1016/j.bbrc.2011.12.099]
 - 132 **Banerjee R**, Mani RS, Russo N, Scanlon CS, Tsodikov A, Jing X, Cao Q, Palanisamy N, Metwally T, Inglehart RC, Tomlins S, Bradford C, Carey T, Wolf G, Kalyana-Sundaram S, Chinnaiyan AM, Varambally S, D'Silva NJ. The tumor suppressor gene rap1GAP is silenced by miR-101-mediated EZH2 overexpression in invasive squamous cell carcinoma. *Oncogene* 2011; **30**: 4339-4349 [PMID: 21532618 DOI: 10.1038/onc.2011.141]
 - 133 **Li CH**, To KF, Tong JH, Xiao Z, Xia T, Lai PB, Chow SC, Zhu YX, Chan SL, Marquez VE, Chen Y. Enhancer of zeste homolog 2 silences microRNA-218 in human pancreatic ductal adenocarcinoma cells by inducing formation of heterochromatin. *Gastroenterology* 2013; **144**: 1086-1097.e9 [PMID: 23395645 DOI: 10.1053/j.gastro.2013.01.058]
 - 134 **Deng J**, He M, Chen L, Chen C, Zheng J, Cai Z. The loss of miR-26a-mediated post-transcriptional regulation of cyclin E2 in pancreatic cancer cell proliferation and decreased patient survival. *PLoS One* 2013; **8**: e76450 [PMID: 24116110 DOI: 10.1371/journal.pone.0076450]
 - 135 **Avan A**, Crea F, Paolicchi E, Funel N, Galvani E, Marquez VE, Honeywell RJ, Danesi R, Peters GJ, Giovannetti E. Molecular mechanisms involved in the synergistic interaction of the EZH2 inhibitor 3-deazaneplanocin A with gemcitabine in pancreatic cancer cells. *Mol Cancer Ther* 2012; **11**: 1735-1746 [PMID: 22622284 DOI: 10.1158/1535-7163.MCT-12-0037]
 - 136 **Lasfargues C**, Pyronnet S. EZH2 links pancreatitis to tissue regeneration and pancreatic cancer. *Clin Res Hepatol Gastroenterol* 2012; **36**: 323-324 [PMID: 22766150 DOI: 10.1016/j.clinre.2012.05.010]
 - 137 **Aguirre AJ**, Bardeesy N, Sinha M, Lopez L, Tuveson DA, Horner J, Redston MS, DePinho RA. Activated Kras and Ink4a/Arf deficiency cooperate to produce metastatic pancreatic ductal adenocarcinoma. *Genes Dev* 2003; **17**: 3112-3126 [PMID: 14681207 DOI: 10.1101/gad.1158703]
 - 138 **Pietras A**. Cancer stem cells in tumor heterogeneity. *Adv Cancer Res* 2011; **112**: 255-281 [PMID: 21925307 DOI: 10.1016/B978-0-12-387688-1.00009-0]
 - 139 **Dalerba P**, Clarke MF. Cancer stem cells and tumor metastasis: first steps into uncharted territory. *Cell Stem Cell* 2007; **1**: 241-242 [PMID: 18371356 DOI: 10.1016/j.stem.2007.08.012]
 - 140 **Rasheed ZA**, Kowalski J, Smith BD, Matsui W. Concise review: Emerging concepts in clinical targeting of cancer stem cells. *Stem Cells* 2011; **29**: 883-887 [PMID: 21509907 DOI: 10.1002/stem.648]
 - 141 **Xia J**, Chen C, Chen Z, Miele L, Sarkar FH, Wang Z. Targeting pancreatic cancer stem cells for cancer therapy. *Biochim Biophys Acta* 2012; **1826**: 385-399 [PMID: 22728049 DOI: 10.1016/j.bbcan.2012.06.002]
 - 142 **Thayer SP**, di Magliano MP, Heiser PW, Nielsen CM, Roberts DJ, Lauwers GY, Qi YP, Gysin S, Fernández-del Castillo C, Yajnik V, Antoniu B, McMahon M, Warshaw AL, Hebrok M. Hedgehog is an early and late mediator of pancreatic cancer tumorigenesis. *Nature* 2003; **425**: 851-856 [PMID: 14520413 DOI: 10.1038/nature02009]
 - 143 **Singh BN**, Fu J, Srivastava RK, Shankar S. Hedgehog signaling antagonist GDC-0449 (Vismodegib) inhibits pancreatic cancer stem cell characteristics: molecular mechanisms. *PLoS One* 2011; **6**: e27306 [PMID: 22087285 DOI: 10.1371/journal.pone.0027306]
 - 144 **Bao B**, Wang Z, Ali S, Kong D, Li Y, Ahmad A, Banerjee S, Azmi AS, Miele L, Sarkar FH. Notch-1 induces epithelial-mesenchymal transition consistent with cancer stem cell phenotype in pancreatic cancer cells. *Cancer Lett* 2011; **307**: 26-36 [PMID: 21463919 DOI: 10.1016/j.canlet.2011.03.012]
 - 145 **Yabuuchi S**, Pai SG, Campbell NR, de Wilde RF, De Oliveira E, Korangath P, Strepel MM, Rasheed ZA, Hidalgo M, Maitra A, Rajeshkumar NV. Notch signaling pathway targeted therapy suppresses tumor progression and metastatic spread in pancreatic cancer. *Cancer Lett* 2013; **335**: 41-51 [PMID: 23402814 DOI: 10.1016/j.canlet.2013.01.054]
 - 146 **Hermann PC**, Huber SL, Herrler T, Aicher A, Ellwart JW, Guba M, Bruns CJ, Heeschen C. Distinct populations of cancer stem cells determine tumor growth and metastatic activity in human pancreatic cancer. *Cell Stem Cell* 2007; **1**: 313-323 [PMID: 18371365 DOI: 10.1016/j.stem.2007.06.002]
 - 147 **Kalatskaya I**, Berchiche YA, Gravel S, Limberg BJ, Rosenbaum JS, Heveker N. AMD3100 is a CXCR7 ligand with allosteric agonist properties. *Mol Pharmacol* 2009; **75**: 1240-1247 [PMID: 19255243 DOI: 10.1124/mol.108.053389]
 - 148 **Bao B**, Wang Z, Ali S, Kong D, Banerjee S, Ahmad A, Li Y, Azmi AS, Miele L, Sarkar FH. Over-expression of FoxM1 leads to epithelial-mesenchymal transition and cancer stem cell phenotype in pancreatic cancer cells. *J Cell Biochem* 2011; **112**: 2296-2306 [PMID: 21503965 DOI: 10.1002/jcb.23150]
 - 149 **Christiansen JJ**, Rajasekaran AK. Reassessing epithelial to mesenchymal transition as a prerequisite for carcinoma invasion and metastasis. *Cancer Res* 2006; **66**: 8319-8326 [PMID: 16951136 DOI: 10.1158/0008-5472.CAN-06-0410]
 - 150 **Wang Z**, Ahmad A, Banerjee S, Azmi A, Kong D, Li Y, Sarkar FH. FoxM1 is a novel target of a natural agent in pancreatic cancer. *Pharm Res* 2010; **27**: 1159-1168 [PMID: 20354770 DOI: 10.1007/s11095-010-0106-x]
 - 151 **Li C**, Lee CJ, Simeone DM. Identification of human pancre-

- atic cancer stem cells. *Methods Mol Biol* 2009; **568**: 161-173 [PMID: 19582426 DOI: 10.1007/978-1-59745-280-9_10]
- 152 **Li C**, Wu JJ, Hynes M, Dosch J, Sarkar B, Welling TH, Pasca di Magliano M, Simeone DM. c-Met is a marker of pancreatic cancer stem cells and therapeutic target. *Gastroenterology* 2011; **141**: 2218-2227.e5 [PMID: 21864475 DOI: 10.1053/j.gastro.2011.08.009]
- 153 **Hage C**, Rausch V, Giese N, Giese T, Schönsiegel F, Labsch S, Nwaeburu C, Mattern J, Gladkich J, Herr I. The novel c-Met inhibitor cabozantinib overcomes gemcitabine resistance and stem cell signaling in pancreatic cancer. *Cell Death Dis* 2013; **4**: e627 [PMID: 23661005 DOI: 10.1038/cddis.2013.158]
- 154 **Murphy G**, Nagase H. Progress in matrix metalloproteinase research. *Mol Aspects Med* 2008; **29**: 290-308 [PMID: 18619669 DOI: 10.1016/j.mam.2008.05.002]
- 155 **Roy R**, Yang J, Moses MA. Matrix metalloproteinases as novel biomarkers and potential therapeutic targets in human cancer. *J Clin Oncol* 2009; **27**: 5287-5297 [PMID: 19738110 DOI: 10.1200/JCO.2009.23.5556]
- 156 **Koshikawa N**, Giannelli G, Cirulli V, Miyazaki K, Quaranta V. Role of cell surface metalloprotease MT1-MMP in epithelial cell migration over laminin-5. *J Cell Biol* 2000; **148**: 615-624 [PMID: 10662785 DOI: 10.1083/jcb.148.3.615]
- 157 **Haro H**, Crawford HC, Fingleton B, Shinomiya K, Spengler DM, Matrisian LM. Matrix metalloproteinase-7-dependent release of tumor necrosis factor- α in a model of herniated disc resorption. *J Clin Invest* 2000; **105**: 143-150 [PMID: 10642592 DOI: 10.1172/JCI7091]
- 158 **Maretzky T**, Reiss K, Ludwig A, Buchholz J, Scholz F, Proksch E, de Strooper B, Hartmann D, Saftig P. ADAM10 mediates E-cadherin shedding and regulates epithelial cell-cell adhesion, migration, and beta-catenin translocation. *Proc Natl Acad Sci USA* 2005; **102**: 9182-9187 [PMID: 15958533 DOI: 10.1073/pnas.0500918102]
- 159 **Lamoreaux WJ**, Fitzgerald ME, Reiner A, Hasty KA, Charles ST. Vascular endothelial growth factor increases release of gelatinase A and decreases release of tissue inhibitor of metalloproteinases by microvascular endothelial cells in vitro. *Microvasc Res* 1998; **55**: 29-42 [PMID: 9473407 DOI: 10.1006/mvre.1997.2056]
- 160 **Thiery JP**. Epithelial-mesenchymal transitions in tumour progression. *Nat Rev Cancer* 2002; **2**: 442-454 [PMID: 12189386 DOI: 10.1038/nrc822]
- 161 **Gaida MM**, Haag N, Günther F, Tschaharganeh DF, Schirmacher P, Friess H, Giese NA, Schmidt J, Wente MN. Expression of A disintegrin and metalloprotease 10 in pancreatic carcinoma. *Int J Mol Med* 2010; **26**: 281-288 [PMID: 20596609 DOI: 10.3892/ijmm.00000463]
- 162 **Krüger A**, Arlt MJ, Gerg M, Kopitz C, Bernardo MM, Chang M, Mobashery S, Fridman R. Antimetastatic activity of a novel mechanism-based gelatinase inhibitor. *Cancer Res* 2005; **65**: 3523-3526 [PMID: 15867341 DOI: 10.1158/0008-5472.CAN-04-3570]
- 163 **Stetler-Stevenson WG**. Tissue inhibitors of metalloproteinases in cell signaling: metalloproteinase-independent biological activities. *Sci Signal* 2008; **1**: re6 [PMID: 18612141 DOI: 10.1126/scisignal.127re6]
- 164 **Bogaczewicz J**, Jasielski P, Mosiewicz A, Trojanowski T, Suchozebrska-Jesioneck D, Stryjecka-Zimmer M. [The role of matrix metalloproteinases and tissue inhibitors of metalloproteinases in invasion of tumours of neuroepithelial tissue]. *Neurol Neurochir Pol* 2006; **40**: 404-412 [PMID: 17103354 DOI: 10.1080/10408360801973244]
- 165 **Won JH**, Zhang Y, Ji B, Logsdon CD, Yule DI. Phenotypic changes in mouse pancreatic stellate cell Ca²⁺ signaling events following activation in culture and in a disease model of pancreatitis. *Mol Biol Cell* 2011; **22**: 421-436 [PMID: 21148289 DOI: 10.1091/mbc.e10-10-0807]
- 166 **Lonardo E**, Frias-Aldeguer J, Hermann PC, Heeschen C. Pancreatic stellate cells form a niche for cancer stem cells and promote their self-renewal and invasiveness. *Cell Cycle* 2012; **11**: 1282-1290 [PMID: 22421149 DOI: 10.4161/cc.19679]
- 167 **Bissell DM**, Friedman SL, Maher JJ, Roll FJ. Connective tissue biology and hepatic fibrosis: report of a conference. *Hepatology* 1990; **11**: 488-498 [PMID: 2179098 DOI: 10.1002/hep.1840110322]
- 168 **Schneider E**, Schmid-Kotsas A, Zhao J, Weidenbach H, Schmid RM, Menke A, Adler G, Waltenberger J, Grünert A, Bachem MG. Identification of mediators stimulating proliferation and matrix synthesis of rat pancreatic stellate cells. *Am J Physiol Cell Physiol* 2001; **281**: C532-C543 [PMID: 11443052]
- 169 **Pandolf S**, Gukovskaya A, Edderkaoui M, Dawson D, Eibl G, Lugea A. Epidemiology, risk factors, and the promotion of pancreatic cancer: role of the stellate cell. *J Gastroenterol Hepatol* 2012; **27** Suppl 2: 127-134 [PMID: 22320930 DOI: 10.1111/j.1440-1746.2011.07013.x]
- 170 **Jacobetz MA**, Chan DS, Neesse A, Bapiro TE, Cook N, Frese KK, Feig C, Nakagawa T, Caldwell ME, Zecchini HI, Lolkema MP, Jiang P, Kultti A, Thompson CB, Maneval DC, Jodrell DL, Frost GL, Shepard HM, Skepper JN, Tuveson DA. Hyaluronan impairs vascular function and drug delivery in a mouse model of pancreatic cancer. *Gut* 2013; **62**: 112-120 [PMID: 22466618 DOI: 10.1136/gutjnl-2012-302529]
- 171 **Apte MV**, Yang L, Phillips PA, Xu Z, Kaplan W, Cowley M, Pirola RC, Wilson JS. Extracellular matrix composition significantly influences pancreatic stellate cell gene expression pattern: role of transgelin in PSC function. *Am J Physiol Gastrointest Liver Physiol* 2013; **305**: G408-G417 [PMID: 23868411 DOI: 10.1152/ajpgi.00016.2013]
- 172 **Walter K**, Omura N, Hong SM, Griffith M, Vincent A, Borges M, Goggins M. Overexpression of smoothed activates the sonic hedgehog signaling pathway in pancreatic cancer-associated fibroblasts. *Clin Cancer Res* 2010; **16**: 1781-1789 [PMID: 20215540 DOI: 10.1158/1078-0432.CCR-09-1913]
- 173 **Hebrok M**. Hedgehog signaling in pancreas development. *Mech Dev* 2003; **120**: 45-57 [PMID: 12490295 DOI: 10.1016/S0925-4773(02)00331-3]
- 174 **Katoh Y**, Katoh M. Hedgehog signaling pathway and gastrointestinal stem cell signaling network (review). *Int J Mol Med* 2006; **18**: 1019-1023 [PMID: 17089004]
- 175 **Chowdhury S**, Pradhan RN, Sarkar RR. Structural and logical analysis of a comprehensive hedgehog signaling pathway to identify alternative drug targets for glioma, colon and pancreatic cancer. *PLoS One* 2013; **8**: e69132 [PMID: 23935937 DOI: 10.1371/journal.pone.0069132]
- 176 **Rodova M**, Fu J, Watkins DN, Srivastava RK, Shankar S. Sonic hedgehog signaling inhibition provides opportunities for targeted therapy by sulforaphane in regulating pancreatic cancer stem cell self-renewal. *PLoS One* 2012; **7**: e46083 [PMID: 23029396 DOI: 10.1371/journal.pone.0046083]
- 177 **Hao K**, Tian XD, Qin CF, Xie XH, Yang YM. Hedgehog signaling pathway regulates human pancreatic cancer cell proliferation and metastasis. *Oncol Rep* 2013; **29**: 1124-1132 [PMID: 23292285 DOI: 10.3892/or.2012.2210]
- 178 **Michl P**, Gress TM. Improving drug delivery to pancreatic cancer: breaching the stromal fortress by targeting hyaluronic acid. *Gut* 2012; **61**: 1377-1379 [PMID: 22661496 DOI: 10.1136/gutjnl-2012-302604]
- 179 **Neesse A**, Michl P, Frese KK, Feig C, Cook N, Jacobetz MA, Lolkema MP, Buchholz M, Olive KP, Gress TM, Tuveson DA. Stromal biology and therapy in pancreatic cancer. *Gut* 2011; **60**: 861-868 [PMID: 20966025 DOI: 10.1136/gut.2010.226092]
- 180 **Mahadevan D**, Von Hoff DD. Tumor-stroma interactions in pancreatic ductal adenocarcinoma. *Mol Cancer Ther* 2007; **6**: 1186-1197 [PMID: 17406031 DOI: 10.1158/1535-7163.MCT-06-0686]
- 181 **Mahlbacher V**, Sewing A, Elsässer HP, Kern HF. Hyaluronan is a secretory product of human pancreatic adenocarcinoma cells. *Eur J Cell Biol* 1992; **58**: 28-34 [PMID: 1644063]

- 182 **Toole BP**, Slomiany MG. Hyaluronan: a constitutive regulator of chemoresistance and malignancy in cancer cells. *Semin Cancer Biol* 2008; **18**: 244-250 [PMID: 18534864 DOI: 10.1016/j.semcancer.2008.03.009]
- 183 **Provenzano PP**, Cuevas C, Chang AE, Goel VK, Von Hoff DD, Hingorani SR. Enzymatic targeting of the stroma ablates physical barriers to treatment of pancreatic ductal adenocarcinoma. *Cancer Cell* 2012; **21**: 418-429 [PMID: 22439937 DOI: 10.1016/j.ccr.2012.01.007]
- 184 **Wang Y**, Ma J, Chow SC, Li CH, Xiao Z, Feng R, Fu J, Chen Y. A potential antitumor ellagitannin, davidiin, inhibited hepatocellular tumor growth by targeting EZH2. *Tumour Biol* 2014; **35**: 205-212 [PMID: 23897557 DOI: 10.1007/s13277-013-1025-3]

P- Reviewer: Chunyi H, Mishra PK **S- Editor:** Qi Y
L- Editor: A **E- Editor:** Liu XM





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>



ISSN 1007-9327

