

Ultrasound-guided continuous fascia iliaca compartment ...

<https://europepmc.org/article/MED/30186422> ▾

Jul 06, 2018 · Another study suggested that both **ultrasound-guided continuous femoral nerve block and fascia iliaca compartment block** with a novel cannula-over-needle provide effective **anesthesia** and...

Ultrasound-guided supra-inguinal fascia iliaca block: a ...

<https://onlinelibrary.wiley.com/doi/10.1111/j.1365-2044.2011.06628.x>

Feb 24, 2011 · Chen Ling, Xing-Qing Liu, Yi-Qun Li, Xian-Jie Wen, Xu-Dong Hu, Kai Yang, **Ultrasound-guided fascia iliaca compartment block combined** with general **anesthesia** for amputation in an acute...

Cited by: 83

Author: P. Hebbard, J. Ivanusic, S. Sha

Publish Year: 2011

PEOPLE ALSO ASK

Is the fascia iliaca compartment block safe to use? ▾

How is FIC used to treat hip fractures? ▾

What are pain protocols for elderly hip fracture patients? ▾

Are there guidelines for anaesthesia for hip fracture patients? ▾

[Combination of high-dose dexmedetomidine sedation and ...](#)

<https://www.sciencedirect.com/science/article/pii/S0952818010000073>

May 01, 2010 · Two patients with hip fracture and with severe cardiac disease, who were anesthetized using a combination of dexmedetomidine sedation and fascia iliaca compartment block, are reported. The block covered almost the entire skin incision area and the major area of the fractured bone.

Cited by: 5 Author: Takayuki Kunisawa, Misaki Ota, Akihiro Su...

Publish Year: 2010

[A continuous infusion fascia iliaca compartment block in ...](#)

europepmc.org/articles/PMC3279500

Jan 17, 2012 · All patients who presented to our institution with a hip fracture were given the option of having a continuous fascia iliaca compartment block for pain control versus usual pain management (non-opioids, opioids, and ice therapy). The block was monitored by the pain service until the day of discharge from the hospital.

Estimated Reading Time: 11 mins


PEOPLE ALSO ASK

What are pain protocols for elderly hip fracture patients? ▾

Can a fascia iliaca block be administered under ultrasound? ▾

How long does it take to get analgesia after a hip fracture? ▾

Why is early anaesthetic important for hip fractures? ▾



Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 68979

Manuscript Type: CASE REPORT

Combined fascia iliaca compartment block and monitored anesthesia care for the geriatric patients with hip fracture: Two case reports

Zhan L. *et al.* Combined FICB and MAC for elderly hip fracture

Li Zhan, Yu-Jie Zhang, Jing-Xian Wang

Match Overview



1	Crossref 18 words Carole Lin, Curtis Darling, Brian C. H. Tsui. "Practical Regional Anesthesia Guide for Elderly Patients", <i>Drugs & Aging</i> , 2019	1%
2	Internet 17 words crawled on 09-May-2020 journals.sagepub.com	1%
3	Crossref 18 words Takayuki Kunitzawa, Misaki Ota, Akihiro Suzuki, Osamu Takahata, Hiroshi Iwasaki. "Combination of high-dose dexmedetomidine ..."	1%
4	Crossref 15 words Nizar H. Amin, Hank L. Hutchinson, Anthony G. Sarzone. "Infiltration Techniques for Local Infiltration Analgesia With Liposo ..."	1%

国内版

国际版

Combined fascia iliaca compartment block and monitored anesthe



ALL

IMAGES

VIDEOS

4,360 Results

Any time ▾

Effectiveness of Fascia Iliaca Compartment Block after ...

<https://europepmc.org/article/PMC/PMC8124308> ▾

May 04, 2021 · An assessment of the feasibility of fascia iliaca compartment block (FICB) combined with nonopioid analgesics and patient controlled analgesia (PCA), oxycodone, in the perioperative anaesthetic management for elective total hip replacement (THR). Design A randomised, single-center, open-label study. Setting A single hospital.

Author: Wojciech Gola, Szymon Bialka, Alek... Publish Year: 2021

Effects of fascia iliaca compartment block combined with ...

<https://www.dovepress.com/effects-of-fascia-iliaca...> ▾

Nov 08, 2018 · Effects of fascia iliaca compartment block combined with general laryngeal mask airway anesthesia in children undergoing femoral fracture surgery: a randomized trial. Background: Postoperative agitation after general anesthesia is a common complication in children; however, pain or uncomfortable feeling is the main reason of emergence agitation ...

Cited by: 1 Author: HY Zhong, XB Deng, Z Wang

Publish Year: 2018

PEOPLE ALSO ASK

How long does it take to get analgesia after a hip fracture? ▾

What are the barriers to regional anesthesia for trauma patients? ▾

How is local anesthesia used in intensive care? ▾

Which is better for anesthesia lidocaine or mepivacaine? ▾

Feedback

Combination of high-dose dexmedetomidine sedation and ...

<https://www.sciencedirect.com/science/article/pii/S0952818010000073>

May 01, 2010 · Two patients with hip fracture and with severe cardiac disease, who were anesthetized using a combination of dexmedetomidine sedation and fascia iliaca compartment block, are reported. The block covered almost the entire skin incision area and the major area of the fractured bone.