

## PEER-REVIEW REPORT

**Name of journal:** World Journal of Clinical Cases

**Manuscript NO:** 68979

**Title:** Combined fascia iliaca compartment block and monitored anesthesia care for geriatric patients with hip fracture: Two case reports

**Reviewer's code:** 05928806

**Position:** Peer Reviewer

**Academic degree:** MD

**Professional title:** Doctor

**Reviewer's Country/Territory:** Japan

**Author's Country/Territory:** China

**Manuscript submission date:** 2021-06-13

**Reviewer chosen by:** AI Technique

**Reviewer accepted review:** 2021-06-16 11:09

**Reviewer performed review:** 2021-06-16 11:29

**Review time:** 1 Hour

|                                 |   |
|---------------------------------|---|
| <b>Scientific quality</b>       | <input type="checkbox"/> Grade A: Excellent <input type="checkbox"/> Grade B: Very good <input checked="" type="checkbox"/> Grade C: Good<br><input type="checkbox"/> Grade D: Fair <input type="checkbox"/> Grade E: Do not publish            |
| <b>Language quality</b>         | <input checked="" type="checkbox"/> Grade A: Priority publishing <input type="checkbox"/> Grade B: Minor language polishing<br><input type="checkbox"/> Grade C: A great deal of language polishing <input type="checkbox"/> Grade D: Rejection |
| <b>Conclusion</b>               | <input type="checkbox"/> Accept (High priority) <input checked="" type="checkbox"/> Accept (General priority)<br><input type="checkbox"/> Minor revision <input type="checkbox"/> Major revision <input type="checkbox"/> Rejection             |
| <b>Re-review</b>                | <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No   |
| <b>Peer-reviewer statements</b> | Peer-Review: <input checked="" type="checkbox"/> Anonymous <input type="checkbox"/> Onymous<br>Conflicts-of-Interest: <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No   |

#### **SPECIFIC COMMENTS TO AUTHORS**

Osteosynthesis of the hip fracture by PFNA is a very short procedure. I have performed more than 10 cases under local anaesthesia in patients in a poor general condition. THA is quite challenging. Do you have a backup plan in case the anaesthetic wears off or is not effective enough? Or did you consider stopping the operation if that happened? It would be helpful to specify the necessary preparations for the reader.