

World Journal of *Clinical Cases*

World J Clin Cases 2021 November 6; 9(31): 9320-9698



FRONTIER

- 9320 Gut-liver axis in cirrhosis: Are hemodynamic changes a missing link?
Maslennikov R, Ivashkin V, Efremova I, Poluektova E, Shirokova E

REVIEW

- 9333 Pharmaconutrition strategy to resolve SARS-CoV-2-induced inflammatory cytokine storm in non-alcoholic fatty liver disease: Omega-3 long-chain polyunsaturated fatty acids
Jeyakumar SM, Vajreswari A
- 9350 Major depressive disorder: Validated treatments and future challenges
Karrouri R, Hammani Z, Benjelloun R, Otheman Y

MINIREVIEWS

- 9368 Gene × environment interaction in major depressive disorder
Zhao MZ, Song XS, Ma JS
- 9376 Deep learning driven colorectal lesion detection in gastrointestinal endoscopic and pathological imaging
Cai YW, Dong FF, Shi YH, Lu LY, Chen C, Lin P, Xue YS, Chen JH, Chen SY, Luo XB

ORIGINAL ARTICLE**Case Control Study**

- 9386 Cognitive behavioral therapy on personality characteristics of cancer patients
Yuan XH, Peng J, Hu SW, Yang Y, Bai YJ

Retrospective Cohort Study

- 9395 Extrapneumonic necrosis volume: A new tool in acute pancreatitis severity assessment?
Cucuteanu B, Negru D, Gavrilescu O, Popa IV, Floria M, Mihai C, Cijevschi Prelipcean C, Dranga M
- 9406 Establishment of a risk assessment score for deep vein thrombosis after artificial liver support system treatment
Ye Y, Li X, Zhu L, Yang C, Tan YW

Retrospective Study

- 9417 Clinical management and susceptibility of primary hepatic lymphoma: A cases-based retrospective study
Hai T, Zou LQ
- 9431 Association of serum pepsinogen with degree of gastric mucosal atrophy in an asymptomatic population
Cai HL, Tong YL

- 9440** Risk factors for relapse and nomogram for relapse probability prediction in patients with minor ischemic stroke

Yu XF, Yin WW, Huang CJ, Yuan X, Xia Y, Zhang W, Zhou X, Sun ZW

- 9452** Incidence, prognosis, and risk factors of sepsis-induced cardiomyopathy

Liang YW, Zhu YF, Zhang R, Zhang M, Ye XL, Wei JR

- 9469** Associations with pancreatic exocrine insufficiency: An United Kingdom single-centre study

Shandro BM, Chen J, Ritehnia J, Poullis A

- 9481** Retrospective analysis of influencing factors on the efficacy of mechanical ventilation in severe and critical COVID-19 patients

Zeng J, Qi XX, Cai WW, Pan YP, Xie Y

Observational Study

- 9491** Vitamin D deficiency, functional status, and balance in older adults with osteoarthritis

Montemor CN, Fernandes MTP, Marquez AS, Poli-Frederico RC, da Silva RA, Fernandes KBP

- 9500** Psychological impact of the COVID-19 pandemic on Chinese population: An online survey

Shah T, Shah Z, Yasmeen N, Ma ZR

- 9509** Outcomes of different minimally invasive surgical treatments for vertebral compression fractures: An observational study

Yeh KL, Wu SH, Liaw CK, Hou SM, Wu SS

META-ANALYSIS

- 9520** Glycated albumin as a biomarker for diagnosis of diabetes mellitus: A systematic review and meta-analysis

Xiong JY, Wang JM, Zhao XL, Yang C, Jiang XS, Chen YM, Chen CQ, Li ZY

CASE REPORT

- 9535** Rapid response to radiotherapy in unresectable tracheal adenoid cystic carcinoma: A case report

Wu Q, Xu F

- 9542** Clinical observation of pediatric-type follicular lymphomas in adult: Two case reports

Liu Y, Xing H, Liu YP

- 9549** Malignant adenomyoepithelioma of the breast: Two case reports and review of the literature

Zhai DY, Zhen TT, Zhang XL, Luo J, Shi HJ, Shi YW, Shao N

- 9557** Validation of diagnostic strategies of autoimmune atrophic gastritis: A case report

Sun WJ, Ma Q, Liang RZ, Ran YM, Zhang L, Xiao J, Peng YM, Zhan B

- 9564** Characteristics of primary giant cell tumor in soft tissue on magnetic resonance imaging: A case report

Kang JY, Zhang K, Liu AL, Wang HL, Zhang LN, Liu WV

- 9571** Acute esophageal necrosis as a complication of diabetic ketoacidosis: A case report
Moss K, Mahmood T, Spaziani R
- 9577** Simultaneous embolization of a spontaneous porto-systemic shunt and intrahepatic arterioportal fistula: A case report
Liu GF, Wang XZ, Luo XF
- 9584** Ureteroscopic holmium laser to transect the greater omentum to remove an abdominal drain: Four case reports
Liu HM, Luo GH, Yang XF, Chu ZG, Ye T, Su ZY, Kai L, Yang XS, Wang Z
- 9592** Forearm compartment syndrome due to acquired hemophilia that required massive blood transfusions after fasciotomy: A case report
Kameda T, Yokota T, Ejiri S, Konno SI
- 9598** Transforaminal endoscopic excision of bi-segmental non-communicating spinal extradural arachnoid cysts: A case report and literature review
Yun ZH, Zhang J, Wu JP, Yu T, Liu QY
- 9607** T-cell lymphoblastic lymphoma with extensive thrombi and cardiac thrombosis: A case report and review of literature
Ma YY, Zhang QC, Tan X, Zhang X, Zhang C
- 9617** Perfect pair, scopes unite – laparoscopic-assisted transumbilical gastroscopy for gallbladder-preserving polypectomy: A case report
Zheng Q, Zhang G, Yu XH, Zhao ZF, Lu L, Han J, Zhang JZ, Zhang JK, Xiong Y
- 9623** Bilateral hematoma after tubeless percutaneous nephrolithotomy for unilateral horseshoe kidney stones: A case report
Zhou C, Yan ZJ, Cheng Y, Jiang JH
- 9629** Atypical endometrial hyperplasia in a 35-year-old woman: A case report and literature review
Wu X, Luo J, Wu F, Li N, Tang AQ, Li A, Tang XL, Chen M
- 9635** Clinical features and literature review related to the material differences in thread rhinoplasty: Two case reports
Lee DW, Ryu H, Jang SH, Kim JH
- 9645** Concurrent tuberculous transverse myelitis and asymptomatic neurosyphilis: A case report
Gu LY, Tian J, Yan YP
- 9652** Diagnostic value of contrast-enhanced ultrasonography in mediastinal leiomyosarcoma mimicking aortic hematoma: A case report and review of literature
Xie XJ, Jiang TA, Zhao QY
- 9662** Misidentification of hepatic tuberculosis as cholangiocarcinoma: A case report
Li W, Tang YF, Yang XF, Huang XY

- 9670** Brunner's gland hyperplasia associated with lipomatous pseudohypertrophy of the pancreas presenting with gastrointestinal bleeding: A case report
Nguyen LC, Vu KT, Vo TTT, Trinh CH, Do TD, Pham NTV, Pham TV, Nguyen TT, Nguyen HC, Byeon JS
- 9680** Metachronous squamous cell carcinoma of pancreas and stomach in an elderly female patient: A case report
Kim JH, Kang CD, Lee K, Lim KH
- 9686** Iatrogenic giant pseudomeningocele of the cervical spine: A case report
Kim KW, Cho JH
- 9691** Traditional Chinese medicine for gait disturbance in adrenoleukodystrophy: A case report and review of literature
Kim H, Kim T, Cho W, Chang H, Chung WS

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Takeo Furuya, MD, PhD, Assistant Professor, Department of Orthopaedic Surgery, Chiba University Graduate School of Medicine, Chiba 2608670, Japan. furuya-takeo@chiba-u.jp

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for *WJCC* as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The *WJCC*'s CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yan-Xia Xing; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Sandro Vento, Bao-Gan Peng

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

November 6, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Observational Study

Psychological impact of the COVID-19 pandemic on Chinese population: An online survey

Taif Shah, Zahir Shah, Nafeesa Yasmeen, Zhong-Ren Ma

ORCID number: Taif Shah 0000-0002-0217-2215; Zahir Shah 0000-0003-0368-4455; Nafeesa Yasmeen 0000-0002-7194-6026; Zhong-Ren Ma 0000-0002-3249-9083.

Author contributions: All the authors contributed to the concept of this study; Shah T, Shah Z and Ma ZR designed the study; Shah T and Yasmeen N acquired and analyzed data; Shah T, Shah Z and Yasmeen N wrote the manuscript; and all the authors reviewed and approved the final manuscript.

Institutional review board

statement: The Institutional Review Ethical Committee of Northwest Minzu University reviewed and approved the protocol used in this study.

Conflict-of-interest statement:

There are no conflicts of interest to report.

Data sharing statement: The aggregate data supporting the findings within this manuscript will be shared upon request submitted to the corresponding author.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in

Taif Shah, Faculty of Life Science and Technology, Kunming University of Science and Technology, Kunming 650500, Yunnan Province, China

Zahir Shah, College of Veterinary Sciences, The University of Agriculture Peshawar, Peshawar 25120, KP, Pakistan

Nafeesa Yasmeen, College of Veterinary Medicine, South China Agricultural University, Guangzhou 510642, Guangdong Province, China

Zhong-Ren Ma, Biomedical Research Center, Northwest Minzu University Gansu China, Lanzhou 730030, Gansu Province, China

Corresponding author: Zhong-Ren Ma, PhD, Professor, Biomedical Research Center, Northwest Minzu University, No. 1 Northwest New Village, Lanzhou 730030, Gansu Province, China. mzr@xbmu.edu.cn

Abstract

BACKGROUND

The ongoing coronavirus disease 2019 (COVID-19) pandemic infiltrates every aspect of our life, including the psychological impact. China has experienced the first wave of this epidemic, and it is now affecting the global population.

AIM

This study aimed to investigate the prevalence and associated factors of post-traumatic stress disorder (PTSD) among the general Chinese population.

METHODS

A detailed questionnaire, comprising of 38 questions designed in both English and Chinese, was developed. The survey was conducted *via* WeChat, a multi-purpose messaging, social media, and mobile payment app, which is widely used by the Chinese population.

RESULTS

In total, 1082 individuals from 31 provinces, autonomous regions, and municipalities participated in the survey by filling the questionnaires through the WeChat app. 97.8% of the participants had an Impact of Event-Scale-Revised (IES-R) total score above 20, which is an indicator of PTSD. The IES-R total and all the three subscales, including intrusion, avoidance, and hyperarousal, are significantly

accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Specialty type: Medicine, research and experimental

Country/Territory of origin: China

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B
Grade C (Good): 0
Grade D (Fair): D
Grade E (Poor): 0

Received: July 5, 2021

Peer-review started: July 5, 2021

First decision: July 26, 2021

Revised: July 28, 2021

Accepted: September 19, 2021

Article in press: September 19, 2021

Published online: November 6, 2021

P-Reviewer: Lozada-Martinez I, Nakaji K, Shrestha DB

S-Editor: Wang JJ

L-Editor: A

P-Editor: Yuan YY



correlated with age. In addition, age, profession, marital status, and level of education are significantly correlated with the degree of PTSD symptoms.

CONCLUSION

The COVID-19 epidemic has widely caused PTSD among the general Chinese population. These results bear important implications for regions struggling with the pandemic to implement effective interventions to cope with these mental health problems.

Key Words: COVID-19; Post-traumatic stress disorder; Impact of Event-Scale-Revised; China

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This study investigated the prevalence and associated factors of post-traumatic stress disorder (PTSD) in the general Chinese population. Overall, coronavirus disease 2019 has caused PTSD in the Chinese population. In addition, the severity of PTSD symptoms is related to age, profession, marital status, and level of education. These findings have significant implications for pandemic-affected regions in terms of implementing effective interventions to address these mental health issues.

Citation: Shah T, Shah Z, Yasmeen N, Ma ZR. Psychological impact of the COVID-19 pandemic on Chinese population: An online survey. *World J Clin Cases* 2021; 9(31): 9500-9508

URL: <https://www.wjgnet.com/2307-8960/full/v9/i31/9500.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v9.i31.9500>

INTRODUCTION

Outbreaks of infectious diseases have constantly threatened human beings, affecting many aspects of society. Compared to other diseases, infectious diseases are more prone to causing public panic[1], and long-term exposure to such events bears a high risk of developing post-traumatic stress disorder (PTSD)[2-4]. The ongoing coronavirus disease 2019 (COVID-19) pandemic infiltrates every aspect of our life and threatens the global population. Control measures are universally implemented to mitigate the pandemic, whereas psychological problems emerge in infected patients and the general population[5].

The COVID-19 outbreak was sparked in December 2019 in Wuhan, China[6] and quickly spread to the rest of China and now all over the world. By implementing strict control measures, including city lockdown, travel ban, and within-population quarantine, the epidemic in China was finally under control by March. This epidemic and the implementation of control measures, in particular quarantine, are expected to have adverse effects on the general population's mental health[7], but primary research in this respect remains limited. Therefore, this study assessed the prevalence and associated factors of PTSD among the general Chinese population who have just experienced the first wave of the COVID-19 epidemic.

MATERIALS AND METHODS

Study population and sample size estimation

Chinese from the general public were invited to participate in this study. The study was conducted from March 26th to March 31st, 2021. The sample size was determined by calculating a representative sample size using the Raosoft website. In 2019, the population of China was estimated to be 1433783686, with a median age of 37. We used a 3% margin of error and a 95% confidence interval to estimate the sample size. Therefore, a total of 1068 participants are needed.

Study design

A detailed questionnaire comprising of 38 questions was designed in both the English and Chinese languages. The survey was conducted *via* WeChat, a Chinese multi-purpose messaging, social media, and mobile payment app. It is widely used in China, with monthly active users estimated to be one billion. In total, 1082 individuals from 31 provinces, autonomous regions, and municipalities participated in the survey by filling the questionnaires (Supplementary Table 1).

The psychological impact was evaluated using a validated scale, *i.e.*, Impact of Event-Scale-Revised (IES-R)[8,9]. IES is a self-report measure used to assess the frequency of intrusive and avoidant phenomena in response to a specific stressful life event[10]. Thus, IES-R is a measurement tool for PTSD to measure subjective response to traumatic events in response to: (1) Intrusion (intrusive thoughts, nightmares, intrusive feelings and imagery, and re-experiencing); (2) Avoidance (numbing of responsiveness, avoidance of feelings, situations, and ideas); and (3) Hyperarousal (anger, irritability, hypervigilance, difficulty concentrating, heightened startle).

It comprises 22 items, each with a Likert rating scale from 0 to 4 (0 not at all; 1 a little bit; 2 moderately; 3 quite a bit; 4 extremely). The maximum score is 88. A higher score indicates a greater concern for PTSD. The IES-R has been translated into Chinese, demonstrated extensive reliability and validity, and is used frequently in trauma research worldwide[11,12].

Statistical analysis

Descriptive and inferential statistics were calculated using SPSS version 20.0 for Windows (SPSS Inc, Chicago IL). Means were calculated to summarize continuous variables. For categorical variables, group proportions were calculated. The one-way analysis of variance (ANOVA) and independent *t*-test were employed to identify the effect of demographic factors on the psychological outcome measure (IES-R). A total score of IES-R > 20 was used to estimate the prevalence of PTSD symptoms. ANOVA was used for age, level of education, profession, marital status, and monthly income, whilst *t*-test was used for gender, ethnicity, and residence. All statistical tests were two-sided and a *P* value < 0.05 was considered statistically significant.

RESULTS

Demographic characteristics

A total of 1082 individuals completed the online survey. Except for 0.3% of overseas Chinese, 99.7% of the respondents are from 31 provinces and autonomous regions and municipalities across China, and they have presumably been placed in quarantine or confined to their homes during the COVID-19 epidemic. Slightly more participants were female (57%). In terms of ethnicity, 67.8% are Hans, and 32.2% are from other ethnicities. 47.6% of the participants are aged younger than 30. 60.9% live in urban areas, and 39.1% are from rural backgrounds. The detailed socio-demographic characteristics of the participants are presented in Table 1.

Furthermore, 103 out of 1082 respondents have reported health problems such as diabetes, cardiovascular diseases, hypertension, chronic respiratory disease, viral hepatitis, or cancer. In addition, 44.4% of respondents are experiencing economic losses, and 6.3% are physically affected due to the COVID-19 epidemic. Only 40.57% of participants reported that they have returned to the normal routine after control of COVID-19 in China (Table 1).

Psychological impacts of the COVID-19 epidemic

The total IES-R scores are significantly correlated with age, level of education, profession, and marital status, but not with gender, ethnicity, residence, and monthly income. Next, we assessed the three subscales of IES-R. The intrusion subscale is significantly correlated with age and marital status. Avoidance is significantly correlated with age, level of education, profession, and marital status. Hyperarousal is correlated with age and level of education (Table 2).

The mean IES-R total score is 42.48 (SD = 12.13). A score of > 20 was used to estimate the prevalence of PTSD. 97.8% of the participants have an IES-R score above 20 (Table 3). The IES-R total scores are significantly associated with age, profession, marital status, and income, but not with gender, ethnicity, residence, or education. The groups with older age (above 50) have significantly higher IES-R scores. The group over 60 years has a mean score of 47.82, followed by the group 50-59 years with a

Table 1 Demographic characteristics of the participants (n = 1082)

Characteristics	Frequency	Percentage (%)
Gender		
Male	463	42.80
Female	619	57.20
Age (yr)		
< 30	515	47.60
31-39	262	24.20
40-49	157	14.50
50-59	131	12.10
> 60	17	1.60
Ethnicity		
Han	734	67.80
Others	348	32.20
Residence		
Rural	423	39.10
Urban	659	60.90
Education		
Illiterate	1	0.10
Primary	12	1.1
Middle	169	15.60
Bachelor	675	62.40
Master	120	11.10
PhD	105	9.70
Profession		
Farmer	40	3.70
Teacher	185	17.10
Doctor	78	7.20
Government	84	7.80
Private	28	2.60
Students	408	37.70
Business	20	1.80
Others	239	22.10
Marital Status		
Married	568	52.50
Divorced	25	2.30
Single	489	45.20
Monthly income (RMB)		
< 3000	496	45.80
3001-5000	251	23.20
5001-10000	235	21.70
> 10000	100	9.20
Positive responses to different questions		

Have you, your family members, or your friends affected by COVID-19?	5	0.46
Are you back to your routine after the control of COVID-19 epidemic?	439	40.57
COVID-19 affected you physically	68	6.3
COVID-19 affected you economically	480	44.4
Diabetes	13	1.2
Cardiovascular diseases	10	0.9
Hypertension	31	2.9
Chronic respiratory disease	4	0.4
Viral hepatitis	4	0.4
Cancer	1	0.1
Other comorbidities	40	3.7

COVID-19: Coronavirus disease 2019.

Table 2 Correlation between Impact of Event-Scale-Revised subscales and demographic characteristics

		Gender	Age	Ethnicity	Residence	Education	Profession	Marital	Income
IES-R total	Pearson <i>r</i>	-0.02	0.17	-0.02	-0.04	-0.09	-0.09	-0.09	0.02
	Sig (2-tailed)	0.47	0.01 ^a	0.53	0.17	0.05 ^a	0.01 ^a	0.01 ^a	0.54
Intrusion	Pearson <i>r</i>	-0.01	0.15	-0.02	-0.03	-0.04	-0.06	-0.10	0.04
	Sig (2-tailed)	0.82	0.01 ^a	0.50	0.34	0.17	0.06	0.01 ^a	0.22
Avoidance	Pearson <i>r</i>	-0.04	-0.20	0.01	-0.04	-0.10	-0.13	-0.08	0.03
	Sig (2-tailed)	0.23	0.01 ^a	0.88	0.16	0.01 ^a	0.01 ^a	0.01 ^a	0.29
Hyperarousal	Pearson <i>r</i>	-0.1	0.10	-0.04	-0.04	-0.08	-0.05	-0.05	-0.03
	Sig (2-tailed)	0.71	0.01 ^a	0.17	0.18	0.01 ^a	0.09	0.10	0.32

^a*P* < 0.05.

IES-R: Impact of Event-Scale-Revised.

mean value of 47.56. The IES-R scores vary tremendously (21-87) among different professions, and farmers, government officers, and people who work in private sectors have higher scores compared to other professions. Government employees have the highest IES-R scores (46.51 ± 12.02; mean ± SD; *P* = 0.001) (Table 3).

DISCUSSION

This study comprehensively assessed the prevalence and degree of PTSD among the general Chinese population affected by the COVID-19 epidemic by conducting an online survey. We further identified important factors associated with the development of PTSD. Mechanistically, we think both the devastation of the current pandemic and the implementation of stringent control measures significantly impact the general population's mental health. The negative consequences of the COVID-19 pandemic are penetrating every aspect of society, not only health, but also the economy, education, religion, and politics. These have immediate psychological effects on the general public. On the other hand, the implementation of heavy control measures, although desperately needed, has profound side effects on the population's physical and mental health. For example, psychological problems such as anxiety, panic disorder, and depression are frequently observed in people under quarantine or confined to their homes[7,13-15].

In this study, an IES-R score higher than 20 was used to estimate the prevalence of PTSD. The prevalence of PTSD is universal among the Chinese population, reaching 97.8%. The IES-R score is significantly higher among the older age groups. This

Table 3 Prevalence of post-traumatic stress symptoms according to participant demographics

Characteristics	Frequency (%) (n = 1082)			
	Range	mean \pm SD	t	P value
Prevalence				
IES-R < 20		24 \pm 2.2		
IES-R > 20		1058 \pm 97.8		
Gender			0.71	0.48
Male	18-88	42.78 \pm 12.48		
Female	18-87	42.25 \pm 11.87		
Ethnicity			0.63	0.53
Han	18-88	42.64 \pm 11.91		
Others	18-87	42.14 \pm 12.59		
Residence			1.37	0.17
Rural	19-87	43.11 \pm 12.52		
Urban	18-88	42.08 \pm 11.87		
Age (yr)			9.81	0.001 ^a
< 30	18-85	40.70 \pm 12.69		
31-39	21-88	43.08 \pm 11.99		
40-49	18-87	42.52 \pm 11.19		
50-59	20-74	47.56 \pm 9.36		
> 60	26-62	47.82 \pm 11.78		
Education			2.1	0.06
Illiterate	34-34	34		
Primary	31-62	44.75 \pm 8.84		
Middle	20-87	44.59 \pm 12.19		
Bachelor	18-88	42.34 \pm 12.23		
Master	18-85	42.28 \pm 11.86		
PhD	21-84	40.02 \pm 11.63		
Profession			4.63	0.001 ^a
Farmer	21-87	46.25 \pm 11.83		
Teacher	19-82	44.41 \pm 10.68		
Doctor	18-74	40.42 \pm 12.92		
Government	22-87	46.51 \pm 12.02		
Private	28-87	46.00 \pm 12.02		
Student	18-85	40.83 \pm 12.62		
Business	18-59	39.00 \pm 12.67		
Others	18-88	42.31 \pm 11.48		
Marital status			5.25	0.005 ^a
Married	18-88	43.42 \pm 11.63		
Divorced	24-63	45.76 \pm 11.51		
Single	18-85	41.23 \pm 12.62		
Monthly income (RMB)			6.29	0.001 ^a
< 3000	18-88	41.47 \pm 12.64		

3001-5000	19-87	44.26 ± 11.34
5001-10000	20-87	43.99 ± 11.48
> 10000	18-78	39.47 ± 12.04

^a*P* < 0.05.

IES-R: Impact of Event-Scale-Revised.

probably reflects that the Chinese populations are well-aware and have adequate epidemiological and clinical knowledge of COVID-19[16]. Because older people, especially when having comorbidities, are vulnerable and have a high COVID-19 fatality rate[17], government employees also experience significantly higher levels of distress compared to other professions. The Chinese central government has mobilized a strong force to contain the epidemic, and government workers have played essential roles. They are heavily involved in the implementation but are also responsible for the effectiveness and outcome of the control measures. Therefore, they face high risks of encountering COVID-19 and political reckoning, explaining their high degree of distress.

Respondents with lower education levels or lower income tend to have more distress. The possible explanation is that this population immediately suffers from economic losses and will face challenges in earning a livelihood. Compared with unmarried participants, the levels of psychological distress are significantly higher among married and divorced participants. Participants with family, especially children, to take care of are likely to worry more about the well-being of their family members.

Of note, this study has some limitations. Firstly, this study used self-reporting instead of diagnostic interviews for assessing psychiatric morbidity. Secondly, the use of online questionnaires has intrinsic limitations as recall, and social desirability biases cannot be excluded. Finally, in addition to PTSD, it would be beneficial to examine depressive disorder, adjustment disorder, psychosomatic disorder, substance use disorder, and antisocial behavior among the psychiatric problems associated with trauma and disasters[18,19].

CONCLUSION

In summary, the COVID-19 epidemic has universally caused PTSD among the general Chinese population. Furthermore, the degree of psychological distress is associated with age, profession, and socio-economic status. Therefore, our results alert the regions struggling with the pandemic to pay attention to the psychological impact and call on the authorities to implement effective interventions to cope with these mental health problems.

ARTICLE HIGHLIGHTS

Research background

The coronavirus disease 2019 (COVID-19) outbreak began in December 2019 in Wuhan, China, and quickly spread to the rest of China and, eventually, the rest of the world. China has been implementing strict control measures such as city lockdowns, travel bans, and within-population quarantine, and the epidemic has been brought under control by March. This epidemic and the implementation of control measures, particularly quarantine, are expected to impact the general population's mental health, but primary research in this area is limited.

Research motivation

COVID-19 has spread to become a pandemic and may become endemic. Unfortunately, knowledge gaps always exist about disease epidemics, potential risks, and the clinical spectrum.

Research objectives

This survey was designed and conducted to investigate the prevalence and associated risk factors of post-traumatic stress disorder (PTSD) among the general Chinese population.

Research methods

A detailed questionnaire, comprising of 38 questions written in both English and Chinese, was developed. The survey was conducted *via* WeChat, a multi-purpose messaging, social media, and mobile payment app widely used by the Chinese population. The 1082 people who participated in this survey belonged to 31 provinces, autonomous regions, and municipalities.

Research results

In total, 1082 people from 31 provinces, autonomous regions, and municipalities participated in the survey by filling questionnaires *via* the WeChat app. 97.8% of the participants had an Impact of Event-Scale-Revised (IES-R) total score above 20, which is the indicator of PTSD. The IES-R total and all three subscales, intrusion, avoidance, and hyperarousal, are significantly correlated with age. In addition, the degree of PTSD symptoms is correlated with age, profession, marital status, and level of education.

Research conclusions

We assessed epidemiological and clinical knowledge of COVID-19 among the general Chinese population and found that the epidemic has widely caused PTSD among the general Chinese population. These results have important implications for regions dealing with the pandemic to implement effective interventions to address these mental health issues.

Research perspectives

We emphasize the importance of launching health promotion programs to educate the general public and healthcare workers about infectious diseases in general to better prepare for future epidemics and pandemics.

ACKNOWLEDGEMENTS

We sincerely thank all the participants who took part in this study.

REFERENCES

- 1 **Pappas G**, Kiriaze IJ, Giannakis P, Falagas ME. Psychosocial consequences of infectious diseases. *Clin Microbiol Infect* 2009; **15**: 743-747 [PMID: 19754730 DOI: 10.1111/j.1469-0691.2009.02947.x]
- 2 **Marshall RD**, Bryant RA, Amsel L, Suh EJ, Cook JM, Neria Y. The psychology of ongoing threat: relative risk appraisal, the September 11 attacks, and terrorism-related fears. *Am Psychol* 2007; **62**: 304-316 [PMID: 17516775 DOI: 10.1037/0003-066x.62.4.304]
- 3 **Xiao S**, Luo D, Xiao Y. Survivors of COVID-19 are at high risk of posttraumatic stress disorder. *Glob Health Res Policy* 2020; **5**: 29 [PMID: 32514428 DOI: 10.1186/s41256-020-00155-2]
- 4 **Roberts T**, Miguel Esponda G, Krupchanka D, Shidhaye R, Patel V, Rathod S. Factors associated with health service utilisation for common mental disorders: a systematic review. *BMC Psychiatry* 2018; **18**: 262 [PMID: 30134869 DOI: 10.1186/s12888-018-1837-1]
- 5 **Xiang YT**, Yang Y, Li W, Zhang L, Zhang Q, Cheung T, Ng CH. Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *Lancet Psychiatry* 2020; **7**: 228-229 [PMID: 32032543 DOI: 10.1016/S2215-0366(20)30046-8]
- 6 **Zhu N**, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R, Niu P, Zhan F, Ma X, Wang D, Xu W, Wu G, Gao GF, Tan W; China Novel Coronavirus Investigating and Research Team. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* 2020; **382**: 727-733 [PMID: 31978945 DOI: 10.1056/NEJMoa2001017]
- 7 **Brooks SK**, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, Rubin GJ. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020; **395**: 912-920 [PMID: 32112714 DOI: 10.1016/S0140-6736(20)30460-8]
- 8 **Beck JG**, Grant DM, Read JP, Clapp JD, Coffey SF, Miller LM, Palyo SA. The impact of event scale-revised: psychometric properties in a sample of motor vehicle accident survivors. *J Anxiety Disord* 2008; **22**: 187-198 [PMID: 17369016 DOI: 10.1016/j.janxdis.2007.02.007]

- 9 **Wilson JP**, Tang CS-k. Cross-cultural assessment of psychological trauma and PTSD. New York: Springer, 2007: 405 [DOI: [10.1007/978-0-387-70990-1](https://doi.org/10.1007/978-0-387-70990-1)]
- 10 **Salsman JM**, Schalet BD, Andrykowski MA, Cella D. The impact of events scale: a comparison of frequency versus severity approaches to measuring cancer-specific distress. *Psychooncology* 2015; **24**: 1738-1745 [PMID: [25773193](https://pubmed.ncbi.nlm.nih.gov/25773193/) DOI: [10.1002/pon.3784](https://doi.org/10.1002/pon.3784)]
- 11 **Chong MY**, Wang WC, Hsieh WC, Lee CY, Chiu NM, Yeh WC, Huang OL, Wen JK, Chen CL. Psychological impact of severe acute respiratory syndrome on health workers in a tertiary hospital. *Br J Psychiatry* 2004; **185**: 127-133 [PMID: [15286063](https://pubmed.ncbi.nlm.nih.gov/15286063/) DOI: [10.1192/bjp.185.2.127](https://doi.org/10.1192/bjp.185.2.127)]
- 12 **Hsu CC**, Chong MY, Yang P, Yen CF. Posttraumatic stress disorder among adolescent earthquake victims in Taiwan. *J Am Acad Child Adolesc Psychiatry* 2002; **41**: 875-881 [PMID: [12108814](https://pubmed.ncbi.nlm.nih.gov/12108814/) DOI: [10.1097/00004583-200207000-00022](https://doi.org/10.1097/00004583-200207000-00022)]
- 13 **Qiu J**, Shen B, Zhao M, Wang Z, Xie B, Xu Y. A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy recommendations. *Gen Psychiatr* 2020; **33**: e100213 [PMID: [32215365](https://pubmed.ncbi.nlm.nih.gov/32215365/) DOI: [10.1136/gpsych-2020-100213](https://doi.org/10.1136/gpsych-2020-100213)]
- 14 **Ma Z**, Idris S, Zhang Y, Zewen L, Wali A, Ji Y, Pan Q, Baloch Z. The impact of COVID-19 pandemic outbreak on education and mental health of Chinese children aged 7-15 years: an online survey. *BMC Pediatr* 2021; **21**: 95 [PMID: [33627089](https://pubmed.ncbi.nlm.nih.gov/33627089/) DOI: [10.1186/s12887-021-02550-1](https://doi.org/10.1186/s12887-021-02550-1)]
- 15 **Ma ZR**, Idris S, Pan QW, Baloch Z. COVID-19 knowledge, risk perception, and information sources among Chinese population. *World J Psychiatry* 2021; **11**: 181-200 [PMID: [34046314](https://pubmed.ncbi.nlm.nih.gov/34046314/) DOI: [10.5498/wjp.v11.i5.181](https://doi.org/10.5498/wjp.v11.i5.181)]
- 16 **Zhong BL**, Luo W, Li HM, Zhang QQ, Liu XG, Li WT, Li Y. Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey. *Int J Biol Sci* 2020; **16**: 1745-1752 [PMID: [32226294](https://pubmed.ncbi.nlm.nih.gov/32226294/) DOI: [10.7150/ijbs.45221](https://doi.org/10.7150/ijbs.45221)]
- 17 **Verity R**, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, Cuomo-Dannenburg G, Thompson H, Walker PGT, Fu H, Dighe A, Griffin JT, Baguelin M, Bhatia S, Boonyasiri A, Cori A, Cucunubá Z, FitzJohn R, Gaythorpe K, Green W, Hamlet A, Hinsley W, Laydon D, Nedjati-Gilani G, Riley S, van Elsland S, Volz E, Wang H, Wang Y, Xi X, Donnelly CA, Ghani AC, Ferguson NM. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis* 2020; **20**: 669-677 [PMID: [32240634](https://pubmed.ncbi.nlm.nih.gov/32240634/) DOI: [10.1016/S1473-3099\(20\)30243-7](https://doi.org/10.1016/S1473-3099(20)30243-7)]
- 18 **McFarlane AC**. Posttraumatic stress disorder: a model of the longitudinal course and the role of risk factors. *J Clin Psychiatry* 2000; **61** Suppl 5: 15-20; discussion 21-23 [PMID: [10761675](https://pubmed.ncbi.nlm.nih.gov/10761675/) DOI: [10.1007/978-981-10-4358-1_10](https://doi.org/10.1007/978-981-10-4358-1_10)]
- 19 **Streb M**, Conway MA, Michael T. Conditioned responses to trauma reminders: How durable are they over time and does memory integration reduce them? *J Behav Ther Exp Psychiatry* 2017; **57**: 88-95 [PMID: [28477531](https://pubmed.ncbi.nlm.nih.gov/28477531/) DOI: [10.1016/j.jbtep.2017.04.005](https://doi.org/10.1016/j.jbtep.2017.04.005)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

