

World Journal of *Clinical Cases*

World J Clin Cases 2022 February 26; 10(6): 1754-2052



Contents

Thrice Monthly Volume 10 Number 6 February 26, 2022

OPINION REVIEW

- 1754 Gut-brain axis: Focus on gut metabolites short-chain fatty acids
Guo C, Huo YJ, Li Y, Han Y, Zhou D

MINIREVIEWS

- 1764 Association between direct-acting antiviral agents in hepatitis C virus treatment and hepatocellular carcinoma occurrence and recurrence: The endless debate
Kamal A, Elsheaita A, Abdelnabi M

ORIGINAL ARTICLE

Retrospective Cohort Study

- 1775 Effects of bilirubin on perioperative myocardial infarction and its long-term prognosis in patients undergoing percutaneous coronary intervention
Li Y, Li DB, Zhao LD, Lv QB, Wang Y, Ren YF, Zhang WB
- 1787 Disease exacerbation is common in inflammatory bowel disease patients treated with immune checkpoint inhibitors for malignancy
Rubin SJS, Balabanis T, Gubatan J, Habtezion A
- 1795 Multidrug-resistant organisms in intensive care units and logistic analysis of risk factors
Han Y, Zhang J, Zhang HZ, Zhang XY, Wang YM

Retrospective Study

- 1806 Change and impact of left ventricular global longitudinal strain during transcatheter aortic valve implantation
Zhang H, Xie JJ, Li RJ, Wang YL, Niu BR, Song L, Li J, Yang Y

Observational Study

- 1815 Early detection of noise-induced hearing loss
Meng ZL, Chen F, Zhao F, Gu HL, Zheng Y
- 1826 Empathetic nursing with mindful cognitive therapy for fatigue, depression, and negative emotions in leukemia patients undergoing long-term chemotherapy
Lu YY, Lu XM, Shao CY, Wang CC, Xu TT, Zhang BL

Prospective Study

- 1834 Superior pancreatic lymphadenectomy with portal vein priority *via* posterior common hepatic artery approach in laparoscopic radical gastrectomy
Zhang YJ, Xiang RC, Li J, Liu Y, Xie SM, An L, Li HL, Mai G

Randomized Controlled Trial

- 1843** Systematic nursing interventions in gastric cancer: A randomized controlled study
He F, He RX

META-ANALYSIS

- 1852** Impact of adding opioids to paravertebral blocks in breast cancer surgery patients: A systematic review and meta-analysis
Chen MH, Chen Z, Zhao D

CASE REPORT

- 1863** Multiple different remote epidural hematomas after craniotomy: A case report
He Q, Tao CY, Fu RH, You C
- 1869** Tuberculous pericarditis-a silent and challenging disease: A case report
Lucero OD, Bustos MM, Ariza Rodríguez DJ, Perez JC
- 1876** Transileocolic endovascular treatment by a hybrid approach for severe acute portal vein thrombosis with bowel necrosis: Two case reports
Shirai S, Ueda T, Sugihara F, Yasui D, Saito H, Furuki H, Kim S, Yoshida H, Yokobori S, Hayashi H, Kumita SI
- 1883** Efficacy of EGFR-TKI sequential therapy in patients with EGFR exon 19 insertion-positive non-small-cell lung cancer: A case report
Shan BB, Li Y, Zhao C, An XQ, Zhang QM
- 1889** Novel compound heterozygous variants in the TAF6 gene in a patient with Alazami-Yuan syndrome: A case report
Lin SZ, Feng JH, Sun LP, Ma HW, Wang WQ, Li JY
- 1896** Asymmetric limb weakness in Guillain-Barré syndrome: Three case reports
Hu M, Li X, Wong HY, Feng XG, Wang YZ, Zhang GR
- 1903** Modified treatment of knee osteoarthritis complicated with femoral varus deformity: A case report
Xu SM, Li W, Zhang DB, Bi HY, Gu GS
- 1909** Novel HNF1A gene mutation in maturity-onset diabetes of the young: A case report
Xu Q, Kan CX, Hou NN, Sun XD
- 1914** Cerebral corridor creator for resection of trigone ventricular tumors: Two case reports
Liu XW, Lu WR, Zhang TY, Hou XS, Fa ZQ, Zhang SZ
- 1922** Left abdominal wall proliferative myositis resection and patch repair: A case report
Xing RW, Nie HQ, Zhou XF, Zhang FF, Mou YH
- 1929** Concurrent ankylosing spondylitis and myelodysplastic syndrome: A case report
Xu GH, Lin J, Chen WQ

- 1937** Life-threatening subclavian artery bleeding following percutaneous coronary intervention with stent implantation: A case report and review of literature
Shi F, Zhang Y, Sun LX, Long S
- 1946** Cryptogenic organizing pneumonia associated with pregnancy: A case report
Lee YJ, Kim YS
- 1952** Eosinophilia complicated with venous thromboembolism: A case report
Su WQ, Fu YZ, Liu SY, Cao MJ, Xue YB, Suo FF, Liu WC
- 1961** Neck and mediastinal hematoma caused by a foreign body in the esophagus with diagnostic difficulties: A case report
Wang LP, Zhou ZY, Huang XP, Bai YJ, Shi HX, Sheng D
- 1966** Therapeutic endoscopy of a Dieulafoy lesion in a 10-year-old girl: A case report
Chen Y, Sun M, Teng X
- 1973** Cavernous hemangioma of an intrapancreatic accessory spleen mimicking a pancreatic tumor: A case report
Huang JY, Yang R, Li JW, Lu Q, Luo Y
- 1981** Surgery and antibiotics for the treatment of lupus nephritis with cerebral abscesses: A case report
Hu QD, Liao LS, Zhang Y, Zhang Q, Liu J
- 1991** Median arcuate ligamentum syndrome: Four case reports
Kim JE, Rhee PL
- 1998** Novel ABCB4 mutations in an infertile female with progressive familial intrahepatic cholestasis type 3: A case report
Liu TF, He JJ, Wang L, Zhang LY
- 2007** Primary duodenal dedifferentiated liposarcoma: A case report and literature review
Kim NI, Lee JS, Choi C, Nam JH, Choi YD, Kim HJ, Kim SS
- 2015** Implant site development using titanium plate and platelet-rich fibrin for congenitally missed maxillary lateral incisors: A case report
Zhang TS, Mudalal M, Ren SC, Zhou YM
- 2023** Successful embolization of an intrahepatic portosystemic shunt using balloon-occluded retrograde transvenous obliteration: A case report
Saito H, Murata S, Sugihara F, Ueda T, Yasui D, Miki I, Hayashi H, Kumita SI
- 2030** Bilateral pneumothorax and pneumomediastinum during colonoscopy in a patient with intestinal Behcet's disease: A case report
Mu T, Feng H
- 2036** Acute kidney injury due to intravenous detergent poisoning: A case report
Park S, Ryu HS, Lee JK, Park SS, Kwon SJ, Hwang WM, Yun SR, Park MH, Park Y

- 2045** Vaginal enterocele after cystectomy: A case report

Liu SH, Zhang YH, Niu HT, Tian DX, Qin F, Jiao W

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Navdeep Singh, MBBS, MS, Assistant Professor, Division of Transplantation, Department of Surgery, The Ohio State University, Columbus, OH 43210, United States.
navdeep.singh@osumc.edu

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJCC as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The WJCC's CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*, Production Department Director: *Xu Guo*, Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

February 26, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Impact of adding opioids to paravertebral blocks in breast cancer surgery patients: A systematic review and meta-analysis

Meng-Hua Chen, Zheng Chen, Da Zhao

Specialty type: Surgery

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Lepot A, Valencia GA

Received: July 19, 2021

Peer-review started: July 19, 2021

First decision: October 16, 2021

Revised: October 21, 2021

Accepted: January 19, 2022

Article in press: January 19, 2022

Published online: February 26, 2022



Meng-Hua Chen, Lanzhou University Medical College, Lanzhou 730000, Gansu Province, China

Zheng Chen, Department of Breast, Shandong Second Provincial General Hospital, Jinan 250021, Shandong Province, China

Da Zhao, Department of Oncology, The First Hospital of Lanzhou University, Lanzhou 730000, Gansu Province, China

Corresponding author: Da Zhao, MD, Doctor, Department of Oncology, The First Hospital of Lanzhou University, No. 1 Donggang West Road, Lanzhou 730000, Gansu Province, China. zhaoda_dr@163.com

Abstract

BACKGROUND

Several breast cancer studies have reported the use of adjuvant opioids with the paravertebral block (PVB) to improve outcomes. However, there is no level-1 evidence justifying its use.

AIM

To elucidate if the addition of opioids to PVB improves pain control in breast cancer surgery patients.

METHODS

We conducted an electronic literature search across PubMed, Embase, Scopus, and Google Scholar databases up to October 20, 2020. Only randomized controlled trials (RCTs) comparing the addition of opioids to PVB with placebo for breast cancer surgery patients were included.

RESULTS

Six RCTs were included. Our meta-analysis indicated significantly reduced 24-h total analgesic consumption with the addition of opioids to PVB as compared to placebo [standardized mean difference (SMD) -1.57, 95% confidence interval (CI): -2.93, -0.21, $I^2 = 94\%$]. However, on subgroup analysis, the results were non-significant for studies using single PVB (SMD: -1.76, 95%CI: -3.65, 0.13 $I^2 = 95.09\%$) and studies using PVB infusion (SMD: -1.30, 95%CI: -4.26, 1.65, $I^2 = 95.49\%$). Analysis of single PVB studies indicated no significant difference in the time to first analgesic request between opioid and placebo groups (mean difference -11.28, 95%CI: -42.00, 19.43, $I^2 = 99.39\%$). Pain scores at 24 h were marginally lower

in the opioid group (mean difference -1.10, 95% CI: -2.20, 0.00, $P = 0\%$). There was no difference in the incidence of postoperative nausea and vomiting between the two groups.

CONCLUSION

Current evidence suggests a limited role of adjuvant opioids with PVB for breast cancer surgery patients. Further homogenous RCTs with a large sample size are needed to clarify the beneficial role of opioids with PVB.

Key Words: Opioids; Pain; Surgery; Breast cancer; Nerve block; Paravertebral block

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The use of opioids as adjuvants for nerve blocks has been increasing. However, it is not clear if the addition of opioids to paravertebral block (PVB) improves outcomes for breast cancer surgery patients. In our first systematic review and meta-analysis, we pooled data from six randomized controlled trials assessing the role of adjuvant opioids for PVB in patients undergoing breast cancer surgery. Our results indicate that the addition of opioids to PVB has a limited role in reducing 24-h analgesic consumption, time to first analgesic request, and pain scores as compared to placebo. Further high-quality studies are required to strengthen the evidence.

Citation: Chen MH, Chen Z, Zhao D. Impact of adding opioids to paravertebral blocks in breast cancer surgery patients: A systematic review and meta-analysis. *World J Clin Cases* 2022; 10(6): 1852-1862

URL: <https://www.wjgnet.com/2307-8960/full/v10/i6/1852.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i6.1852>

INTRODUCTION

Breast cancer is the most common malignancy in females worldwide, and surgical intervention is the primary mode of management even in advanced cases (basically palliative in selected populations)[1]. Studies have shown improved survival with mastectomy in patients with breast cancer[1,2]. While general anesthesia is the standard-setting used for surgical interventions for these patients, a substantial number of individuals encounter significant postoperative pain[3,4]. Inadequate analgesia in the immediate postoperative period can lead to a prolonged hospital stay, increased healthcare cost, and reduced patient satisfaction[5]. Optimal management of acute pain in breast cancer survivors can also reduce the development of chronic post-surgical pain[6].

Over the last decade, several quality-improvement protocols have been described to improve peri-operative management and optimize pain control in breast cancer patients[7]. One such method is the use of locoregional anesthetic techniques like the paravertebral block (PVB), intercostal nerve block, erector spinae plane block, and pectoral block[5,8,9]. Of these, the PVB has been widely used to provide better analgesia after surgery in breast cancer patients. The clinical efficacy of PVB has also been demonstrated by several studies[10,11]. Terkawi *et al*[12] in a meta-analysis of 24 studies demonstrated that the use of PVB decreased opioid consumption and postoperative pain scores at 2, 24, 48, and 72 h after surgery. However, the effects of PVB were found to be modest with a limited beneficial effect on postoperative recovery. In this context, several researchers have evaluated the addition of adjuvants to PVB to improve its efficacy. It is hypothesized that the addition of drugs like opioids, clonidine, and dexmedetomidine would lead to better analgesic efficacy of PVB[13,14]. Opioids have been used in combination with local anesthetics for several locoregional anesthetic techniques leading to better pain control in the immediate postoperative period[15,16]. However, it is not known if the addition of opioids to PVB would lead to better outcomes in breast cancer patients. Despite several studies reporting the use of adjuvant opioids with PVB, there is a lack of pooled evidence to guide clinical practice. Thus, this systematic review and meta-analysis aimed to answer the following clinical question: Does the addition of opioids to PVB lead to improved pain control in the immediate postoperative period in patients undergoing surgery for breast cancer?

MATERIALS AND METHODS

Search strategy

The authors planned and executed this study conforming to the recommendations of the PRISMA

statement (Preferred Reporting Items for Systematic Reviews and Meta-analyses)[17] and the Cochrane Handbook for Systematic Reviews of Intervention[18]. Protocol registration was, however, not carried out. We conducted an electronic literature search across PubMed, Embase, Scopus, and Google Scholar databases. Two reviewers independently carried out the literature search. Search limits were from the inception of databases to October 20, 2020. The search terms included “breast surgery”, “mastectomy”, “paravertebral block”, “opioid”, “morphine”, “fentanyl”, “buprenorphine”, and “tramadol”. **Supplementary Table 1** presents the search strategy and the result of the PubMed database. At first, the search records were reviewed by their titles and abstracts. Relevant articles to the review were identified, and full texts of the articles were extracted. Both the reviewers assessed individual articles based on the inclusion and exclusion criteria. Any disagreements were resolved by discussion. The bibliography of studies meeting the inclusion criteria was also hand-searched for any missed references.

Inclusion criteria

We defined the inclusion and exclusion criteria of the review based on the PICOS (Population, Intervention, Comparison, Outcome, Study type) framework *a priori*. Population: Studies conducted on patients undergoing breast cancer surgery and receiving PVB before general anesthesia. The Intervention was to be the addition of an opioid to the PVB. The comparison was the addition of placebo or no drug to the PVB. Studies were to report at least one of the following outcomes: 24 h total analgesic consumption, pain scores, time to the first analgesic, and/or incidence of Postoperative nausea and vomiting (PONV). Only randomized controlled trials (RCTs) were eligible to be included in the review. No language restriction was placed. Studies comparing opioids with any other active drugs were excluded. We also excluded studies using opioids not as an addition to PVB but *via* other routes like intravenous, subcutaneous, *etc.* Furthermore, non-RCTs, retrospective studies, single-arm studies, and studies not reporting relevant data were also excluded.

Data extraction and quality assessment

Data were extracted using a data extraction form by two reviewers independently. Name of the first author, publication year, study type, study location, age of patients, surgery type, sample size, intervention drug and dose, PVB protocol, use of other analgesics, and study outcomes were extracted. The primary outcome of the interest of our analysis was 24-h total analgesic consumption. The secondary outcomes were time to first analgesic demand, pain scores, and incidence of PONV. Furthermore, a descriptive analysis of other outcomes reported by the included studies was also performed.

Two reviewers assessed the quality of each RCT using the Cochrane Collaboration risk assessment tool[18]. Each study was assessed for bias in random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting, and other bias. The study was judged to have a “high”, “unclear”, or “low” risk of bias for each domain. Any disagreements were resolved by discussion.

Statistical analysis

The software “Open MetaAnalyst” was used for the meta-analysis[19]. Meta-analysis was conducted only if at least three trials reported similar outcomes. Owing to the methodological heterogeneity of the included studies, we used a random-effects model to calculate the pooled effect size for all analyses. Continuous data not reported on the same scale were summarized using standardized mean difference (SMD) with 95% confidence interval (CI) or else mean difference (MD) was used. Specifically, different analgesics were used by the individual studies for the outcome of ‘total analgesic consumption’, hence we used SMD to pool this variable. For studies not reporting mean and standard deviation (SD) scores of continuous variables, the method described by Wan *et al*[20] was used to calculate data from the median and interquartile range. For the incidence of PONV, we calculated odds ratios (OR) with 95% CI. Sub-group analysis was conducted for single and continuous PVB. Heterogeneity was assessed using the *I*² statistic. *I*² values of 25%-50% represented low, values of 50%-75% represented medium, and more than 75% represented substantial heterogeneity. As < 10 studies were included per meta-analysis, funnel plots were not used to assess publication bias.

RESULTS

The PRISMA flowchart of the review is presented in **Figure 1**. Of the 11 studies assessed by the full-texts, five were excluded with reasons, and a total of six RCTs fulfilled the inclusion criteria[21-26]. Characteristics of studies included in the review are presented in **Table 1**. Three trials[22,23,25] were conducted exclusively on modified radical mastectomy patients, while the remaining three included other breast cancer surgeries as well. The sample size of the included studies was small, ranging from 12-20 patients per group. The opioids used as an adjunct to PVB were fentanyl in three trials, while morphine, tramadol, and buprenorphine were used in one study each. Two trials[22,26] used continuous PVB infusions, while the remaining used single PVB.

Table 1 Details of included studies

Ref.	Yr	Study location	Surgery type	Sample size		Age (yr)	Common protocol for PVB	Opioid added to PVB in intervention group)	Drug added to PVB in control group	Post-operative analgesics used
				Study	Control					
Mostafa <i>et al</i> [21]	2018	Egypt	MRM or breast conservation surgery with axillary node dissection	20	20	18-78	Nalbuphine 10 mg	PVB at the level of T4 with bupivacaine 0.5% 0.3 mL/kg	No drug	Tramadol as PCA
Pushparajan <i>et al</i> [22]	2017	India	MRM	20	20	18-60	Fentanyl 2 µg/mL at 0.1 mL/kg/h for 24 h	Continuous PVB at the level of T4 with 0.2% ropivacaine for 24 h	No drug	Fentanyl as PCA. Paracetamol or tramadol or fentanyl for breakthrough pain
Morsy <i>et al</i> [23]	2017	Egypt	MRM	15	15	NR	Morphine 2 mg	PVB at the level of T3 with 20 mL of bupivacaine 0.25%	No drug	Meperidine for breakthrough pain
Bhuvaneshwari <i>et al</i> [24]	2012	India	Total mastectomy and axillary lymph node dissection	12	12	Study: 49.1 ± 7.1; Control: 50.7 ± 11	Fentanyl 2 µg/mL	PVB at the level of T3 with bupivacaine 0.25% and epinephrine 5 µg/mL	No drug	Morphine for breakthrough pain
Omar <i>et al</i> [25]	2011	Egypt	MRM	19	20	Study: 47.5 ± 9.3; Control: 49.3 ± 10.5	Tramadol 1.5 mg/kg (maximum of 150 mg)	PVB at the level of T1 (1/3 rd of the dose) and T4 (2/3 rd of the dose) with bupivacaine 0.5% 2 mg/kg	No drug	Fentanyl as PCA. Paracetamol 1 g thrice daily and ibuprofen 400-600 mg thrice daily
Burlacu <i>et al</i> [26]	2006	Ireland	Wide local excisions (at least one breast quadrant), mastectomies, and mastectomies with reconstruction	13	13	Study: 54 ± NR; Control: 51 ± NR	Fentanyl 50 µg with bolus followed by 4 µg/mL infusion	Continuous PVB at the level of T3 with initial bolus of 19 mL levobupivacaine 0.25% followed by continuous infusion of 0.1% solution for 24 h	Saline	Morphine as PCA

PVB: Paravertebral block; MRM: Modified radical mastectomy; PCA: Patient controlled analgesia; T: Thoracic vertebral level.

Outcomes

Table 2 presents the results of the outcomes reported by the included studies. For the primary outcome, data were reported by all six studies. Our meta-analysis indicated significantly reduced 24-h total analgesic consumption with the addition of opioids to PVB as compared to placebo (SMD: -1.57, 95%CI: -2.93, -0.21, $I^2 = 94\%$) (Figure 2). However, on subgroup analysis, the results were non-significant for studies using single PVB (SMD: -1.76, 95%CI: -3.65, 0.13, $I^2 = 95.09\%$) and studies using PVB infusion (SMD: -1.30, 95%CI: -4.26, 1.65, $I^2 = 95.49\%$). Data on time to first analgesic request were reported by four studies using a single PVB. Our analysis demonstrated no significant difference in the time to first analgesic request between opioid and placebo groups in hours (MD -11.28, 95%CI: -42.00, 19.43, $I^2 = 99.39\%$) (Figure 3).

Data for a meta-analysis on pain scores on the Visual Analog Scale were available only from three studies. The study of Bhuvaneshwari *et al*[24] reported cumulative 24-h pain scores, while SD values of pain scores were not reported by Mostafa *et al*[21]. Both these studies reported significantly lower pain scores in the opioid group. For the remaining studies, our pooled analysis indicated pain scores at 24 h were marginally lower in the opioid group (MD -1.10, 95%CI: -2.20, 0.00, $I^2 = 0\%$) (Figure 4). Sub-group

Table 2 Outcomes reported by included studies

Ref.	Outcome	Results
Mostafa <i>et al</i> [21]	Time to first analgesic request	Significantly longer in the opioid group
	Post-operative analgesic time	Significantly longer in the opioid group
	24 h total analgesic consumption	Significantly lower in the opioid group
	Pain scores up to 24 h	Significantly lower in the opioid group
	HR, SBP, DBP	No difference between the two groups
	Ramsay sedation scores	Patients in the control group were more agitated then opioid group in the first four hours after the operation. No difference between the two groups after four hours
Pushparajan <i>et al</i> [22]	Pain scores up to discharge	Significantly lower scores in the opioid group only at 24 h and not at other time periods
	24 h total analgesic consumption	No difference between the two groups
	PONV	No difference between the two groups
	Urinary retention, pruritis	No difference between the two groups
	Patient satisfaction	No difference between the two groups
Morsy <i>et al</i> [23]	24 h total analgesic consumption	No difference between the two groups
	Time to first analgesic request	No difference between the two groups
	Ramsay sedation scores	No difference between the two groups
	PONV	No difference between the two groups
	HR, SBP, DBP	No difference between the two groups
Bhuvaneshwari <i>et al</i> [24]	Time to first analgesic request	Significantly longer in the opioid group
	24 h total analgesic consumption	Significantly lower in the opioid group
	Cumulative pain scores at 24 h	Significantly lower in the opioid group
	PONV	No difference between the two groups
	Patient satisfaction	Significantly higher in the opioid group
Omar <i>et al</i> [25]	Time to first analgesic request	No difference between the two groups
	24 h total analgesic consumption	No difference between the two groups
	Pain scores up to 24 h	No difference between the two groups
	PONV	No difference between the two groups
Burlacu <i>et al</i> [26]	Total analgesic consumption	Significantly lower in the opioid group
	Pain scores up to 24 h	No difference between the two groups
	Nausea scores	Significantly higher in the opioid group
	SBP	No difference between the two groups
	Patient satisfaction	Significantly higher in the opioid group

PONV: Postoperative nausea and vomiting; HR: Heart rate; SBP: Systolic blood pressure; DBP: Diastolic blood pressure.




Figure 1 Study flow chart.




Figure 2 Forest plot of 24-h total analgesic consumption with sub-group analysis based on type of paravertebral block.



analysis demonstrated marginal difference favoring the opioid group in the studies using PVB infusion (MD -1.30, 95%CI: -2.61, 0.00, $I^2 = 0\%$); however, for the study using single PVB no difference was noted (MD -0.60, 95%CI: -2.65, 1.45). Data on the incidence of PONV were reported by four studies. Our analysis indicated no statistically significant difference between opioid and placebo groups (OR 0.87, 95%CI: 0.39, 1.93, $I^2 = 0\%$) (Figure 5). Results were non-significant on subgroup analysis as well.

Risk of bias analysis

Table 3 presents the risk of bias assessment of included studies. The majority of studies (5/6) were of good quality with a low risk of bias across six of the seven domains. Only the trial of Morsy *et al*[23] did not provide adequate information on randomization, allocation concealment, and blinding.

Table 3 Risk of bias in included studies

Ref.	Random sequence generation	Allocation concealment	Blinding of participants and personnel	Blinding of outcome assessment	Incomplete outcome data	Selective reporting	Other bias
Mostafa <i>et al</i> [21]	Low risk	Low risk	Low risk	Low risk	Low risk	High risk	Low risk
Pushparajan <i>et al</i> [22]	Low risk	Unclear risk	Low risk	Low risk	Low risk	Low risk	Low risk
Morsy <i>et al</i> [23]	Unclear risk	Unclear risk	Unclear risk	Unclear risk	Low risk	Low risk	Low risk
Bhuvaneshwari <i>et al</i> [24]	Low risk	Unclear risk	Low risk	Low risk	Low risk	Low risk	Low risk
Omar <i>et al</i> [25]	Low risk	Unclear risk	Low risk	Low risk	Low risk	Low risk	Low risk
Burlacu <i>et al</i> [26]	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk

**Figure 3 Forest plot of time to first analgesic request in hours.****Figure 4 Forest plot of 24-h visual analog scale pain score with sub-group analysis based on type of paravertebral block.**

DISCUSSION

The results of our review assessing the role of adjuvant opioids to PVB for breast cancer surgery patients indicate that: (1) Total analgesic consumption in the immediate postoperative period may not be reduced with the addition of opioids to a single PVB or PVB infusion; (2) Time to first analgesic request is not increased with the addition of opioids to a single PVB; (3) There is only a marginal difference in pain score at 24 h with the addition of opioids; and (4) Adjuvant opioids do not increase the incidence of PONV.

PVB as a technique of regional anesthesia has gained popularity for pain control in patients undergoing breast surgery. The procedure involves the deposition of local anesthetic just lateral to the spinous process of the vertebrae in the area where spinal nerves emerge out of the intervertebral foramina. The local anesthetic blocks the somatic and sympathetic nerve supply of the dermatome of interest thereby providing postsurgical analgesia[27]. Its efficacy has been tested in a randomized setting by numerous authors not only for breast surgery but also for thoracic surgeries[12,28]. However, one important limitation of nerve blocks, in general, is the short duration of action of the local anesthetic. Even with agents with a longer duration of action, like bupivacaine and ropivacaine, the duration of analgesia is often inadequately sustained in the postoperative period[29]. Prolonging the




Figure 5 Forest plot of incidence of postoperative nausea and vomiting with sub-group analysis based on type of paravertebral block.

duration of analgesia by increasing the dose of local anesthetic entails the risk of adverse events involving the cardiovascular and central nervous systems[30]. To overcome this issue, several adjuvants have been used with local anesthetics to prolong the analgesic effect while maintaining the safety of regional anesthesia.

One of the earliest adjuvants used with local anesthetics was opioids. Their use with local anesthetics has accelerated since studies reported the presence of peripheral opioid receptors in the primary afferent neurons and peripheral sensory nerves[31]. It is thought that opioids exhibit a local anesthetic-like action causing hyperpolarization of the afferent sensory neuron through G protein-coupled receptor mechanism[32]. Nishikawa *et al*[33] have suggested that adjuvant opioids with local anesthetics can lead to the increased duration and improved quality of local anesthetic blockade. However, the results of our review indicate that this mechanism may have a limited role especially for PVB in breast cancer surgery patients. For the primary outcome of 24-h total analgesic consumption, our analysis demonstrated a statistically significant difference in favor of the opioid group, but no such difference was noted on sub-group analysis of single PVB and PVB infusions studies. It is important to note that the lower end of 95% CI of the meta-analysis was on the higher side (-3.65 for single PVB sub-group and -4.26 for PVB infusion group) and the upper end of the 95% CI close to zero (for single PVB sub-group) in the meta-analysis. Similar was the case in the analysis of 24-h pain scores with the upper end of 95% CI at 0, albeit with a very limited number of studies. The addition of opioids was found to be safe with no increase in the incidence of PONV. Therefore, while our results do not demonstrate a clear advantage of adding opioids to PVB they are suggestive of a probable role of adjuvant opioids in PVB for breast cancer surgery patients, which needs further research.

Such inconclusive result was also evident on descriptive analysis of the included studies with three RCTs[22,23,25] reporting no benefit of adding opioids to PVB, while the remaining three[21,24,26] reporting significant advantage in favor of adjuvant opioids. Such varied results with local anesthetic adjuvants have been reported with other nerve blocks as well. Soulioti *et al*[34] in an RCT reported improved outcomes with the addition of 100 mg of tramadol to brachial plexus block with ropivacaine. In contrast, no such benefit was noted by Kesimci *et al*[35] using a similar dose of tramadol with ropivacaine. Several other RCTs have reported the limited benefit of adding opioids to peripheral nerve blocks. In a recently published trial, Kim *et al*[36] failed to prove any beneficial effect of adding fentanyl to continuous femoral nerve blocks for knee arthroplasty patients. Two RCTs have shown that the addition of fentanyl to transverse abdominus plane block did not improve postoperative outcomes in patients undergoing cesarean section and gynecological surgeries[37,38]. The contrasting results on the role of adjuvant opioids in literature as well as in our review can be attributed to several factors, like the type of surgery, type of nerve block, type and dosage of local anesthetic, type and dose of opioid, pain threshold of patients, the post-operative pain control protocol, *etc.* In our review, while the type of surgery and nerve block were the same, four different opioids were used with varying doses in the included studies. Postoperative pain control protocol was different across studies, which could also have contributed to the heterogeneous results. However, such heterogeneity is expected in clinical trials conducted in different geographical settings. Several other meta-analyses in literature have also pooled outcomes of different opioids in a single analysis[39-41].

In addition to the inter-study heterogeneity, other limitations of our review should also be taken into account while interpreting the results. Firstly, only six RCTs were available for analysis in this review, all with limited sample size. Thus, our review may not have been statistically powered to detect significant differences between the groups. Secondly, data on pain scores at different periods were not available from included studies for a meta-analysis. Furthermore, data as mean and SD were not presented by all studies, further restricting our analysis. Thirdly, pain score and analgesic consumption after surgery can depend on several surgical and patient-dependent factors, like duration and complexity of the surgical procedure, the pain threshold of the patient, *etc.* Such uncontrolled factors could have also influenced the outcomes of the included trials. Lastly, we were unable to register the review protocol on any online database, and this is a significant limitation of our review. Nevertheless, our study has certain novelties. We have presented the first systematic review and meta-analysis assessing the role of adjuvant opioids in PVB for breast cancer surgery patients. The overall quality of

the included RCTs was high, and this lends credibility to the overall review.

CONCLUSION

To conclude, there is a limited role of adjuvant opioids with PVB for breast cancer surgery patients. The addition of opioids had no significant effect on 24-h total analgesic consumption, time to first analgesic, and 24-h pain score. Further homogenous RCTs with a large sample size are needed to clarify the beneficial role of opioids with PVB.

ARTICLE HIGHLIGHTS

Research background

Opioids have been used in combination with local anesthetics for several locoregional anesthetic techniques, leading to better pain control in the immediate postoperative period. However, it is not known if the addition of opioids to paravertebral block (PVB) would lead to better outcomes in breast cancer patients.

Research motivation

No meta-analysis has summarized evidence to assess the value of adding opioids to PVB in breast cancer patients undergoing surgical intervention.

Research objectives

To compare total analgesic consumption, time to first analgesic request, and pain scores with and without the addition of opioids to PVB in breast cancer surgery patients.

Research methods

We conducted an electronic literature search across PubMed, Embase, Scopus, and Google Scholar databases up to October 20, 2020 for randomized controlled trials (RCTs) comparing the addition of opioids to PVB with placebo for breast cancer surgery patients.

Research results

Analysis of six RCTs demonstrated that the addition of opioids to PVB significantly reduced 24-h total analgesic consumption but had no impact on the time to first analgesic request. Pain scores at 24 h were marginally lower with the addition of opioids.

Research conclusions

Current evidence suggests a limited role of adjuvant opioids with PVB for breast cancer surgery patients.

Research perspectives

Further homogenous RCTs with a large sample size are needed to clarify the beneficial role of opioids with PVB.

FOOTNOTES

Author contributions: Chen MH conceived and designed the study, analyzed the data, and wrote the paper; Chen Z and Zhao D were involved in literature search and data collection and reviewed and edited the manuscript; all authors read and approved the final manuscript.

Conflict-of-interest statement: The authors deny any conflict of interest.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Meng-Hua Chen 0000-0002-9174-7198; Zheng Chen 0000-0001-9636-7891; Da Zhao 0000-0001-7994-5215.

S-Editor: Wu YXJ

L-Editor: Filipodia

P-Editor: Wu YXJ

REFERENCES

- Harris E, Barry M, Kell MR. Meta-analysis to determine if surgical resection of the primary tumour in the setting of stage IV breast cancer impacts on survival. *Ann Surg Oncol* 2013; **20**: 2828-2834 [PMID: 23653043 DOI: 10.1245/s10434-013-2998-2]
- Arciero C, Liu Y, Gillespie T, Subhedar P. Surgery and survival in patients with stage IV breast cancer. *Breast J* 2019; **25**: 644-653 [PMID: 31087448 DOI: 10.1111/tbj.13296]
- Sada A, Thiels CA, Britain MK, Dudakovic A, Bergquist WJ, Nickel SR, Moran MJ, Martinez-Jorge J, Jakub JW. Optimizing Discharge Opioid Prescribing Practices After Mastectomy With Immediate Reconstruction. *Mayo Clin Proc Innov Qual Outcomes* 2019; **3**: 183-188 [PMID: 31193969 DOI: 10.1016/j.mayocpiqo.2019.03.001]
- Baron RH. Surgical management of breast cancer. *Semin Oncol Nurs* 2007; **23**: 10-19 [PMID: 17303512 DOI: 10.1016/j.soncn.2006.11.003]
- Cali Cassi L, Biffoli F, Francesconi D, Petrella G, Buonomo O. Anesthesia and analgesia in breast surgery: the benefits of peripheral nerve block. *Eur Rev Med Pharmacol Sci* 2017; **21**: 1341-1345 [PMID: 28387892]
- Poleshuck EL, Katz J, Andrus CH, Hogan LA, Jung BF, Kulick DI, Dworkin RH. Risk factors for chronic pain following breast cancer surgery: a prospective study. *J Pain* 2006; **7**: 626-634 [PMID: 16942948 DOI: 10.1016/j.jpain.2006.02.007]
- Temple-Oberle C, Shea-Budgell MA, Tan M, Semple JL, Schrag C, Barreto M, Blondeel P, Hamming J, Dayan J, Ljungqvist O; ERAS Society. Consensus Review of Optimal Perioperative Care in Breast Reconstruction: Enhanced Recovery after Surgery (ERAS) Society Recommendations. *Plast Reconstr Surg* 2017; **139**: 1056e-1071e [PMID: 28445352 DOI: 10.1097/PRS.00000000000003242]
- Jin Z, Durrands T, Li R, Gan TJ, Lin J. Pectoral block vs paravertebral block: a systematic review, meta-analysis and trial sequential analysis. *Reg Anesth Pain Med* 2020; **45**: 727-732 [PMID: 32727818 DOI: 10.1136/rapm-2020-101512]
- Leong RW, Tan ESJ, Wong SN, Tan KH, Liu CW. Efficacy of erector spinae plane block for analgesia in breast surgery: a systematic review and meta-analysis. *Anaesthesia* 2021; **76**: 404-413 [PMID: 32609389 DOI: 10.1111/anae.15164]
- Kasimahanti R, Arora S, Bhatia N, Singh G. Ultrasound-guided single- vs double-level thoracic paravertebral block for postoperative analgesia in total mastectomy with axillary clearance. *J Clin Anesth* 2016; **33**: 414-421 [PMID: 27555203 DOI: 10.1016/j.jclinane.2016.01.027]
- Salviz EA, Sivriköz N, Ozonur A, Orhan-Sungur M, Savran-Karadeniz M, Altun D, Hocaoglu E, Celet-Ozden B, Tugrul KM. Ultrasound-Guided Bilateral Thoracic Paravertebral Blocks as an Adjunct to General Anesthesia in Patients Undergoing Reduction Mammoplasty: A Historical Cohort Study. *Plast Reconstr Surg* 2017; **139**: 20e-28e [PMID: 28027224 DOI: 10.1097/PRS.00000000000002842]
- Terkawi AS, Tsang S, Sessler DI, Terkawi RS, Nunemaker MS, Durieux ME, Shilling A. Improving Analgesic Efficacy and Safety of Thoracic Paravertebral Block for Breast Surgery: A Mixed-Effects Meta-Analysis. *Pain Physician* 2015; **18**: E757-E780 [PMID: 26431130]
- Priya S, Bamba C. Comparison of Morphine and Clonidine as Adjuvants in Paravertebral Block. *Anesth Essays Res* 2018; **12**: 459-463 [PMID: 29962616 DOI: 10.4103/aer.AER_27_18]
- Mohta M, Kalra B, Sethi AK, Kaur N. Efficacy of dexmedetomidine as an adjuvant in paravertebral block in breast cancer surgery. *J Anesth* 2016; **30**: 252-260 [PMID: 26694929 DOI: 10.1007/s00540-015-2123-8]
- Hamed MA, Ghaber S, Reda A. Dexmedetomidine and Fentanyl as an Adjunct to Bupivacaine 0.5% in Supraclavicular Nerve Block: A Randomized Controlled Study. *Anesth Essays Res* 2018; **12**: 475-479 [PMID: 29962619 DOI: 10.4103/aer.AER_50_18]
- Akhondzadeh R, Rashidi M, Gousheh M, Olapour A, Tasbihi B. Comparison of the Ketamine-Lidocaine and Fentanyl-Lidocaine in Postoperative Analgesia in Axillary Block in Upper Limb Fractures By Ultrasound Guidance. *Anesth Pain Med* 2019; **9**: e92695 [PMID: 32280613 DOI: 10.5812/aapm.92695]
- Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**: e1000097 [PMID: 19621072 DOI: 10.1371/journal.pmed.1000097]
- Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page M, Welch VA. Cochrane Handbook for Systematic Reviews of Interventions. Version 6. Cochrane; 2019 [DOI: 10.1002/9781119536604]
- Wallace BC, Schmid CH, Lau J, Trikalinos TA. Meta-Analyst: software for meta-analysis of binary, continuous and diagnostic data. *BMC Med Res Methodol* 2009; **9**: 80 [PMID: 19961608 DOI: 10.1186/1471-2288-9-80]
- Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014; **14**: 135 [PMID: 25524443 DOI: 10.1186/1471-2288-14-135]
- Omar Mostafa M, Makram Botros J, Sayed Khaleel AM. Effect of Dexmedetomidine Versus Nalbuphine as an Adjuvant on Paravertebral Block to Manage Postoperative Pain After Mastectomies. *Anesth Pain Med* 2018; **8**: e13308 [PMID: 30027066 DOI: 10.5812/aapm.13308]
- Pushparajan HK, Punj J, Pandey R, Darlong V, Srivastava A, Batra RK, Bhan A. Addition of Fentanyl to Ropivacaine Infusion in Continuous Thoracic Paravertebral Infusion Does Not Improve Its Analgesic Effect Following Modified Radical Mastectomy: A Randomized Controlled Trial. *AANA J* 2017; **85**: 352-356 [PMID: 31566535]

- 23 **Morsy A**, Abd-elmaksoud M, Abdel Aziz R, Metwally M. Comparison between dexmedetomidine vs morphine added to bupivacaine in paravertebral block in patients scheduled for modified radical mastectomy. *Res Opin Anesth Intensive Care* 2017; **4**: 59-64 [DOI: [10.4103/roaic.roaic_10_16](https://doi.org/10.4103/roaic.roaic_10_16)]
- 24 **Bhuvaneswari V**, Wig J, Mathew PJ, Singh G. Post-operative pain and analgesic requirements after paravertebral block for mastectomy: A randomized controlled trial of different concentrations of bupivacaine and fentanyl. *Indian J Anaesth* 2012; **56**: 34-39 [PMID: [22529417](https://pubmed.ncbi.nlm.nih.gov/22529417/) DOI: [10.4103/0019-5049.93341](https://doi.org/10.4103/0019-5049.93341)]
- 25 **Omar AM**, Mansour MA, Abdelwahab HH, Aboushanab OH. Role of ketamine and tramadol as adjuncts to bupivacaine 0.5% in paravertebral block for breast surgery: A randomized double-blind study. *Egypt J Anaesth* 2011; **27**: 101-105 [DOI: [10.1016/j.egja.2011.04.002](https://doi.org/10.1016/j.egja.2011.04.002)]
- 26 **Burlacu CL**, Frizelle HP, Moriarty DC, Buggy DJ. Fentanyl and clonidine as adjunctive analgesics with levobupivacaine in paravertebral analgesia for breast surgery. *Anaesthesia* 2006; **61**: 932-937 [PMID: [16978305](https://pubmed.ncbi.nlm.nih.gov/16978305/) DOI: [10.1111/j.1365-2044.2006.04793.x](https://doi.org/10.1111/j.1365-2044.2006.04793.x)]
- 27 **Vila H Jr**, Liu J, Kavasmaneck D. Paravertebral block: new benefits from an old procedure. *Curr Opin Anaesthesiol* 2007; **20**: 316-318 [PMID: [17620838](https://pubmed.ncbi.nlm.nih.gov/17620838/) DOI: [10.1097/ACO.0b013e328166780e](https://doi.org/10.1097/ACO.0b013e328166780e)]
- 28 **Yang J**, Hao Z, Li W, Duan C, Fan X, Xin J, Ren C. The Efficacy and Safety of Paravertebral Block Combined with Parecoxib During Video-Assisted Thoracic Surgery: A Randomized Controlled Trial. *J Pain Res* 2020; **13**: 355-366 [PMID: [32104057](https://pubmed.ncbi.nlm.nih.gov/32104057/) DOI: [10.2147/JPR.S244787](https://doi.org/10.2147/JPR.S244787)]
- 29 **Mansour NA**. Ropivacaine vs Bupivacaine in Postoperative Pain Control. *J Biotechnol Biomater* 2012; **02**: 1-8 [DOI: [10.4172/2155-952X.1000137](https://doi.org/10.4172/2155-952X.1000137)]
- 30 **Krishna Prasad GV**, Khanna S, Jaishree SV. Review of adjuvants to local anesthetics in peripheral nerve blocks: Current and future trends. *Saudi J Anaesth* 2020; **14**: 77-84 [PMID: [31998024](https://pubmed.ncbi.nlm.nih.gov/31998024/) DOI: [10.4103/sja.SJA_423_19](https://doi.org/10.4103/sja.SJA_423_19)]
- 31 **Stein C**. The control of pain in peripheral tissue by opioids. *N Engl J Med* 1995; **332**: 1685-1690 [PMID: [7760870](https://pubmed.ncbi.nlm.nih.gov/7760870/) DOI: [10.1056/nejm199506223322506](https://doi.org/10.1056/nejm199506223322506)]
- 32 **Gissen AJ**, Gugino LD, Datta S, Miller J, Covino BG. Effects of fentanyl and sufentanil on peripheral mammalian nerves. *Anesth Analg* 1987; **66**: 1272-1276 [PMID: [2961289](https://pubmed.ncbi.nlm.nih.gov/2961289/) DOI: [10.1213/00000539-198712000-00012](https://doi.org/10.1213/00000539-198712000-00012)]
- 33 **Nishikawa K**, Kanaya N, Nakayama M, Igarashi M, Tsunoda K, Namiki A. Fentanyl improves analgesia but prolongs the onset of axillary brachial plexus block by peripheral mechanism. *Anesth Analg* 2000; **91**: 384-387 [PMID: [10910853](https://pubmed.ncbi.nlm.nih.gov/10910853/) DOI: [10.1097/00000539-200008000-00028](https://doi.org/10.1097/00000539-200008000-00028)]
- 34 **Soulioti E**, Tsaroucha A, Makris A, Koutsaki M, Sklika E, Mela A, Megaloikononimos PD, Mavrogenis AF, Fassoulaki A. Addition of 100 mg of Tramadol to 40 mL of 0.5% Ropivacaine for Interscalene Brachial Plexus Block Improves Postoperative Analgesia in Patients Undergoing Shoulder Surgeries as Compared to Ropivacaine Alone-A Randomized Controlled Study. *Medicina (Kaunas)* 2019; **55** [PMID: [31340565](https://pubmed.ncbi.nlm.nih.gov/31340565/) DOI: [10.3390/medicina55070399](https://doi.org/10.3390/medicina55070399)]
- 35 **Kesimci E**, Izdes S, Gozdemir M, Kanbak O. Tramadol does not prolong the effect of ropivacaine 7.5 mg/mL for axillary brachial plexus block. *Acta Anaesthesiol Scand* 2007; **51**: 736-741 [PMID: [17425616](https://pubmed.ncbi.nlm.nih.gov/17425616/) DOI: [10.1111/j.1399-6576.2007.01308.x](https://doi.org/10.1111/j.1399-6576.2007.01308.x)]
- 36 **Kim GH**, Lee JW, Kim GE, Lee SS, Son SL, Kim BU, Cho HN, Kwon MY, Koo MS, Kim JE, Yun MJ. Analgesic effect of ropivacaine with fentanyl in comparison with ropivacaine alone for continuous femoral nerve block after knee replacement arthroplasty: a prospective, randomized, double-blinded study. *Anesth Pain Med (Seoul)* 2020; **15**: 209-216 [PMID: [33329816](https://pubmed.ncbi.nlm.nih.gov/33329816/) DOI: [10.17085/apm.2020.15.2.209](https://doi.org/10.17085/apm.2020.15.2.209)]
- 37 **Chen Q**, Liu X, Zhong X, Yang B. Addition of dexmedetomidine or fentanyl to ropivacaine for transversus abdominis plane block: evaluation of effect on postoperative pain and quality of recovery in gynecological surgery. *J Pain Res* 2018; **11**: 2897-2903 [PMID: [30532583](https://pubmed.ncbi.nlm.nih.gov/30532583/) DOI: [10.2147/JPR.S178516](https://doi.org/10.2147/JPR.S178516)]
- 38 **Wang LZ**, Liu X, Zhang YF, Hu XX, Zhang XM. Addition of fentanyl to the ultrasound-guided transversus abdominis plane block does not improve analgesia following cesarean delivery. *Exp Ther Med* 2016; **11**: 1441-1446 [PMID: [27073462](https://pubmed.ncbi.nlm.nih.gov/27073462/) DOI: [10.3892/etm.2016.3053](https://doi.org/10.3892/etm.2016.3053)]
- 39 **Frauenknecht J**, Kirkham KR, Jacot-Guillarmod A, Albrecht E. Analgesic impact of intra-operative opioids vs. opioid-free anaesthesia: a systematic review and meta-analysis. *Anaesthesia* 2019; **74**: 651-662 [PMID: [30802933](https://pubmed.ncbi.nlm.nih.gov/30802933/) DOI: [10.1111/anae.14582](https://doi.org/10.1111/anae.14582)]
- 40 **Wang J**, Li QB, Wu YY, Wang BN, Kang JL, Xu XW. Efficacy and Safety of Opioids for the Prevention of Etomidate-Induced Myoclonus: A Meta-Analysis. *Am J Ther* 2018; **25**: e517-e523 [PMID: [26840341](https://pubmed.ncbi.nlm.nih.gov/26840341/) DOI: [10.1097/MJT.0000000000000404](https://doi.org/10.1097/MJT.0000000000000404)]
- 41 **Luo N**, Tan S, Li X, Singh S, Liu S, Chen C, Huang Z, Feng S, Lin Y, Cen H, Liang M, Chen M. Efficacy and Safety of Opioids in Treating Cancer-Related Dyspnea: A Systematic Review and Meta-Analysis Based on Randomized Controlled Trials. *J Pain Symptom Manage* 2021; **61**: 198-210.e1 [PMID: [32730950](https://pubmed.ncbi.nlm.nih.gov/32730950/) DOI: [10.1016/j.jpainsymman.2020.07.021](https://doi.org/10.1016/j.jpainsymman.2020.07.021)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

