

World Journal of *Clinical Cases*

World J Clin Cases 2022 June 16; 10(17): 5518-5933



MINIREVIEWS

- 5518 Occult hepatitis B – the result of the host immune response interaction with different genomic expressions of the virus
Gherlan GS
- 5531 Pulmonary complications of portal hypertension: The overlooked decompensation
Craciun R, Mocan T, Procopet B, Nemes A, Tefas C, Sparchez M, Mocan LP, Sparchez Z
- 5541 Ethical review of off-label drugs during the COVID-19 pandemic
Li QY, Lv Y, An ZY, Dai NN, Hong X, Zhang Y, Liang LJ

ORIGINAL ARTICLE**Case Control Study**

- 5551 Gut peptide changes in patients with obstructive jaundice undergoing biliary drainage: A prospective case control study
Pavić T, Pelajić S, Blažević N, Kralj D, Milošević M, Mikolasevic I, Lerotic I, Hrabar D

Retrospective Cohort Study

- 5566 Longitudinal assessment of liver stiffness by transient elastography for chronic hepatitis C patients
Mezina A, Krishnan A, Woreta TA, Rubenstein KB, Watson E, Chen PH, Rodriguez-Watson C

Retrospective Study

- 5577 Clinical evaluation of prone position ventilation in the treatment of acute respiratory distress syndrome induced by sepsis
Xia WH, Yang CL, Chen Z, Ouyang CH, Ouyang GQ, Li QG
- 5586 Three-dimensional arterial spin labeling and diffusion kurtosis imaging in evaluating perfusion and infarct area size in acute cerebral ischemia
Jiang YY, Zhong ZL, Zuo M
- 5595 Intrathecal methotrexate in combination with systemic chemotherapy in glioblastoma patients with leptomeningeal dissemination: A retrospective analysis
Kang X, Chen F, Yang SB, Wang YL, Qian ZH, Li Y, Lin H, Li P, Peng YC, Wang XM, Li WB
- 5606 Hepatic epithelioid hemangioendothelioma: Clinical characteristics, diagnosis, treatment, and prognosis
Zhao M, Yin F
- 5620 Difference between type 2 gastroesophageal varices and isolated fundic varices in clinical profiles and portosystemic collaterals
Song YH, Xiang HY, Si KK, Wang ZH, Zhang Y, Liu C, Xu KS, Li X

- 5634** Assessment of incidental focal colorectal uptake by analysis of fluorine-18 fluorodeoxyglucose positron emission tomography parameters

Lee H, Hwang KH, Kwon KA

Observational Study

- 5646** "Zero ischemia" laparoscopic partial nephrectomy by high-power GreenLight laser enucleation for renal carcinoma: A single-center experience

Zhang XM, Xu JD, Lv JM, Pan XW, Cao JW, Chu J, Cui XG

- 5655** High Eckardt score and previous treatment were associated with poor postperoral endoscopic myotomy pain control: A retrospective study

Chen WN, Xu YL, Zhang XG

- 5667** Higher volume growth rate is associated with development of worrisome features in patients with branch duct-intraductal papillary mucinous neoplasms

Innocenti T, Danti G, Lynch EN, Dragoni G, Gottin M, Fedeli F, Palatresi D, Biagini MR, Milani S, Miele V, Galli A

Prospective Study

- 5680** Application of a new anatomic hook-rod-pedicle screw system in young patients with lumbar spondylolysis: A pilot study

Li DM, Li YC, Jiang W, Peng BG

META-ANALYSIS

- 5690** Systematic review of Yougui pills combined with levothyroxine sodium in the treatment of hypothyroidism

Liu XP, Zhou YN, Tan CE

CASE REPORT

- 5702** Allogeneic stem cell transplantation-A curative treatment for paroxysmal nocturnal hemoglobinuria with PIGT mutation: A case report

Schenone L, Notarantonio AB, Latger-Cannard V, Fremaux-Bacchi V, De Carvalho-Bittencourt M, Rubio MT, Muller M, D'Aveni M

- 5708** Gray zone lymphoma effectively treated with cyclophosphamide, doxorubicin, vincristine, prednisolone, and rituximab chemotherapy: A case report

Hojo N, Nagasaki M, Mihara Y

- 5717** Diagnosis of spontaneous isolated superior mesenteric artery dissection with ultrasound: A case report

Zhang Y, Zhou JY, Liu J, Bai C

- 5723** Adrenocorticotrophic hormone-secreting pancreatic neuroendocrine carcinoma with multiple organ infections and widespread thrombosis: A case report

Yoshihara A, Nishihama K, Inoue C, Okano Y, Eguchi K, Tanaka S, Maki K, Fridman D'Alessandro V, Takeshita A, Yasuma T, Uemura M, Suzuki T, Gabazza EC, Yano Y

- 5732** Management of the palato-radicular groove with a periodontal regenerative procedure and prosthodontic treatment: A case report

Ling DH, Shi WP, Wang YH, Lai DP, Zhang YZ

- 5741** Combined thoracic paravertebral block and interscalene brachial plexus block for modified radical mastectomy: A case report
Hu ZT, Sun G, Wang ST, Li K
- 5748** Chondromyxoid fibroma of the cervical spine: A case report
Li C, Li S, Hu W
- 5756** Preterm neonate with a large congenital hemangioma on maxillofacial site causing thrombocytopenia and heart failure: A case report
Ren N, Jin CS, Zhao XQ, Gao WH, Gao YX, Wang Y, Zhang YF
- 5764** Simultaneous multiple primary malignancies diagnosed by endoscopic ultrasound-guided fine-needle aspiration: A case report
Yang J, Zeng Y, Zhang JW
- 5770** Neuroendocrine tumour of the descending part of the duodenum complicated with schwannoma: A case report
Zhang L, Zhang C, Feng SY, Ma PP, Zhang S, Wang QQ
- 5776** Massive hemothorax following internal jugular vein catheterization under ultrasound guidance: A case report
Kang H, Cho SY, Suk EH, Ju W, Choi JY
- 5783** Unilateral adrenal tuberculosis whose computed tomography imaging characteristics mimic a malignant tumor: A case report
Liu H, Tang TJ, An ZM, Yu YR
- 5789** Modified membrane fixation technique in a severe continuous horizontal bone defect: A case report
Wang LH, Ruan Y, Zhao WY, Chen JP, Yang F
- 5798** Surgical repair of an emergent giant hepatic aneurysm with an abdominal aortic dissection: A case report
Wen X, Yao ZY, Zhang Q, Wei W, Chen XY, Huang B
- 5805** Heterotopic ossification beneath the upper abdominal incision after radical gastrectomy: Two case reports
Zhang X, Xia PT, Ma YC, Dai Y, Wang YL
- 5810** Non-alcoholic Wernicke encephalopathy in an esophageal cancer patient receiving radiotherapy: A case report
Zhang Y, Wang L, Jiang J, Chen WY
- 5816** New approach for the treatment of vertical root fracture of teeth: A case report and review of literature
Zhong X, Yan P, Fan W
- 5825** Ultrasound-guided microwave ablation as a palliative treatment for mycosis fungoides eyelid involvement: A case report
Chen YW, Yang HZ, Zhao SS, Zhang Z, Chen ZM, Feng HH, An MH, Wang KK, Duan R, Chen BD
- 5833** Pulp revascularization on an adult mandibular right second premolar: A case report
Yang YQ, Wu BL, Zeng JK, Jiang C, Chen M

- 5841** Barrett's esophagus in a patient with bulimia nervosa: A case report
Gouda A, El-Kassas M
- 5846** Spontaneous gallbladder perforation and colon fistula in hypertriglyceridemia-related severe acute pancreatitis: A case report
Wang QP, Chen YJ, Sun MX, Dai JY, Cao J, Xu Q, Zhang GN, Zhang SY
- 5854** Beware of gastric tube in esophagectomy after gastric radiotherapy: A case report
Yurttas C, Wichmann D, Gani C, Bongers MN, Singer S, Thiel C, Koehngrainer A, Thiel K
- 5861** Transition from minimal change disease to focal segmental glomerulosclerosis related to occupational exposure: A case report
Tang L, Cai Z, Wang SX, Zhao WJ
- 5869** Lung adenocarcinoma metastasis to paranasal sinus: A case report
Li WJ, Xue HX, You JQ, Chao CJ
- 5877** Follicular lymphoma presenting like marginal zone lymphoma: A case report
Peng HY, Xiu YJ, Chen WH, Gu QL, Du X
- 5884** Primary renal small cell carcinoma: A case report
Xie K, Li XY, Liao BJ, Wu SC, Chen WM
- 5893** Gitelman syndrome: A case report
Chen SY, Jie N
- 5899** High-frame-rate contrast-enhanced ultrasound findings of liver metastasis of duodenal gastrointestinal stromal tumor: A case report and literature review
Chen JH, Huang Y
- 5910** Tumor-like disorder of the brachial plexus region in a patient with hemophilia: A case report
Guo EQ, Yang XD, Lu HR
- 5916** Response to dacomitinib in advanced non-small-cell lung cancer harboring the rare delE709_T710insD mutation: A case report
Xu F, Xia ML, Pan HY, Pan JW, Shen YH
- 5923** Loss of human epidermal receptor-2 in human epidermal receptor-2+ breast cancer after neoadjuvant treatment: A case report
Yu J, Li NL

LETTER TO THE EDITOR

- 5929** Repetitive transcranial magnetic stimulation for post-traumatic stress disorder: Lights and shadows
Concerto C, Lanza G, Fisticaro F, Pennisi M, Rodolico A, Torrisi G, Bella R, Aguglia E

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Raden Andri Primadhi, MD, PhD, Assistant Professor, Surgeon, Department of Orthopaedics and Traumatology, Universitas Padjadjaran Medical School, Hasan Sadikin Hospital, Bandung 40161, Indonesia. randri@unpad.ac.id

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for *WJCC* as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The *WJCC*'s CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Hua-Ge Yan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jim-Lai Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

June 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Cohort Study

Longitudinal assessment of liver stiffness by transient elastography for chronic hepatitis C patients

Anya Mezina, Arunkumar Krishnan, Tinsay A Woreta, Kevin B Rubenstein, Eric Watson, Po-Hung Chen, Carla Rodriguez-Watson

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Kishida Y

Received: August 10, 2021

Peer-review started: August 10, 2021

First decision: November 17, 2021

Revised: December 16, 2021

Accepted: April 21, 2022

Article in press: April 21, 2022

Published online: June 16, 2022



Anya Mezina, Department of Medicine, Johns Hopkins School of Medicine, Baltimore, MD 21287, United States

Arunkumar Krishnan, Tinsay A Woreta, Po-Hung Chen, Division of Gastroenterology and Hepatology, Johns Hopkins School of Medicine, Baltimore, MD 21231, United States

Kevin B Rubenstein, Eric Watson, Carla Rodriguez-Watson, Mid-Atlantic Permanente Research Institute, Kaiser Permanente Mid-Atlantic States, Rockville 20852, United States

Carla Rodriguez-Watson, Department of Health Policy and Management, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD 21287, United States

Carla Rodriguez-Watson, Innovation in Medical Evidence Development and Surveillance (IMEDS) Program, Reagan-Udall Foundation for the FDA, Washington, 20036, United States

Corresponding author: Carla Rodriguez-Watson, PhD, Senior Researcher, Mid-Atlantic Permanente Research Institute, Kaiser Permanente Mid-Atlantic States, 2101 E Jefferson Street, Rockville 20852, United States. crodriguezwatson@reaganudall.org

Abstract

BACKGROUND

Liver fibrosis is a common pathway of liver injury and is a feature of most chronic liver diseases. Fibrosis progression varies markedly in patients with hepatitis C virus (HCV). Liver stiffness has been recommended as a parameter of fibrosis progression/regression in patients with HCV.

AIM

To investigate changes in liver stiffness measured by transient elastography (TE) in a large, racially diverse cohort of United States patients with chronic hepatitis C (CHC).

METHODS

We evaluated the differences in liver stiffness between patients treated with direct-acting antiviral (DAA) therapy and untreated patients. Patients had ≥ 2 TE measurements and no prior DAA exposure. We used linear regression to measure the change in liver stiffness between first and last TE in response to treatment, controlling for age, sex, race, diabetes, smoking status, human immunodeficiency

virus status, baseline alanine aminotransferase, and baseline liver stiffness. Separate regression models analyzed the change in liver stiffness as measured by kPa, stratified by cirrhosis status.

RESULTS

Of 813 patients, 419 (52%) initiated DAA treatment. Baseline liver stiffness was 12 kPa in 127 (16%). Median time between first and last TE was 11.7 and 12.7 mo among treated and untreated patients, respectively. There was no significant change in liver stiffness observed over time in either the group initiating DAA treatment (0.016 kPa/month; CI: -0.051, 0.084) or in the untreated group (0.001 kPa/mo; CI: -0.090, 0.092), controlling for covariates. A higher baseline kPa score was independently associated with decreased liver stiffness.

CONCLUSION

DAA treatment was not associated with a differential change in liver stiffness over time in patients with CHC compared to untreated patients.

Key Words: Chronic hepatitis C; Liver stiffness; Cirrhosis; Transient elastography; Direct-acting antiviral therapy

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: We evaluated changes in liver stiffness measured by transient elastography (TE) in a large, racially diverse cohort of United States patients with chronic hepatitis C (CHC). We retrospectively evaluated differences in liver stiffness between patients treated with direct-acting antiviral (DAA) therapy and untreated patients. Our study shows a higher baseline kPa score was independently associated with decreased liver stiffness, and that differences in liver stiffness may be observed on serial TE measurements in patients with higher baseline scores, irrespective of treatment effect. DAA treatment was not associated with a differential change in liver stiffness over time in patients with CHC compared to untreated patients.

Citation: Mezina A, Krishnan A, Woreta TA, Rubenstein KB, Watson E, Chen PH, Rodriguez-Watson C. Longitudinal assessment of liver stiffness by transient elastography for chronic hepatitis C patients. *World J Clin Cases* 2022; 10(17): 5566-5576

URL: <https://www.wjgnet.com/2307-8960/full/v10/i17/5566.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i17.5566>

INTRODUCTION

Chronic hepatitis C (CHC) infection, affecting an estimated 71.1 million people worldwide, is a major cause of liver cirrhosis [1,2]. The degree of liver fibrosis is a chief determinant of complications related to end-stage liver disease and is associated with the risk of hepatic decompensation, hepatocellular carcinoma, and mortality[3]. The level of CHC-related hepatic fibrosis is detected through histology by the meta-analysis of histological data in viral hepatitis (METAVIR) scoring system (F0 indicating no fibrosis to F4 indicating cirrhosis)[4]. The temporal evolution of fibrosis in patients with CHC is non-linear, with variable progression observed between different stages of disease specified by METAVIR fibrosis scores[5]. The degree of progression of liver fibrosis is further influenced by variables including age, gender, alcohol consumption, insulin resistance, high body mass index, and co-infection with hepatitis B virus (HBV) or human immunodeficiency virus (HIV)[6-9]. Histological assessment of the liver is considered the optimal approach to evaluate changes in fibrosis over time and remains the gold standard for the diagnosis of cirrhosis[10]. However, reliance on liver biopsy is limited by the intra- and interobserver variability of fibrosis scoring, the sampling variability, and finally, the potential for complications from an invasive procedure, which renders serial histopathological assessments impractical in routine clinical care[11-13].

Transient elastography (TE) has become a widely used, noninvasive ultrasound-based technique to measure liver stiffness and accurately assess severe fibrosis and cirrhosis without the morbidity and mortality associated with liver biopsy. TE is validated in patients with CHC with 89% sensitivity and 91% specificity for the diagnosis of cirrhosis[14]. Achieving sustained virological response (SVR) after treatment of CHC has been shown to reduce the rate of liver fibrosis progression and even partially reverse fibrosis as assessed by histology as well as TE[15-17]. Eradication of HCV is associated with a reduction in all-cause and liver-related mortality and a lower risk of progression to cirrhosis[15,18,19]. However, despite achieving SVR, some patients will develop progressively higher liver stiffness measurements over time, and up to 60% of those with cirrhosis at the time of treatment with direct-

acting antiviral agents (DAA) will continue to exhibit liver stiffness measurements in the cirrhotic range, increasing the risk for hepatic decompensation and adverse clinical outcomes[20-22]. Due to the broad implementation of guideline-directed treatment of CHC, understanding the factors leading to differential changes in liver stiffness will be integral to refining clinical decisions regarding management and surveillance for complications of end-stage liver disease[17,23]. Many studies utilizing TE to evaluate longitudinal changes in liver stiffness have been limited to the duration of DAA therapy or the 12-24 wk post-treatment when SVR is ascertained[24,25-27]. Other studies with long-term follow-up periods have been conducted outside of the United States and may not reflect the same racially diverse patient population and genotypic distribution of HCV, limiting the generalizability of findings[22,28].

The American Gastroenterological Association has recognized the need for large, long-term investigations of outcomes in patients with CHC treated with DAA therapy and indicated that current evidence is insufficient to recommend a universal policy of noninvasive testing of liver fibrosis in successfully treated patients[14,29]. To address this need, we performed a longitudinal, retrospective observational study investigating changes in liver stiffness measured by TE in a racially diverse cohort of United States patients with CHC.

MATERIALS AND METHODS

Study design: We conducted a longitudinal retrospective study of patients with confirmed CHC infection seen at Johns Hopkins Health System (JHHS) and Kaiser Permanente Mid-Atlantic States (KPMAS) between April 1, 2014 and March 31, 2018. JHHS is an academic health system primarily serving Baltimore, MD, and the surrounding area. KPMAS is an integrated healthcare system serving more than 750000 members in a region that includes the District of Columbia, Maryland, and Virginia. The Institutional Review Board approved the study at JHHS and KPMAS.

Study population: Patients at JHHS were identified by querying the electronic health record (EHR) system for the International Classification of Disease, Ninth (ICD-9), or Tenth Revision (ICD-10) diagnosis codes for hepatitis C viral (HCV) infection or a positive HCV RNA test. At KPMAS, HCV patients were identified by positive HCV RNA, HCV genotype, ≥ 2 refills of interferon-based HCV therapy within 1 year, or positive HCV antibody test plus ≥ 1 HCV-coded visit. Included patients were at least 18 years of age, treatment naïve to DAA at the time of the first liver stiffness measurement, and had at least two liver stiffness measurements. Patients were excluded if the date of the first liver stiffness measurement was more than 30 d before the date on which our HCV definition is met. We reason that liver stiffness measurements performed fewer than 30 d before HCV recognition was likely due to clinically suspected HCV, but those performed more than 30 d before were due to other chronic liver conditions. We further excluded any patients found to have an undetectable HCV viral load within 180 d prior to the first liver stiffness measurement indicating clearance of HCV infection. The baseline was defined as the date of the first liver stiffness measurement. Treatment was defined by pharmacy dispensation dates in the EHR. Only those patients initiating treatment between the date of the first and last TE were included in the treatment group. TE results and biochemical, clinical, and demographic characteristics were extracted from the EHR. We collected data on the following: age, sex, race, diabetes, chronic kidney disease (CKD), body mass index (BMI), smoking status, alcohol consumption, HIV status, HBV status, hepatocellular carcinoma, aspartate aminotransferase (AST), alanine aminotransferase (ALT), HCV RNA viral load, bilirubin, platelet count, and TE results. The most proximal laboratory and clinical data were collected from a window that spanned 180 days prior to the baseline TE scan and up to 30 d after.

Liver stiffness measurements: Liver stiffness measurements were obtained using transient elastography (Fibroscan 502, Echosens, Paris, France) by trained operators at JHHS and KPMAS. Participants were instructed to fast for a minimum of 2 h prior to testing. The liver stiffness measurement was considered reliable only if a minimum of 10 successful acquisitions were obtained, with an interquartile range $\leq 30\%$ of the median liver stiffness measurement and success rate $\geq 60\%$. The success rate was calculated as the ratio of the number of successful acquisitions over the total number of acquisitions. The result was reported as a median of liver stiffness expressed in kilopascals (kPa).

Statistical analysis: All statistical analyses were conducted using R version 3.6.0, Stata version 14 (StataCorp LP., College Station, TX, United States), and SAS version 9.4 (SAS Institute Inc., Cary, NC, United States). Descriptive statistics were computed, including counts and percentages for categorical variables, and either means and standard deviations, medians with quartiles, or medians with minimum and maximum values for continuous variables. A *t*-test was used to compare mean baseline liver stiffness measurements (kPa) between treatment groups. We used multivariable linear regression models to assess the association between treatment and the difference in liver stiffness measurements between first and last TE, adjusting for baseline kPa, and the covariates mentioned above. We included an interaction term between time and treatment to directly compare the change in kPa between treated and untreated subjects. This model can be considered equivalent to one with only the last TE

measurement as the outcome, provided that baseline kPa (also referred to as first TE measurement) is included along with all of the same covariates[30]. The analysis was further stratified by baseline cirrhosis status (kPa \geq 12). A *P* value $<$ 0.05 was considered statistically significant.

Sensitivity analysis: We found that some patients had two liver stiffness measurements obtained in rapid succession within 7 d of one another. We chose to omit patients with only two measurements taken within 7 d because differences in liver stiffness during a brief time interval are likely to reflect measurement error rather than meaningful physiological changes in the liver. Therefore, as a sensitivity analysis, we repeated all analyses, including patients with liver stiffness measurements taken within 7 d of one another. We will describe results, both including and excluding pairs of liver stiffness measurements taken within 7 d, to assess whether these data impact our conclusions.

RESULTS

Of 813 patients, 84% were at least 50 years of age, 79% were Black, 79% were current or former smokers, 37% were coinfecting with HIV, 3% were coinfecting with HBV, 19% had diabetes, and 52% initiated treatment with a DAA (Table 1). The median (Q1, Q3) time between first and last liver stiffness measurement was 11.73 (7.41, 17.87) months among those initiating treatment and 12.68 (8.71, 16.94) months among those who were untreated (Table 2). 22% ($n = 91$) of treated subjects had a kPa score \geq 12, compared with 9% ($n = 36$) of untreated patients. The mean (SD) baseline liver stiffness among treated patients was 10.69 (9.48) kPa, significantly higher compared to the mean (SD) of 7.32 (6.10) kPa among untreated patients ($P <$ 0.001). The HCV genotype was genotype 1 in 60% of patients (comprising 91% of patients with known HCV genotype); however, HCV genotype data are not included elsewhere in our analyses because of missing data in 34% of patients. Additionally, due to missing data regarding sustained virologic response (SVR) rates in treated patients, alcohol consumption, and body mass index, these variables were also excluded from the analyses. We also omitted 38 patients (4.7%) who did not have an ALT result up to 180 d before the first liver stiffness measurement, as well as 3 patients (0.4%) with unknown race and 2 patients (0.2%) with unknown smoking status. After excluding these 43 total patients with incomplete information, 770 total patients were included in the regression models. In the multivariable linear regression model examining the change in kPa as the dependent variable, the baseline kPa score was independently associated with a reduction in liver stiffness between the first and last TE (-0.557 kPa; CI: -0.699, -0.415), controlling for age, race (Black, White, or other), sex, HIV status, smoking status, baseline ALT score, and treatment status (Table 3). We observed no significant change in liver stiffness over time among patients who were treated (0.016 kPa/mo; CI: -0.051, 0.084) or untreated (0.001 kPa/mo; CI: -0.090, 0.092); as such, there was no significant difference in the change in liver stiffness among treated patients compared to untreated patients (0.015 kPa/mo, CI: -0.106, 0.136). The difference in the kPa score between the first and last TE, stratified by treatment status, is shown in Figure 1. Next, the analysis was stratified to examine the association of these covariates with liver stiffness in patients with cirrhosis (kPa \geq 12, $n = 119$) as well as non-cirrhotic patients (kPa $<$ 12, $n = 651$). There was not a significant effect of DAA treatment on the change in liver stiffness over time in either cohort, and the difference in liver stiffness between treated and untreated patients was similar. A higher baseline kPa was an independent predictor of a reduction in liver stiffness measurement in both groups. Among patients without cirrhosis, age \geq 50 was independently associated with increased liver stiffness (0.949 kPa; CI: 0.336, 1.563).

Sensitivity analysis results: We repeated the analyses described above with the inclusion of patients who had two liver stiffness measurements obtained within 7 d ($n = 7$). We did not observe a significant change over time in liver stiffness in the treated (0.017 kPa/mo; CI: -0.051, 0.086) or untreated patients (-0.027 kPa/mo; CI: -0.120, 0.066), and the rate of change was not different when comparing treated and untreated patients. Results stratified by cirrhosis status (kPa $<$ 12 and kPa \geq 12) were similar to the original analysis (Supplementary Table 1 and Supplementary Figure 1). Estimates for all other covariates yielded qualitatively comparable results to the main analysis.

DISCUSSION

The widespread application of TE as a noninvasive tool to estimate liver stiffness in patients with CHC has raised important questions regarding the clinical benefit of repeating TE following DAA treatment and the evolution of fibrosis on post-treatment measurements. Our retrospective longitudinal cohort study employed TE to investigate changes in liver stiffness over time in a large, racially diverse patient population with CHC infection and assess treatment response.

In our multivariable regression models, we compared the rate of change of liver stiffness over time between DAA-treated and untreated subjects, controlling for race, age, gender, diabetes, smoking status, HIV co-infection, baseline ALT, and baseline kPa that may be independently associated with liver

Table 1 Demographic and clinical variables in patients with chronic hepatitis C stratified by treatment status (n = 813)

Variables	HCV treated	HCV untreated
	(n = 419 (51.5%))	[n = 394 (48.5%)]
Age (yr) [mean (SD)]	56.54 (8.26)	56.14 (9.02)
Male sex [n (%)]	285 (68.02)	234 (59.39)
Race [n (%)]		
Black	339 (80.91)	303 (77.49)
Other	18 (4.30)	20 (5.12)
White	62 (14.80)	68 (17.39)
Smoking history		
Non-smoker	70 (16.71)	98 (25.00)
Current smoker	216 (51.55)	183 (46.68)
Former smoker	133 (31.74)	111 (28.32)
Co-infections [n (%)]		
HIV	184 (43.91)	113 (28.68)
HBV	9 (2.15)	15 (3.81)
HCV RNA at baseline, IU/mL [median (Q1, Q3)]	2220000 (803000, 6606934)	2340000 (693015, 5924209)
Biochemistry		
AST, IU/L, mean (SD)	57.91 (44.75)	46.38 (31.13)
ALT, IU/L, mean (SD)	56.63 (51.54)	51.52 (59.04)
Bilirubin < 2 mg/dL [n (%)]	393 (97.52)	359 (98.09)
Platelets, K/uL, median (Q1, Q3)	196.00 (155.50, 238.00)	209.00 (170.50, 255.00)
Hepatocellular carcinoma [n (%)]	4 (0.95)	2 (0.51)
Baseline diabetes [n (%)]	82 (19.57)	73 (18.53)
Baseline chronic kidney disease [n (%)]	41 (9.79)	26 (6.60)

HCV: Hepatitis C virus; AST: Aspartate aminotransferase; ALT: Alanine aminotransferase; HBV: Hepatitis B virus; HIV: human immunodeficiency virus.

Table 2 Transient elastography characteristics in patients with chronic hepatitis C according to treatment status

Variables	HCV treated (n = 419)	HCV untreated (n = 394)
Baseline liver stiffness (kPa) [mean (SD); range]	10.69 (9.48); 1.30-75.00	7.32 (6.10); 0.00-75.00
Change in kPa score [mean (SD)]	-2.78 (7.10)	0.51 (5.14)
Months between first and last TE [median (Q1, Q3)]	11.73 (7.41, 17.87)	12.68 (8.71, 16.94)
Months between first TE and DAA start [median (Q1, Q3)]	1.64 (0.95, 3.84)	-
Months between DAA end and last TE [median (Q1, Q3)]	4.17 (3.07, 10.96)	-

DAA: Direct-acting antiviral therapy; kPa: kilopascals; TE: Transient elastography

fibrosis. Interestingly, estimated liver stiffness as measured in kPa did not improve over a median of 11.7 mo among patients treated with DAA therapy. There was no significant difference in the change in liver stiffness between treated and untreated patients when stratifying patients based on underlying cirrhosis. Patients with longer times between TE measurements did not exhibit more pronounced differences in liver stiffness.

Our findings were unexpected due to abundant clinical evidence that initiation of antiviral treatment is associated with rapid improvement in liver stiffness scores as well as long-term attenuation of liver fibrosis in patients who achieve SVR[17]. It is possible that our study was underpowered to detect a

Table 3 Variables associated with the change in liver stiffness measurements in patients with chronic hepatitis C stratified by baseline kilopascals score

Variables	All patients (n = 770)			Baseline kPa < 12 (n = 651)			Baseline kPa ≥ 12 (n = 119)		
	Effect estimate	95%CI	P	Effect estimate	95%CI	P value	Effect estimate	95%CI	P value
Antiviral treatment effect	-1.821	(-3.452, -0.191)	0.029	-1.487	(-2.820, -0.154)	0.029	-2.071	(-7.477, 3.335)	0.453
Baseline kPa	-0.557	(-0.699, -0.415)	< 0.001	-0.677	(-0.875, -0.480)	< 0.001	-0.600	(-0.818, -0.383)	< 0.001
Black race ¹	-0.797	(-1.775, 0.181)	0.110	-0.654	(-1.362, 0.054)	0.070	-2.747	(-7.304, 1.810)	0.237
Other race ¹	-0.497	(-1.901, 0.907)	0.488	-0.687	(-1.893, 0.520)	0.264	1.071	(-6.423, 8.566)	0.779
Age ≥ 50 ²	0.663	(-0.165, 1.490)	0.117	0.949	(0.336, 1.563)	0.002	-0.204	(-4.089, 3.682)	0.918
Female gender	-0.210	(-0.843, 0.424)	0.517	-0.269	(-0.772, 0.235)	0.295	0.383	(-3.330, 4.096)	0.840
Diabetes	0.758	(-0.189, 1.705)	0.117	0.751	(-0.097, 1.599)	0.082	1.721	(-2.034, 5.475)	0.369
Current smoker	0.617	(-0.390, 1.624)	0.230	0.199	(-0.467, 0.865)	0.558	2.266	(-3.371, 7.903)	0.431
Former smoker	-0.258	(-1.155, 0.638)	0.572	-0.236	(-0.867, 0.395)	0.464	-0.554	(-6.741, 5.633)	0.861
HIV co-infected	0.468	(-0.230, 1.166)	0.189	0.254	(-0.286, 0.794)	0.356	2.040	(-1.292, 5.372)	0.230
Baseline ALT (per 10 units)	0.023	(-0.043, 0.089)	0.499	0.018	(-0.039, 0.075)	0.532	0.060	(-0.250, 0.370)	0.705
Time between first and last TE (untreated group) (mo)	0.001	(-0.090, 0.092)	0.975	-0.012	(-0.103, 0.078)	0.787	0.129	(-0.229, 0.488)	0.480
Time between first and last TE (treated group) (mo) ³	0.016	(-0.051, 0.084)	0.634	0.008	(-0.044, 0.059)	0.772	0.004	(-0.215, 0.223)	0.972
Interaction of treatment and time	0.015	(-0.106, 0.136)	0.807	0.020	(-0.086, 0.126)	0.711	-0.125	(-0.523, 0.273)	0.537
Intercept	4.146	(2.365, 5.927)	< 0.001	4.781	(2.478, 7.084)	< 0.001	5.852	(-0.994, 12.698)	0.094

kPa: kilopascals; ALT: Alanine aminotransferase; TE: Transient elastography; HIV: Human immunodeficiency virus;


¹Effect of race estimated relative to white race as a reference.

²Reference ages 18-49.

³Estimated effect of time between first and last TE (treated group) is not directly included as a covariate in the model. Estimates for this row are derived from the antiviral treatment, time between first and last TE, and interaction terms.

change in liver stiffness during a median follow-up time period of 4.2 mo after treatment. Since liver stiffness measurements were not always available at the initiation of DAA therapy, the effect of treatment may have been underestimated due to the inclusion of a median of 1.6 mo of untreated time in the treated group. Our study may have been underpowered to detect a modest effect of treatment on cirrhosis regression, as the analysis subsumed only 119 patients with cirrhosis at baseline. The apparent lack of improvement in hepatic stiffness among treated patients may be due to continued liver injury from other, potentially modifiable causes, including non-alcoholic fatty liver disease, alcohol use and drug use, which were not fully captured in our dataset. Despite successful treatment, a subset of patients cured of CHC, particularly those with advanced fibrosis at baseline, will exhibit a progression of liver fibrosis and are at risk of developing adverse liver-related outcomes[21,31]. Although we did not capture clinical data indicative of portal hypertension or decompensated cirrhosis in our study population, it has been postulated that decompensated cirrhosis represents an irreversible threshold that precludes regression to a noncirrhotic state[31,32].

In all analyses, higher baseline liver stiffness was independently associated with a reduction in kPa measurements between first and last TE, which is consistent with results from other investigations. In one systematic review and meta-analysis by Singh *et al*[17] of studies examining liver stiffness measurements after antiviral therapy, a greater absolute decrease in stiffness over time was observed



DOI: 10.12998/wjcc.v10.i17.5566 Copyright ©The Author(s) 2022.

Figure 1 Change in liver stiffness between first and last transient elastography measurement in patients with chronic hepatitis C infection. Scatter plot illustrating the difference in liver stiffness (kPa) between the first and last transient elastography measurement for patients who were treated ($n = 405$) and not treated ($n = 365$) with direct-acting antiviral therapy for chronic hepatitis C infection during the study period. Circles correspond to values for individual patients. kPa: Kilopascals; TE: Transient elastography.

among studies that included a larger proportion of patients with advanced fibrosis and cirrhosis at baseline. We found that antiviral treatment status was associated with a decline in kPa when analyzing the entire study population as well as patients without baseline cirrhosis. However, this estimated effect of treatment was independent of time, therefore, this does not represent the incremental decrease in stiffness that may be expected in successfully treated patients, and may instead reflect another confounder in our population. We cannot rule out the possibility that this observed decline in kPa is an artifact of regression towards the mean[33,34]. If there is random variability in the measurement of liver stiffness, then misclassifying patients with cirrhosis based on high baseline liver stiffness may result in observed declines in subsequent stiffness measurements that do not reflect a biological change in the liver[34,35]. We further found that treated patients have significantly higher baseline liver stiffness than untreated patients on average, which may make this effect stronger in the treated group than the untreated group. We included baseline kPa in our models to mitigate the impact of regression to the mean on the change over time in liver stiffness, but we cannot be certain that we have entirely accounted for this effect[35].

Based on our eligibility criteria, we identified 7 subjects who underwent serial TE measurements within 7 d or fewer. We performed a sensitivity analysis to investigate a potential bias introduced by our decision to exclude these subjects from the main analysis, based on the hypothesis that changes in liver stiffness observed in a short time period would be more likely to reflect measurement error than changes in inflammation or fibrosis due to DAA treatment. The sensitivity analysis demonstrated that DAA treatment did not significantly change liver stiffness when comparing treated *vs* untreated patients.

We found that age ≥ 50 was independently associated with increased liver stiffness, consistent with the natural history of fibrosis progression in CHC infection[36]. Baseline pre-treatment ALT levels were not predictive of a change in liver stiffness. However, other reports have identified a positive relationship between elevated ALT levels and improvement in liver stiffness over time as measured by TE, which may be partially due to a decline in necroinflammation[25,37].

Limitations of our study include missing data regarding SVR rates in patients initiating treatment with DAA. Although reported SVR rates are greater than 95% with DAA combinations, this approximation may dilute the effect of treatment on liver stiffness measurements in our study[38]. We were unable to assess the relationship between DAA treatment and liver-related outcomes in our cohort as these data were not collected as part of our study, however, achieving SVR has been demonstrated to lower the risk of liver-related morbidity and mortality[19,36,37].

The results from our study contribute to the clinical evidence regarding the real-world utility of post-treatment evaluation for fibrosis regression using TE[14,39]. We found that differences in liver stiffness may be observed on serial TE measurements in patients with higher baseline scores, irrespective of treatment effect, which suggests that false positive results on TE are not uncommon in clinical practice. The Baveno VI Consensus criteria includes a weak recommendation for obtaining two fasting TE measurements for patients undergoing screening for compensated advanced chronic liver disease[40]. Since liver stiffness measurements may be taken into account by both clinicians and patients as part of the decision to pursue a liver biopsy for definitive histopathologic assessment, and may impact clinical

management with respect to variceal bleeding surveillance, we advocate that a repeat confirmatory TE measurement should be considered for patients with an elevated kPa without other clinical or radiographic signs of advanced fibrosis.

CONCLUSION

Our study underscores the imperfect characteristics of TE for the assessment of liver fibrosis and cirrhosis, compared with the gold standard metrics of hepatic venous pressure gradient measurement and liver histopathology, despite the impetus to avoid invasive testing as part of routine clinical care of patients with CHC[41]. Ultimately, the question remains whether a decline in liver stiffness measurements after successful treatment with DAA is correlated with a reduced risk of liver-related complications, particularly among patients with advanced liver disease at baseline. Further long-term prospective studies must be performed to better understand the clinical utility of obtaining TE measurements in patients with CHC who have achieved SVR and to evaluate liver fibrosis progression in diverse populations, with incorporation of histological, hemodynamic, and clinical data on liver-related outcomes.

ARTICLE HIGHLIGHTS

Research background

Liver fibrosis is a common pathway of liver injury and is a feature of most chronic liver diseases. Fibrosis progression varies markedly in patients with hepatitis C virus (HCV), and the severity of liver fibrosis is associated with the prognosis of liver disease. Liver stiffness has been recommended as a parameter of fibrosis progression/regression in patients with HCV.

Research motivation

To investigate the changes in liver stiffness measured by transient elastography (TE) in a large, racially diverse cohort of U.S. patients with chronic hepatitis C (CHC).

Research objectives

We evaluated the differences in liver stiffness between patients treated with direct-acting antiviral (DAA) therapy and untreated patients. In addition, we performed a longitudinal, retrospective observational study investigating changes in liver stiffness measured by TE in a racially diverse cohort of United States patients with CHC.

Research methods

We conducted a longitudinal retrospective study of patients with confirmed CHC infection seen at Johns Hopkins Health System (JHHS) and Kaiser Permanente Mid-Atlantic States (KPMAS). Patients had ≥ 2 TE measurements and no prior DAA exposure. We used linear regression to measure the change in liver stiffness between first and last TE in response to treatment, controlling for age, sex, race, diabetes, smoking status, HIV status, baseline ALT, and baseline liver stiffness.

Research results

Of 813 patients, 84% were at least 50 years of age, 79% were Black, 79% were current or former smokers, 37% were coinfecting with HIV, 3% were coinfecting with HBV, 19% had diabetes, and 52% initiated treatment with a DAA. The median time between first and last TE was 11.7 and 12.7 mo among treated and untreated patients, respectively. There was no significant change in liver stiffness observed over time in either the group initiating DAA treatment (0.016 kPa/month; CI: -0.051, 0.084) or in the untreated group (0.001 kPa/month; CI: -0.090, 0.092), controlling for covariates. A higher baseline kPa score was independently associated with decreased liver stiffness.

Research conclusions

DAA treatment was not associated with a differential change in liver stiffness over time, as measured by TE, in patients with CHC compared to untreated patients. Our study underscores the imperfect characteristics of any single noninvasive test for assessing liver fibrosis, which continues to be compared to the gold standard of liver biopsy and histopathology, despite the impetus to avoid invasive testing for CHC infection in clinical practice.

Research perspectives

Direct-acting antiviral therapy was not associated with a differential change in liver stiffness over time in patients with CHC compared to untreated patients. Further longitudinal prospective studies are

needed to evaluate the clinical utility of obtaining TE measurements in patients with CHC who have achieved sustained virologic response and assess liver fibrosis progression in diverse populations.

FOOTNOTES

Author contributions: Tinsay A Woreta and Po-Hung Chen conceptualized the study; Anya Mezina, Tinsay A Woreta and Po-Hung Chen investigated the study; Anya Mezina, Tinsay A Woreta, Kevin B Rubenstein Po-Hung Chen and Carla Rodriguez-Watson did the methodology; Anya Mezina did the writing - original draft; Arunkumar Krishnan, Tinsay A Woreta, Kevin B Rubenstein Po-Hung Chen and Carla Rodriguez-Watson did the writing-review and editing; Kevin B Rubenstein did the software and visualization; Tinsay A Woreta and Eric Watson did the data curation; Arunkumar Krishnan did the validation.

Supported by the National Center for Advancing Translational Sciences, No. 5KL2TR001077-05 (to Po-Hung Chen).

Institutional review board statement: The study was reviewed and approved for publication by the Institutional Review Board of Johns Hopkins Medicine and Kaiser Permanente Mid-Atlantic States.

Informed consent statement: The need for informed consent was waived due to the retrospective nature of the study.

Conflict-of-interest statement: All the Authors have no conflict of interest related to the manuscript.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: United States

ORCID number: Anya Mezina [0000-0002-1515-4358](https://orcid.org/0000-0002-1515-4358); Arunkumar Krishnan [0000-0002-9452-7377](https://orcid.org/0000-0002-9452-7377); Tinsay A Woreta [0000-0001-7292-4518](https://orcid.org/0000-0001-7292-4518); Kevin B Rubenstein [0000-0003-3516-5932](https://orcid.org/0000-0003-3516-5932); Eric Watson [0000-0002-1917-2748](https://orcid.org/0000-0002-1917-2748); Po-Hung Chen [0000-0002-9865-1009](https://orcid.org/0000-0002-9865-1009); Carla Rodriguez-Watson [0000-0003-4898-1613](https://orcid.org/0000-0003-4898-1613).

S-Editor: Xing YX

L-Editor: A

P-Editor: Xing YX

REFERENCES

- 1 **Perz JF**, Armstrong GL, Farrington LA, Hutin YJ, Bell BP. The contributions of hepatitis B virus and hepatitis C virus infections to cirrhosis and primary liver cancer worldwide. *J Hepatol* 2006; **45**: 529-538 [PMID: [16879891](https://pubmed.ncbi.nlm.nih.gov/16879891/) DOI: [10.1016/j.jhep.2006.05.013](https://doi.org/10.1016/j.jhep.2006.05.013)]
- 2 **Polaris Observatory HCV Collaborators**. Global prevalence and genotype distribution of hepatitis C virus infection in 2015: a modelling study. *Lancet Gastroenterol Hepatol* 2017; **2**: 161-176 [PMID: [28404132](https://pubmed.ncbi.nlm.nih.gov/28404132/) DOI: [10.1016/S2468-1253\(16\)30181-9](https://doi.org/10.1016/S2468-1253(16)30181-9)]
- 3 **Singh S**, Fujii LL, Murad MH, Wang Z, Asrani SK, Ehman RL, Kamath PS, Talwalkar JA. Liver stiffness is associated with risk of decompensation, liver cancer, and death in patients with chronic liver diseases: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2013; **11**: 1573-84.e1 [PMID: [23954643](https://pubmed.ncbi.nlm.nih.gov/23954643/) DOI: [10.1016/j.cgh.2013.07.034](https://doi.org/10.1016/j.cgh.2013.07.034)]
- 4 **Bedossa P**, Poynard T. An algorithm for the grading of activity in chronic hepatitis C. The METAVIR Cooperative Study Group. *Hepatology* 1996; **24**: 289-293 [PMID: [8690394](https://pubmed.ncbi.nlm.nih.gov/8690394/) DOI: [10.1002/hep.510240201](https://doi.org/10.1002/hep.510240201)]
- 5 **Thein HH**, Yi Q, Dore GJ, Krahn MD. Estimation of stage-specific fibrosis progression rates in chronic hepatitis C virus infection: a meta-analysis and meta-regression. *Hepatology* 2008; **48**: 418-431 [PMID: [18563841](https://pubmed.ncbi.nlm.nih.gov/18563841/) DOI: [10.1002/hep.22375](https://doi.org/10.1002/hep.22375)]
- 6 **Marcellin P**, Asselah T, Boyer N. Fibrosis and disease progression in hepatitis C. *Hepatology* 2002; **36**: S47-S56 [PMID: [12407576](https://pubmed.ncbi.nlm.nih.gov/12407576/) DOI: [10.1053/jhep.2002.36993](https://doi.org/10.1053/jhep.2002.36993)]
- 7 **Hoefs JC**, Shiffman ML, Goodman ZD, Kleiner DE, Dienstag JL, Stoddard AM; HALT-C Trial Group. Rate of progression of hepatic fibrosis in patients with chronic hepatitis C: results from the HALT-C Trial. *Gastroenterology* 2011; **141**: 900-908.e1 [PMID: [21699796](https://pubmed.ncbi.nlm.nih.gov/21699796/) DOI: [10.1053/j.gastro.2011.06.007](https://doi.org/10.1053/j.gastro.2011.06.007)]
- 8 **Konerman MA**, Mehta SH, Sutcliffe CG, Vu T, Higgins Y, Torbenson MS, Moore RD, Thomas DL, Sulkowski MS. Fibrosis progression in human immunodeficiency virus/hepatitis C virus coinfecting adults: prospective analysis of 435 liver biopsy pairs. *Hepatology* 2014; **59**: 767-775 [PMID: [24436062](https://pubmed.ncbi.nlm.nih.gov/24436062/) DOI: [10.1002/hep.26741](https://doi.org/10.1002/hep.26741)]
- 9 **Patel S**, Jinjuvadia R, Patel R, Liangpunsakul S. Insulin Resistance is Associated With Significant Liver Fibrosis in Chronic Hepatitis C Patients: A Systemic Review and Meta-Analysis. *J Clin Gastroenterol* 2016; **50**: 80-84 [PMID: [26302498](https://pubmed.ncbi.nlm.nih.gov/26302498/) DOI: [10.1097/MCG.0000000000000400](https://doi.org/10.1097/MCG.0000000000000400)]

- 10 **McHutchison J**, Poynard T, Afdhal N. Fibrosis as an end point for clinical trials in liver disease: a report of the international fibrosis group. *Clin Gastroenterol Hepatol* 2006; **4**: 1214-1220 [PMID: 16979947 DOI: 10.1016/j.cgh.2006.07.006]
- 11 **Maharaj B**, Maharaj RJ, Leary WP, Cooppan RM, Naran AD, Pirie D, Pudifin DJ. Sampling variability and its influence on the diagnostic yield of percutaneous needle biopsy of the liver. *Lancet* 1986; **1**: 523-525 [PMID: 2869260 DOI: 10.1016/s0140-6736(86)90883-4]
- 12 **Regev A**, Berho M, Jeffers LJ, Milikowski C, Molina EG, Pyrsopoulos NT, Feng ZZ, Reddy KR, Schiff ER. Sampling error and intraobserver variation in liver biopsy in patients with chronic HCV infection. *Am J Gastroenterol* 2002; **97**: 2614-2618 [PMID: 12385448 DOI: 10.1111/j.1572-0241.2002.06038.x]
- 13 **Colloredo G**, Guido M, Sonzogni A, Leandro G. Impact of liver biopsy size on histological evaluation of chronic viral hepatitis: the smaller the sample, the milder the disease. *J Hepatol* 2003; **39**: 239-244 [PMID: 12873821 DOI: 10.1016/s0168-8278(03)00191-0]
- 14 **Lim JK**, Flamm SL, Singh S, Falck-Ytter YT; Clinical Guidelines Committee of the American Gastroenterological Association. American Gastroenterological Association Institute Guideline on the Role of Elastography in the Evaluation of Liver Fibrosis. *Gastroenterology* 2017; **152**: 1536-1543 [PMID: 28442119 DOI: 10.1053/j.gastro.2017.03.017]
- 15 **Poynard T**, McHutchison J, Manns M, Trepo C, Lindsay K, Goodman Z, Ling MH, Albrecht J. Impact of pegylated interferon alfa-2b and ribavirin on liver fibrosis in patients with chronic hepatitis C. *Gastroenterology* 2002; **122**: 1303-1313 [PMID: 11984517 DOI: 10.1053/gast.2002.33023]
- 16 **D'Ambrosio R**, Aghemo A, Fraquelli M, Rumi MG, Donato MF, Paradis V, Bedossa P, Colombo M. The diagnostic accuracy of Fibroscan for cirrhosis is influenced by liver morphometry in HCV patients with a sustained virological response. *J Hepatol* 2013; **59**: 251-256 [PMID: 23528378 DOI: 10.1016/j.jhep.2013.03.013]
- 17 **Singh S**, Facciorusso A, Loomba R, Falck-Ytter YT. Magnitude and Kinetics of Decrease in Liver Stiffness After Antiviral Therapy in Patients With Chronic Hepatitis C: A Systematic Review and Meta-analysis. *Clin Gastroenterol Hepatol* 2018; **16**: 27-38.e4 [PMID: 28479504 DOI: 10.1016/j.cgh.2017.04.038]
- 18 **Singal AG**, Volk ML, Jensen D, Di Bisceglie AM, Schoenfeld PS. A sustained viral response is associated with reduced liver-related morbidity and mortality in patients with hepatitis C virus. *Clin Gastroenterol Hepatol* 2010; **8**: 280-288, 288.e1 [PMID: 19948249 DOI: 10.1016/j.cgh.2009.11.018]
- 19 **Backus LI**, Boothroyd DB, Phillips BR, Belperio P, Halloran J, Mole LA. A sustained virologic response reduces risk of all-cause mortality in patients with hepatitis C. *Clin Gastroenterol Hepatol* 2011; **9**: 509-516.e1 [PMID: 21397729 DOI: 10.1016/j.cgh.2011.03.004]
- 20 **Chekuri S**, Nickerson J, Bichoupan K, Sefcik R, Doobay K, Chang S, DelBello D, Harty A, Dieterich DT, Perumalswami PV, Branch AD. Liver Stiffness Decreases Rapidly in Response to Successful Hepatitis C Treatment and Then Plateaus. *PLoS One* 2016; **11**: e0159413 [PMID: 27442255 DOI: 10.1371/journal.pone.0159413]
- 21 **van der Meer AJ**, Feld JJ, Hofer H, Almasio PL, Calvaruso V, Fernández-Rodríguez CM, Aleman S, Ganne-Carrié N, D'Ambrosio R, Pol S, Trapero-Marugan M, Maan R, Moreno-Otero R, Mallet V, Hultcrantz R, Weiland O, Rutter K, Di Marco V, Alonso S, Bruno S, Colombo M, de Knecht RJ, Veldt BJ, Hansen BE, Janssen HLA. Risk of cirrhosis-related complications in patients with advanced fibrosis following hepatitis C virus eradication. *J Hepatol* 2017; **66**: 485-493 [PMID: 27780714 DOI: 10.1016/j.jhep.2016.10.017]
- 22 **Pietsch V**, Deterding K, Attia D, Ringe KI, Heidrich B, Cornberg M, Gebel M, Manns MP, Wedemeyer H, Potthoff A. Long-term changes in liver elasticity in hepatitis C virus-infected patients with sustained virologic response after treatment with direct-acting antivirals. *United European Gastroenterol J* 2018; **6**: 1188-1198 [PMID: 30288281 DOI: 10.1177/2050640618786067]
- 23 **AASLD-IDSAs**. HCV testing and linkage to care. Recommendations for testing, managing, and treating hepatitis C. [cited 2018 Nov 21] Available from: <https://www.hcvguidelines.org/evaluate/when-whom>
- 24 **Bernuth S**, Yagmur E, Schuppan D, Sprinzl MF, Zimmermann A, Schad A, Kittner JM, Weyer V, Knapstein J, Schattenberg JM, Wörms MA, Galle PR, Zimmermann T. Early changes in dynamic biomarkers of liver fibrosis in hepatitis C virus-infected patients treated with sofosbuvir. *Dig Liver Dis* 2016; **48**: 291-297 [PMID: 26514736 DOI: 10.1016/j.dld.2015.09.015]
- 25 **Sporea I**, Lupuşoru R, Mare R, Popescu A, Gheorghe L, Iacob S, Şirli R. Dynamics of liver stiffness values by means of transient elastography in patients with HCV liver cirrhosis undergoing interferon free treatment. *J Gastrointest Liver Dis* 2017; **26**: 145-150 [PMID: 28617884 DOI: 10.15403/jgld.2014.1121.262.dyn]
- 26 **Knop V**, Hoppe D, Welzel T, Vermehren J, Herrmann E, Vermehren A, Friedrich-Rust M, Sarrazin C, Zeuzem S, Welker MW. Regression of fibrosis and portal hypertension in HCV-associated cirrhosis and sustained virologic response after interferon-free antiviral therapy. *J Viral Hepat* 2016; **23**: 994-1002 [PMID: 27500382 DOI: 10.1111/jvh.12578]
- 27 **Elsharkawy A**, Alem SA, Fouad R, El Raziky M, El Akel W, Abdo M, Tantawi O, AbdAllah M, Bourliere M, Esmat G. Changes in liver stiffness measurements and fibrosis scores following sofosbuvir based treatment regimens without interferon. *J Gastroenterol Hepatol* 2017; **32**: 1624-1630 [PMID: 28177543 DOI: 10.1111/jgh.13758]
- 28 **Bachofner JA**, Valli PV, Kröger A, Bergamin I, Künzler P, Baserga A, Braun D, Seifert B, Moncsek A, Fehr J, Semela D, Magenta L, Müllhaupt B, Terziroli Beretta-Piccoli B, Mertens JC. Direct antiviral agent treatment of chronic hepatitis C results in rapid regression of transient elastography and fibrosis markers fibrosis-4 score and aspartate aminotransferase-platelet ratio index. *Liver Int* 2017; **37**: 369-376 [PMID: 27678216 DOI: 10.1111/liv.13256]
- 29 **Jacobson IM**, Lim JK, Fried MW. American Gastroenterological Association Institute Clinical Practice Update-Expert Review: Care of Patients Who Have Achieved a Sustained Virologic Response After Antiviral Therapy for Chronic Hepatitis C Infection. *Gastroenterology* 2017; **152**: 1578-1587 [PMID: 28344022 DOI: 10.1053/j.gastro.2017.03.018]
- 30 **Garcia-Tsao G**. Regression of HCV cirrhosis: Time will tell. *Hepatology* 2018; **67**: 1651-1653 [PMID: 29222930 DOI: 10.1002/hep.29720]
- 31 **Labarga P**, Fernandez-Montero JV, de Mendoza C, Barreiro P, Pinilla J, Soriano V. Liver fibrosis progression despite HCV cure with antiviral therapy in HIV-HCV-coinfected patients. *Antivir Ther* 2015; **20**: 329-334 [PMID: 25372299 DOI: 10.3851/IMP2909]

- 32 Garcia-Tsao G. Regression of HCV cirrhosis: Time will tell. *Hepatology* 2018; **67**: 1651-1653 [PMID: 29222930 DOI: 10.1002/hep.29720]
- 33 Joy KP, Agha AK. Seasonal effects of administration of melatonin and 5-methoxytryptophol on ovarian activity in the catfish *Heteropneustes fossilis* (Bloch). *J Pineal Res* 1991; **10**: 65-70 [PMID: 2056435 DOI: 10.1002/hep.23744]
- 34 Fleming S. Statistical notes. Regression towards the mean. *BMJ* 1994; **309**: 539 [PMID: 8086922]
- 35 Barnett AG, van der Pols JC, Dobson AJ. Regression to the mean: what it is and how to deal with it. *Int J Epidemiol* 2005; **34**: 215-220 [PMID: 15333621 DOI: 10.1093/ije/dyh299]
- 36 Lingala S, Ghany MG. Natural History of Hepatitis C. *Gastroenterol Clin North Am* 2015; **44**: 717-734 [PMID: 26600216 DOI: 10.1016/j.gtc.2015.07.003]
- 37 Martinez SM, Foucher J, Combis JM, Métivier S, Brunetto M, Capron D, Bourlière M, Bronowicki JP, Dao T, Maynard-Muet M, Lucidarme D, Merrouche W, Forns X, de Ledinghen V. Longitudinal liver stiffness assessment in patients with chronic hepatitis C undergoing antiviral therapy. *PLoS One* 2012; **7**: e47715 [PMID: 23082200 DOI: 10.1371/journal.pone.0047715]
- 38 Nahon P, Bourcier V, Layese R, Audureau E, Cagnot C, Marcellin P, Guyader D, Fontaine H, Larrey D, De Ledinghen V, Ouzan D, Zoulim F, Roulot D, Tran A, Bronowicki JP, Zarski JP, Leroy V, Riachi G, Calès P, Péron JM, Alric L, Bourlière M, Mathurin P, Dharancy S, Blanc JF, Abergel A, Serfaty L, Mallat A, Grangé JD, Attali P, Bacq Y, Wartelle C, Dao T, Benhamou Y, Pilette C, Silvain C, Christidis C, Capron D, Bernard-Chabert B, Zucman D, Di Martino V, Thibaut V, Salmon D, Ziol M, Sutton A, Pol S, Roudot-Thoraval F; ANRS CO12 CirVir Group. Eradication of Hepatitis C Virus Infection in Patients With Cirrhosis Reduces Risk of Liver and Non-Liver Complications. *Gastroenterology* 2017; **152**: 142-156.e2 [PMID: 27641509 DOI: 10.1053/j.gastro.2016.09.009]
- 39 Cheng PN, Chiu HC, Chiu YC, Chen SC, Chen Y. Comparison of FIB-4 and transient elastography in evaluating liver fibrosis of chronic hepatitis C subjects in community. *PLoS One* 2018; **13**: e0206947 [PMID: 30403744 DOI: 10.1371/journal.pone.0206947]
- 40 de Franchis R; Baveno VI Faculty. Expanding consensus in portal hypertension: Report of the Baveno VI Consensus Workshop: Stratifying risk and individualizing care for portal hypertension. *J Hepatol* 2015; **63**: 743-752 [PMID: 26047908 DOI: 10.1016/j.jhep.2015.05.022]
- 41 Castera L, Pinzani M, Bosch J. Non invasive evaluation of portal hypertension using transient elastography. *J Hepatol* 2012; **56**: 696-703 [PMID: 21767510 DOI: 10.1016/j.jhep.2011.07.005]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

