

World Journal of *Clinical Cases*

World J Clin Cases 2022 February 16; 10(5): 1457-1753



Contents

Thrice Monthly Volume 10 Number 5 February 16, 2022

REVIEW

- 1457 Nonalcoholic fatty liver disease shows significant sex dimorphism
Chen XY, Wang C, Huang YZ, Zhang LL

MINIREVIEWS

- 1473 Management of procedural pain in the intensive care unit
Guo NN, Wang HL, Zhao MY, Li JG, Liu HT, Zhang TX, Zhang XY, Chu YJ, Yu KJ, Wang CS

ORIGINAL ARTICLE

Clinical and Translational Research

- 1485 Effect of prior malignancy on the prognosis of gastric cancer and somatic mutation
Yin X, He XK, Wu LY, Yan SX

Retrospective Cohort Study

- 1498 Elemene-containing hyperthermic intraperitoneal chemotherapy combined with chemotherapy for elderly patients with peritoneal metastatic advanced gastric cancer
Chen ZX, Li J, Liu WB, Zhang SR, Sun H

Retrospective Study

- 1508 Timing theory continuous nursing, resistance training: Rehabilitation and mental health of caregivers and stroke patients with traumatic fractures
Shen YL, Zhang ZQ, Zhu LJ, Liu JH
- 1517 Effect of precise nursing service mode on postoperative urinary incontinence prevention in patients with prostate disease
Zheng XC, Luo TT, Cao DD, Cai WZ

- 1527 Significance of serum glucagon-like peptide-1 and matrix Gla protein levels in patients with diabetes and osteoporosis
Xie FF, Zhang YF, Hu YF, Xie YY, Wang XY, Wang SZ, Xie BQ

- 1536 Castleman disease and TAFRO syndrome: To improve the diagnostic consciousness is the key
Zhou QY

Observational Study

- 1548 Correlation of myopia onset and progression with corneal biomechanical parameters in children
Lu LL, Hu XJ, Yang Y, Xu S, Yang SY, Zhang CY, Zhao QY

META-ANALYSIS

- 1557** Intensive *vs* non-intensive statin pretreatment before percutaneous coronary intervention in Chinese patients: A meta-analysis of randomized controlled trials

Yang X, Lan X, Zhang XL, Han ZL, Yan SM, Wang WX, Xu B, Ge WH

CASE REPORT

- 1572** Giant nodular fasciitis originating from the humeral periosteum: A case report
Yu SL, Sun PL, Li J, Jia M, Gao HW
- 1580** Tumor-related cytokine release syndrome in a treatment-naïve patient with lung adenocarcinoma: A case report
Deng PB, Jiang J, Hu CP, Cao LM, Li M
- 1586** Submucosal protuberance caused by a fish bone in the absence of preoperative positive signs: A case report
Du WW, Huang T, Yang GD, Zhang J, Chen J, Wang YB
- 1592** Misdiagnosis of unroofed coronary sinus syndrome as an ostium primum atrial septal defect by echocardiography: A case report
Chen JL, Yu CG, Wang DJ, Chen HB
- 1598** Uncommon complication of nasoenteral feeding tube: A case report
Jiang YP, Zhang S, Lin RH
- 1602** Treatment of extracranial internal carotid artery dissecting aneurysm with SUPERA stent implantation: Two case reports
Qiu MJ, Zhang BR, Song SJ
- 1609** Combination of atezolizumab and chidamide to maintain long-term remission in refractory metastatic extranodal natural killer/T-cell lymphoma: A case report
Wang J, Gao YS, Xu K, Li XD
- 1617** Hemangioma in the lower labial vestibule of an eleven-year-old girl: A case report
Aloyouny AY, Alfaifi AJ, Aladhyani SM, Alshalan AA, Alfayadh HM, Salem HM
- 1623** Primary orbital monophasic synovial sarcoma with calcification: A case report
Ren MY, Li J, Li RM, Wu YX, Han RJ, Zhang C
- 1630** Small-cell carcinoma of the prostate with negative CD56, NSE, Syn, and CgA indicators: A case report
Shi HJ, Fan ZN, Zhang JS, Xiong BB, Wang HF, Wang JS
- 1639** Disseminated peritoneal leiomyomatosis with malignant transformation involving right ureter: A case report
Wen CY, Lee HS, Lin JT, Yu CC

- 1645** Arthroscopic surgery for synovial chondroma of the subacromial bursa with non-traumatic shoulder subluxation complications: Two case reports
Tang XF, Qin YG, Shen XY, Chen B, Li YZ
- 1654** Wilkie's syndrome as a cause of anxiety-depressive disorder: A case report and review of literature
Apostu RC, Chira L, Colcear D, Lebovici A, Nagy G, Scurtu RR, Drasovean R
- 1667** Gastric schwannoma misdiagnosed as gastrointestinal stromal tumor by ultrasonography before surgery: A case report
Li QQ, Liu D
- 1675** Giant retroperitoneal lipoma presenting with abdominal distention: A case report and review of the literature
Chen ZY, Chen XL, Yu Q, Fan QB
- 1684** Pneumothorax during retroperitoneal laparoscopic partial nephrectomy in a lupus nephritis patient: A case report
Zhao Y, Xue XQ, Xia D, Xu WF, Liu GH, Xie Y, Ji ZG
- 1689** Bulbar conjunctival vascular lesion combined with spontaneous retrobulbar hematoma: A case report
Lei JY, Wang H
- 1697** Hepatitis B virus in cerebrospinal fluid of a patient with purulent bacterial meningitis detected by multiplex-PCR: A case report
Gao DQ, Hu YQ, Wang X, Zhang YZ
- 1702** Aseptic abscess in the abdominal wall accompanied by monoclonal gammopathy simulating the local recurrence of rectal cancer: A case report
Yu Y, Feng YD, Zhang C, Li R, Tian DA, Huang HJ
- 1709** Tacrolimus treatment for relapsing-remitting chronic inflammatory demyelinating polyradiculoneuropathy: Two case reports
Zhu WJ, Da YW, Chen H, Xu M, Lu Y, Di L, Duo JY
- 1716** Vedolizumab-associated diffuse interstitial lung disease in patients with ulcerative colitis: A case report
Zhang J, Liu MH, Gao X, Dong C, Li YX
- 1723** Unusual magnetic resonance imaging findings of brain and leptomeningeal metastasis in lung adenocarcinoma: A case report
Li N, Wang YJ, Zhu FM, Deng ST
- 1729** Diffuse invasive signet ring cell carcinoma in total colorectum caused by ulcerative colitis: A case report and review of literature
Zhang Z, Yu PF, Gu GL, Zhang YH, Wang YM, Dong ZW, Yang HR
- 1738** Neurothekeoma located in the hallux and axilla: Two case reports
Huang WY, Zhang YQ, Yang XH

- 1747 Subclavian artery stenting *via* bilateral radial artery access: Four case reports

Qiu T, Fu SQ, Deng XY, Chen M, Dai XY

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Prashanth Panta, MDS, Reader (Associate Professor), Department of Oral Medicine and Radiology, Malla Reddy Institute of Dental Sciences, Suraram 500055, Telangana, India. maithreya.prashanth@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJCC as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The WJCC's CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Lin-YuTong Wang; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

February 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Giant retroperitoneal lipoma presenting with abdominal distention: A case report and review of the literature

Zhi-Yan Chen, Xian-Long Chen, Qi Yu, Qing-Bo Fan

ORCID number: Zhi-Yan Chen 0000-0002-1695-1455; Xian-Long Chen 0000-0002-3677-4281; Qi Yu 0000-0001-9737-5957; Qing-Bo Fan 0000-0002-8745-1132.

Author contributions: Chen ZY wrote and revised the manuscript; Chen ZY and Fan QB were part of the clinical team that treated the patient; Chen XL participated in the review of the pathology; Yu Q and Fan QB revised the manuscript and supervised the study.

Informed consent statement:

Informed written consent was obtained from the patient for publication of this case report and any accompanying images.

Conflict-of-interest statement: No conflict of interests exists to any of the authors.

CARE Checklist (2016) statement:

The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Supported by the National Key Research and Development Program, No. 2018YFC1002105.

Country/Territory of origin: China

Specialty type: Obstetrics and gynecology

Zhi-Yan Chen, Qi Yu, Qing-Bo Fan, Department of Obstetrics and Gynecology, Peking Union Medical College Hospital, Beijing 100730, China

Xian-Long Chen, Department of Pathology, Peking Union Medical College Hospital, Beijing 100730, China

Corresponding author: Qing-Bo Fan, MD, Professor, Department of Obstetrics and Gynecology, Peking Union Medical College Hospital, No. 1 Shuaifuyuan, Wangfujing, Dongcheng District, Beijing 100730, China. qbfan@sohu.com

Abstract

BACKGROUND

Retroperitoneal lipomas are extremely rare tumors and tend to be large in size (> 10 cm) when diagnosed, causing various clinical manifestations. Preoperative diagnosis of retroperitoneal lipomas is difficult. There is a lack of relevant information about the management and prognosis of these benign tumors due to limited reports.

CASE SUMMARY

A 53-year-old woman who complained about progressive abdominal distention and aggravating satiety was referred to the gynecological outpatient department of Peking Union Medical College Hospital. Computerized tomography (CT) revealed an immense mass with fat density, measuring 28.6 cm× 16.6 cm in size. Adjacent organs, including the intestinal tract and uterus, were squeezed to the right side of the abdomen. An exploratory laparotomy was performed with suspicion of liposarcoma. Intraoperatively, a giant yellowish lobulated mass was found occupying the retroperitoneum and it was removed by tumor debulking. Postoperative histopathological results confirmed the diagnosis of retroperitoneal lipoma.

CONCLUSION

Retroperitoneal lipoma is a very rare condition and is difficult to differentiate from well-differentiated liposarcoma. Radiographic investigations, especially CT and magnetic resonance imaging, are important for preoperative diagnosis. Surgical resection is the fundamental treatment, which is difficult due to its size and relation to neighboring structures.

Key Words: Retroperitoneal lipoma; Well-differentiated liposarcoma; Retroperitoneal

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind**Peer-review report's scientific quality classification**

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Received: August 21, 2021

Peer-review started: August 21, 2021

First decision: November 17, 2021

Revised: December 8, 2021

Accepted: January 8, 2022

Article in press: January 8, 2022

Published online: February 16, 2022

P-Reviewer: Dhali A, Meglio LD

S-Editor: Wang JL

L-Editor: Wang TQ

P-Editor: Wang JL



tumors; Treatment; Prognosis; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Retroperitoneal lipomas are rare benign tumors originating from adipose tissues and they tend to have large sizes. Imaging examinations, especially computerized tomography and magnetic resonance imaging, are fundamental diagnostic tools for these tumors. Surgical resection is the main treatment method. *En bloc* resection is commonly required. Postoperative histopathology determines the final diagnosis, and immunohistochemical analysis could be useful in the differentiation of liposarcomas. Regular follow-ups are also required for the patients.

Citation: Chen ZY, Chen XL, Yu Q, Fan QB. Giant retroperitoneal lipoma presenting with abdominal distention: A case report and review of the literature. *World J Clin Cases* 2022; 10(5): 1675-1683

URL: <https://www.wjgnet.com/2307-8960/full/v10/i5/1675.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i5.1675>

INTRODUCTION

Lipomas are a category of benign tumors originating from well-differentiated adipocytes. Their predilection sites are subdermal tissues of the trunk and extremities [1,2]. Retroperitoneal lipomas are a rare condition, with just 22 case reports describing the tumour in adults in the previous literature of PubMed since 1970[3-22]. They account for 2.9% of primary retroperitoneal tumors, approximately 80% of which are known as malignant[6,23,24]. Unlike subcutaneous lipomas, which are related to obesity, hyperlipidaemia, and injuries, retroperitoneal lipomas have an unknown aetiology[5,25-27]. On account of the rarity and limited knowledge of these tumors, further reports and investigations are necessary. In this report, we describe the case of a 53-year-old postmenopausal woman who presented with a massive fatty retroperitoneal mass measuring 28.6 cm× 16.6 cm and weighing 7.126 kg.

CASE PRESENTATION

Chief complaints

A 53-year-old postmenopausal woman who complained about progressive abdominal distention and aggravating satiety was referred to the department of gynaecology in our center.

History of present illness

The patient started to feel intensifying abdominal distention and satiety for the last 2 mo. She also found a significant increase in abdominal circumference and thinning of the limbs. The patient denied other discomforts, including fever, abdominal pain, nausea, and vomiting. The patient had been postmenopausal for 5 years and did not report abnormal vaginal bleeding.

History of past illness

The patient had a 10-year history of hypertension and took Loxone once per day, with stable control of blood pressure. The patient denied any history of diabetes, coronary heart disease, or malignancy. She also reported no drug allergy or other physical impairment. Additionally, the patient did not receive regular physical examinations, and the last medical examination had occurred more than 10 years prior.

Personal and family history

No noteworthy personal or family history was reported by the patient.

Physical examination

The patient's height was 161 cm, and her weight was 60 kg (body mass index: 23.3, within the normal range). The physical examination revealed a palpable giant abdominal mass reaching the xiphisternum with a rubbery consistency. Other clinical symptoms, including tenderness, rebound tenderness, and mobile turbid sounds, were found.

Laboratory examinations

After hospitalization, the patient received a series of laboratory examinations for testing liver and kidney function, faecal occult blood, blood coagulation factors, electrolyte panel, and tumor biomarkers. The laboratory findings fell within the normal range.

Imaging examinations

The patient had received a computed tomography (CT) plain scan at another hospital, and was re-evaluated by ultrasonography at our hospital. A massive hyperechoic mass, approximately 30 cm × 17 cm in size, was visualized *via* ultrasound. The mass was clearly defined and had internal echogenicity, filled with stripe-like structures. Minimal blood signals were detected by colour Doppler ultrasound (Figure 1). A CT plain scan demonstrated a giant homogeneous mass mainly consisting of fatty tissue and thin septa. It measured 16.6 cm × 28.6 cm in volume and pushed the peritoneal contents, such as the bowel loops and uterus, to the right part of the abdomen (Figure 2).

FINAL DIAGNOSIS

Based on the clinical manifestations, normal laboratory examinations, and imaging examinations indicating its adipose origin, the clinicians considered the mass to be a giant retroperitoneal lipoma. However, the possibility of malignancy cannot be overlooked due to its large size.

TREATMENT

After completing the examinations and preoperative assessments, the patient underwent an exploratory laparotomy with the suspicion of malignancy, most likely retroperitoneal liposarcoma. During the operation, a bulky yellowish tumor originating from perirenal fatty tissues in the left retroperitoneal region was found to occupy the retroperitoneum. The uterus and adnexa were displaced by the mass. The mass adhered to the left psoas major muscle and wrapped around the left ureter, making it unfeasible to perform *en bloc* resection. After carefully separating the left ureter, we performed tumor debulking and resection of the left adnexa, which was also tightly adhered to the tumor. The total weight of the mass was 7.126 kg (Figure 3). The frozen pathological results suggested that the mass mainly consisted of adipose tissues, and a retroperitoneal lipomatous tumor was considered.

After 1 wk of uneventful hospitalization, the patient was discharged from the hospital with full recovery from her clinical symptoms. The final paraffin pathology showed that the tumor was composed of mature adipose tissues and hematopoietic cells, without cytologic atypia, and confirmed the diagnosis of multiple lipomas and multiple myelolipomas (Figure 4).

OUTCOME AND FOLLOW-UP

The patient reported no relevant clinical symptoms after the operation. During a series of follow-ups for 18 mo, the laboratory tests and imaging examinations were normal and indicated no signs of relapse.

DISCUSSION

We herein report a massive retroperitoneal lipoma, which consisted of multiple




Figure 1 Abdominal ultrasonography of the mass. A giant hyperechoic mass filling the abdomen was presented on grey-scale ultrasound. The mass had a relative clear margin and internal septas.




Figure 2 Abdominal computed tomography in the axial plane. Computed tomography imaging showed a giant homogenous mass, mainly consisting of fatty tissue measuring 16.6 cm × 28.6 cm with thin septa, pushing the peritoneal containing such as bowel loops and uterus to the right part of abdomen.

conventional lipomas and multiple myelolipomas. Retroperitoneal lipomas are rare mesenchymal-originated tumors. It was first reported in 1947[22], and since then, a total of 22 cases have been reported in adults sporadically (Table 1). The peak incidence of adult retroperitoneal lipomas occurs between the ages of 40 and 60, with no discernible gender predisposition. According to the morphologic characteristics, lipomas can be subdivided into conventional lipoma, fibrolipoma, angioliipoma, spindle cell lipoma, pleomorphic lipoma, and myelolipoma[7], of which, almost all myelolipomas have been identified inside the adrenal gland, with just about 50 cases of myelolipomas being identified in extra-adrenal locations, such as the retroperitoneum[28]. The exact underlying aetiology of retroperitoneal lipomas is not well understood. Seeding after fibroid excision, exogenous hormone treatment, or chronic abnormalities in glucose homeostasis have all been blamed for these benign tumors. And genetic factors are thought to have an important role in adipocyte proliferation[5].

Retroperitoneal tumors are often asymptomatic for a long period of time throughout their early clinical course, owing to the vast potential spaces in the retroperitoneum. Local compression of surrounding organs and tissues, which can manifest obstructive urinary/bowel symptoms such as stomach pain, fullness, early satiety, or lower extremity oedema, may occur once the tumors have grown to gigantic sizes. The

Table 1 Summary of all case reports describing retroperitoneal lipomas resected in adults

Ref.	Age	Sex	Symptoms	Imaging methods	Tumor size (cm)	Tumor weight	Follow-up
Cattell <i>et al</i> [22], 1947	55	Female			10 cm in diameter		
Cattell <i>et al</i> [22], 1947	61	Female	Epigastric distress and bloating			12700 g	
Deppe <i>et al</i> [21], 1985	24	Female		Barium enema, CT	11 × 8 × 3	NA	
Zhang <i>et al</i> [20], 1987	65	Male	Weight gain, leg edema	NA	50 cm in diameter	19500 g	4 yr
Acheson <i>et al</i> [19], 1997	76	Female	Swollen leg	CT, MRI	20 × 20 × 12	576 g	
Matsubara <i>et al</i> [18], 2000	65	Male		NA	12 × 13	NA	
Marshall <i>et al</i> [17], 2001	47	Male		CT	NA	4990 g	
Forte <i>et al</i> [15], 2002	61	Male	Urinary frequency, urgency and nocturia	CT			
Foa <i>et al</i> [16], 2002	52	Male			10.5 × 9.5 × 2	145 g	
Raftopoulos <i>et al</i> [14], 2002	62	Male	Abdominal pain	CT	20 × 15 × 10	790 g	
Martinez <i>et al</i> [12], 2003	32	Female	Abdominal pain	US, barium enema	20 × 13 × 10	3400 g	17 yr
Drop <i>et al</i> [13], 2003	60	Female	Abdominal pain, gastrointestinal symptoms	US, CT	13 × 12		
Drop <i>et al</i> [13], 2003	72	Female	Abdominal pain, sickness	US, CT	12 × 9 × 4		
Ida <i>et al</i> [11], 2008	65	Male	Painless swelling in left inguinal region		22 × 14 × 5		18 mo
Ukita <i>et al</i> [10], 2009	61	Female	Gluteal pain	MRI	25 × 15		
Singh <i>et al</i> [9], 2011	65	Male	Inguinal pain		15.6 cm in diameter	NA	
Chander <i>et al</i> [8], 2012	36	Female			13.6 × 11.2 × 9.1	1300 g	
Wei <i>et al</i> [6], 2013	25	Female		US	20 × 12 × 10	1650 g	6 mo
Saito <i>et al</i> [7], 2013	65	Male	Flank pain	US, CT	30 cm in diameter	NA	
Weniger <i>et al</i> [5], 2015	73	Female	Abdominal swelling, pain, and obstipation	CT	55 × 40 × 10	8950 g	
Al-Ali <i>et al</i> [4], 2019	34	Female	Abdominal distention and back pain	US, CT	45 × 48 × 13	1200 g	6 mo
Mitchell <i>et al</i> [3], 2020	29	Female	Abdominal pain, distention, orthopnea	MRI	28 × 14 × 6		

CT: Computerized tomography; MRI: Magnetic resonance imaging; US: Ultrasonography; NA: Not available.

clinical presentations tend to be variable and nonspecific[4]. Hence, imaging examinations play an essential role in the diagnosis of these lesions.

Ultrasound is generally used for the initial diagnosis and screening of abdominal masses. Radiography, especially CT and magnetic resonance imaging (MRI), is a crucial diagnostic tool for further evaluation of retroperitoneal tumors. The characteristics of adipose tissues are consistent on CT and MRI, but they differ on ultrasonography depending on the physical properties and histologic types. The fatty content is the fundamental feature to identify fat-containing retroperitoneal tumors during imaging examinations. Typical lipomas appear as extensive hyperechoic lesions on ultrasound, while they appear as homogeneous fat-containing masses with thin septa on CT and MRI. Retroperitoneal lipomas are difficult to identify preoperatively since they mimic liposarcomas, which account for the majority of fat-containing retroperitoneal tumors. Liposarcomas present heterogeneous signal intensity and variable appearances on MRI and CT due to the varying subtypes, which included well-differentiated liposarcoma (WDLPS), dedifferentiated liposarcoma, myxoid/round cell liposarcoma, pleomorphic liposarcoma, and mixed liposarcoma. The increased vascularity in liposarcomas that present as low-intensity signals on T1-weighted images can be used for differentiation. However, both lipomas and WDLPS are accompanied by a large amount of fat and minimal soft tissue and have identical appearances on CT and MRI, making it hard to distinguish lipomas from well-differentiated liposarcomas preoperatively.




Figure 3 Macroscopic view of extracted retroperitoneal lipoma. During the operation, a bulky yellowish tumor, originating from the left retroperitoneal region, was found to occupy the retroperitoneum. The mass weighted 7.126 kg.

Due to the large size, the measurement of retroperitoneal tumors by preoperative imaging examinations can be inaccurate. Despite their typical presentations on CT and MRI, both imaging modalities may not rule out the possibility of WDLPS[29]. Approximately 80% of retroperitoneal tumors appear to be malignant, most of which are soft-tissue sarcomas, a category of very uncommon neoplasms, with an overall incidence of 0.3% to 0.4% per 100000 people[10]. Liposarcomas account for 41% of sarcomas, and the majority of the cases are malignant from the start. A few outliers arise from benign lipomas in the early stages[30]. The final diagnosis of lipomas depends on histopathology. Tissue for pathology can be acquired by fine-needle aspiration or core-needle biopsy, but it is nearly impossible to distinguish lipoma-like WDLPS and lipomas due to the limited tissue sample obtained by these methods for detecting atypia and hyperchromatic cells. Postoperative histopathology remains the gold standard for diagnosis. Histologic characteristics for WDLPS include mature adipocytes punctuated with big atypical hyperchromatic cells. However, WDLPS are likely to be misdiagnosed, because atypia may be localized, especially in deep lesions with tiny samples. Murine double minute (*MDM2*, located at 12q14-15) and cyclin-dependent kinase 4 gene are regularly amplified in WDLPS, which cannot be observed in benign lipomas. Hence, fluorescence *in situ* hybridization has emerged as a promising method for differential diagnosis[11,16,18].

It is of great importance to discern tumor characteristics intraoperatively and make decisions about the resection extent subsequently. In cases of the pathological diagnosis of liposarcoma, resection with negative margins (R0) is crucial. If infiltrative growth is detected by frozen pathology, a broad excision should be performed. Surgeons should also tailor personalized surgical strategies for patients with important involved adjacent structures who are unsuitable for an entire resection. Commonly, *en*




Figure 4 Microscopical picture of the extracted tumors (H&E, 20 ×). A: Myelolipoma was composed of mature adipose adipocytes and hematopoietic cells, without necrosis, atypia, and hyperchromatic cells; B: Conventional lipoma was composed of mature adipocytes.

en bloc removal of the involved structures is required[3,5]. The prognosis and recurrence risk for patients with benign retroperitoneal lipomas are unclear due to the limited number of case reports. Patients are often recommended to receive regular clinical and radiologic follow-ups.

CONCLUSION

Retroperitoneal lipomas are rare benign tumors originating from adipose tissues and they tend to have large sizes. Imaging examinations, especially CT and MRI, are fundamental diagnostic tools for these tumors. Surgical resection is the main treatment method. *En bloc* resection is commonly required. Postoperative histopathology determines the final diagnosis, and immunohistochemical analysis could be useful in the differentiation of liposarcomas. Regular follow-ups are also required for the patients.

ACKNOWLEDGEMENTS

The authors are deeply grateful to Dr. Zhao H and Dr. Shi J for their help in reviewing the postoperative pathology.

REFERENCES

- 1 **Terada T.** Giant fibrolipoma of the spermatic cord. *Pathol Int* 2010; **60**: 330-332 [PMID: 20403037 DOI: 10.1111/j.1440-1827.2010.02521.x]
- 2 **Johnson CN, Ha AS, Chen E, Davidson D.** Lipomatous Soft-tissue Tumors. *J Am Acad Orthop Surg* 2018; **26**: 779-788 [PMID: 30192249 DOI: 10.5435/JAAOS-D-17-00045]
- 3 **Mitchell K, Fuller K, Thomay A, Shapiro R.** Diagnosis and Surgical Management of a Retroperitoneal Lipoma in Pregnancy. *Case Rep Obstet Gynecol* 2020; **2020**: 6309417 [PMID: 32724687 DOI: 10.1155/2020/6309417]
- 4 **Al-Ali MHM, Salih AM, Ahmed OF, Kakamad FH, Mohammed SH, Hassan MN, Sidiq SH, Mustafa MQ, Najar KA, Abdullah IY.** Retroperitoneal lipoma; a benign condition with frightening presentation. *Int J Surg Case Rep* 2019; **57**: 63-66 [PMID: 30904820 DOI: 10.1016/j.ijscr.2019.02.044]
- 5 **Weniger M, D'Haese JG, Kunz W, Pratschke S, Guba M, Werner J, Angele MK.** En-bloc resection of a giant retroperitoneal lipoma: a case report and review of the literature. *BMC Res Notes* 2015; **8**: 75 [PMID: 25890295 DOI: 10.1186/s13104-015-1038-7]
- 6 **Wei D, Shen L, Yang K, Fang F.** Giant retroperitoneal lipoma in a pregnant patient. *J Obstet Gynaecol* 2013; **33**: 522 [PMID: 23815212 DOI: 10.3109/01443615.2013.788621]
- 7 **Saito S.** Retroperitoneal lipoma presenting with nutcracker-like phenomenon. *Case Rep Urol* 2013; **2013**: 893242 [PMID: 24349819 DOI: 10.1155/2013/893242]
- 8 **Chander B, Krishna M, Thakur S, Mahajan N, Vij A, Diwakaran J.** Extremely rare giant retroperitoneal fibrolipoma: a case report. *J Cancer Res Ther* 2012; **8**: 314-316 [PMID: 22842386 DOI: 10.4103/0973-1482.99002]
- 9 **Singh G, Bharadwaj RN, Purandare SN, Gore CR, Dubhashi SP, Vaidya S, Patil A, Kompally GR.** Giant retroperitoneal lipoma presenting as inguinal hernia. *Indian J Surg* 2011; **73**: 187-189 [PMID: 22654328 DOI: 10.1007/s12262-010-0210-5]
- 10 **Ukita S, Koshiyama M, Ohnaka M, Miyagawa N, Yamanishi Y, Nishimura F, Nagura M, Kim T, Hirose M, Shirase T, Kobayashi H, Ozasa H.** Retroperitoneal lipoma arising from the urinary bladder. *Rare Tumors* 2009; **1**: e13 [PMID: 21139884 DOI: 10.4081/rt.2009.e13]
- 11 **Ida CM, Wang X, Erickson-Johnson MR, Wenger DE, Blute ML, Nascimento AG, Oliveira AM.** Primary retroperitoneal lipoma: a soft tissue pathology heresy? *Am J Surg Pathol* 2008; **32**: 951-954 [PMID: 18551755 DOI: 10.1097/pas.0b013e318160c6bf]
- 12 **Martinez CA, Palma RT, Waisberg J.** Giant retroperitoneal lipoma: a case report. *Arq Gastroenterol* 2003; **40**: 251-255 [PMID: 15264048 DOI: 10.1590/s0004-28032003000400010]
- 13 **Drop A, Czekajka-Chehab E, Maciejewski R.** Giant retroperitoneal lipomas--radiological case report. *Ann Univ Mariae Curie Skłodowska Med* 2003; **58**: 142-146 [PMID: 15323181]
- 14 **Raftopoulos I, Lee T, Byrne MP.** Image of the month: retroperitoneal lipoma. *Arch Surg* 2002; **137**: 865-866 [PMID: 12093348 DOI: 10.1001/archsurg.137.7.865]
- 15 **Forte F, Maturo G, Catania A, Sorrenti S, Gemma D, Foti N, Vanni B, Virgili G, Vespasiani G, De Antoni E.** Retroperitoneal lipoma. Unusual presentation with detrusor instability. *Minerva Urol Nefrol* 2002; **54**: 131-133 [PMID: 12070462]
- 16 **Foa C, Mainguéné C, Dupré F, Coindre JM, Huguet C, Kober C, Pedetour F.** Rearrangement involving chromosomes 1 and 8 in a retroperitoneal lipoma. *Cancer Genet Cytogenet* 2002; **133**: 156-159. [PMID: 11943344 DOI: 10.1016/s0165-4608(01)00573-8]
- 17 **Marshall MT, Rosen P, Berlin R, Greenson N.** Appendicitis masquerading as tumor: a case of two diagnoses. *J Emerg Med* 2001; **21**: 397-399 [PMID: 11728767 DOI: 10.1016/s0736-4679(01)00422-x]
- 18 **Matsubara N, Yoshitaka T, Matsuno T, Ikeda M, Isozaki H, Tanaka N, Shimizu K.** Multiple tumors and a novel E2F-4 mutation. A case report. *Digestion* 2000; **62**: 213-216 [PMID: 11025371 DOI: 10.1159/000007816]
- 19 **Acheson A, McIlrath E, Barros D'Sa AA.** Pelvic lipoma causing venous obstruction syndrome. *Eur J Vasc Endovasc Surg* 1997; **14**: 149-150 [PMID: 9314859 DOI: 10.1016/s1078-5884(97)80213-4]
- 20 **Zhang SZ, Yue XH, Liu XM, Lo SL, Wang XZ.** Giant retroperitoneal pleomorphic lipoma. *Am J Surg Pathol* 1987; **11**: 557-562 [PMID: 3605490]
- 21 **Deppe G, Malviya VK, Hercule J, Gleicher N.** Retroperitoneal pelvic lipoma. *J Natl Med Assoc* 1985; **77**: 574-576 [PMID: 4046056]
- 22 **Cattell RB, Warren KW.** Retroperitoneal lipoma. *Surg Clin North Am* 1947; **27**: 659-665 [PMID: 20242255 DOI: 10.1016/s0039-6109(16)32148-x]
- 23 **Pai MR, Naik R, Raghuveer CV.** Primary retroperitoneal tumors a 25 year study. *Indian J Med Sci* 1995; **49**: 139-141 [DOI: 10.1111/j.1744-1633.2010.00485.x]
- 24 **Armstrong JR, Cohn I.** Primary malignant retroperitoneal tumors. *Am J Surg* 1965; **110**: 937-943 [DOI: 10.1016/0002-9610(65)90181-9]
- 25 **Rubinstein A, Goor Y, Gazit E, Cabili S.** Non-symmetric subcutaneous lipomatosis associated with familial combined hyperlipidaemia. *Br J Dermatol* 1989; **120**: 689-694 [PMID: 2757931 DOI: 10.1111/j.1365-2133.1989.tb01357.x]
- 26 **Self TH, Akins D.** Dramatic reduction in lipoma associated with statin therapy. *J Am Acad Dermatol* 2008; **58**: S30-S31 [PMID: 18191694 DOI: 10.1016/j.jaad.2007.08.034]
- 27 **Pires Botelho da Costa JS, Reis JC, Valença-Filipe R.** Giant atypical lipoma of the thigh. *Dermatol*

- Surg* 2014; **40**: 213-214 [PMID: [24354663](#) DOI: [10.1111/dsu.12397](#)]
- 28 **Cho J**, Kinsey D, Kimchi ET, O'Carroll KS, Nguyen V, Alsabbagh M, Gaballah A. Retroperitoneal extra-adrenal myelolipoma misdiagnosed as liposarcoma: A case report. *Radiol Case Rep* 2021; **16**: 364-368 [PMID: [33532014](#) DOI: [10.1016/j.radcr.2020.11.045](#)]
 - 29 **Shaaban AM**, Rezvani M, Tubay M, Elsayes KM, Woodward PJ, Menias CO. Fat-containing Retroperitoneal Lesions: Imaging Characteristics, Localization, and Differential Diagnosis. *Radiographics* 2016; **36**: 710-734 [PMID: [27163589](#) DOI: [10.1148/rg.2016150149](#)]
 - 30 **Vijay A**, Ram L. Retroperitoneal liposarcoma: a comprehensive review. *Am J Clin Oncol* 2015; **38**: 213-219 [PMID: [24136142](#) DOI: [10.1097/COC.0b013e31829b5667](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

