

World Journal of *Clinical Cases*

World J Clin Cases 2022 April 6; 10(10): 2976-3320



Contents

Thrice Monthly Volume 10 Number 10 April 6, 2022

REVIEW

- 2976 Gut microbiota in gastrointestinal diseases during pregnancy
Liu ZZ, Sun JH, Wang WJ
- 2990 Targeting metabolism: A potential strategy for hematological cancer therapy
Tang X, Chen F, Xie LC, Liu SX, Mai HR

MINIREVIEWS

- 3005 Elevated intra-abdominal pressure: A review of current knowledge
Łagosz P, Sokolski M, Biegus J, Tycinska A, Zymlinski R

ORIGINAL ARTICLE

Case Control Study

- 3014 Changes in corneal nerve morphology and function in patients with dry eyes having type 2 diabetes
Fang W, Lin ZX, Yang HQ, Zhao L, Liu DC, Pan ZQ
- 3027 Combined sevoflurane-dexmedetomidine and nerve blockade on post-surgical serum oxidative stress biomarker levels in thyroid cancer patients
Du D, Qiao Q, Guan Z, Gao YF, Wang Q

Retrospective Cohort Study

- 3035 Early warning prevention and control strategies to reduce perioperative venous thromboembolism in patients with gastrointestinal cancer
Lu Y, Chen FY, Cai L, Huang CX, Shen XF, Cai LQ, Li XT, Fu YY, Wei J
- 3047 Dose-response relationship between risk factors and incidence of COVID-19 in 325 hospitalized patients: A multicenter retrospective cohort study
Zhao SC, Yu XQ, Lai XF, Duan R, Guo DL, Zhu Q

Retrospective Study

- 3060 Preventive online and offline health management intervention in polycystic ovary syndrome
Liu R, Li M, Wang P, Yu M, Wang Z, Zhang GZ
- 3069 Evidence-based intervention on postoperative fear, compliance, and self-efficacy in elderly patients with hip fracture
Fu Y, Zhu LJ, Li DC, Yan JL, Zhang HT, Xuan YH, Meng CL, Sun YH
- 3078 Significance of dysplasia in bile duct resection margin in patients with extrahepatic cholangiocarcinoma: A retrospective analysis
Choe JW, Kim HJ, Kim JS

- 3088** Diagnostic value and safety of medical thoracoscopy for pleural effusion of different causes

Liu XT, Dong XL, Zhang Y, Fang P, Shi HY, Ming ZJ

Observational Study

- 3101** Oxaliplatin-induced neuropathy and colo-rectal cancer patient's quality of life: Practical lessons from a prospective cross-sectional, real-world study

Prutianu I, Alexa-Stratulat T, Cristea EO, Nicolau A, Moisuc DC, Covrig AA, Ivanov K, Croitoru AE, Miron MI, Dinu MI, Ivanov AV, Marinca MV, Radu I, Gafton B

- 3113** Breast-conserving surgery and sentinel lymph node biopsy for breast cancer and their correlation with the expression of polyligand proteoglycan-1

Li FM, Xu DY, Xu Q, Yuan Y

SYSTEMATIC REVIEWS

- 3121** Clinical significance of aberrant left hepatic artery during gastrectomy: A systematic review

Tao W, Peng D, Cheng YX, Zhang W

META-ANALYSIS

- 3131** Betel quid chewing and oral potential malignant disorders and the impact of smoking and drinking: A meta-analysis

Lin HJ, Wang XL, Tian MY, Li XL, Tan HZ

- 3143** Effects of physical exercise on the quality-of-life of patients with haematological malignancies and thrombocytopenia: A systematic review and meta-analysis

Yang YP, Pan SJ, Qiu SL, Tung TH

CASE REPORT

- 3156** Primary malignant peritoneal mesothelioma mimicking tuberculous peritonitis: A case report

Lin LC, Kuan WY, Shiu BH, Wang YT, Chao WR, Wang CC

- 3164** Endoscopic submucosal dissection combined with adjuvant chemotherapy for early-stage neuroendocrine carcinoma of the esophagus: A case report

Tang N, Feng Z

- 3170** Lymph-node-first presentation of Kawasaki disease in a 12-year-old girl with cervical lymphadenitis caused by *Mycoplasma pneumoniae*: A case report

Kim N, Choi YJ, Na JY, Oh JW

- 3178** Tuberculosis-associated hemophagocytic lymphohistiocytosis misdiagnosed as systemic lupus erythematosus: A case report

Chen WT, Liu ZC, Li MS, Zhou Y, Liang SJ, Yang Y

- 3188** Migration of a Hem-o-Lok clip to the renal pelvis after laparoscopic partial nephrectomy: A case report

Sun J, Zhao LW, Wang XL, Huang JG, Fan Y

- 3194** Ectopic intrauterine device in the bladder causing cystolithiasis: A case report
Yu HT, Chen Y, Xie YP, Gan TB, Gou X
- 3200** Giant tumor resection under ultrasound-guided nerve block in a patient with severe asthma: A case report
Liu Q, Zhong Q, Zhou NN, Ye L
- 3206** Myomatous erythrocytosis syndrome: A case report
Shu XY, Chen N, Chen BY, Yang HX, Bi H
- 3213** Middle thyroid vein tumor thrombus in metastatic papillary thyroid microcarcinoma: A case report and review of literature
Gui Y, Wang JY, Wei XD
- 3222** Severe pneumonia and acute myocardial infarction complicated with pericarditis after percutaneous coronary intervention: A case report
Liu WC, Li SB, Zhang CF, Cui XH
- 3232** IgA nephropathy treatment with traditional Chinese medicine: A case report
Zhang YY, Chen YL, Yi L, Gao K
- 3241** Appendico-vesicocolonic fistula: A case report and review of literature
Yan H, Wu YC, Wang X, Liu YC, Zuo S, Wang PY
- 3251** *Scedosporium apiospermum* infection of the lumbar vertebrae: A case report
Shi XW, Li ST, Lou JP, Xu B, Wang J, Wang X, Liu H, Li SK, Zhen P, Zhang T
- 3261** Woman diagnosed with obsessive-compulsive disorder became delusional after childbirth: A case report
Lin SS, Gao JF
- 3268** Emphysematous pyelonephritis: Six case reports and review of literature
Ma LP, Zhou N, Fu Y, Liu Y, Wang C, Zhao B
- 3278** Atypical infantile-onset Pompe disease with good prognosis from mainland China: A case report
Zhang Y, Zhang C, Shu JB, Zhang F
- 3284** *Mycobacterium tuberculosis* bacteremia in a human immunodeficiency virus-negative patient with liver cirrhosis: A case report
Lin ZZ, Chen D, Liu S, Yu JH, Liu SR, Zhu ML
- 3291** Cervical aortic arch with aneurysm formation and an anomalous right subclavian artery and left vertebral artery: A case report
Wu YK, Mao Q, Zhou MT, Liu N, Yu X, Peng JC, Tao YY, Gong XQ, Yang L, Zhang XM
- 3297** Dedifferentiated chondrosarcoma of the middle finger arising from a solitary enchondroma: A case report
Yonezawa H, Yamamoto N, Hayashi K, Takeuchi A, Miwa S, Igarashi K, Morinaga S, Asano Y, Saito S, Tome Y, Ikeda H, Nojima T, Tsuchiya H

- 3306** Endoscopic-catheter-directed infusion of diluted (-)-noradrenaline for atypical hemobilia caused by liver abscess: A case report
Zou H, Wen Y, Pang Y, Zhang H, Zhang L, Tang LJ, Wu H
- 3313** *Pneumocystis jiroveci* pneumonia after total hip arthroplasty in a dermatomyositis patient: A case report
Hong M, Zhang ZY, Sun XW, Wang WG, Zhang QD, Guo WS

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Hui-Jeong Hwang, MD, PhD, Associate Professor, Department of Cardiology, Kyung Hee University Hospital at Gangdong, Kyung Hee University College of Medicine, Seoul 05278, South Korea. neonic7749@hanmail.net

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJCC as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The WJCC's CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Xu Guo; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lai Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

April 6, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Appendico-vesicocolonic fistula: A case report and review of literature

Han Yan, Ying-Chao Wu, Xin Wang, Yu-Cun Liu, Shuai Zuo, Peng-Yuan Wang

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Corvino A, Silano F

Received: October 10, 2021

Peer-review started: October 10, 2021

First decision: December 12, 2021

Revised: December 30, 2021

Accepted: February 23, 2022

Article in press: February 23, 2022

Published online: April 6, 2022



Han Yan, Ying-Chao Wu, Xin Wang, Yu-Cun Liu, Shuai Zuo, Peng-Yuan Wang, Department of General Surgery, Peking University First Hospital, Beijing 100032, China

Corresponding author: Peng-Yuan Wang, MD, Doctor, Department of General Surgery, Peking University First Hospital, No. 8 Xishiku Street, Xicheng District, Beijing 100032, China.
wangpengyuan2014@126.com

Abstract

BACKGROUND

Appendico-vesicocolonic fistulas and appendiceal-colonic fistulas are two kinds of intestinal and bladder diseases that are rarely seen in the clinic. To our knowledge, no more than 4 cases of appendico-vesicocolonic fistulas have been publicly reported throughout the world, and no more than 100 cases of appendiceal-colonic fistulas have been reported. Although the overall incidence is low, an early diagnosis is difficult due to their atypical initial symptoms, but these diseases still require our attention.

CASE SUMMARY

Here, we report a case of a 77-year-old male patient diagnosed with an appendico-vesicocolonic fistula combined with an appendiceal-colonic fistula. The main manifestations were diarrhea and urine that contained fecal material. The diagnosis was confirmed by multiple laboratory and imaging examinations. A routine urinalysis showed red blood cells and white blood cells. Abdominal and pelvic computed tomography scans showed close adhesions between the bowels and the bladder, and fistulas could be seen. Colonoscopy and cystoscopy and some other imaging examinations clearly showed fistulas. The preoperative diagnoses were a colovesical fistula and an appendiceal-colonic fistula. The fistulas were repaired by laparoscopic surgical treatment. The diseased bowel and part of the bladder wall were removed, followed by a protective ileostomy. The postoperative diagnosis was an appendico-vesicocolonic fistula combined with an appendiceal-colonic fistula, and the pathology suggested inflammatory changes. The patient recovered well after surgery, and all his symptoms resolved.

CONCLUSION

The final diagnosis in this case was a double fistula consisting of an appendico-vesicocolonic fistula combined with an appendiceal-colonic fistula.

Key Words: Appendico-vesicocolonic; Appendiceal-colonic; Enterovesical; Fistula;

Laparoscopic surgery; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Appendico-vesicocolonic fistulas and appendicovesical fistulas are rare diseases involving the intestine and urinary bladder. These conditions have a very low morbidity even in patients with appendicitis after conservative treatment or with malignant tumors. Early symptoms and signs are usually atypical, making early diagnosis and treatment more difficult. Here, we present a case of an appendico-vesicocolonic fistula combined with an appendiceal-colonic fistula and provide some recommendations for diagnosis and treatment along with a review of the literature.

Citation: Yan H, Wu YC, Wang X, Liu YC, Zuo S, Wang PY. Appendico-vesicocolonic fistula: A case report and review of literature. *World J Clin Cases* 2022; 10(10): 3241-3250

URL: <https://www.wjgnet.com/2307-8960/full/v10/i10/3241.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i10.3241>

INTRODUCTION

Appendicovesical fistulas are commonly seen as complications after conservative treatment of acute and chronic appendicitis and benign or malignant appendiceal neoplasms[1]. Appendicovesical fistulas account for approximately 5% of all enterovesical fistulas[2]. Appendiceal-colonic fistulas are commonly seen in diseases such as a vermiform appendix combined with acute or chronic appendicitis or benign or malignant appendiceal neoplasms. It also occurs in the sigmoid colon combined with colonic diverticulitis or colonic malignancy[3]. Colovesical fistulas are more commonly seen in sigmoid diverticulitis or colonic malignancy disease and are rarely seen with bladder malignancies originating on the left side of the bladder that invade the sigmoid colon. Sigmoid diverticulitis is the most common cause[4]. It has been reported that the incidence of colovesical fistula is 2%-23% in diverticular disease [5].

Appendico-vesicocolonic fistulas are composed of the above three fistulas. It is a rare disease, with only 3 cases reported thus far[6-8]. A case of simultaneous double fistulas, "appendico-vesicocolonic fistula combined with appendiceal-colonic fistula", has not been reported thus far throughout the world. Here, we report a rare case of an appendico-vesicocolonic fistula with an appendiceal-colonic fistula.

CASE PRESENTATION

Chief complaints

A 77-year-old male patient was admitted to the hospital because of "diarrhea and urine having fecal material for 7 years".

History of present illness

Seven years before presentation, the patient experienced diarrhea and frequent and urgent urination, accompanied by urine having fecal material with no obvious cause. He was admitted to the local hospital. Colonoscopy showed a fistula 15 cm away from the anus, and a colovesical fistula was considered. Surgical treatment was suggested, but the patient refused surgery because the symptoms did not significantly affect his quality of life at that time. Recently, the patient came to our hospital due to worsening of the symptoms of frequent and urgent urination. The patient did not receive any other trauma or treatment during the period between refusing the surgery and its acceptance.

History of past illness

The patient had hypertension and diabetes for many years but had no history of trauma or surgery.

Personal and family history

There was no specific family history.

Physical examination

The patient had some mild tenderness in the lower abdomen. Percussion of the abdomen revealed dull

percussion sounds.

Laboratory examinations

Routine blood test: White blood cell (WBC) $7.9 \times 10^9/L$, hemoglobin (HBG) 127 g/L; Stool routine test (-); Urine routine test: WBC (+++), red blood cell 945/H, WBC 1277/HP.

Imaging examinations

Colonoscopy: A fistula approximately 1.5 cm in size could be seen involving the sigmoid colon 20 cm from the anus. The endoscope could pass through the fistula, and the bladder mucosa could be seen. Another fistula with a size of approximately 1.2 cm could be seen 15 cm away from the anus. This suggested multiple colovesical fistulas (Figure 1A). Cystoscopy showed a high turbidity of the urine in the bladder, with a large amount of feces visible, and the wall of the bladder was edematous. A fistula could be seen in the lateral wall of the bladder. Both ureteral orifices were fissured, but the flow of urine was clearly visible (Figure 1B). Abdominal pelvic computed tomography (CT) showed that the appendix was not clearly visible. The distal sigmoid colon-cecum, right posterior to the top wall of the bladder-sigmoid colon, and middle sigmoid colon-cecal wall contained adhesions, and fistulas could be seen among these structures. A gas density shadow could be seen in the bladder. The involved intestinal wall and bladder wall were thickened, and a contrast-enhanced CT scan showed enhancements (Figure 2). In view of the previous consideration of a possible colovesical fistula and the complexity of the patient's condition, we considered a CT examination directly rather than a plain film or ultrasound. Plain films, ultrasound, fistulography and magnetic resonance imaging (MRI) were not performed later because the CT scan provided comprehensive information.

FINAL DIAGNOSIS

Preoperative diagnosis was a colovesical fistula and an appendiceal-colonic fistula. Postoperative diagnosis was an appendico-vesicocolonic fistula with an appendiceal-colonic fistula.


TREATMENT

After completing the preoperative preparation and signing the informed consent, the surgeries, which consisted of "cystoscopy, bilateral Double J (D-J) ureteral tube placement, laparoscopic exploration, appendectomy, resection and anastomosis of sigmoid colon, double-lumen protective ileostomy, bladder fistula neoplasty, and drainage tube placement" were performed. First, for the operation, bilateral D-J tubes were placed with cystoscopy. A zebra guidewire was set through the sinus tract. A catheter was inserted through the guide wire, and the catheter water bag was filled with water for subsequent guidance to find the sinus canal. Using laparoscopy, the water sac was found and located in the cecum. After the appendix was severed at its root from the cecum, the sinus tract was cut open, and the urinary duct was withdrawn. The urinary duct was withdrawn into the colon lumen and then into the bladder lumen. Then, it was confirmed that there was a three-way fistula of the appendix, colon and bladder. After that, sigmoidectomy and anastomosis were performed, and the other fistula revealed by colonoscopy was not found during the operation. The rectum was dissected in the upper segment, and a double-lumen protective ileostomy and bladder neoplasty were performed. The operation was successful with a minimal amount of bleeding.

The specimen was opened and revealed a fistula of approximately 1.5 cm in diameter in the middle of the appendix and leading to the colon and bladder.


Another fistula was seen approximately 5 cm away from the tip of the appendix, which was confirmed as a case of a double fistula consisting of an appendico-vesicocolonic fistula combined with an appendiceal-colonic fistula (Figure 3).

In the course of postoperative recovery, a urinary tract infection occurred, and the urinary culture grew *Candida albicans*. The infection was cured after antibiotics and antifungal agents were given. The bilateral D-J tubes were removed from the bladder under cystoscopy guidance one month after the operation, and the procedure was successful. Postoperative pathology of the resected rectum, appendix, and part of bladder showed the following: a segment of large intestine was evaluated with a length of 8 cm, and acute or chronic inflammation was present in the large intestinal wall. There was local mucosal erosion and infiltration of some eosinophils in the lamina propria. Two fistulas were observed in the wall of the large intestine. One fistula was adhered to part of the appendiceal wall, and the appendiceal wall was dilated, with mucosal erosion and degeneration. The other fistula was adhered to a small amount of hyperplastic fibers and smooth muscle tissue, and no transitional epithelium was observed. One lymph node around the large intestine showed reactive hyperplasia.



DOI: 10.12998/wjcc.v10.i10.3241 Copyright ©The Author(s) 2022.

Figure 1 Endoscope. A: Colonoscopy. A fistula approximately 1.5 cm in size could be seen on the sigmoid colon 20 cm from the anus. The endoscope could pass through the fistula, and the bladder mucosa could be seen. Another fistula with a size of approximately 1.2 cm could be seen 15 cm away from the anus. This suggested multiple colovesical fistulas; B: Cystoscopy. Turbidity of the urine in the bladder, with a large amount of feces visible, and the wall of the bladder was edematous. A fistula could be seen in the lateral wall of the bladder. Both ureteral orifices were fissured, and the release of urine was adequate.



DOI: 10.12998/wjcc.v10.i10.3241 Copyright ©The Author(s) 2022.

Figure 2 Contrast-enhanced computed tomography of the abdominal pelvis: the appendix was not clearly visualized. A: The distal sigmoid colon-cecum, right posterior to the top wall of the bladder - sigmoid colon, and middle sigmoid colon-cecal wall were adhered, and fistulas could be seen among these structures. The correlative intestinal wall was thickened, and a contrast-enhanced computed tomography (CT) scan showed enhancement; B: A gas density shadow could be seen in the bladder. The correlative bladder wall was thickened, and a contrast-enhanced CT scan showed enhancement.


OUTCOME AND FOLLOW-UP

The patient has recovered well after surgery. In addition, the symptoms above have completely resolved.

DISCUSSION

Enterovesical fistulas were first described by Cripps[9] in 1888. Appendiceal-colic fistulas were first reported by Cherigie *et al*[10] in 1953.

The common causes of appendico-vesicocolonic fistulas mainly include inflammation, tumors, trauma, and iatrogenic and congenital diseases. Appendiceal-colonic fistulas often begin when acute or chronic appendicitis is combined with the formation of a local abscess, especially after conservative treatment. When the appendix is long and in the pelvic position, local inflammation and abscesses are more likely to be close to and invade the lengthy sigmoid colonic wall, leading to the formation of an appendiceal-colonic fistula. In addition, malignancy of the appendix, neuroma, cystic fibrosis, villous adenoma and parasites are also common causes[2]. Colovesical fistulas are more common in patients with colonic diverticulitis and colonic malignancy, especially in the sigmoid colon, which may be due to its high mobility and anatomical characteristics. According to the data, the incidence of diverticula developing into fistulas was 4% to 23%. Enterovesical fistulas are the most common type, accounting for 33% to 65% of all diagnosed diverticular fistulas[11]. According to the report, inflammatory bowel



DOI: 10.12998/wjcc.v10.i10.3241 Copyright ©The Author(s) 2022.

Figure 3 A schematic drawing depicting the fistulous paths between the various organs. The fistula 20 cm away from the anus was an appendico-vesicocolonic fistula with a diameter of 1.5 cm, and the fistula 15 cm away from the anus was an appendiceal-colonic fistula with a diameter of 1.2 cm.

disease, colon radiotherapy, or bladder malignancy are also rare causes of colovesical fistula[12]. Dangle *et al*[13] reported that, rarely, enterovesical fistulas were the result of inflammatory and necrotic responses to the intravesical injection of mitomycin. The underlying etiologies of 66 patients diagnosed with enterovesical fistulas, as reported by Qiu *et al*[14], were intestinal cancer (46.97%), Crohn's disease (16.67%), colonic diverticulitis (15.15%), bladder cancer (12.12%), appendicitis and other inflammatory diseases (7.58%), and intraoperative injury (1.52%).

The clinical manifestations of appendico-vesicocolonic fistulas are diverse, and the common manifestations are frequent urination, pneumaturia, urine containing fecal material complex and persistent urinary tract infection, dysuria, diarrhea, abdominal discomfort and other symptoms. Sometimes the symptoms, such as urinary tract irritation, are relieved because the fistula is completely blocked by a bezoar, but temporary abdominal pain may occur when the bezoar in the fistula falls into the bladder [15]. Different patients have different clinical manifestations, and the above clinical symptoms do not always occur simultaneously. Therefore, the early diagnosis of an appendico-vesicocolonic fistula is much more difficult. In this case, the main clinical manifestations of the patient were diarrhea, fecal contamination, and frequent and urgent urination. Melchior *et al*[16] reported 49 patients with colovesical fistulas due to sigmoid diverticulitis, 71.4% of patients had pneumaturia, and 51% had fecal material within urine. Yagi *et al*[17] had previously reported on a patient with a sigmoid carcinoma within the colonic diverticulum, which invaded the bladder and resulted in a colovesical fistula, the main symptom of which was hematuria. In addition, Keane and Tebala[18] reported a rare case of hypokalemia and hyperchloremic metabolic acidosis. Most of the patients with appendico-vesicocolonic fistulas were male, which may be because the uterus plays a protective role in the formation of enterovesical fistulas[4]. Melchior *et al*[16] reported 49 patients with colovesical fistulas, only 7 patients (14.3%) were female, and 5 of them had undergone a hysterectomy. However, the clinical manifestations of appendiceal-colic fistulas are relatively vague and have a poor specificity, and patients have nonspecific symptoms such as discomfort in the lower abdomen, constipation, diarrhea and so on. Due to its nonspecific manifestations, a preoperative diagnosis is often difficult. The postoperative diagnosis is usually made through histopathological examination of resected specimens after exploratory surgery for other suspected diseases[19].

The diagnoses of appendico-vesicocolonic fistulas and appendiceal-colonic fistulas are often difficult. Although there are various diagnostic methods, different methods have different specificities and sensitivities, so it is often necessary to apply multiple simultaneous diagnostic methods to aid in diagnosis and to improve the detection rates.

In terms of laboratory examinations in patients with appendico-vesicocolonic fistulas, routine urine tests and urine cultures had the best results. Routine urine tests are commonly characterized by turbid urine with feces containing a large number of white blood cells and red blood cells. *Escherichia coli* and *Klebsiella* are usually found in the urine cultures[20].

In terms of imaging examinations in patients with appendico-vesicocolonic fistulas, compared with other diagnostic tools, abdominal CT is considered to be one of the most sensitive detection tools[21]. The main manifestations are as follows: the intestine and bladder are closely adhered, fistulas can be seen in some cases, gas density shadows can be seen in the bladder, there is corresponding intestinal and bladder wall thickening, and some other manifestations can be present. It has been reported that the

detection rate of colovesical fistulas using abdominal spiral CT is between 60% and 100%[22]. In the study from Melchior *et al*[16], the detection rate using CT was 40%. Although the detection rate of appendico-vesicocolonic fistulas by CT is highly variable, CT is useful to exclude other organic diseases in the abdominal cavity, such as malignant tumors.

Colonoscopy and cystoscopy have a limited diagnostic value for appendico-vesicocolonic fistulas. According to relevant reports, the detection rate of enterovesical fistulas by colonoscopy is highly variable, ranging from 11% to 89%[22]. The low colonoscopy detection rate was mainly because some fistulas with stenosis in the neck were easily ignored or mistaken for colonic diverticula. Although cystoscopy is a routine test, it has been reported to diagnose no more than 50% of enterovesical fistulas [22]. Cystoscopy often presents some nonspecific findings, such as diffuse edema of the bladder wall and mucosal bulging. Only mature fistulas with obvious openings can be easily diagnosed with endoscopy. In the study from Melchior *et al*[16], the detection rate using colonoscopy was 8.5%, and the detection rate using cystoscopy was 10.2%. Although the diagnostic value of colonoscopy for enterovesical fistulas is limited, it helps to rule out inflammatory bowel disease and colon, appendix, and bladder tumors.

Detection of poppy seeds in urine is considered a positive confirmatory test for enterovesical fistulas. In some studies, the sensitivity of the poppy seed test to detect fistulas was 94.8%[23]. The poppy seed test involves taking 50 g of poppy seeds orally and visually testing urine for the presence of poppy seeds within 48 h. This experiment is simple and effective, with a high diagnostic rate and almost no side effects. However, it has not been carried out in our country (China) or in our center.

MRI is widely used in the diagnosis of appendico-vesicocolonic fistulas because of its quality imaging of soft tissues. The main radiographic findings were tubular hyperintensities on T2-weighted images and focal disruptions of the muscular layer of the bladder wall. A study by Ravichandran *et al*[24] showed that enterovesical fistulas were diagnosed by MRI in 18 out of 19 patients, and the related causes were analyzed. The patient reported in this paper did not receive an MRI examination because the diagnosis was confirmed by CT and endoscopy. In addition, barium enema studies, total alimentary tract contrast studies, cystography, and retrograde urography also play auxiliary roles in the diagnosis. Among them, the Bourne test had a higher detection rate, urine specimens were collected immediately after a barium enema, and the sediment was examined by radiography after centrifugation. If the sediment contained the barium agent, the diagnosis could be confirmed. However, few institutions have carried out such test.

For appendiceal-colonic fistulas, the main manifestations on CT are as follows: the distal appendix may be closely adhered to the lengthy sigmoid colon, local inflammation or abscesses can be seen, and the fistula can be seen in some cases. Fluorodeoxyglucose (FDG) positron emission tomography is helpful to distinguish between benign and malignant diseases[25]. Barium enemas are useful for the confirmation of appendiceal-colonic fistulas[26].

Given the atypical symptoms of enterovesical and appendiceal-colonic fistulas, differential diagnoses need to be considered. Because the common symptoms are abdominal pain in the hypogastrium and right/left iliac fossa, they are often differentiated from the following conditions: inflammatory and infectious conditions involving the ileocecal region, diverticulitis occurring in different parts of the digestive tract, malignancies of the appendix and colorectum, and other rare diseases[27]. Interestingly, van Breda Vriesman AC *et al*[28] found that infarction of the epiploic appendage or omentum can cause similar localized pain and can easily produce clinical symptoms that resemble appendicitis and diverticulitis, leading to unnecessary surgery. Trovato *et al*[29] reported a case of epiploic appendagitis with left lower abdominal pain, whose symptoms were similar to diseases such as diverticulitis and could be treated with conservative treatment. These possible differential diagnoses should all be taken into account.

For appendico-vesicocolonic fistulas, if they are secondary to a benign disease and the symptoms are mild, conservative treatment such as antibiotics can be attempted. However, relevant literature shows that the closure rate of the fistula remains low and there is a significant risk of infectious complications after conservative treatment[30]. Surgical treatment is recommended for appendico-vesicocolonic fistulas caused by malignant diseases or with more severe symptoms that affect normal life. The preferred surgical treatment strategy is a one-stage resection and anastomosis of the disordered bowel segment with or without prophylactic ileum or colon neostomy[16]. However, Hsieh *et al*[31] showed that the one-stage operation of enterovesical fistulas should be limited to patients with good nutrition and no severe inflammation, radiation injuries, intestinal obstructions, or advanced malignant tumors present and should not be pursued in elderly patients. In the study by Smeek *et al*[32], 53% of the patients underwent secondary surgery due to the advanced age of the patients in the study and the high rate of complications. For the bladder, if the lesion is caused by a benign disease, it is feasible to resect and suture the bladder wall. If malignant tumors involve the trigone of the bladder, a partial resection of the bladder wall with the preservation of the bladder may not be safe. If located outside the trigone of the bladder, partial bladder wall resection with bladder preservation should be considered[17]. There is a marked difference between benign and malignant disease in the treatment of appendiceal-colonic fistulas. If the lesion is benign, resection of the appendix, the fistula and part of the colon should be considered[33]. If the lesion is malignant, a right hemicolectomy, lymph node dissection with resection of the fistula and part of colon should be considered[34]. Specific methods of surgery include open

Table 1 Comparison of 4 cases of appendico-vesicocolonic fistula

Ref.	Age and gender (man-M/woman-F)	Etiology	Clinical manifestation	Library examinations	Imaging examinations	Treatment
Marsha <i>et al</i> [6], 1975	55/M	Chronic appendicitis	Frequent urination, diarrhea	Urine culture: <i>E. coli</i>	Barium enema: fistula	Open surgery (lesion resection + colostomy)
Blalock[7], 1981	45/M	Appendicular abscess	Diarrhea, urinary tract infections, gas urine, urine fecal stain	Urine culture: <i>E. coli</i>	Excretory imaging, barium enema, cystoscopy: Nonspecific changes	Open operation (lesion resection)
Kathie <i>et al</i> [8], 1983	15/M	Appendicular abscess, trauma	Abdominal pain, nausea→trauma→abdominal pain, diarrhea, frequent urination, dysuria	Urine culture: <i>E. coli</i> Urine routine: WBC	Colonoscopy, cystoscopy, barium enema: nonspecific findings. Excretory imaging: Extravasation	Open surgery (removal of lesion)
Our 2022	77/M	Acute or chronic inflammation of the colon and inflammation of the appendix	Diarrhea, urine and feces, frequent and urgent urination	Urine routine: WBC, RBC (+)	CT: Fistula visible cystoscopy & colonoscopy: Fistula visible	Laparoscopic surgery (lesion resection + ileostomy)

E. coli: *Escherichia coli*; CT: Computed tomography; WBC: White blood cell; RBC: Red blood cell.

surgery, laparoscopic surgery and robot-assisted surgery. Open surgery is considered the traditional method of treatment, and laparoscopy has been widely used with the maturity of technology in recent decades. Chung *et al*[35], Albrecht *et al*[1], Lee *et al*[36] and others reported successful cases using laparoscopic surgery. In our case, the laparoscopic technique was also used to successfully remove the diseased intestinal segment and repair the fistula. Kibar *et al*[37] reported a successful case using robot-assisted enterovesical fistula surgery. Postoperative complications include anastomotic fistulas and urinary tract infections. Our patient also developed a urinary tract infection after surgery. The main influencing factors for poor anastomotic healing include radiotherapy, the American Society of Anaesthesiologists (ASA) score, hormone therapy, emergency surgery, lack of surgical experience and a higher body mass index (BMI)[38].

The case presented here is the fourth reported case of an appendico-vesicocolonic fistula to date worldwide (Table 1). Marsha *et al*[6] previously reported a case in 1975 involving a 55-year-old male patient with bladder irritation and diarrhea as the main manifestations. A urine culture showed an *E. coli* infection, and a barium enema showed that the contrast agent quickly entered the bladder and filled the colon. Intraoperative exploration revealed an appendico-vesicocolonic fistula. Colostomy was performed after the resection of the lesion, and the stoma was closed 3 mo after surgery. Postoperative pathology showed chronic inflammatory changes. Blalock[7] reported a case in 1981 involving a 45-year-old male patient with diarrhea, urinary tract infection, pneumaturia and fecal contamination as the main manifestations. A urine culture showed an *Escherichia coli* infection. The excretory radiography, barium enema, and cystoscopy showed nonspecific inflammatory changes. During laparotomy, the necrotic tip of the appendix was found in the abscess cavity connected with the sigmoid colon and bladder, and the lesion was removed. Postoperative pathology suggested severe inflammatory changes. Keane and Tebala[18] reported a 15-year-old male who developed abdominal pain, dysuria, frequent urination and diarrhea after conservative treatment for appendicitis and upper abdominal trauma in

1983. A urinalysis revealed leukocytes present in the urine, and a urine culture grew *Escherichia coli*. Cystourethrography showed an extravasation of contrast medium, but no obvious abnormality was found by colonoscopy and cystoscopy. An exploratory laparotomy revealed fistulas in the appendix, bladder, and rectum, and the lesions were removed. In the 77-year-old man reported in our case, the main manifestations were diarrhea, frequent and urgent urination, and fecal contamination of the urine. Routine urine tests showed a significant increase in white blood cells and red blood cells, the CT showed fistula formation, and cystoscopy and colonoscopy showed fistulas. A laparoscopic resection of the lesion was followed by a protective ileum neostomy. Pathological findings indicated inflammation of the appendix and colon. This is the fourth publicly reported case of an appendico-vesicocolonic fistula. It was also the first double fistula case having an appendico-vesicocolonic fistula combined with an appendiceal-colonic fistula.

CONCLUSION

This case is the first double fistula case involving an appendico-vesicocolonic fistula with an appendiceal-colonic fistula that has been publicly reported in the world, and it is also the first case of an appendico-vesicocolonic fistula treated by laparoscopic surgery. The clinical manifestations of this case varied, including feces in the urine, pneumaturia, complicated urinary tract infection and abdominal discomfort. However, the symptoms of this condition often lack specificity. The sensitivity and specificity of different examination methods vary greatly, and the combined application of multiple examinations is often required. The main treatment was the excision of the diseased intestine and bladder. Laparoscopic surgery can be performed by a skilled surgeon with satisfactory results.

FOOTNOTES

Author contributions: Yan H reviewed the literature and contributed to data collection and manuscript drafting; Wu YC and Zuo S were the patient's surgeons and were responsible for the revision of the manuscript; all other members equally contributed to the medical treatment and manuscript drafting.

Informed consent statement: Informed written consent was obtained from the patient's parents for publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Han Yan 0000-0003-1341-3667; Ying-Chao Wu 0000-0002-1831-1056; Xin Wang 0000-0002-0381-5497; Yu-Cun Liu 0000-0002-6158-755X; Shuai Zuo 0000-0003-2187-1246; Peng-Yuan Wang 0000-0003-3884-0227.

S-Editor: Chen YL

L-Editor: A

P-Editor: Chen YL

REFERENCES

- 1 Albrecht K, Schumann R, Peitgen K, Walz MK. [Laparoscopic therapy of appendicovesical fistula -- two case reports]. *Zentralbl Chir* 2004; **129**: 396-398 [PMID: 15486792 DOI: 10.1055/s-2004-820387]
- 2 Wang W, Wang L, Xu J, Shi S, Tian Y, Zhang Y. Combination of CT imaging and endoscopy in diagnosis of appendicovesical fistula caused by appendiceal adenocarcinoma. *J Xray Sci Technol* 2014; **22**: 493-501 [PMID: 25080116 DOI: 10.3233/XST-140440]
- 3 Agarwal S, Sharma A, Aggarwal A, Sankhwar S. Air in the urine: a mystified patient and a concerned surgeon. *BMJ Case Rep* 2019; **12** [PMID: 30635317 DOI: 10.1136/bcr-2018-227334]
- 4 Garcea G, Majid I, Sutton CD, Pattenden CJ, Thomas WM. Diagnosis and management of colovesical fistulae; six-year

- experience of 90 consecutive cases. *Colorectal Dis* 2006; **8**: 347-352 [PMID: 16630242 DOI: 10.1111/j.1463-1318.2005.00928.x]
- 5 **Small WP**, Smith AN. Fistula and conditions associated with diverticular disease of the colon. *Clin Gastroenterol* 1975; **4**: 171-199 [PMID: 1167350]
 - 6 **Marshall VR**, Molland E, Bland JP. Appendico-vesico-colic fistula. *Br J Urol* 1975; **47**: 544 [PMID: 1191924 DOI: 10.1111/j.1464-410x.1975.tb06258.x]
 - 7 **Blalock ARK**. Appendico-vesicocolonic fistula. *Br J Urol* 1981; **53**: 190
 - 8 **Dalessandri KM**, Swafford GR. Appendico-vesicocolonic fistula. *J Urol* 1983; **130**: 777 [PMID: 6887418 DOI: 10.1016/s0022-5347(17)51457-3]
 - 9 **Cripps H**. Passage of air and feces from the urethra. *Lancet* 1888; **2**: 619 [DOI: 10.1016/s0140-6736(02)24860-6]
 - 10 **Cherigie**, Proux, Bourdon. [Sigmoid pseudocancer, appendiculo-sigmoid fistula, resection of the fistula, postoperative obstruction, study of the small intestine]. *Arch Mal Appar Dig Mal Nutr* 1953; **42**: 989-991 [PMID: 13114966]
 - 11 **Eglinton T**, Nguyen T, Raniga S, Dixon L, Dobbs B, Frizelle FA. Patterns of recurrence in patients with acute diverticulitis. *Br J Surg* 2010; **97**: 952-957 [PMID: 20474006 DOI: 10.1002/bjs.7035]
 - 12 **Najjar SF**, Jamal MK, Savas JF, Miller TA. The spectrum of colovesical fistula and diagnostic paradigm. *Am J Surg* 2004; **188**: 617-621 [PMID: 15546583 DOI: 10.1016/j.amjsurg.2004.08.016]
 - 13 **Dangle PP**, Wang WP, Pohar KS. Vesicoenteric, vesicovaginal, vesicocutaneous fistula -an unusual complication with intravesical mitomycin. *Can J Urol* 2008; **15**: 4269-4272 [PMID: 18814818]
 - 14 **Qiu SL**, Yu JW, Liu XH, Wang HY. Diagnosis and treatment of 66 cases with vesicoenteric fistula. *Chin J Postgrad Med* 2019; **42** [DOI: 10.1007/s11684-008-0011-4]
 - 15 **Abubakar AM**, Pindiga UH, Chinda JY, Nggada HA. Appendicovesical fistula associated with Hirschsprung's disease. *Pediatr Surg Int* 2006; **22**: 617-618 [PMID: 16733721 DOI: 10.1007/s00383-006-1691-y]
 - 16 **Melchior S**, Cudovic D, Jones J, Thomas C, Gillitzer R, Thüroff J. Diagnosis and surgical management of colovesical fistulas due to sigmoid diverticulitis. *J Urol* 2009; **182**: 978-982 [PMID: 19616793 DOI: 10.1016/j.juro.2009.05.022]
 - 17 **Yagi Y**, Shoji Y, Sasaki S, Yoshikawa A, Tsukioka Y, Fukushima W, Hirosawa H, Izumi R, Saito K. Sigmoid colon cancer arising in a diverticulum of the colon with involvement of the urinary bladder: a case report and review of the literature. *BMC Gastroenterol* 2014; **14**: 90 [PMID: 24884743 DOI: 10.1186/1471-230X-14-90]
 - 18 **Keane S**, Tebala GD. Appendicovesical fistula presenting as hypokalaemic hyperchloraemic metabolic acidosis: a case report. *Ann R Coll Surg Engl* 2019; **101**: e131-e132 [PMID: 31155884 DOI: 10.1308/rcsann.2019.0047]
 - 19 **Hakim S**, Amin M, Cappell MS. Limited, local, extracolonic spread of mucinous appendiceal adenocarcinoma after perforation with formation of a malignant appendix-to-sigmoid fistula: Case report and literature review. *World J Gastroenterol* 2016; **22**: 8624-8630 [PMID: 27784975 DOI: 10.3748/wjg.v22.i38.8624]
 - 20 **Trinavarat P**, Mahayosnond A, Riccabona M. Appendico-vesical fistula: A clinical and imaging challenge—A case report based discussion of imaging implication. *Eur J Radiol Extra* 2009; **71**: e101-e103 [DOI: 10.1016/j.ejrex.2009.04.002]
 - 21 **Scozzari G**, Arezzo A, Morino M. Enterovesical fistulas: diagnosis and management. *Tech Coloproctol* 2010; **14**: 293-300 [PMID: 20617353 DOI: 10.1007/s10151-010-0602-3]
 - 22 **Tsivian A**, Kyzer S, Shtricker A, Benjamin S, Sidi AA. Laparoscopic treatment of colovesical fistulas: technique and review of the literature. *Int J Urol* 2006; **13**: 664-667 [PMID: 16771754 DOI: 10.1111/j.1442-2042.2006.01382.x]
 - 23 **Leicht W**, Thomas C, Thüroff J, Roos F. [Colovesical fistula caused by diverticulitis of the sigmoid colon: diagnosis and treatment]. *Urologe A* 2012; **51**: 971-974 [PMID: 22772496 DOI: 10.1007/s00120-012-2910-x]
 - 24 **Ravichandran S**, Ahmed HU, Matanhelia SS, Dobson M. Is there a role for magnetic resonance imaging in diagnosing colovesical fistulas? *Urology* 2008; **72**: 832-837 [PMID: 18718640 DOI: 10.1016/j.urology.2008.06.036]
 - 25 **Nakamoto H**, Yokota R, Namba H, Ishikawa T, Yamada K, Hosoda M, Taguchi K. A Benign Appendiceal-Colonic Fistula, Diagnosed and Managed Laparoscopically: A Case Report. *Am J Case Rep* 2020; **21**: e925946 [PMID: 33162549 DOI: 10.12659/AJCR.925946]
 - 26 **Morris-Stiff GJ**, Islam KA. Appendico-colic fistula complicating appendicitis in cystic fibrosis. *BMJ Case Rep* 2010; **2010** [PMID: 22778186 DOI: 10.1136/bcr.02.2010.2714]
 - 27 **Purysko AS**, Remer EM, Filho HM, Bittencourt LK, Lima RV, Racy DJ. Beyond appendicitis: common and uncommon gastrointestinal causes of right lower quadrant abdominal pain at multidetector CT. *Radiographics* 2011; **31**: 927-947 [PMID: 21768232 DOI: 10.1148/rg.314105065]
 - 28 **van Breda Vriesman AC**, Lohle PN, Coerkamp EG, Puylaert JB. Infarction of omentum and epiploic appendage: diagnosis, epidemiology and natural history. *Eur Radiol* 1999; **9**: 1886-1892 [PMID: 10602970 DOI: 10.1007/s003300050942]
 - 29 **Trovato P**, Simonetti I, Verde F, Lomoro P, Vinci G, Tarotto L, Corvino F, Corvino A. Acute epiploic appendagitis: ultrasound and computed tomography findings of a rare case of acute abdominal pain and the role of other imaging techniques. *Pol J Radiol* 2020; **85**: e178-e182 [PMID: 32419882 DOI: 10.5114/pjr.2020.94335]
 - 30 **Granieri S**, Sessa F, Bonomi A, Paleino S, Bruno F, Chierici A, Sciannamea IM, Germini A, Campi R, Talso M, Facciorusso A, Deiana G, Serni S, Cotsoglou C. Indications and outcomes of enterovesical and colovesical fistulas: systematic review of the literature and meta-analysis of prevalence. *BMC Surg* 2021; **21**: 265 [PMID: 34044862 DOI: 10.1186/s12893-021-01272-6]
 - 31 **Hsieh JH**, Chen WS, Jiang JK, Lin TC, Lin JK, Hsu H. Enterovesical fistula: 10 years experience. *Zhonghua Yi Xue Za Zhi (Taipei)* 1997; **59**: 283-288 [PMID: 9248120]
 - 32 **Smeenk RM**, Plaisier PW, van der Hoeven JA, Hesp WL. Outcome of surgery for colovesical and colovaginal fistulas of diverticular origin in 40 patients. *J Gastrointest Surg* 2012; **16**: 1559-1565 [PMID: 22653331 DOI: 10.1007/s11605-012-1919-1]
 - 33 **Joo JS**, Agachan F, Wexner SD. Laparoscopic surgery for lower gastrointestinal fistulas. *Surg Endosc* 1997; **11**: 116-118 [PMID: 9069139 DOI: 10.1007/s004649900310]
 - 34 **Knigge U**, Hansen CP. [Appendiceal carcinoid tumors and goblet cell carcinoids]. *Ugeskr Laeger* 2010; **172**: 1678-1681 [PMID: 20525467]

- 35 **Chung CW**, Kim KA, Chung JS, Park DS, Hong JY, Hong YK. Laparoscopic treatment of appendicovesical fistula. *Yonsei Med J* 2010; **51**: 463-465 [PMID: [20376905](#) DOI: [10.3349/ymj.2010.51.3.463](#)]
- 36 **Lee JW**, Ahn JH, Ha HK. Vesico-Appendiceal Fistula Misdiagnosed as Meckel's Diverticulum: A Laparoscopic Approach. *World J Mens Health* 2012; **30**: 195-197 [PMID: [23596613](#) DOI: [10.5534/wjmh.2012.30.3.195](#)]
- 37 **Kibar Y**, Yalcin S, Kopru B, Kaya E, Topuz B, Ebiloglu T. Robot-Assisted Laparoscopic Repair of Spontaneous Appendicovesical Fistula. *J Endourol Case Rep* 2016; **2**: 111-113 [PMID: [27579435](#) DOI: [10.1089/cren.2016.0063](#)]
- 38 **Alves A**, Panis Y, Trancart D, Regimbeau JM, Pocard M, Valleur P. Factors associated with clinically significant anastomotic leakage after large bowel resection: multivariate analysis of 707 patients. *World J Surg* 2002; **26**: 499-502 [PMID: [11910487](#) DOI: [10.1007/s00268-001-0256-4](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

