

# World Journal of *Clinical Cases*

*World J Clin Cases* 2022 July 6; 10(19): 6341-6758



## Contents

Thrice Monthly Volume 10 Number 19 July 6, 2022

## MINIREVIEWS

- 6341** Review of clinical characteristics, immune responses and regulatory mechanisms of hepatitis E-associated liver failure  
*Chen C, Zhang SY, Chen L*
- 6349** Current guidelines for *Helicobacter pylori* treatment in East Asia 2022: Differences among China, Japan, and South Korea  
*Cho JH, Jin SY*
- 6360** Review of epidermal growth factor receptor-tyrosine kinase inhibitors administration to non-small-cell lung cancer patients undergoing hemodialysis  
*Lan CC, Hsieh PC, Huang CY, Yang MC, Su WL, Wu CW, Wu YK*

## ORIGINAL ARTICLE

## Case Control Study

- 6370** Pregnancy-related psychopathology: A comparison between pre-COVID-19 and COVID-19-related social restriction periods  
*Chieffo D, Avallone C, Serio A, Kotzalidis GD, Balocchi M, De Luca I, Hirsch D, Gonzalez del Castillo A, Lanzotti P, Marano G, Rinaldi L, Lanzone A, Mercuri E, Mazza M, Sani G*
- 6385** Intestinal mucosal barrier in functional constipation: Does it change?  
*Wang JK, Wei W, Zhao DY, Wang HF, Zhang YL, Lei JP, Yao SK*

## Retrospective Cohort Study

- 6399** Identification of risk factors for surgical site infection after type II and type III tibial pilon fracture surgery  
*Hu H, Zhang J, Xie XG, Dai YK, Huang X*

## Retrospective Study

- 6406** Total knee arthroplasty in Ranawat II valgus deformity with enlarged femoral valgus cut angle: A new technique to achieve balanced gap  
*Lv SJ, Wang XJ, Huang JF, Mao Q, He BJ, Tong PJ*
- 6417** Preliminary evidence in treatment of eosinophilic gastroenteritis in children: A case series  
*Chen Y, Sun M*
- 6428** Self-made wire loop snare successfully treats gastric persimmon stone under endoscopy  
*Xu W, Liu XB, Li SB, Deng WP, Tong Q*
- 6437** Neoadjuvant transcatheter arterial chemoembolization and systemic chemotherapy for the treatment of undifferentiated embryonal sarcoma of the liver in children  
*He M, Cai JB, Lai C, Mao JQ, Xiong JN, Guan ZH, Li LJ, Shu Q, Ying MD, Wang JH*

- 6446** Effect of cold snare polypectomy for small colorectal polyps

*Meng QQ, Rao M, Gao PJ*

- 6456** Field evaluation of COVID-19 rapid antigen test: Are rapid antigen tests less reliable among the elderly?

*Tabain I, Cucevic D, Skreb N, Mrzljak A, Ferencak I, Hruskar Z, Misic A, Kuzle J, Skoda AM, Jankovic H, Vilibic-Cavlek T*

### Observational Study

- 6464** Tracheobronchial intubation using flexible bronchoscopy in children with Pierre Robin sequence: Nursing considerations for complications

*Ye YL, Zhang CF, Xu LZ, Fan HF, Peng JZ, Lu G, Hu XY*

- 6472** Family relationship of nurses in COVID-19 pandemic: A qualitative study

*Çelik MY, Kiliç M*

### META-ANALYSIS

- 6483** Diagnostic accuracy of  $\geq 16$ -slice spiral computed tomography for local staging of colon cancer: A systematic review and meta-analysis

*Liu D, Sun LM, Liang JH, Song L, Liu XP*

### CASE REPORT

- 6496** Delayed-onset endophthalmitis associated with *Achromobacter* species developed in acute form several months after cataract surgery: Three case reports

*Kim TH, Lee SJ, Nam KY*

- 6501** Sustained dialysis with misplaced peritoneal dialysis catheter outside peritoneum: A case report

*Shen QQ, Behera TR, Chen LL, Attia D, Han F*

- 6507** Arteriovenous thrombotic events in a patient with advanced lung cancer following bevacizumab plus chemotherapy: A case report

*Kong Y, Xu XC, Hong L*

- 6514** Endoscopic ultrasound radiofrequency ablation of pancreatic insulinoma in elderly patients: Three case reports

*Rossi G, Petrone MC, Capurso G, Partelli S, Falconi M, Arcidiacono PG*

- 6520** Acute choroidal involvement in lupus nephritis: A case report and review of literature

*Yao Y, Wang HX, Liu LW, Ding YL, Sheng JE, Deng XH, Liu B*

- 6529** Triple A syndrome-related achalasia treated by per-oral endoscopic myotomy: Three case reports

*Liu FC, Feng YL, Yang AM, Guo T*

- 6536** Choroidal thickening with serous retinal detachment in BRAF/MEK inhibitor-induced uveitis: A case report

*Kiraly P, Groznik AL, Valentinčič NV, Mekjavić PJ, Urbančič M, Ocvirk J, Mesti T*

- 6543** Esophageal granular cell tumor: A case report

*Chen YL, Zhou J, Yu HL*

- 6548** Hem-o-lok clip migration to the common bile duct after laparoscopic common bile duct exploration: A case report  
*Liu DR, Wu JH, Shi JT, Zhu HB, Li C*
- 6555** Chidamide and sintilimab combination in diffuse large B-cell lymphoma progressing after chimeric antigen receptor T therapy  
*Hao YY, Chen PP, Yuan XG, Zhao AQ, Liang Y, Liu H, Qian WB*
- 6563** Relapsing polychondritis with isolated tracheobronchial involvement complicated with Sjogren's syndrome: A case report  
*Chen JY, Li XY, Zong C*
- 6571** Acute methanol poisoning with bilateral diffuse cerebral hemorrhage: A case report  
*Li J, Feng ZJ, Liu L, Ma YJ*
- 6580** Immunoabsorption therapy for Klinefelter syndrome with antiphospholipid syndrome in a patient: A case report  
*Song Y, Xiao YZ, Wang C, Du R*
- 6587** Roxadustat for treatment of anemia in a cancer patient with end-stage renal disease: A case report  
*Zhou QQ, Li J, Liu B, Wang CL*
- 6595** Imaging-based diagnosis for extraskeletal Ewing sarcoma in pediatrics: A case report  
*Chen ZH, Guo HQ, Chen JJ, Zhang Y, Zhao L*
- 6602** Unusual course of congenital complete heart block in an adult: A case report  
*Su LN, Wu MY, Cui YX, Lee CY, Song JX, Chen H*
- 6609** Penile metastasis from rectal carcinoma: A case report  
*Sun JJ, Zhang SY, Tian JJ, Jin BY*
- 6617** Isolated cryptococcal osteomyelitis of the ulna in an immunocompetent patient: A case report  
*Ma JL, Liao L, Wan T, Yang FC*
- 6626** Magnetic resonance imaging features of intrahepatic extramedullary hematopoiesis: Three case reports  
*Luo M, Chen JW, Xie CM*
- 6636** Giant retroperitoneal liposarcoma treated with radical conservative surgery: A case report and review of literature  
*Lieto E, Cardella F, Erario S, Del Sorbo G, Reginelli A, Galizia G, Urraro F, Panarese I, Auricchio A*
- 6647** Transplanted kidney loss during colorectal cancer chemotherapy: A case report  
*Pośpiech M, Kolonko A, Nieszporek T, Kozak S, Kozaczka A, Karkoszka H, Winder M, Chudek J*
- 6656** Massive gastrointestinal bleeding after endoscopic rubber band ligation of internal hemorrhoids: A case report  
*Jiang YD, Liu Y, Wu JD, Li GP, Liu J, Hou XH, Song J*

- 6664** Mills' syndrome is a unique entity of upper motor neuron disease with N-shaped progression: Three case reports  
*Zhang ZY, Ouyang ZY, Zhao GH, Fang JJ*
- 6672** Entire process of electrocardiogram recording of Wellens syndrome: A case report  
*Tang N, Li YH, Kang L, Li R, Chu QM*
- 6679** Retroperitoneal tumor finally diagnosed as a bronchogenic cyst: A case report and review of literature  
*Gong YY, Qian X, Liang B, Jiang MD, Liu J, Tao X, Luo J, Liu HJ, Feng YG*
- 6688** Successful treatment of Morbihan disease with total glucosides of paeony: A case report  
*Zhou LF, Lu R*
- 6695** Ant sting-induced whole-body pustules in an inebriated male: A case report  
*Chen SQ, Yang T, Lan LF, Chen XM, Huang DB, Zeng ZL, Ye XY, Wan CL, Li LN*
- 6702** Plastic surgery for giant metastatic endometrioid adenocarcinoma in the abdominal wall: A case report and review of literature  
*Wang JY, Wang ZQ, Liang SC, Li GX, Shi JL, Wang JL*
- 6710** Delayed-release oral mesalamine tablet mimicking a small jejunal gastrointestinal stromal tumor: A case report  
*Frosio F, Rausa E, Marra P, Boutron-Ruault MC, Lucianetti A*
- 6716** Concurrent alcoholic cirrhosis and malignant peritoneal mesothelioma in a patient: A case report  
*Liu L, Zhu XY, Zong WJ, Chu CL, Zhu JY, Shen XJ*
- 6722** Two smoking-related lesions in the same pulmonary lobe of squamous cell carcinoma and pulmonary Langerhans cell histiocytosis: A case report  
*Gencer A, Ozcibik G, Karakas FG, Sarbay I, Batur S, Borekci S, Turna A*
- 6728** Proprotein convertase subtilisin/kexin type 9 inhibitor non responses in an adult with a history of coronary revascularization: A case report  
*Yang L, Xiao YY, Shao L, Ouyang CS, Hu Y, Li B, Lei LF, Wang H*
- 6736** Multimodal imaging study of lipemia retinalis with diabetic retinopathy: A case report  
*Zhang SJ, Yan ZY, Yuan LF, Wang YH, Wang LF*
- 6744** Primary squamous cell carcinoma of the liver: A case report  
*Kang LM, Yu DP, Zheng Y, Zhou YH*
- 6750** Tumor-to-tumor metastasis of clear cell renal cell carcinoma to contralateral synchronous pheochromocytoma: A case report  
*Wen HY, Hou J, Zeng H, Zhou Q, Chen N*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Abdulqadir Jeprel Naswhan, MSc, RN, Director, Research Scientist, Senior Lecturer, Senior Researcher, Nursing for Education and Practice Development, Hamad Medical Corporation, Doha 576214, Qatar. anashwan@hamad.qa

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJCC as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The WJCC's CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Xu Guo; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lai Wang.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

July 6, 2022

**COPYRIGHT**

© 2022 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Roxadustat for treatment of anemia in a cancer patient with end-stage renal disease: A case report

Qiao-Qiao Zhou, Jing Li, Bin Liu, Chun-Li Wang

**Specialty type:** Medicine, research and experimental

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

**P-Reviewer:** Cabezu AS, Spain

**Received:** November 23, 2021

**Peer-review started:** November 23, 2021

**First decision:** April 8, 2022

**Revised:** April 16, 2022

**Accepted:** May 8, 2022

**Article in press:** May 8, 2022

**Published online:** July 6, 2022



**Qiao-Qiao Zhou, Jing Li, Chun-Li Wang,** Hemodialysis Room, Sichuan Cancer Hospital and Institute, Sichuan Cancer Center, School of Medicine, University of Electronic Science and Technology of China, Chengdu 610042, Sichuan Province, China

**Bin Liu,** Department of Internal Medicine-Oncology, Sichuan Cancer Hospital and Institute, Sichuan Cancer Center, School of Medicine, University of Electronic Science and Technology of China, Chengdu 610042, Sichuan Province, China

**Corresponding author:** Jing Li, PhD, Professor, Hemodialysis Room, Sichuan Cancer Hospital and Institute, Sichuan Cancer Center, School of Medicine, University of Electronic Science and Technology of China, No. 55 Section 4, Renmin South Road, Chengdu 610042, Sichuan Province, China. [24978201@qq.com](mailto:24978201@qq.com)

### Abstract

#### BACKGROUND

Most cancer patients are accompanied by anemia, which will be more serious when combined with end-stage renal disease (ESRD). At present, cancer-related anemia and renal anemia treatments mainly include erythropoiesis-stimulating agents (ESAs), iron supplementation, and blood transfusion, but their effects are often poor with several safety concerns. We have used roxadustat to treat anemia in a cancer patient with ESRD and achieved a successful outcome for the first time.

#### CASE SUMMARY

A 64-year-old man was diagnosed with right renal cancer (clear cell renal cell carcinoma). He did not receive surgery or radiotherapy before admission. He was treated with oral sunitinib (sunitinib malate) on April 18, 2017. During oral chemotherapy, he had numerous complications, including anemia, hypertension, thyroid hypofunction, skin pigment loss, and renal function deterioration. At last, he progressed to ESRD and began hemodialysis treatment. We initially treated the patient with high-dose ESAs, iron supplementation, adequate dialysis, and even blood transfusion, but his anemia did not improve. Roxadustat is a newly developed drug for renal anemia treatment, but not for cancer-related anemia, let alone to treat anemia in cancer patients with ESRD. We prescribed oral roxadustat to the patient. After a period, his hemoglobin gradually increased. He did not have obvious discomfort symptoms, and his tumor did not progress significantly.

#### CONCLUSION

Oral roxadustat could achieve good results in treating anemia in cancer patients



with ESRD.

**Key Words:** Cancer; End-stage renal disease; Hemodialysis; Anemia; Roxadustat; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Currently, erythropoiesis-stimulating agents (ESAs) are the cornerstones for the treatment of renal anemia or cancer-related anemia. Roxadustat is a newly developed drug for renal anemia treatment, but not for cancer-related anemia, let alone to treat anemia in cancer patients with end-stage renal disease (ESRD). At the beginning, the patient was treated with high-dose ESAs, iron supplementation, and even blood transfusion, but his anemia did not improve. However, after orally taking roxadustat, his hemoglobin gradually increased without significant tumor progression. This is the first case using roxadustat to successfully improve anemia in a cancer patient with ESRD.

**Citation:** Zhou QQ, Li J, Liu B, Wang CL. Roxadustat for treatment of anemia in a cancer patient with end-stage renal disease: A case report. *World J Clin Cases* 2022; 10(19): 6587-6594

**URL:** <https://www.wjgnet.com/2307-8960/full/v10/i19/6587.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v10.i19.6587>

## INTRODUCTION

Most cancer patients have anemia symptoms. If they progress to end-stage renal disease (ESRD) and need maintenance hemodialysis treatment, anemia will be more difficult to correct. At present, cancer-related anemia and renal anemia treatments mainly include erythropoiesis-stimulating agents (ESAs), iron supplementation, and blood transfusion. However, ESAs are often used at high dosage and accompanied by thrombosis, hypertension, and other adverse events. Hypoxia-inducible factor prolyl hydroxylase inhibitors (HIF-PHI) are newly developed, oral-based small molecule drugs for the treatment of renal anemia, of which roxadustat (FG-4592, FibroGen, Astellas, AstraZeneca) has been listed in China and Japan[1]. To our best knowledge, it has not been used to treat cancer-related anemia. In this case report, we, for the first time, used roxadustat to treat anemia in a cancer patient with ESRD and achieved a successful outcome.

## CASE PRESENTATION

### Chief complaints

A 64-year-old man was admitted to our hospital because of general fatigue and tight breath. He had been on maintenance hemodialysis for 1 year.

### History of present illness

The patient developed ESRD 2 years ago due to oral chemotherapy for right renal cancer and began hemodialysis on May 23, 2019. At that time, his hemoglobin (Hb) level was 79 g/L. He was given 10000 U of recombinant human erythropoietin (rHuEPO) (weight 50 kg) twice a week *via* intravenous injection. However, after 2 mo of treatment, his Hb level was still 77 g/L. He was then intravenously injected with 100 mg iron sucrose once a week for 10 wk. His Hb gradually rose to about 100 g/L for a short period and began to decline. During that time, we actively looked for the causes of anemia. The patient had no gastrointestinal bleeding, evident inflammation, hemolytic anemia, erythropoietin (EPO) resistance, and other factors. Therefore, he was given 10000 U rHuEPO three times a week and subjected to adequate hemodialysis three times *per* week and hemodiafiltration once every 2 wk. In addition, he was prescribed oral ketoacids, intravenous infusion of human albumin, and other nutritional support treatments. However, these measures did not work, and his Hb dropped from 100 g/L to 65 g/L within 5 mo. Meanwhile, he developed discomfort symptoms such as increased heart rate, tight breath, and general fatigue.

### History of past illness

The patient had been diagnosed with right renal cancer (clear cell renal carcinoma) in 2017. He did not receive surgery or radiotherapy. He was orally given 50 mg of soltan (sunitinib malate) daily for 14 d since April 18, 2017. He stopped taking it for 7 d and began to retake it. During oral chemotherapy, he had numerous complications, including anemia, hypertension, positive urine protein, thyroid



hypofunction, skin pigment loss, and renal function deterioration. At last, he progressed to ESRD.

He took amlodipine besylate tablets 5 mg daily to control blood pressure and levothyroxine sodium tablets 150 mg to treat hypothyroidism.

### Personal and family history

The patient had no history of smoking or drinking and no family history of similar diseases.

### Physical examination

The vital signs of the patient were as follows: Body temperature, 36.5 °C; blood pressure, 168/82 mmHg; pulse rate, 98 beats/min; and respiratory rate, 26 breaths/min. He was in a semi-recumbent position, and his eyelid conjunctiva was pale. His bilateral thoracic movement weakened, palpation speech fibrillation weakened, percussion sounded dull, and auscultation respiratory sound disappeared. His heart rate was 98 beats/min, his rhythm was neat, and there was no obvious murmur. His abdominal examination showed no obvious abnormalities, and his lower limbs had slight edema.

### Laboratory examinations

The blood examination of the patient after admission showed an Hb level of 65 g/L, erythrocyte count of  $2.29 \times 10^{12}/L$ , reticulocyte count of  $0.079 \times 10^{12}/L$ , serum ferritin level of 345 mg/L, transferrin saturation of 19.9%, EPO level of 28.6 mIU/mL, C-reactive protein level of 15.41 mg/L, albumin level of 24.4 g/L, and intact parathyroid hormone level of 27.3 pg/mL. The direct Coombs test was negative, and anti-EPO antibodies were negative (Table 1).

### Imaging examinations

The plain chest computed tomography (CT) scan showed a large amount of effusion in the bilateral thorax, marks on the right, and partial compression atelectasis of both lungs. The plain abdominal CT scan (Figure 1A) showed an enlarged right kidney, irregular soft tissue occupying, unclear boundary, an about 59.4 mm × 49.7 mm larger layer, and thickened perirenal fascia. In addition, nodules with soft tissue density were seen at the lower pole of the left kidney, with a larger layer of about 26.0 mm × 22.1 mm.

## FINAL DIAGNOSIS

The patient was diagnosed with chronic kidney disease (CKD) stage 5, renal anemia, maintenance hemodialysis, right renal cancer, cancer-related anemia, bilateral pleural effusion, very high-risk grade 3 hypertension, and hypoproteinemia.

## TREATMENT

During outpatient dialysis, to correct anemia, we increased the patient's rHuEPO dosage from 20000 U *per* week to 30000 U *per* week and gave him a total of 1 g of iron sucrose intravenously. Therefore, he was treated with 1.5 units of red blood cell transfusion after hospitalization. To alleviate the tight breath symptoms, he was subjected to pleural puncture. Examination of the collected chest fluid identified some cancer cells. He was continuously treated with hemodialysis three times a week and hemodiafiltration once every 2 wk. He was also given oral ketoacids, intravenous infusion of human albumin, and other nutritional support treatments. After that, his Hb rose to 87 g/L, but his serum ferritin was as high as 1252.2 mg/L (Figure 2). His general fatigue symptoms were markedly relieved. After a series of symptomatic treatments, the patient's condition improved, and he was discharged from the hospital.

Because the patient was given a very high dosage of rHuEPO, we were worried that further increasing rHuEPO dosage would lead to severe complications. At the same time, the patient had an iron overload. Therefore, we recommended oral roxadustat to improve his anemia. However, the patient was only willing to accept 20 mg rather than the standard starting dose of 100 mg roxadustat treatment three times a week because of the financial burden since August 1, 2020.

## OUTCOME AND FOLLOW-UP

During the 4 mo of follow-up, the patient's Hb remained stable at about 90 g/L. We repeatedly advised him to increase roxadustat dosage. Finally, he agreed to take 50 mg oral roxadustat three times a week from December 8, 2020. Two months later, his Hb level rose to about 100 g/L (Figure 2). During the course of oral administration, he was regularly monitored for blood parameters and subjected to abdominal CT. His iron index was further improved (Figure 3). The plain abdomen CT scan (Figure 1B)

**Table 1 Laboratory data before blood transfusion**


Parameter	Recorded value	Reference range
White blood cell count ( $\times 10^9/L$ )	5.22	4.0-10.0
Red blood cell count ( $\times 10^{12}/L$ )	2.29	4.0-5.5
Hemoglobin (g/L)	65	120-160
Platelet count ( $\times 10^9/L$ )	146	100-300
Hematocrit (%)	22	38.0-50.8
Mean corpuscular volume (fl)	95.9	80-100
Mean corpuscular hemoglobin (pg)	29.1	27-31
Mean corpuscular hemoglobin concentration (g/L)	304	320-360
Reticulocyte count ( $\times 10^{12}/L$ )	0.079	0.024-0.084
Reticulocyte percentage (%)	3.45	0.5-1.5
Iron ( $\mu\text{mol}/L$ )	7.68	10.6-36.7
Serum ferritin ( $\mu\text{g}/L$ )	345	30-300
Total iron binding capacity ( $\mu\text{mol}/L$ )	26.51	45-75
Transferrin saturation (%)	19.9	20-50
C-reactive protein (mg/L)	15.41	< 10
Albumin (g/L)	24.4	40.0-55.0
Calcium (mmol/L)	2.01	2.11-2.52
Phosphonium (mmol/L)	1.58	0.85-1.51
Intact parathyroid hormone (pg/ml)	27.3	150-300
Blood urea nitrogen (mmol/L)	21.66	3.6-9.5
Serum creatinine ( $\mu\text{mol}/L$ )	556	57.0-111.0
EPO concentration (mIU/ml)	28.6	4.3-29
Direct Coombs (IgG, C3) tests	Negative	Negative
Anti-EPO antibodies	Negative	Negative
Fecal occult blood test	Negative	Negative

EPO: Erythropoietin.

showed an enlarged right kidney, irregular soft tissue occupying, unclear boundary, and an about 59.2 mm  $\times$  50.6 mm larger layer. Compared with previous CT, there was no significant change in tumor size. Meanwhile, he did not have obvious discomfort.


## DISCUSSION

To our best knowledge, this is the first case report of roxadustat for the treatment of anemia in a cancer patient with ESRD. Many studies have shown that 30% to 90% of cancer patients have anemia. The incidence and severity of anemia are related to the age, tumor type, tumor stage, disease progression, treatment plan, dosage of chemotherapy drugs, inflammation, and other factors[2,3]. Based on the data from Chinese clinical trials, the prevalence of anemia is significantly higher in patients with CKD than in the general population and increases with the progression of CKD, reaching as high as 91.6%-98.2% in dialysis patients[4-6]. A bidirectional relationship has been observed between kidney disease and cancer. On the one hand, cancer is an important complication noted in kidney disease and a major cause of morbidity and mortality in this group. On the other hand, cancer patients may develop CKD directly or indirectly due to the toxicity of tumors to the kidney itself or treatment-related factors. Currently, ESAs are the cornerstones for the treatment of renal anemia or cancer-related anemia[7] besides iron supplementation and blood transfusion.



DOI: 10.12998/wjcc.v10.i19.6587 Copyright ©The Author(s) 2022.

**Figure 1 Plain abdomen computed tomography scan.** A: Plain abdomen computed tomography scan at admission showed an enlarged right kidney, irregular soft tissue occupying, unclear boundary, and an about 59.4 mm×49.7 mm larger layer. Compared with previous computed tomography, there was no significant change in tumor size; B: Plain abdomen computed tomography scan on February 2, 2021 showed an enlarged right kidney, irregular soft tissue occupying, unclear boundary, and an about 59.2 mm × 50.6 mm larger layer.




DOI: 10.12998/wjcc.v10.i19.6587 Copyright ©The Author(s) 2022.

**Figure 2 Dynamic changes of hemoglobin and serum ferritin.** The patient was admitted to our hospital on July 24, 2020. He was treated with 1.5 units of red blood cell transfusion on that day. After discharge, he began to take 20 mg oral roxadustat three times a week on August 1, 2020. Then he began to take 50 mg oral roxadustat three times a week on December 8, 2020. Hb: Hemoglobin; SF: Serum ferritin.

ESAs are the analogs of EPO with characteristics of good tolerance and ease to use. They have been used to treat anemia patients with normal physiology and can significantly improve the quality of life of patients. However, ESAs can increase blood pressure, promote tumor growth[8], and increase the risk of stroke and thromboembolism in patients with malignant tumors[9-11]. According to the summary of the American Society of Clinical Oncology/American Society of Hematology Clinical Practice Guideline Update, ESAs should not be offered to patients with chemotherapy-associated anemia whose cancer treatment is curative in intent. Still, it could be provided to patients with chemotherapy-associated anemia whose cancer treatment is not curative in intent and whose Hb is < 10 g/dL[7].

For patients with renal failure caused by tumors or chemotherapy, continuous use of ESAs can lead to functional iron deficiency. At present, the main iron supplements are oral and parenteral[12]. However, due to the close relationship between unstable iron and oxidative stress, bacterial growth, gastro-



**Figure 3 Changes of serum ferritin and transferrin saturation.** The patient's serum ferritin was as high as 1252.2 mg/L on the second day after blood transfusion on July 25, 2020. His iron index was further improved after taking roxadustat orally in the following months. SF: Serum ferritin; TAST: Transferrin saturation.

intestinal reaction, and hypersensitivity, iron supplementation (especially intravenous iron) increases the risk of infection and death of CKD patients[13,14].

Transfusion of red blood cells or whole blood can rapidly increase Hb concentration. Thus, it is used for tumor patients with severe anemia, patients with anemia caused by acute hemorrhage, and asymptomatic anemia patients with heart disease, chronic lung disease, and cerebrovascular disease. However, transfusion may lead to a series of risks such as transfusion-related response, blood transfusion-related cycle overload, virus transmission, bacterial infection, iron overload, and red blood cell allogeneic immunity[15].

HIF, a dimer composed of  $\alpha$  and  $\beta$  subunits, is the key transcription factor regulating the body's physiological response to the change of oxygen concentration. There are three subtypes of HIF- $\alpha$ : HIF-1 $\alpha$ , HIF-2 $\alpha$ , and HIF-3 $\alpha$ . HIF-2 $\alpha$  is mainly distributed in the kidney and participates in the production of red blood cells. HIF is regulated by the prolyl hydroxylase (PHD) family. PHD mainly regulates HIF by sensing the changes of intracellular oxygen partial pressure. Under normal oxygen conditions, HIF- $\alpha$  is recognized and hydroxylated by PHD and then degraded by proteasome. Under hypoxia, PHD activity decreases, and HIF- $\alpha$  is freed from degradation. HIF- $\alpha$  and HIF- $\beta$  combine to form a dimer, promoting the upregulation of a series of genes. HIF-PHI drugs weaken the effect of PHD on HIF mainly by replacing the necessary synergistic substrate of PHD or by blocking the interference between enzyme catalytic sites and PHD substrate- $\alpha$  so that HIF can be stably expressed to regulate the transcription and expression of target genes downstream of the HIF signaling pathway[1]. HIF-PHI could promote the expression of iron metabolism-related proteins by promoting EPO production and its receptor expression and reducing hepcidin levels, thereby comprehensively regulating the production of erythrocytes[16,17]. Roxadustat (FG-4592, FibroGen, Astellas, AstraZeneca) is one of HIF-PHI and has been listed in China and Japan. On December 17, 2018, roxadustat was officially approved by the State Drug Administration of China to treat CKD anemia in dialysis patients. But it has not been used to treat cancer-related anemia, let alone to treat anemia in cancer patients with ESRD.

In our case report, the patient had both renal cancer and ESRD and was receiving hemodialysis. His anemia was caused by various factors, including insufficient EPO production, reduced EPO activity, iron deficiency, metabolic disorders, malnutrition, inflammatory state, uremic toxin, and the tumor itself. Because the patient's cancer treatment was not curative in intent, he was given ESAs to treat anemia at first. Later, to maintain Hb stability, we adjusted ESA dose and supplemented iron to ensure dialysis adequacy and improve nutritional status. However, the patient's Hb decreased rapidly (Table 1). Although the patient's tumor was incurable, he still hoped to prolong his survival time and improve his quality of life as much as possible. At this time, a large dose of ESAs could not maintain Hb stability. Although blood transfusion temporarily increased Hb, he was still in a state of iron overload and inflammation.

Many studies have shown that roxadustat can effectively treat renal anemia[18,19]. However, it has not been reported to treat cancer-related anemia, cancer complicated with renal anemia, and anemia with a combination of roxadustat and ESAs. Based on the above reasons, we advised the patient to take oral roxadustat on the basis of maintaining the original treatment regimen. Unfortunately, he received 20 mg rather than the initial 100 mg of roxadustat three times a week. He was also given 30000 U of rHuEPO *via* intravenous injections *per* week. During the subsequent follow-up months, his Hb remained at about 90 g/L, his BP slightly increased but was controllable by drugs, and his iron index was improved (Figure 3). With our repeated efforts, the patient accepted treatment with 50 mg roxadustat three times a week, and his condition further improved: His Hb gradually rose to about 100 g/L. The patient's abdominal CT (Figure 1B) revealed no apparent progression of the renal tumor. Over the period of oral roxadustat, the patient had no obvious discomfort symptoms. Therefore, we believe that roxadustat was effective for anemia treatment in cancer patients with ESRD and should be recognized as a routine treatment.

Some studies have shown that increased HIF expression in cancers is associated with a poor prognosis[20]. It is unclear whether HIF is a pro-tumor factor or whether an increased HIF level reflects the hypoxic environment in faster-growing and more aggressive tumors. Currently, its causative relationship with primary tumors has not been clinically proven. Since our report is just an individual case, more clinical trials are needed to observe the therapeutic effect and mechanism of roxadustat for anemia treatment in cancer patients with ESRD.

## CONCLUSION

Oral roxadustat is safe and effective for anemia treatment of cancer patients with ESRD undergoing hemodialysis. A combination of roxadustat and ESAs may be an additional treatment option for these patients. However, more studies are needed to assess the risk and adverse effects of roxadustat.

## FOOTNOTES

**Author contributions:** Li J designed the study; Zhou QQ collected the patient's clinical data and contributed to drafting the manuscript; Wang CL collected the patient's clinical data; Liu B provided this case; and all authors issued final approval for this version to be submitted.

**Informed consent statement:** Informed written consent was obtained from the patient for publication of this report and any accompanying images.

**Conflict-of-interest statement:** The authors declare that they have no conflict of interest to report.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Qiao-Qiao Zhou 0000-0001-7314-6746; Jing Li 0000-0003-2153-4055; Bin Liu 0000-0002-0881-0265; Chun-Li Wang 0000-0002-8910-3967.

**S-Editor:** Fan JR

**L-Editor:** Wang TQ

**P-Editor:** Fan JR

## REFERENCES

- 1 **Working group of guidelines for renal anemia of nephrologist branch of Chinese Medical Association.** Clinical practice guidelines for diagnosis and treatment of renal anemia in China. *Zhonghua Yixue Zazhi* 2021; **101**: 1463-502 [DOI: [10.3760/cma.j.cn112137-20210201-00309](https://doi.org/10.3760/cma.j.cn112137-20210201-00309)]
- 2 **Knight K, Wade S, Balducci L.** Prevalence and outcomes of anemia in cancer: a systematic review of the literature. *Am J Med* 2004; **116** Suppl 7A: 11S-26S [PMID: [15050883](https://pubmed.ncbi.nlm.nih.gov/15050883/) DOI: [10.1016/j.amjmed.2003.12.008](https://doi.org/10.1016/j.amjmed.2003.12.008)]



- 3 Li Y, Shi H, Wang WM, Peng A, Jiang GR, Zhang JY, Ni ZH, He LQ, Niu JY, Wang NS, Mei CL, Xu XD, Guo ZY, Yuan WJ, Yan HD, Deng YY, Yu C, Cen J, Zhang Y, Chen N. Prevalence, awareness, and treatment of anemia in Chinese patients with nondialysis chronic kidney disease: First multicenter, cross-sectional study. *Medicine (Baltimore)* 2016; **95**: e3872 [PMID: 27310973 DOI: 10.1097/MD.0000000000003872]
- 4 Wang Y, Wei RB, Su TY, Huang MJ, Li P, Chen XM. Clinical and pathological factors of renal anaemia in patients with IgA nephropathy in Chinese adults: a cross-sectional study. *BMJ Open* 2019; **9**: e023479 [PMID: 30647035 DOI: 10.1136/bmjopen-2018-023479]
- 5 Li Y, Zhang W, Ren H, Wang W, Shi H, Li X, Chen X, Shen P, Wu X, Xie J, Chen N. Evaluation of anemia and serum iPTH, calcium, and phosphorus in patients with primary glomerulonephritis. *Contrib Nephrol* 2013; **181**: 31-40 [PMID: 23689565 DOI: 10.1159/000348636]
- 6 Huang Z, Song T, Fu L, Rao Z, Zeng D, Qiu Y, Wang X, Xie L, Wei Q, Wang L, Lin T. Post-renal transplantation anemia at 12 mo: prevalence, risk factors, and impact on clinical outcomes. *Int Urol Nephrol* 2015; **47**: 1577-1585 [PMID: 26246037 DOI: 10.1007/s11255-015-1069-y]
- 7 Latcha S. Anemia management in cancer patients with chronic kidney disease. *Semin Dial* 2019; **32**: 513-519 [PMID: 31596520 DOI: 10.1111/sdi.12841]
- 8 Henke M, Laszig R, Rübe C, Schäfer U, Haase KD, Schilcher B, Mose S, Beer KT, Burger U, Dougherty C, Frommhold H. Erythropoietin to treat head and neck cancer patients with anaemia undergoing radiotherapy: randomised, double-blind, placebo-controlled trial. *Lancet* 2003; **362**: 1255-1260 [PMID: 14575968 DOI: 10.1016/S0140-6736(03)14567-9]
- 9 Bohlius J, Schmidlin K, Brillant C, Schwarzer G, Trelle S, Seidenfeld J, Zwahlen M, Clarke M, Weingart O, Kluge S, Piper M, Rades D, Steensma DP, Djulbegovic B, Fey MF, Ray-Coquard I, Machtay M, Moebus V, Thomas G, Untch M, Schumacher M, Egger M, Engert A. Recombinant human erythropoiesis-stimulating agents and mortality in patients with cancer: a meta-analysis of randomised trials. *Lancet* 2009; **373**: 1532-1542 [PMID: 19410717 DOI: 10.1016/S0140-6736(09)60502-X]
- 10 Wright JR, Ung YC, Julian JA, Pritchard KI, Whelan TJ, Smith C, Szechtman B, Roa W, Mulroy L, Rudinskas L, Gagnon B, Okawara GS, Levine MN. Randomized, double-blind, placebo-controlled trial of erythropoietin in non-small-cell lung cancer with disease-related anemia. *J Clin Oncol* 2007; **25**: 1027-1032 [PMID: 17312332 DOI: 10.1200/JCO.2006.07.1514]
- 11 Leyland-Jones B, Semiglazov V, Pawlicki M, Pienkowski T, Tjulandini S, Manikhas G, Makhson A, Roth A, Dodwell D, Baselga J, Biakhov M, Valuckas K, Voznyi E, Liu X, Vercammen E. Maintaining normal hemoglobin levels with epoetin alfa in mainly nonanemic patients with metastatic breast cancer receiving first-line chemotherapy: a survival study. *J Clin Oncol* 2005; **23**: 5960-5972 [PMID: 16087945 DOI: 10.1200/JCO.2005.06.150]
- 12 The society of chemotherapy; Chinese Anti-Cancer Association; Committee of neoplastic supportive-care (CONS); China Anti-Cancer Association. A consensus on the clinical diagnosis, treatment, and prevention of cancer and chemotherapy-related anemia in China (2019 edition). *Chin J Clin Oncol* 2019; **11**: 78-85
- 13 Li X, Cole SR, Kshirsagar AV, Fine JP, Stürmer T, Brookhart MA. Safety of Dynamic Intravenous Iron Administration Strategies in Hemodialysis Patients. *Clin J Am Soc Nephrol* 2019; **14**: 728-737 [PMID: 30988164 DOI: 10.2215/CJN.03970318]
- 14 Bailie GR, Larkina M, Goodkin DA, Li Y, Pisoni RL, Bieber B, Mason N, Tong L, Locatelli F, Marshall MR, Inaba M, Robinson BM. Data from the Dialysis Outcomes and Practice Patterns Study validate an association between high intravenous iron doses and mortality. *Kidney Int* 2015; **87**: 162-168 [PMID: 25075769 DOI: 10.1038/ki.2014.275]
- 15 Spivak JL, Gascón P, Ludwig H. Anemia management in oncology and hematology. *Oncologist* 2009; **14** Suppl 1: 43-56 [PMID: 19762516 DOI: 10.1634/theoncologist.2009-S1-43]
- 16 Locatelli F, Del Vecchio L, De Nicola L, Minutolo R. Are all erythropoiesis-stimulating agents created equal? *Nephrol Dial Transplant* 2021; **36**: 1369-1377 [PMID: 32206785 DOI: 10.1093/ndt/gfaa034]
- 17 Maxwell PH, Eckardt KU. HIF prolyl hydroxylase inhibitors for the treatment of renal anaemia and beyond. *Nat Rev Nephrol* 2016; **12**: 157-168 [PMID: 26656456 DOI: 10.1038/nrneph.2015.193]
- 18 Chen N, Qian J, Chen J, Yu X, Mei C, Hao C, Jiang G, Lin H, Zhang X, Zuo L, He Q, Fu P, Li X, Ni D, Hemmerich S, Liu C, Szczec L, Besarab A, Neff TB, Peony Yu KH, Valone FH. Phase 2 studies of oral hypoxia-inducible factor prolyl hydroxylase inhibitor FG-4592 for treatment of anemia in China. *Nephrol Dial Transplant* 2017; **32**: 1373-1386 [PMID: 28371815 DOI: 10.1093/ndt/gfx011]
- 19 Akizawa T, Ueno M, Shiga T, Reusch M. Oral roxadustat three times weekly in ESA-naïve and ESA-converted patients with anemia of chronic kidney disease on hemodialysis: Results from two phase 3 studies. *Ther Apher Dial* 2020; **24**: 628-641 [PMID: 31891449 DOI: 10.1111/1744-9987.13468]
- 20 Semenza GL. Defining the role of hypoxia-inducible factor 1 in cancer biology and therapeutics. *Oncogene* 2010; **29**: 625-634 [PMID: 19946328 DOI: 10.1038/onc.2009.441]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

