

World Journal of *Clinical Cases*

World J Clin Cases 2022 June 16; 10(17): 5518-5933



MINIREVIEWS

- 5518 Occult hepatitis B – the result of the host immune response interaction with different genomic expressions of the virus
Gherlan GS
- 5531 Pulmonary complications of portal hypertension: The overlooked decompensation
Craciun R, Mocan T, Procopet B, Nemes A, Tefas C, Sparchez M, Mocan LP, Sparchez Z
- 5541 Ethical review of off-label drugs during the COVID-19 pandemic
Li QY, Lv Y, An ZY, Dai NN, Hong X, Zhang Y, Liang LJ

ORIGINAL ARTICLE**Case Control Study**

- 5551 Gut peptide changes in patients with obstructive jaundice undergoing biliary drainage: A prospective case control study
Pavić T, Pelajić S, Blažević N, Kralj D, Milošević M, Mikolasevic I, Lerotic I, Hrabar D

Retrospective Cohort Study

- 5566 Longitudinal assessment of liver stiffness by transient elastography for chronic hepatitis C patients
Mezina A, Krishnan A, Woreta TA, Rubenstein KB, Watson E, Chen PH, Rodriguez-Watson C

Retrospective Study

- 5577 Clinical evaluation of prone position ventilation in the treatment of acute respiratory distress syndrome induced by sepsis
Xia WH, Yang CL, Chen Z, Ouyang CH, Ouyang GQ, Li QG
- 5586 Three-dimensional arterial spin labeling and diffusion kurtosis imaging in evaluating perfusion and infarct area size in acute cerebral ischemia
Jiang YY, Zhong ZL, Zuo M
- 5595 Intrathecal methotrexate in combination with systemic chemotherapy in glioblastoma patients with leptomeningeal dissemination: A retrospective analysis
Kang X, Chen F, Yang SB, Wang YL, Qian ZH, Li Y, Lin H, Li P, Peng YC, Wang XM, Li WB
- 5606 Hepatic epithelioid hemangioendothelioma: Clinical characteristics, diagnosis, treatment, and prognosis
Zhao M, Yin F
- 5620 Difference between type 2 gastroesophageal varices and isolated fundic varices in clinical profiles and portosystemic collaterals
Song YH, Xiang HY, Si KK, Wang ZH, Zhang Y, Liu C, Xu KS, Li X

- 5634** Assessment of incidental focal colorectal uptake by analysis of fluorine-18 fluorodeoxyglucose positron emission tomography parameters

Lee H, Hwang KH, Kwon KA

Observational Study

- 5646** "Zero ischemia" laparoscopic partial nephrectomy by high-power GreenLight laser enucleation for renal carcinoma: A single-center experience

Zhang XM, Xu JD, Lv JM, Pan XW, Cao JW, Chu J, Cui XG

- 5655** High Eckardt score and previous treatment were associated with poor postperoral endoscopic myotomy pain control: A retrospective study

Chen WN, Xu YL, Zhang XG

- 5667** Higher volume growth rate is associated with development of worrisome features in patients with branch duct-intraductal papillary mucinous neoplasms

Innocenti T, Danti G, Lynch EN, Dragoni G, Gottin M, Fedeli F, Palatresi D, Biagini MR, Milani S, Miele V, Galli A

Prospective Study

- 5680** Application of a new anatomic hook-rod-pedicle screw system in young patients with lumbar spondylolysis: A pilot study

Li DM, Li YC, Jiang W, Peng BG

META-ANALYSIS

- 5690** Systematic review of Yougui pills combined with levothyroxine sodium in the treatment of hypothyroidism

Liu XP, Zhou YN, Tan CE

CASE REPORT

- 5702** Allogeneic stem cell transplantation-A curative treatment for paroxysmal nocturnal hemoglobinuria with PIGT mutation: A case report

Schenone L, Notarantonio AB, Latger-Cannard V, Fremeaux-Bacchi V, De Carvalho-Bittencourt M, Rubio MT, Muller M, D'Aveni M

- 5708** Gray zone lymphoma effectively treated with cyclophosphamide, doxorubicin, vincristine, prednisolone, and rituximab chemotherapy: A case report

Hojo N, Nagasaki M, Mihara Y

- 5717** Diagnosis of spontaneous isolated superior mesenteric artery dissection with ultrasound: A case report

Zhang Y, Zhou JY, Liu J, Bai C

- 5723** Adrenocorticotrophic hormone-secreting pancreatic neuroendocrine carcinoma with multiple organ infections and widespread thrombosis: A case report

Yoshihara A, Nishihama K, Inoue C, Okano Y, Eguchi K, Tanaka S, Maki K, Fridman D'Alessandro V, Takeshita A, Yasuma T, Uemura M, Suzuki T, Gabazza EC, Yano Y

- 5732** Management of the palato-radicular groove with a periodontal regenerative procedure and prosthodontic treatment: A case report

Ling DH, Shi WP, Wang YH, Lai DP, Zhang YZ

- 5741** Combined thoracic paravertebral block and interscalene brachial plexus block for modified radical mastectomy: A case report
Hu ZT, Sun G, Wang ST, Li K
- 5748** Chondromyxoid fibroma of the cervical spine: A case report
Li C, Li S, Hu W
- 5756** Preterm neonate with a large congenital hemangioma on maxillofacial site causing thrombocytopenia and heart failure: A case report
Ren N, Jin CS, Zhao XQ, Gao WH, Gao YX, Wang Y, Zhang YF
- 5764** Simultaneous multiple primary malignancies diagnosed by endoscopic ultrasound-guided fine-needle aspiration: A case report
Yang J, Zeng Y, Zhang JW
- 5770** Neuroendocrine tumour of the descending part of the duodenum complicated with schwannoma: A case report
Zhang L, Zhang C, Feng SY, Ma PP, Zhang S, Wang QQ
- 5776** Massive hemothorax following internal jugular vein catheterization under ultrasound guidance: A case report
Kang H, Cho SY, Suk EH, Ju W, Choi JY
- 5783** Unilateral adrenal tuberculosis whose computed tomography imaging characteristics mimic a malignant tumor: A case report
Liu H, Tang TJ, An ZM, Yu YR
- 5789** Modified membrane fixation technique in a severe continuous horizontal bone defect: A case report
Wang LH, Ruan Y, Zhao WY, Chen JP, Yang F
- 5798** Surgical repair of an emergent giant hepatic aneurysm with an abdominal aortic dissection: A case report
Wen X, Yao ZY, Zhang Q, Wei W, Chen XY, Huang B
- 5805** Heterotopic ossification beneath the upper abdominal incision after radical gastrectomy: Two case reports
Zhang X, Xia PT, Ma YC, Dai Y, Wang YL
- 5810** Non-alcoholic Wernicke encephalopathy in an esophageal cancer patient receiving radiotherapy: A case report
Zhang Y, Wang L, Jiang J, Chen WY
- 5816** New approach for the treatment of vertical root fracture of teeth: A case report and review of literature
Zhong X, Yan P, Fan W
- 5825** Ultrasound-guided microwave ablation as a palliative treatment for mycosis fungoides eyelid involvement: A case report
Chen YW, Yang HZ, Zhao SS, Zhang Z, Chen ZM, Feng HH, An MH, Wang KK, Duan R, Chen BD
- 5833** Pulp revascularization on an adult mandibular right second premolar: A case report
Yang YQ, Wu BL, Zeng JK, Jiang C, Chen M

- 5841** Barrett's esophagus in a patient with bulimia nervosa: A case report
Gouda A, El-Kassas M
- 5846** Spontaneous gallbladder perforation and colon fistula in hypertriglyceridemia-related severe acute pancreatitis: A case report
Wang QP, Chen YJ, Sun MX, Dai JY, Cao J, Xu Q, Zhang GN, Zhang SY
- 5854** Beware of gastric tube in esophagectomy after gastric radiotherapy: A case report
Yurttas C, Wichmann D, Gani C, Bongers MN, Singer S, Thiel C, Koehngrainer A, Thiel K
- 5861** Transition from minimal change disease to focal segmental glomerulosclerosis related to occupational exposure: A case report
Tang L, Cai Z, Wang SX, Zhao WJ
- 5869** Lung adenocarcinoma metastasis to paranasal sinus: A case report
Li WJ, Xue HX, You JQ, Chao CJ
- 5877** Follicular lymphoma presenting like marginal zone lymphoma: A case report
Peng HY, Xiu YJ, Chen WH, Gu QL, Du X
- 5884** Primary renal small cell carcinoma: A case report
Xie K, Li XY, Liao BJ, Wu SC, Chen WM
- 5893** Gitelman syndrome: A case report
Chen SY, Jie N
- 5899** High-frame-rate contrast-enhanced ultrasound findings of liver metastasis of duodenal gastrointestinal stromal tumor: A case report and literature review
Chen JH, Huang Y
- 5910** Tumor-like disorder of the brachial plexus region in a patient with hemophilia: A case report
Guo EQ, Yang XD, Lu HR
- 5916** Response to dacomitinib in advanced non-small-cell lung cancer harboring the rare delE709_T710insD mutation: A case report
Xu F, Xia ML, Pan HY, Pan JW, Shen YH
- 5923** Loss of human epidermal receptor-2 in human epidermal receptor-2+ breast cancer after neoadjuvant treatment: A case report
Yu J, Li NL

LETTER TO THE EDITOR

- 5929** Repetitive transcranial magnetic stimulation for post-traumatic stress disorder: Lights and shadows
Concerto C, Lanza G, Fisticaro F, Pennisi M, Rodolico A, Torrisi G, Bella R, Aguglia E

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Raden Andri Primadhi, MD, PhD, Assistant Professor, Surgeon, Department of Orthopaedics and Traumatology, Universitas Padjadjaran Medical School, Hasan Sadikin Hospital, Bandung 40161, Indonesia. randri@unpad.ac.id

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for *WJCC* as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The *WJCC*'s CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Hua-Ge Yan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jim-Lai Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

June 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Beware of gastric tube in esophagectomy after gastric radiotherapy: A case report

Can Yurttas, Doerte Wichmann, Cihan Gani, Malte N Bongers, Stephan Singer, Christian Thiel, Alfred Koenigsrainer, Karolin Thiel

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C
Grade D (Fair): D
Grade E (Poor): 0

P-Reviewer: Di Meglio L, Italy;
Ding X, China

Received: December 27, 2021

Peer-review started: December 27, 2021

First decision: March 10, 2022

Revised: March 11, 2022

Accepted: April 9, 2022

Article in press: April 9, 2022

Published online: June 16, 2022



Can Yurttas, Doerte Wichmann, Christian Thiel, Alfred Koenigsrainer, Karolin Thiel, Department of General, Visceral and Transplant Surgery, University Hospital of Tuebingen, Tuebingen 72076, Germany

Cihan Gani, Department of Radiooncology, University Hospital of Tuebingen, Tuebingen 72076, Germany

Malte N Bongers, Department of Diagnostic and Interventional Radiology, University Hospital of Tuebingen, Tuebingen 72076, Germany

Stephan Singer, Department of Pathology, University Hospital of Tuebingen, Tuebingen 72076, Germany

Corresponding author: Karolin Thiel, MD, Department of General, Visceral and Transplant Surgery, University Hospital of Tuebingen, Hoppe Seyler Str 3, Tuebingen 72076, Germany. karolin.thiel@med.uni-tuebingen.de

Abstract

BACKGROUND

Gastric tube formation and pull-up is the most common technique of reconstruction following esophagectomy for esophageal cancer. If previous treatment with radiotherapy for gastric mucosa-associated lymphoid tissue (MALT)-lymphoma restricts suitability of the stomach for anastomosis to the esophagus is unknown.

CASE SUMMARY

A 57-year-old man underwent sequential chemotherapy and radiotherapy for gastric MALT-lymphoma seven years prior to diagnosis of esophageal adenocarcinoma. Esophagectomy without neoadjuvant treatment was recommended by the multidisciplinary tumor board due to early tumor stage [uT1 (sm2) uN+ cM0 according to TNM-classification of malignant tumors, 8th edition] without lymph node involvement. Minimal invasive esophageal resection with esophagogastrotomy was performed. Due to gastric tube necrosis with anastomotic leakage on the twelfth postoperative day, diverting resection with construction of a cervical salivary fistula was necessary. Rapid recovery facilitated colonic interposition without any complications six months afterwards.

CONCLUSION

This case report may represent the start for further investigation to know if it is reasonable to refrain from esophagogastrectomy in patients with a long interval between gastric radiotherapy and surgery.

Key Words: Esophageal cancer; Mucosa-associated lymphoid tissue lymphoma; Esophagogastrectomy; Cervical fistula; Colonic interposition; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: A patient with previous radiotherapy for gastric mucosa-associated lymphoid tissue lymphoma underwent esophagectomy and esophagogastrectomy for esophageal cancer more than seven years later. Gastric tube necrosis, made diversion surgery with salivary fistula necessary. Six months later, interposition of the transverse colon was performed without occurrence of any complications. The patient fully recovered with unlimited oral intake capability and remains free of tumor recurrence at date of publication. In patients with a long interval between gastric radiotherapy and surgery esophagogastrectomy should be avoided.

Citation: Yurttas C, Wichmann D, Gani C, Bongers MN, Singer S, Thiel C, Koenigsrainer A, Thiel K. Beware of gastric tube in esophagectomy after gastric radiotherapy: A case report. *World J Clin Cases* 2022; 10(17): 5854-5860

URL: <https://www.wjgnet.com/2307-8960/full/v10/i17/5854.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i17.5854>

INTRODUCTION

Esophagectomy, combined with neoadjuvant chemo(radio)therapy in the locally advanced situation, is considered standard treatment with curative intention for carcinomas of the esophagus and the esophagogastric junction[1]. Most commonly, anastomosis of the remnant esophagus to a gastric tube is performed[2]. Whether prior chemoradiotherapy for gastric mucosa-associated lymphoid tissue lymphoma limits the stomach's suitability for reconstruction is unknown. With this case report we provide first evidence for pretreated stomach usage for esophagogastrectomy in esophagectomy.

CASE PRESENTATION

Chief complaints

Due to asymptomatic gastro-esophageal reflux disease with Long Segment Barrett's esophagus C9M13 according to Prague Classification, a 64-year-old patient underwent repetitive esophagogastroduodenoscopy.

History of present illness

In 2020, biopsy of the distal esophagus 34 cm from row of teeth revealed invasive moderately differentiated (G2) adenocarcinoma. Moreover, erythema and atrophy of the gastric mucosa were detected. However, the patient had no disease-specific complaints when he first presented to our department. Oral intake of standard western-diet was unrestricted and body weight was constant at a BMI of 29.1 kg/m².

History of past illness

In 2012, the 57-year-old man was diagnosed with diffuse large B-cell lymphoma (DLBCL) of the stomach in the course of endoscopic treatment of gastric bleeding (2a according to the Forrest Classification of gastrointestinal bleedings). Although there was no detection of *Helicobacter pylori*, eradication therapy was performed. Endosonography proved localization at the posterior gastric wall without infiltration of neighboring tissues, whereas computed tomography (CT) scan and bone marrow biopsy were without evidence of disease equivalent to stage IE according to the Ann Arbor staging system. Following four courses of rituximab, cyclophosphamide, doxorubicin hydrochloride (hydroxydaunorubicin), vincristine sulfate (Oncovin), and prednisone (R-CHOP) with curative intention percutaneous normofractionated radiotherapy of the stomach with a total of 39.6 Gray (Gy) in 20 fractions weekly was performed as consolidating therapy. Both systemic and radiation therapy were well tolerated. Due to

herpes zoster of the left thorax antiviral therapy with aciclovir was introduced.

The patient had a history of herniated vertebral disc, struma nodosa, chronic-venous insufficiency and endoscopic resection of a low-grade adenoma of the sigmoid colon and regularly took metformin, thyroxine and sitagliptin for type 2 diabetes mellitus and hypothyroidism respectively. Hepatic and renal function were not impaired. Follow-up examinations up to five years were without any peculiarities or evidence of tumor recurrence. The patient had skipped drinking and smoking after intake of 60 pack-years.

Personal and family history

Family history was unremarkable and not related to the present case.

Physical examination

The patient was in a normal general state without any evidence of disease or restriction of normal activities.

Laboratory examinations

Preoperative blood examinations were unremarkable. Tumor markers CEA, CA19-9 and CA72-4 were within reference range.

Imaging examinations

Whereas CT scan showed no signs of distant metastases or involvement of locoregional lymph nodes, endosonography described uT1 (sm2) uN+ according to TNM classification of malignant tumors, 8th edition. Positron emission tomography-CT was performed for further clarification, which ruled out involvement of locoregional lymph nodes.

Material and methods

Surgery for esophageal cancer and gastric tube necrosis: Surgery was performed in minimally invasive technique of Ivor Lewis esophagectomy. Access to the abdominal cavity and capnoperitoneum was established with the help of a Veress needle. An optic trocar was introduced under vision with a 30° camera (KARL STORZ SE & Co. KG, Tuttlingen, Germany). The abdominal cavity was inspected to rule out injuries during access and also peritoneal or hepatic metastases. Then, gastric mobilization was performed with an electrosurgical vessel sealer, left gastric artery was clipped whereas the right gastric artery as well as the right gastroepiploic arcade were preserved. Complete D2-lymphadenectomy was performed followed by stapled gastric tube formation of approximately 5 cm in diameter. Esophagectomy including mediastinal lymphadenectomy was operated thoracoscopically with four right-sided intercostal trocars. The resection was completed with formation of a stapled circular end-to-side-esophagogastrostomy.

Emergency thoracotomy was necessary for resection of the necrotic gastric tube, hemithyroidectomy and creation of the salivary glandula. A jejunal feeding tube was inserted after laparotomy. Continuous intestinal passage was reconstructed by colonic interposition. Following laparotomy, the transverse colon was prepared for retrosternal pull-up and formation of an end-to-end esophagocolostomy and an end-to-side colojejunosomy. A side-to-side ascendodendostomy was created.

Endoscopy and endoscopic negative-pressure therapy: Endoscopy was performed with a standard gastroscope with 9.8-mm outer caliber and 3.2-mm working channel (PENTAX Medical, Tokyo, Japan). A thin open-pore film wrapped around a drain (Medicoplast, Illingen, Germany) and fixed with a suture was constructed prior to endoscopically controlled insertion and positioning of the device. Negative pressure of -125 mmHg was established with the use of a vacuum therapy system (KCI medical, Wiesbaden, Germany).

FINAL DIAGNOSIS

Moderately differentiated adenocarcinoma of the distal esophagus with infiltration of the submucosal layer without locoregional lymph node metastases [TNM: pT1b, pN0 (0/17) L0, V0, Pn0, R0, Grading: G2].

TREATMENT

The multidisciplinary tumor board consequently recommended surgical resection without neoadjuvant treatment. Thoracoscopic and laparoscopic abdominal right thoracic esophagectomy with two-field lymphadenectomy (Ivor Lewis) and stapled end-to-side esophagogastrostomy was performed. Histopathological examination confirmed the diagnosis and staging results and complete resection of a

moderately differentiated adenocarcinoma of the distal esophagus. The gastric mucosa showed signs of erosive gastritis with denuded surface epithelium, subepithelial and interstitial hemorrhage, but no recurrent lymphoma infiltrates. The initial postoperative course was regular and without any pathologic findings. Following extubation immediately after surgery, the patient was monitored at the intermediate care unit for one day without requiring cardiocirculatory or respiratory support before transfer to the general ward. Low-dose anticoagulation with unfractionated heparin was initiated six hours after surgery. Amount and quality of drain output were unsuspecting. Seven days after surgery the patient's general state was seen to deteriorate and elevated leukocytes and C-reactive protein were observed, which required endoscopic assessment of the esophagogastronomy to rule out anastomotic leakage. The gastric interposition showed compromised perfusion without evidence of anastomotic insufficiency. Endoscopic negative-pressure therapy was therefore introduced. After vomiting with aspiration during anaesthetization the patient was transferred to the intensive care unit. Despite initiation of calculated antibiotic therapy with meropenem, vancomycin and anidulafungin there was no observable improvement. On day 12 postoperative, endoscopy revealed necrosis of the gastric interposition with a pronounced anastomotic insufficiency prompting surgical resection of the gastric tube interposition, creation of a cervical fistula and insertion of a jejunal feeding catheter (Figure 1). Histopathology confirmed ischemic necrosis of the proximal gastric tube with anastomotic leakage. There was no evidence of residual adenocarcinoma or recurrent lymphoma in the resected esophagogastronomy or gastric tube. Postoperative pleural effusion was treated with a thoracic drain and central venous line-associated blood-stream infection, while paroxysmal tachycardia and delirium necessitated respective therapy. The patient slowly recovered until he was discharged 40 d after esophageal resection. Follow-up care was recommended by the multidisciplinary tumor board.

OUTCOME AND FOLLOW-UP

Six months later, the patient underwent colonoscopy and CT scan in preparation for colonic interposition without any contraindications or signs of tumor recurrence. Retrosternal interposition of the transverse colon creating an end-to-end esophagotransversostomy, end-to-side transversojejunos-tomy and a side-to-side ascendotransversostomy was performed. Postoperative course was normal. Oral intake of food and liquids was without difficulty. Supportive enteral feeding was continued. The patient was discharged home on day 12 postoperative. Nine weeks later, the patient was in an unrestricted general condition with stable body weight so that the jejunal feeding catheter was removed. Table 1 shows information from this case report organized in a time table.

DISCUSSION

When the patient first presented to our out-patient clinic, the suitability of the pretreated stomach for construction of an esophagogastronomy was uncertain because evidence was missing. In the literature, complications of esophagogastronomy in general are reported to occur in 12% and mortality in 4% of all cases[3]. According to the present literature, small bowel or colonic interposition may be considered alternative grafts. Compared to the colon, small bowel grafts require fewer anastomoses, are rarely affected by malignancies and have good peristalsis, but provide no reservoir function. Colonic interposition is complicated by the need for three to four anastomoses and potential metachronous development of adenoma and carcinoma. Nevertheless, longer grafts are available offering reservoir-like function and less reflux[4,5]. However, a retrospective cohort study comparing complex esophageal reconstruction including 44.7% of patients with other than gastric tube formation to non-complex esophagectomy with direct gastric pull-up reported higher morbidity and longer length of stay for patients in the complex therapy group[6]. Jejunal grafts are described as suitable primary alternatives for any scope of esophageal replacement, but are accompanied by up to 36% anastomotic leakage and 10% mortality[7]. In colonic interposition, higher overall morbidity of 45.0%-64.0% and increased risk of anastomotic leakage occurring in 13.0%-30.0% of patients is shown[8-11]. Alternatively, construction of a cervical salivary fistula with secondary gastric tube formation could be an option, but especially patients with cancer were shown to have poor outcome after primary diversion and secondary reconstruction in esophagectomy[12]. Considering our experience with gastric tubes and the lower complication rates as compared to small bowel and colonic interposition, the decision for esophagogastronomy was therefore made together with the patient.


Despite expectable poor outcome following resection of the necrotic gastric tube with diversion[12], creation of a cervical fistula and secondary colonic interposition, our patient fully recovered, has sufficient oral intake capacity and to date remains without signs of any tumor recurrence.

Neoadjuvant radiochemotherapy prior to esophagectomy has been shown to improve overall survival compared to surgery alone with a very favourable toxicity profile. In particular, no increase in anastomotic leakage was reported in the CROSS trial[13], whereas in-field creation of anastomosis following neoadjuvant radiochemotherapy and esophagectomy was identified as a risk factor for

Table 1 Information from this case report organized in a time table

Date	Diagnosis	Intervention
September, 2012	Gastric MALT lymphoma	R-CHOP
February, 2013	Gastric MALT lymphoma	Intensity-modulated radiation therapy up to 39.6 Gy
July, 2020	Barrett's carcinoma	Endoscopic biopsy
November, 2020	Barrett's carcinoma	Abdominal right-thoracic esophagectomy with two-field lymphadenectomy (Ivor Lewis)
December, 2020	Gastric tube necrosis	Diverting resection with creation of cervical salivary fistula
June, 2021	Presence of a cervical salivary fistula	Colonic interposition and insertion of a jejunal feeding catheter
August, 2021	Needless jejunal feeding catheter	Extraction of jejunal feeding catheter

MALT: Mucosa-associated lymphoid tissue; R-CHOP: Rituximab, cyclophosphamide, doxorubicin hydrochloride (hydroxydaunorubicin), vincristine sulfate (Oncovin), and prednisone.



DOI: 10.12998/wjcc.v10.i17.5854 Copyright ©The Author(s) 2022.

Figure 1 Ischemic necrosis of the gastric interposition with anastomotic leakage. A and B: Computed tomography (A) and endoscopy (B). Computed tomography also shows left-sided pleural effusion and inserted nasogastric tube. Endoscopy also revealed the anastomotic dehiscence with cavity and the exposed staples.

anastomotic leakage in a retrospective analysis of 285 patients treated for esophageal cancer[14]. Especially in distal esophageal cancer the celiac lymph nodes and the ones at the lesser gastric curvature are frequently irradiated in the preoperative setting with doses that are comparable to the dose given in the current case presentation resulting in a considerable dose burden to the stomach without causing an excessive rate of anastomotic leakage. A major difference however between preoperative radiotherapy for esophageal cancer and the previous treatment with radiotherapy in the current case is the interval between radiotherapy and surgery. While surgery after planned neoadjuvant therapy is commonly scheduled within a couple of weeks, the interval was seven years in the present case. One can hypothesize that the tissue turned less “flexible” over the time due to fibrosis which might have contributed to anastomotic leakage. However, in the present case radiotherapy was applied to the specimen employed for reconstruction and not to the resected organ.

CONCLUSION

We therefore recommend that stomachs pretreated by radiotherapy should not be utilized for reconstruction in esophagectomy. Although this case report provides little evidence from a single patient only without proven causality, further investigations as to whether stomachs pretreated by radiotherapy in general should not be utilized for reconstruction in esophagectomy are required.

ACKNOWLEDGEMENTS

The authors thank Margreiter MH for her kind contribution to preparation of the manuscript.

FOOTNOTES

Author contributions: Yurttas C, Thiel K and Koenigsrainer A conceived and wrote the manuscript; Thiel K and Koenigsrainer A were responsible for overall treatment; Endoscopic treatment was performed by Wichmann D; Yurttas C and Bongers MN prepared the figures; Yurttas C, Thiel K, Gani C, Thiel C, Singer S and Koenigsrainer A revised the manuscript; all authors read and approved the final version of the manuscript.

Informed consent statement: For all clinical data and personal images written consent for publication was obtained from the patient.

Conflict-of-interest statement: The authors have no competing interests to declare in the context of the current case.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Germany

ORCID number: Can Yurttas 0000-0001-9720-8243; Doerte Wichmann 0000-0002-8832-5031; Cihan Gani 0000-0003-0242-9428; Malte N Bongers 0000-0003-2852-0270; Stephan Singer 0000-0003-4775-6609; Christian Thiel 0000-0001-8880-4860; Alfred Koenigsrainer 0000-0002-2301-4080; Karolin Thiel 0000-0002-7328-7589.

S-Editor: Gao CC

L-Editor: A

P-Editor: Gao CC

REFERENCES

- 1 **Low DE.** Evolution in surgical management of esophageal cancer. *Dig Dis* 2013; **31**: 21-29 [PMID: 23797119 DOI: 10.1159/000343650]
- 2 **Low DE,** Kuppusamy MK, Alderson D, Ceconello I, Chang AC, Darling G, Davies A, D'Journo XB, Gisbertz SS, Griffin SM, Hardwick R, Hoelscher A, Hofstetter W, Jobe B, Kitagawa Y, Law S, Mariette C, Maynard N, Morse CR, Nafteux P, Pera M, Pramesh CS, Puig S, Reynolds JV, Schroeder W, Smithers M, Wijnhoven BPL. Benchmarking Complications Associated with Esophagectomy. *Ann Surg* 2019; **269**: 291-298 [PMID: 29206677 DOI: 10.1097/SLA.0000000000002611]
- 3 **Butskiy O,** Rahmanian R, White RA, Durham S, Anderson DW, Prisman E. Revisiting the gastric pull-up for pharyngoesophageal reconstruction: A systematic review and meta-analysis of mortality and morbidity. *J Surg Oncol* 2016; **114**: 907-914 [PMID: 27774626 DOI: 10.1002/jso.24477]
- 4 **Watanabe M,** Mine S, Nishida K, Kuroguchi T, Okamura A, Imamura Y. Reconstruction after esophagectomy for esophageal cancer patients with a history of gastrectomy. *Gen Thorac Cardiovasc Surg* 2016; **64**: 457-463 [PMID: 27234222 DOI: 10.1007/s11748-016-0661-0]
- 5 **Sohn M,** Agha A, Trum S, Moser C, Gundling F, Hochrein A, Pratschke J, Aigner F, Ritschl P. Frequency of Polyps and Adenocarcinoma in Colon Interposition After Esophagectomy in Adulthood - A Systematic Review. *Anticancer Res* 2019; **39**: 6419-6430 [PMID: 31810906 DOI: 10.21873/anticancer.13856]
- 6 **Moore JM,** Hooker CM, Molena D, Mungo B, Brock MV, Battafarano RJ, Yang SC. Complex Esophageal Reconstruction Procedures Have Acceptable Outcomes Compared With Routine Esophagectomy. *Ann Thorac Surg* 2016; **102**: 215-222 [PMID: 27217296 DOI: 10.1016/j.athoracsur.2016.02.039]
- 7 **Gaur P,** Blackmon SH. Jejunal graft conduits after esophagectomy. *J Thorac Dis* 2014; **6** Suppl 3: S333-S340 [PMID: 24876939 DOI: 10.3978/j.issn.2072-1439.2014.05.07]
- 8 **Hamai Y,** Hihara J, Emi M, Aoki Y, Okada M. Esophageal reconstruction using the terminal ileum and right colon in esophageal cancer surgery. *Surg Today* 2012; **42**: 342-350 [PMID: 22200754 DOI: 10.1007/s00595-011-0103-7]
- 9 **Mine S,** Udagawa H, Tsutsumi K, Kinoshita Y, Ueno M, Ehara K, Haruta S. Colon interposition after esophagectomy with extended lymphadenectomy for esophageal cancer. *Ann Thorac Surg* 2009; **88**: 1647-1653 [PMID: 19853126 DOI: 10.1016/j.athoracsur.2009.05.081]
- 10 **Klink CD,** Binnebösel M, Schneider M, Ophoff K, Schumpelick V, Jansen M. Operative outcome of colon interposition in the treatment of esophageal cancer: a 20-year experience. *Surgery* 2010; **147**: 491-496 [PMID: 20004440 DOI: 10.1016/j.surg.2009.10.045]
- 11 **Davis PA,** Law S, Wong J. Colonic interposition after esophagectomy for cancer. *Arch Surg* 2003; **138**: 303-308 [PMID: 12611579 DOI: 10.1001/archsurg.138.3.303]
- 12 **DiPierro FV,** Rice TW, DeCamp MM, Rybicki LA, Blackstone EH. Esophagectomy and staged reconstruction. *Eur J Cardiothorac Surg* 2000; **17**: 702-709 [PMID: 10856863 DOI: 10.1016/s1010-7940(00)00408-5]
- 13 **van Hagen P,** Hulshof MC, van Lanschot JJ, Steyerberg EW, van Berge Henegouwen MI, Wijnhoven BP, Richel DJ,

- Nieuwenhuijzen GA, Hospers GA, Bonenkamp JJ, Cuesta MA, Blaisse RJ, Busch OR, ten Kate FJ, Creemers GJ, Punt CJ, Plukker JT, Verheul HM, Spillenaar Bilgen EJ, van Dekken H, van der Sagen MJ, Rozema T, Biermann K, Beukema JC, Piet AH, van Rij CM, Reinders JG, Tilanus HW, van der Gaast A; CROSS Group. Preoperative chemoradiotherapy for esophageal or junctional cancer. *N Engl J Med* 2012; **366**: 2074-2084 [PMID: [22646630](#) DOI: [10.1056/NEJMoa1112088](#)]
- 14 **Juloori A**, Tucker SL, Komaki R, Liao Z, Correa AM, Swisher SG, Hofstetter WL, Lin SH. Influence of preoperative radiation field on postoperative leak rates in esophageal cancer patients after trimodality therapy. *J Thorac Oncol* 2014; **9**: 534-540 [PMID: [24736077](#) DOI: [10.1097/JTO.000000000000100](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

