

World Journal of *Clinical Cases*

World J Clin Cases 2023 February 26; 11(6): 1224-1433



Contents

Thrice Monthly Volume 11 Number 6 February 26, 2023

OPINION REVIEW

- 1224 Collagen matrix scaffolds: Future perspectives for the management of chronic liver diseases
Martinez-Castillo M, Altamirano-Mendoza I, Zielinski R, Priebe W, Piña-Barba C, Gutierrez-Reyes G

MINIREVIEWS

- 1236 Sex dimorphism and metabolic profiles in management of metabolic-associated fatty liver disease
Martin-Grau M, Monleon D
- 1245 Epidemiology and etiology of chemical ocular injury: A brief review
Akgun Z, Selver OB
- 1252 Review of the prevalence, diagnostics, and containment measures of the current mpox outbreak
Sanyaolu A, Marinkovic A, Okorie C, Prakash S, Haider N, Dixon Y, Izurieta R, Badaru O, Smith S
- 1261 Clinical and pathophysiological understanding of the hepatorenal syndrome: Still wrong or still not exactly right?
Wilde B, Canbay A, Katsounas A
- 1267 Flare of the silent pandemic in the era of the COVID-19 pandemic: Obstacles and opportunities
Rayan RA
- 1275 Implications of metabolic dysfunction associated fatty liver disease in COVID-19
Chakraborty R, Sharma D, Kapoor DU, Dwivedi A, Khabiya R, Sen S

ORIGINAL ARTICLE

Retrospective Study

- 1287 Hyperglycemia in COVID-19 infection without diabetes mellitus: Association with inflammatory markers
Geetha HS, Singh G, Sekar A, Gogtay M, Singh Y, Abraham GM, Trivedi N

Clinical Trials Study

- 1299 Efficacy of invisible advancement correction for mandibular retraction in adolescents based on Pancherz analysis
Kong L, Liu XQ

Observational Study

- 1310 Survey study of the etiology of non-traumatic altered consciousness in the Emergency Department at Suez Canal University Hospital in Egypt
Moussa BS, Abd Elatiff ZM, Kamal Eldin Elhadary GM

- 1318** Metformin effect on internal carotid artery blood flow assessed by area under the curve of carotid artery Doppler in women with polycystic ovarian syndrome

Akram W, Nori W, Abdul Ghani Zghair M

- 1330** Effect of continuous nursing combined with respiratory exercise nursing on pulmonary function of postoperative patients with lung cancer

Qiu QX, Li WJ, Ma XM, Feng XH

CASE REPORT

- 1341** Functioning gonadotroph adenoma with hyperestrogenemia and ovarian hyperstimulation in a reproductive-aged woman: A case report and review of literature

He Y, Gao YT, Sun L

- 1349** Clinical manifestations of adult hereditary spherocytosis with novel *SPTB* gene mutations and hyperjaundice: A case report

Jiang N, Mao WY, Peng BX, Yang TY, Mao XR

- 1356** Post-traumatic cauda equina nerve calcification: A case report

Liu YD, Deng Q, Li JJ, Yang HY, Han XF, Zhang KD, Peng RD, Xiang QQ

- 1365** Endometriosis-associated endometrioid adenocarcinoma of the fallopian tube synchronized with endometrial adenocarcinoma: A case report

Feng JY, Jiang QP, He H

- 1372** Gemcitabine-induced peripheral vascular disease and prolonged response in a patient with metastatic pancreatic adenocarcinoma: A case report

Fabien MB, Elodie P, Anna S, Addeo P, Meher B

- 1379** Epidemic Japanese B encephalitis combined with contactin-associated protein-like 2 antibody-positive autoimmune encephalitis: A case report

Huang P

- 1385** Acute pancreatitis as initial presentation of acute myeloid leukemia-M2 subtype: A case report

Yang WX, An K, Liu GF, Zhou HY, Gao JC

- 1393** Postoperative jaundice related to *UGT1A1* and *ABCB11* gene mutations: A case report and literature review

Jiang JL, Liu X, Pan ZQ, Jiang XL, Shi JH, Chen Y, Yi Y, Zhong WW, Liu KY, He YH

- 1403** Hidrotic ectodermal dysplasia in a Chinese pedigree: A case report

Liao MY, Peng H, Li LN, Yang T, Xiong SY, Ye XY

- 1410** Hepatitis A virus-associated acute acalculous cholecystitis in an adult-onset Still's disease patient: A case report and review of the literature

Chang CH, Wang YY, Jiao Y

- 1419** Transverse myelitis caused by herpes zoster following COVID-19 vaccination: A case report

Cho SY, Jang BH, Seo JW, Kim SW, Lim KJ, Lee HY, Kim DJ

- 1426** Primary malignant melanoma of the esophagus: A case report

Wang QQ, Li YM, Qin G, Liu F, Xu YY

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Goran Augustin, MD, MSc, PhD, Assistant Professor, Senior Scientist, Surgeon, Department of Surgery, University Hospital Centre Zagreb, Zagreb 10000, Croatia. augustin.goran@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, George Kontogeorgos, Jerzy Tadeusz Chudek, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

February 26, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Hyperglycemia in COVID-19 infection without diabetes mellitus: Association with inflammatory markers

Harinivaas Shanmugavel Geetha, Garima Singh, Abinash Sekar, Maya Gogtay, Yuvaraj Singh, George M Abraham, Nitin Trivedi

Specialty type: Medicine, research and experimental

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Chan ASW, China; Navarro-Alvarez N, Mexico

Received: October 30, 2022

Peer-review started: October 30, 2022

First decision: November 27, 2022

Revised: December 17, 2022

Accepted: February 3, 2023

Article in press: February 3, 2023

Published online: February 26, 2023



Harinivaas Shanmugavel Geetha, Garima Singh, Abinash Sekar, Yuvaraj Singh, George M Abraham, Nitin Trivedi, Department of Internal Medicine, Saint Vincent Hospital, Worcester, MA 01608, United States

Maya Gogtay, Department of Hospice and Palliative Medicine, University of Texas Health, San Antonio, TX 78229, United States

George M Abraham, Department of Internal Medicine, University of Massachusetts Chan Medical School, Worcester, MA 01655, United States

Corresponding author: Yuvaraj Singh, MD, Chief Medical Resident, Researcher, Department of Internal Medicine, Saint Vincent Hospital, No. 123 Summer Street, Worcester, MA 01608, United States. yuvarajmle@gmail.com

Abstract

BACKGROUND

New onset hyperglycemia is common in patients with severe coronavirus disease 2019 (COVID-19) infection. Cytokine storm due to COVID-19 infection is an essential etiology for new-onset hyperglycemia, but factors like direct severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)-induced pancreatic β -cell failure have also been postulated to play a role.

AIM

We plan to investigate further the mechanisms underlying SARS-CoV-2 infection-induced hyperglycemia, particularly the rationale of the cytokine-induced hyperglycemia hypothesis, by evaluating the association between inflammatory markers and new onset hyperglycemia in non-diabetic patients with COVID-19 infection.

METHODS

We conducted a retrospective case-control study on adults without diabetes mellitus hospitalized for COVID-19 infection. The serum levels of glucose and inflammatory markers at presentation before initiation of corticosteroid were collected. Hyperglycemia was defined as glucose levels ≥ 140 mg/dL. C-Reactive protein (CRP) ≥ 100 mg/L, ferritin ≥ 530 ng/mL, lactate dehydrogenase (LDH) ≥ 590 U/L, and D-dimer ≥ 0.5 mg/L were considered elevated. We used the χ^2 test for categorical variables and the Mann-Whitney U test for continuous variables

and calculated the logistic regression for hyperglycemia.

RESULTS

Of the 520 patients screened, 248 met the inclusion criteria. Baseline demographics were equally distributed between patients with hyperglycemia and those who were normoglycemic. Serum inflammatory markers in patients with or without new-onset hyperglycemia were elevated as follows: CRP (58.1% *vs* 65.6%, $P = 0.29$), ferritin (48.4% *vs* 34.9%, $P = 0.14$), D-dimer (37.1% *vs* 37.1%, $P = 0.76$) and LDH (19.4% *vs* 11.8%, $P = 0.02$). Logistic regression analysis showed LDH odds ratio (OR) = 1.623 ($P = 0.256$). We observed significantly higher mortality (24.2% *vs* 9.1%, $P = 0.001$; OR = 2.528, $P = 0.024$) and length of stay (8.89 *vs* 6.69, $P = 0.026$) in patients with hyperglycemia.

CONCLUSION

Our study showed no association between CRP, ferritin, LDH, D-dimer levels, and new-onset hyperglycemia in non-diabetic patients with COVID-19 infection. It also shows an increased mortality risk and length of stay in patients with hyperglycemia. With new-onset hyperglycemia being closely associated with poor prognostic indices, it becomes pivotal to understand the underlying pathophysiological mechanisms behind the SARS-CoV-2 infection-induced hyperglycemia. We conclude that the stress hyperglycemia hypothesis is not the only mechanism of SARS-CoV-2 infection-induced hyperglycemia but rather a multicausal pathogenesis leading to hyperglycemia that requires further research and understanding. This would help us improve not only the clinical outcomes of COVID-19 disease and inpatient hyperglycemia management but also understand the long-term effects of SARS-CoV-2 infection and further management.

Key Words: COVID-19; Inflammatory markers; Hyperglycemia; C-reactive protein; Mortality; Severity; Mechanisms; Diabetes mellitus

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Our study suggests that there is no correlation between the inflammatory marker levels and the presence of hyperglycemia in non-diabetic patients with coronavirus disease 2019 (COVID-19) infection. With an increased need to understand the mechanism underlying severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection-induced hyperglycemia, we assessed the validity of the most accepted COVID-19 infection-induced cytokine storm-related stress hyperglycemia theory. However, our study did not show any correlation between inflammatory marker levels that correlate with the cytokine storm and the level of hyperglycemia. This suggests the possibility of other mechanisms playing a role in the SARS-CoV-2 infection-induced hyperglycemia. Our study also demonstrated that new-onset hyperglycemia was an independent risk factor for higher mortality and length of stay, thereby emphasizing the need to understand the mechanisms leading to hyperglycemia.

Citation: Geetha HS, Singh G, Sekar A, Gogtay M, Singh Y, Abraham GM, Trivedi N. Hyperglycemia in COVID-19 infection without diabetes mellitus: Association with inflammatory markers. *World J Clin Cases* 2023; 11(6): 1287-1298

URL: <https://www.wjgnet.com/2307-8960/full/v11/i6/1287.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i6.1287>

INTRODUCTION

At the end of 2019, a novel coronavirus rapidly spread worldwide, resulting in a global pandemic. The virus was designated severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), and the illness it caused was coronavirus disease 2019 (COVID-19). The spectrum of COVID-19 in adults ranges from asymptomatic infection to mild respiratory tract symptoms to severe pneumonia with acute respiratory distress syndrome and multiorgan dysfunction.

The presence of hyperglycemia was an independent factor associated with poor outcomes. One study of hospitalized, elderly COVID-19 patients in Wuhan reported that 21.6% had a history of diabetes, 20.8% were newly diagnosed with diabetes [fasting admission glucose ≥ 7.0 mmol/L or glycated hemoglobin (HbA1c) $\geq 6.5\%$], and 28.4% were diagnosed with dysglycemia (fasting glucose 5.6–6.9 mmol/L or HbA1c 5.7%–6.4%)[1].

Although there is clear evidence demonstrating the association between COVID-19 disease and hyperglycemia, there exists a lack of sufficient literature explaining the mechanism of action of SARS-CoV-2-induced hyperglycemia. The pathophysiological mechanisms of hyperglycemia in COVID-19 patients remain poorly understood. Several complex processes have been hypothesized, including previously undiagnosed diabetes, stress hyperglycemia, steroid-induced hyperglycemia, and direct or indirect effects of SARS-CoV-2 on the pancreatic β -cell[2].

Studies have shown that patients with newly diagnosed diabetes have higher inflammatory markers such as C-reactive protein (CRP), erythrocyte sedimentation rate, and white blood cells. Acute inflammation seen in cytokine storms may worsen insulin resistance. One study shows that neutrophils, D-dimers, and inflammatory markers are significantly higher in those with hyperglycemia than those with normal glucose[3].

We assessed the rationale of the cytokine-induced hyperglycemia hypothesis by evaluating the association between routinely tested inflammatory markers like CRP, ferritin, Lactate dehydrogenase (LDH), and D-dimer and new onset hyperglycemia in non-diabetic patients with COVID-19 infection.

MATERIALS AND METHODS

Study design and participants

We conducted a retrospective, single-center case-control study of hospitalized patients between Dec 1, 2019, and Jan 1, 2022, at a 329 bed community teaching hospital in central Massachusetts. The study inclusion criteria included: (1) Inpatients diagnosed with SARS-CoV-2 infection; (2) Age > 18 years; and (3) Patients with documented inflammatory markers and glucose levels on admission. The exclusion criteria included: (1) Pregnant patients; (2) Patients with previous diagnoses of Type 1 or Type 2 Diabetes mellitus (DM), and (3) Patients who received steroids before admission. The data were obtained by reviewing the patients' medical records, including demographic information, past medical history, medication, labs, and course during hospitalization. The Institutional Review Board approved this study.

Exposure and outcomes

The primary endpoint measured was hyperglycemia, defined as glucose levels ≥ 140 mg/dL. The patients with hyperglycemia were defined as cases, and the control group included patients with normoglycemia (glucose levels < 140 mg/dL). We assessed the level of inflammatory markers between the two study groups, including CRP, ferritin, LDH, and D-dimer levels. We used prespecified cutoffs for the inflammatory markers described by previous studies[4-6]. The inflammatory markers were categorized as binary variables, *i.e.* either elevated or normal. A CRP ≥ 100 mg/L, ferritin ≥ 530 ng/mL, LDH ≥ 590 U/L, and D-dimer ≥ 0.5 mg/L were considered elevated. Demographic data collected included age, sex, weight, height, body mass index (BMI), and vaccination status. Relevant clinical data that is associated with hyperglycemia in previous studies which included a family history of diabetes, past medical history of prediabetes, American Diabetes Association (ADA) diabetes risk score, hypertension, chronic liver disease (CLD), chronic obstructive pulmonary disease (COPD), congestive heart failure (CHF), and chronic kidney disease (CKD) was collected. We also collected data about the outcomes of these patients that included mortality rate, use of steroids, use of remdesivir, and length of hospital stay. The ADA diabetes risk assessment tool that provides a risk stratification index for developing Type II DM was also used.

Statistical analyses

The data were subjected to the normalcy test (Shapiro-Wilk test) that showed non-normalcy distribution. Hence, non-parametric tests were employed. The baseline demographic characteristics between the two study populations were assessed using χ^2 analysis for categorical variables and the Mann-Whitney *U* test for continuous variables. The association between the covariates and primary endpoints was evaluated using a bivariate logistic regression with hyperglycemia as a dependent variable and other variables significant in the univariate analysis as independent variables. The level of significance was set at 5%.

RESULTS

Patient characteristics

A total of 520 patients were screened, among which 248 met the inclusion criteria, with 186 patients (75%) being normoglycemic on admission and 62 patients (25%) being hyperglycemic on admission. Patients with hyperglycemia were more likely to have a history of hypertension (58.1% *vs* 43.5%, $P = 0.047$) and increased ADA Diabetes risk scoring (66.1% *vs* 51.6%, $P = 0.048$) when compared to patients with normoglycemia (Tables 1 and 2). The demographic data like sex distribution, vaccination status,

Table 1 Demographic information

Patient characteristic	Normoglycemia	Hyperglycemia	Total	P value ²
Male sex	102 54.8%	33 53.2%	135 54.4%	0.83
Vaccinated for COVID	34 18.3%	10 16.1%	44 17.7%	0.87
Hypertension	81 43.5%	36 58.1%	117 47.2%	0.047 ¹
Family history of diabetes	16 8.6%	8 12.9%	24 9.7%	0.32
Prediabetes	5 2.7%	1 1.6%	6 2.4%	1 ³
ADA - At risk	96 51.6%	41 66.1%	137 55.2%	0.05*
Chronic liver disease	7 3.8%	2 3.2%	9 3.6%	1.00
Chronic obstructive pulmonary disease	27 14.5%	7 11.3%	34 13.7%	0.39
Congestive heart failure	33 17.7%	7 11.3%	40 16.1%	0.23
Chronic kidney disease	31 16.7%	11 17.7%	42 16.9%	0.85

¹Significant.² χ^2 test.³Fisher's exact test.

ADA: American Diabetes Association; COVID: Coronavirus disease.

Table 2 Demographic information

Variable	Category	n	Mean	Std dev	Median	IQR	P value ¹
Age, yr	Normoglycemic	186	63.38	18.52	68.50	23.00	0.366
	Hyperglycemic	62	66.32	15.64	65.00	29.00	
Weight, kg	Normoglycemic	186	84.96	26.24	82.50	31.00	0.982
	Hyperglycemic	62	84.71	25.63	82.00	29.00	
Height, cm	Normoglycemic	186	168.48	10.67	166.00	18.00	0.376
	Hyperglycemic	62	167.02	10.99	168.00	17.00	
BMI	Normoglycemic	186	29.62	7.75	28.50	9.00	0.566
	Hyperglycemic	62	30.02	7.52	28.00	9.00	

¹Mann Whitney U test.

BMI: Body mass index; IQR: Interquartile range; Std dev: Standard deviation.

and history of CLD, COPD, CHF, and CKD were similar between the two groups.

Inflammatory markers and hyperglycemia

The association between the levels of inflammatory markers like CRP, ferritin, D-dimer, and LDH in

patients with normoglycemia and patients with hyperglycemia was assessed (Tables 3 and 4). Although there was a difference in the mean levels of CRP (106.56 mg/L *vs* 85.65 mg/L), ferritin (1404.40 ng/mL *vs* 470.38 ng/mL), D dimer (0.15 mg/L *vs* 1.71 mg/L) and LDH levels (319.11 U/L *vs* 224.45 U/L), there were statistically no significant differences between serum inflammatory markers except LDH in patients with or without new-onset hyperglycemia [CRP (58.1% *vs* 65.6%, $P = 0.29$), ferritin (48.4% *vs* 34.9%, $P = 0.14$), D-dimer (37.1% *vs* 37.1%, $P = 0.76$) and LDH (19.4% *vs* 11.8%, $P = 0.02$)]. On further binary logistic regression analysis to predict hyperglycemia, there was no difference in LDH levels between the two groups (OR = 1.623, $P = 0.256$).

Hyperglycemia and prognosis

We further analyzed the prognostic indices like mortality rate and length of stay between the two groups (Tables 5 and 6). There was significantly higher mortality (24.2% *vs* 9.1%, $P = 0.001$) and length of stay (8.89 d *vs* 6.69 d, $P = 0.026$) in patients with hyperglycemia compared to patients with normoglycemia. Further analysis with binary logistic regression shows an increased risk of mortality in patients with hyperglycemia (OR = 2.528, $P = 0.024$). The administration of remdesivir initially showed a statistically significant difference in patients with hyperglycemia (59.7% *vs* 44.6%, $P = 0.04$) but did not show any significant difference in binary logistic regression (OR = 1.620, CI: 0.882 - 2.974) (Table 7). There was also no significant difference in the rate of steroid administration between the two groups.

DISCUSSION

COVID-19 and Hyperglycemia

The SARS-CoV-2 infection that was first noticed in December 2019 has resulted in the worldwide COVID-19 pandemic that has claimed over 6.5 million lives worldwide[7]. Initially thought to be a pathogen that affects only the respiratory system, as we continued to learn more about the virus, its effects on multiple organ systems are being slowly discovered, with several studies assessing its association in different disease states[8,9]. Studies have revealed the association of certain risk factors with increased predisposition and severity of the disease. DM has been shown to increase the risk of morbidity and mortality due to its associated metabolic, microvascular, and macrovascular complications and reactive hyperglycemia being a predictor of severity in previous SARS-CoV-1 and Middle East Respiratory Syndrome coronavirus-CoV infections. We noticed an increased prevalence of hyperglycemia in patients with COVID-19 infection, irrespective of the presence of pre-existing diabetes mellitus. An Italian study of 271 patients admitted for COVID-19 showed that hyperglycemia was independently associated with increased mortality[10]. However, this study failed to elucidate the underlying pathophysiological mechanisms explaining SARS-CoV-2 infection-induced hyperglycemia. In our study, we investigated the association of hyperglycemia at presentation with inflammatory markers and the impact of hyperglycemia on mortality and morbidity in non-diabetic COVID-19 patients. The results showed no association between CRP, ferritin, LDH, and D-dimer levels and new-onset hyperglycemia in non-diabetic patients with COVID-19 infection.

Although age and male sex were the initially associated risk factors, further studies revealed the increased prevalence of hypertension and diabetes mellitus in patients with severe disease compared to those with milder forms of infection[11,12]. Our study showed a similar finding of increased hypertension in patients with hyperglycemia. This was further analyzed by binary logistic regression to account for hypertension as a potential confounding agent, given the increased mortality and length of stay in the hyperglycemia group. However, further analysis demonstrated the absence of a significant difference in the presence of hypertension between the two groups. Other risk factors such as male sex, increased BMI, and vaccination status and comorbidities such as chronic liver disease, congestive heart failure, chronic obstructive pulmonary disease, and chronic kidney disease were similar between the two groups, eliminating potential confounders. Although we accounted for major potential confounders, the demographic prevalence of prediabetes was not accounted for due to the non-availability of baseline HbA1c levels.

Covid and Inflammatory markers

With the advent of the COVID-19 pandemic, the utility of inflammatory markers has been on the rise. Biomarkers are quantitative measurements that reflect the pathophysiology of the disease and help gauge the underlying disease severity. Initial studies demonstrated increased levels of inflammatory markers in COVID-19 patients that directly correlated with the disease severity.

Although a milieu of inflammatory markers like Interleukin(IL)-6, IL-1 β , and IL-8 exist, these are not routinely assessed in labs. It would not be cost-effective to employ them in the routine monitoring of every patient with SARS-CoV-2 infection, thus necessitating the use of other inflammatory markers that are cost-effective and can be utilized to assess the severity of the disease. This led to further research on the utility of markers like C-reactive protein, ferritin, LDH, and D-dimer. Studies have demonstrated the utility of these biomarkers in correlating with the severity of the disease. The meta-analyses by Malik *et al*[13] showed an increased association between the elevated levels of different inflammatory markers

Table 3 Inflammatory marker levels between the two study groups on admission

Variable	Category	n	Mean	Std dev	Median	IQR	P value ²
CRP levels on admission	Normoglycemic	186	85.65	74.12	84.00	149.00	0.278
	Hyperglycemic	62	106.56	94.09	74.00	109.00	
Ferritin levels on admission	Normoglycemic	186	470.38	984.06	385.50	1109.50	0.028 ¹
	Hyperglycemic	62	1404.40	4758.84	124.00	647.50	
LDH levels on admission	Normoglycemic	186	224.45	245.15	276.00	319.50	0.041 ¹
	Hyperglycemic	62	319.11	326.57	231.00	389.00	
D-dimer levels on admission	Normoglycemic	186	1.71	4.63	1.00	1.25	0.470
	Hyperglycemic	62	2.15	6.26	0.00	1.00	

¹Significant.²Mann-Whitney U test.

CRP: C-reactive protein; IQR: Interquartile range; LDH: Lactate dehydrogenase; Std dev: Standard deviation.

Table 4 Inflammatory marker levels between the two study groups

Variable	Normoglycemia	Hyperglycemia	Total	P value ²
CRP ≥ 100 mg/L	122	36	158	0.29
	65.6%	58.1%	63.7%	
Ferritin ≥ 530 ng/mL	65	30	95	0.14
	34.9%	48.4%	38.3%	
LDH ≥ 590 U/L	22	12	34	0.02 ¹
	11.8%	19.4%	13.7%	
D-dimer ≥ 0.5 mg/L	69	23	92	0.76
	37.1%	37.1%	37.1%	

¹Significant.² χ^2 test.

CRP: C-reactive protein; LDH: Lactate dehydrogenase.

Table 5 Analyses of outcomes

Variable	Normoglycemia	Hyperglycemia	Total	P value ²
Mortality	17	15	32	0.001 ¹
	9.1%	24.2%	12.9%	
Received remdesivir	83	37	120	0.04 ¹
	44.6%	59.7%	48.4%	
Received steroids	157	56	213	0.25
	84.4%	90.3%	85.9%	

¹Significant.² χ^2 test.

and the severity of COVID-19 disease. Elevated CRP levels (> 10 mg/L) showed a fourfold increase in poor outcomes. Elevated D-dimer values (\geq 0.5 mg/L) were associated with a threefold higher risk of poor outcomes in COVID-19 patients. Elevated LDH showed a fivefold increased risk of poor outcomes. The Meta-analysis by Huang *et al*[14] demonstrated that patients with poor composite outcome had higher levels of Ferritin. Although various studies used different values of CRP such as Koozi *et al*[4] \geq

Table 6 Analyses of outcomes

Variable	Category	n	Mean	Std dev	Median	IQR	P value ²
Length of stay	Normoglycemic	186	6.69	4.38	6.50	6.00	0.026 ¹
	Hyperglycemic	62	8.89	6.80	5.00	6.00	

¹Significant.²Mann-Whitney *U* test.

IQR: Interquartile range; Std dev: Standard deviation.

Table 7 Binary logistic regression for hyperglycemia

Variable	P value	OR	95%CI for OR	
			Lower	Upper
LDH \geq 590 U/L	0.256	1.623	0.704	3.745
Hypertension	0.375	1.355	0.692	2.651
Mortality	0.024 ¹	2.528	1.129	5.662
ADA	0.256	0.616	0.267	1.421
Received remdesivir	0.120	1.620	0.882	2.974

¹Significant.

ADA American Diabetes Association; CI: Confidence interval; LDH: Lactate dehydrogenase; OR: Odds ratio.

1000 mg/L, Ryoo *et al*[5] \geq 140 mg/L, and Liu *et al*[6] \geq 41.8 mg/L, we uniformly observed increased risk of COVID-19 severity with elevated levels of CRP. Despite its value in predicting a poor outcome in COVID-19, it should be noted that various factors could affect serum CRP levels, including age, sex, smoking status, weight, lipid levels, blood pressure, and liver injury[15].

Mechanisms of new-onset hyperglycemia in COVID-19

As we continued to discover the close association between COVID-19 disease and hyperglycemia, several theories were postulated on the underlying pathophysiology between SARS-CoV-2 infection and the onset of hyperglycemia (Figure 1).

Stress hyperglycemia

The most widely accepted hypothesis explaining the mechanism of new-onset hyperglycemia in SARS-CoV-2 infection is the stress hyperglycemia theory. Previous studies have shown that any acute illness, including myocardial infarction or severe infection, tends to drive lipolysis production and an increased number of free fatty acids in the blood[16] which in turn causes production of proinflammatory cytokines. In COVID-19 infection, we expect a similar mechanism or even greater response due to the profound cytokine storm triggered by the SARS-CoV-2 infection. Since previous studies had established the correlation between the levels of inflammatory markers like C-reactive protein, ferritin, LDH, and D-dimer and the severity of SARS-CoV-2 infection, we employed these markers to assess the presence of hyperglycemia in correlation with the disease severity. Our study showed no difference in the levels of C-reactive protein, ferritin, and D-dimer between the two groups. Although initial results showed a significant difference in the LDH levels between the two groups, further binary logistic regression did not reveal any significant change. The mean levels of C-reactive protein, ferritin, LDH, and D-dimer were higher in patients with hyperglycemia. However, no significant difference was observed when a cutoff for severity was imposed based on the established inflammatory marker levels for severe COVID-19 disease.

Preadmission diabetes

One of the first theories postulated was the presence of undiagnosed preexisting diabetes in the population that was brought to light when such patients were admitted to the hospital for COVID-19 infection. Studies also supported this hypothesis, such as recently increased weight gain and worsening hyperglycemia due to mandatory isolation, reduced physical activities, social distancing, and poor mental health that led to eating disorders[17]. These changes could potentially lead to metabolic syndrome that culminates in insulin resistance resulting in hyperglycemia. However, our study accounted for such changes by calculating the ADA diabetes risk score, a validated tool by the American




Figure 1 Mechanisms of hyperglycemia in patients with severe acute respiratory syndrome–coronavirus–2 infection. ACE: Angiotensin converting enzyme; SARS-CoV-2: Severe acute respiratory syndrome-coronavirus–2.

Diabetes Association, to detect an individual's risk of developing diabetes mellitus[18]. A cutoff score of 5 or greater indicates a greater risk of developing diabetes when compared to the general population. In our study, although the initial χ^2 analysis showed a significantly greater ADA risk score in the hyperglycemic study group, further analysis with binary logistic regression showed no statistically significant difference in the ADA risk score between the two groups. This finding emphasizes the poor validity of the preexisting undiagnosed diabetes hypothesis.

Impairments in insulin secretion

The first association between SARS-CoV-2 infection and pancreatic damage was postulated when the presence of Angiotensin converting enzyme 2 receptors, the binding site for SARS-CoV-2, was identified in the pancreatic islet cells[19,20]. More recent studies have observed that SARS-CoV-2 can infect pancreatic cells permitting entry of the virus, leading to attenuation of pancreatic insulin levels and consequently leading to β -cell apoptosis[21,22].

Steroid-induced diabetes

A meta-analysis of 13 studies showed that 32.3% of patients treated for COVID-19 infection with dexamethasone developed glucocorticoid-induced hyperglycemia and 18.6% of patients developed new-onset diabetes[23]. Since steroid-induced hyperglycemia would confound our results, we negated the effects of glucocorticoids by obtaining blood glucose levels on admission before administering steroids. We also excluded patients who had received steroids at outside facilities or in the ED before collecting their on-admission blood glucose levels.

Clinical outcomes of new-onset hyperglycemia

There is growing evidence that hyperglycemia in COVID-19 patients bears significant prognostic implications. Hyperglycemia in critically ill patients is a common manifestation directly correlated with increased mortality or morbidity[24]. More importantly, hyperglycemia was found to worsen the progression of respiratory failure.

Mortality related to hyperglycemia

Platelet activation incited by Fc glycosylated immune complexes is consistent with platelet hyperactivity in severe COVID-19 patients. Excessive macrophage stimulation by enhanced Fc-glycosylated immune complexes is consistent with the macrophage activation syndrome. The resulting hypercoagulability with compromised microperfusion, pulmonary endothelial fluid leakage, and severe respiratory distress

syndrome can result in death[25]. In our study, we observed a significantly increased rate of mortality in patients with new onset hyperglycemia compared to those without. This correlated with the findings of existing literature, such as that by Bode *et al*[26] (2020), who demonstrated that uncontrolled hyperglycemia without prior diabetes was related to an increase in mortality in comparison to normoglycemic subjects. Our study also observed an increased mortality rate in both groups, patients with hyperglycemia (24.2%) and normoglycemia (9.1%), compared to other studies. Our center is one of the referral hospitals receiving the most severe COVID-19 patients, which might explain the higher mortality rate compared to previous studies from China and the United States.

Impact of new-onset hyperglycemia on length of stay

We also observed a significant increase in the mean length of stay in patients with hyperglycemia compared to patients with normoglycemia. This impact of hyperglycemia in those without an established diagnosis of diabetes is a more concerning matter for clinicians, and research has shown that this could be explained by the acute hyperglycemia causing impairment in innate immunity, leading to a heightened risk of infections and increasing the length of stay (LOS) in hospital. In the study by Bode *et al*[26], among 493 discharged patients, the median LOS was longer in 184 patients with hyperglycemia compared with 386 patients without diabetes or hyperglycemia (5.7 *vs* 4.3 d, $P < 0.001$). Our study results are parallel and support the findings of increased mortality and length of stay in patients with hyperglycemia compared with patients with normoglycemia. These results suggest that disease severity and mortality risk significantly increase with newly diagnosed hyperglycemia after hospital admission.

Limitations

One of the most important limitations of the study is the small study population. Since the study was based on a community hospital in the United States, the generalizability of the study is limited. Although due consideration was provided regarding the possible confounding factors, due to the study's retrospective nature, it is possible that all potential confounding factors were not adequately adjusted for. Furthermore, the current study was performed during a period of travel limitations and limited physical activity, which could have contributed to the increased risk of dysglycemia in the study population. Our study is also limited by the inflammatory markers used since it did not include markers like IL-6, IL-1 β , and IL-8, which better correlate with the level of inflammation. We were also limited by the non-availability of baseline HbA1c levels in the study population in order to negate any pre-existing diabetic conditions.

Future implications

Further studies are needed to analyze the different potential mechanisms underlying SARS-CoV-2 infection-induced hyperglycemia. Basic science research that would help better understand the underlying pathophysiology and further clinical studies that assess the utility of different treatment strategies in managing hyperglycemia are required to improve the clinical outcomes of COVID-19.

CONCLUSION

Our study investigated the association of hyperglycemia at presentation with inflammatory markers and the impact of hyperglycemia on mortality and morbidity in non-diabetic COVID-19 patients. The current study showed no association between CRP, ferritin, LDH, and D-dimer levels and new-onset hyperglycemia in non-diabetic patients with COVID-19 infection. It also shows an increased mortality risk and length of stay in patients with hyperglycemia. With new-onset hyperglycemia being closely associated with poor prognostic indices, it becomes pivotal to understand the underlying pathophysiological mechanisms behind the SARS-CoV-2 infection-induced hyperglycemia. This will help us better control the glycemic status and prevent new-onset hyperglycemia, thereby improving the clinical recovery of patients. Although current guidelines recommend the treatment of hyperglycemia using an Insulin sliding scale, these recommendations are based on the hypothesis that hyperglycemia secondary to COVID-19 infection is due to cytokine storm-induced stress hyperglycemia. However, if hyperglycemia arises due to other possible mechanisms, such as viral infection-induced direct suppression of insulin secretion, treatment modalities of achieving glycemic control have to be adjusted to better manage the disease and improve prognosis. Our study results indicate the low probability of the stress hyperglycemia hypothesis being the sole mechanism of SARS-CoV-2 infection-induced hyperglycemia but rather a multicausal pathogenesis leading to hyperglycemia that requires further research and understanding. This would help us improve not only the clinical outcomes of COVID-19 disease and inpatient hyperglycemia management but also understand the long-term effects of SARS-CoV-2 infection and further management.

ARTICLE HIGHLIGHTS

Research background

New onset hyperglycemia is common in patients with severe coronavirus disease 2019 (COVID-19) infection. Cytokine storm and direct severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)-induced pancreatic β -cell failure have been postulated to play a role in new-onset hyperglycemia.

Research motivation

Based on evidence regarding the "Immune-mediated inflammatory storm" during the COVID-19 illness. It has also been proposed that COVID-19 is likely associated with an increased risk of developing diabetes. This motivated us to study the underlying mechanisms contributing to new onset hyperglycemia in hospitalized COVID-19 cases.

Research objectives

To assess the validity of the cytokine-induced hyperglycemia hypothesis by evaluating the association between inflammatory markers and new onset hyperglycemia in non-diabetic patients with COVID-19 infection.

Research methods

A retrospective case-control study was conducted on adults without diabetes mellitus hospitalized for COVID-19 infection. The serum levels of glucose and inflammatory markers at presentation before initiation of corticosteroid were collected. Hyperglycemia was defined as glucose levels ≥ 140 mg/dL. Prespecified cutoffs were used for the inflammatory markers. Statistical methods of analysis were used to calculate the logistic regression for hyperglycemia.

Research results

Of the 520 patients screened, 248 met the inclusion criteria. Our study showed no association between C-reactive protein, ferritin, Lactate dehydrogenase, D-dimer levels, and new-onset hyperglycemia in non-diabetic patients with COVID-19 infection. We observed significantly higher mortality and length of stay in patients with hyperglycemia.

Research conclusions

With new-onset hyperglycemia being closely associated with poor prognostic indices, it becomes pivotal to understand the underlying pathophysiological mechanisms behind the SARS-CoV-2 infection-induced hyperglycemia. This will help us better control the glycemic status and prevent new-onset hyperglycemia, thereby improving the clinical recovery of patients. Our study results indicate the low probability of the stress hyperglycemia hypothesis being the sole mechanism of SARS-CoV-2 infection-induced hyperglycemia but rather a multicausal pathogenesis leading to hyperglycemia that requires further research and understanding.

Research perspectives

Basic science research that would help better understand the underlying pathophysiology and further clinical studies that assess the utility of different treatment strategies in managing hyperglycemia are required to improve the clinical outcomes of COVID-19.

FOOTNOTES

Author contributions: Geetha HS and Trivedi N conceived the idea for the study; Geetha HS, Gogtay M, Abraham GM, and Trivedi N designed and undertook the literature review; Geetha HS, Singh G, Sekar A, and Gogtay M collected data; Gogtay M and Singh Y performed the statistical analysis, figures, and appendix and analyzed and interpreted the data; Geetha HS, Singh G, Sekar A, Singh Y and Gogtay M wrote the first draft of the manuscript; Geetha HS, Singh G, Sekar A, Gogtay M, Singh Y, Abraham GM, Trivedi N revised the subsequent drafts of the manuscript; All authors reviewed and agreed on the final draft of the manuscript.

Institutional review board statement: The study was reviewed and approved by our local Medical Center Institutional Review Board (Approval No. 2020-035).

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: No additional data is available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license

their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: United States

ORCID number: Harinivaas Shanmugavel Geetha 0000-0001-6293-208X; Maya Gogtay 0000-0001-9955-7121; Yuvaraj Singh 0000-0003-4970-8870; GM Abraham 0000-0003-4296-8362; Nitin Trivedi 0000-0002-2510-8339.

S-Editor: Liu GL

L-Editor: Filipodia

P-Editor: Liu GL

REFERENCES

- 1 Li H, Tian S, Chen T, Cui Z, Shi N, Zhong X, Qiu K, Zhang J, Zeng T, Chen L, Zheng J. Newly diagnosed diabetes is associated with a higher risk of mortality than known diabetes in hospitalized patients with COVID-19. *Diabetes Obes Metab* 2020; **22**: 1897-1906 [PMID: 32469464 DOI: 10.1111/dom.14099]
- 2 Khunti K, Del Prato S, Mathieu C, Kahn SE, Gabbay RA, Buse JB. COVID-19, Hyperglycemia, and New-Onset Diabetes. *Diabetes Care* 2021; **44**: 2645-2655 [PMID: 34625431 DOI: 10.2337/dc21-1318]
- 3 Michalakis K, Ilias I. COVID-19 and hyperglycemia/diabetes. *World J Diabetes* 2021; **12**: 642-650 [PMID: 33995851 DOI: 10.4239/wjd.v12.i5.642]
- 4 Koozi H, Lengquist M, Frigyesi A. C-reactive protein as a prognostic factor in intensive care admissions for sepsis: A Swedish multicenter study. *J Crit Care* 2020; **56**: 73-79 [PMID: 31855709 DOI: 10.1016/j.jcrc.2019.12.009]
- 5 Ryoo SM, Han KS, Ahn S, Shin TG, Hwang SY, Chung SP, Hwang YJ, Park YS, Jo YH, Chang HL, Suh GJ, You KM, Kang GH, Choi SH, Lim TH, Kim WY; Korean Shock Society (KoSS) Investigators. The usefulness of C-reactive protein and procalcitonin to predict prognosis in septic shock patients: A multicenter prospective registry-based observational study. *Sci Rep* 2019; **9**: 6579 [PMID: 31036824 DOI: 10.1038/s41598-019-42972-7]
- 6 Liu F, Li L, Xu M, Wu J, Luo D, Zhu Y, Li B, Song X, Zhou X. Prognostic value of interleukin-6, C-reactive protein, and procalcitonin in patients with COVID-19. *J Clin Virol* 2020; **127**: 104370 [PMID: 32344321 DOI: 10.1016/j.jcv.2020.104370]
- 7 Grauhan A. [Combined session of the International Labor organization and the World Health Organization on job and economic conditions of hospital personnel, from 19-30 November, 1973 in Geneva. 1]. *Dtsch Krankenpflegez* 1975; **28**: 262-264 [PMID: 1038992]
- 8 Tripathi K, Godoy Brewer G, Thu Nguyen M, Singh Y, Saleh Ismail M, Sauk JS, Parian AM, Limketkai BN. COVID-19 and Outcomes in Patients With Inflammatory Bowel Disease: Systematic Review and Meta-Analysis. *Inflamm Bowel Dis* 2022; **28**: 1265-1279 [PMID: 34718595 DOI: 10.1093/ibd/izab236]
- 9 Song D, Geetha HS, Kim A, Seen T, Almas T, Nagarajan VR, Alsaed N, Cheng JH, Lieber J. Transformation of acute cholecystitis to acute choledocholithiasis in COVID-19 patient. *Ann Med Surg (Lond)* 2021; **71**: 102946 [PMID: 34664016 DOI: 10.1016/j.amsu.2021.102946]
- 10 Coppelli A, Giannarelli R, Aragona M, Penno G, Falcone M, Tiseo G, Ghiadoni L, Barbieri G, Monzani F, Virdis A, Menichetti F, Del Prato S; Pisa COVID-19 Study Group. Hyperglycemia at Hospital Admission Is Associated With Severity of the Prognosis in Patients Hospitalized for COVID-19: The Pisa COVID-19 Study. *Diabetes Care* 2020; **43**: 2345-2348 [PMID: 32788285 DOI: 10.2337/dc20-1380]
- 11 Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DSC, Du B, Li LJ, Zeng G, Yuen KY, Chen RC, Tang CL, Wang T, Chen PY, Xiang J, Li SY, Wang JL, Liang ZJ, Peng YX, Wei L, Liu Y, Hu YH, Peng P, Wang JM, Liu JY, Chen Z, Li G, Zheng ZJ, Qiu SQ, Luo J, Ye CJ, Zhu SY, Zhong NS; China Medical Treatment Expert Group for Covid-19. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020; **382**: 1708-1720 [PMID: 32109013 DOI: 10.1056/NEJMoa2002032]
- 12 Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, Wu Y, Zhang L, Yu Z, Fang M, Yu T, Wang Y, Pan S, Zou X, Yuan S, Shang Y. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020; **8**: 475-481 [PMID: 32105632 DOI: 10.1016/S2213-2600(20)30079-5]
- 13 Malik P, Patel U, Mehta D, Patel N, Kelkar R, Akrmah M, Gabrilove JL, Sacks H. Biomarkers and outcomes of COVID-19 hospitalisations: systematic review and meta-analysis. *BMJ Evid Based Med* 2021; **26**: 107-108 [PMID: 32934000 DOI: 10.1136/bmjebm-2020-111536]
- 14 Huang I, Pranata R, Lim MA, Oehadian A, Alisjahbana B. C-reactive protein, procalcitonin, D-dimer, and ferritin in severe coronavirus disease-2019: a meta-analysis. *Ther Adv Respir Dis* 2020; **14**: 1753466620937175 [PMID: 32615866 DOI: 10.1177/1753466620937175]
- 15 Sproston NR, Ashworth JJ. Role of C-Reactive Protein at Sites of Inflammation and Infection. *Front Immunol* 2018; **9**: 754 [PMID: 29706967 DOI: 10.3389/fimmu.2018.00754]
- 16 Capes SE, Hunt D, Malmberg K, Gerstein HC. Stress hyperglycaemia and increased risk of death after myocardial infarction in patients with and without diabetes: a systematic overview. *Lancet* 2000; **355**: 773-778 [PMID: 10711923 DOI: 10.1016/S0140-6736(99)08415-9]
- 17 World Health Organization. COVID-19 significantly impacts health services for noncommunicable diseases. [Internet] [accessed 12 March 2021]. Available from: <https://www.who.int/news/item/01-06-2020-covid-19-significantly-impacts-health-services-for-noncommunicable-diseases>

- 18 **Chima CC**, Anikpezie N, Pongetti LS, Wade BC, Powell T, Beech B. 1443-P: Can the ADA Diabetes Risk Score Be Approximated Using Routine Data in the Electronic Health Record? *Diabetes* 2020; **69** (Supplement_1): 1443-P [DOI: [10.2337/db20-1443-P](https://doi.org/10.2337/db20-1443-P)]
- 19 **Liu F**, Long X, Zhang B, Zhang W, Chen X, Zhang Z. ACE2 Expression in Pancreas May Cause Pancreatic Damage After SARS-CoV-2 Infection. *Clin Gastroenterol Hepatol* 2020; **18**: 2128-2130.e2 [PMID: [32334082](https://pubmed.ncbi.nlm.nih.gov/32334082/) DOI: [10.1016/j.cgh.2020.04.040](https://doi.org/10.1016/j.cgh.2020.04.040)]
- 20 **Figliani D**, Licata G, Brusco N, Nigi L, Grieco GE, Marselli L, Overbergh L, Gysemans C, Colli ML, Marchetti P, Mathieu C, Eizirik DL, Sebastiani G, Dotta F. SARS-CoV-2 Receptor Angiotensin I-Converting Enzyme Type 2 (ACE2) Is Expressed in Human Pancreatic β -Cells and in the Human Pancreas Microvasculature. *Front Endocrinol (Lausanne)* 2020; **11**: 596898 [PMID: [33281748](https://pubmed.ncbi.nlm.nih.gov/33281748/) DOI: [10.3389/fendo.2020.596898](https://doi.org/10.3389/fendo.2020.596898)]
- 21 **Wu CT**, Lidsky PV, Xiao Y, Lee IT, Cheng R, Nakayama T, Jiang S, Demeter J, Bevacqua RJ, Chang CA, Whitener RL, Stalder AK, Zhu B, Chen H, Goltsev Y, Tzankov A, Nayak JV, Nolan GP, Matter MS, Andino R, Jackson PK. SARS-CoV-2 infects human pancreatic β cells and elicits β cell impairment. *Cell Metab* 2021; **33**: 1565-1576.e5 [PMID: [34081912](https://pubmed.ncbi.nlm.nih.gov/34081912/) DOI: [10.1016/j.cmet.2021.05.013](https://doi.org/10.1016/j.cmet.2021.05.013)]
- 22 **Shaharuddin SH**, Wang V, Santos RS, Gross A, Wang Y, Jawanda H, Zhang Y, Hasan W, Garcia G Jr, Arumugaswami V, Sareen D. Deleterious Effects of SARS-CoV-2 Infection on Human Pancreatic Cells. *Front Cell Infect Microbiol* 2021; **11**: 678482 [PMID: [34282405](https://pubmed.ncbi.nlm.nih.gov/34282405/) DOI: [10.3389/fcimb.2021.678482](https://doi.org/10.3389/fcimb.2021.678482)]
- 23 **Liu XX**, Zhu XM, Miao Q, Ye HY, Zhang ZY, Li YM. Hyperglycemia induced by glucocorticoids in nondiabetic patients: a meta-analysis. *Ann Nutr Metab* 2014; **65**: 324-332 [PMID: [25402408](https://pubmed.ncbi.nlm.nih.gov/25402408/) DOI: [10.1159/000365892](https://doi.org/10.1159/000365892)]
- 24 **Plummer MP**, Bellomo R, Cousins CE, Annink CE, Sundararajan K, Reddi BA, Raj JP, Chapman MJ, Horowitz M, Deane AM. Dysglycaemia in the critically ill and the interaction of chronic and acute glycaemia with mortality. *Intensive Care Med* 2014; **40**: 973-980 [PMID: [24760120](https://pubmed.ncbi.nlm.nih.gov/24760120/) DOI: [10.1007/s00134-014-3287-7](https://doi.org/10.1007/s00134-014-3287-7)]
- 25 **Mamtani M**, Athavale AM, Abraham M, Vernik J, Amarah AR, Ruiz JP, Joshi AJ, Itteera M, Zhukovski SD, Madaiah RP, White BC, Hart P, Kulkarni H. Association of hyperglycaemia with hospital mortality in nondiabetic COVID-19 patients: A cohort study. *Diabetes Metab* 2021; **47**: 101254 [PMID: [33781926](https://pubmed.ncbi.nlm.nih.gov/33781926/) DOI: [10.1016/j.diabet.2021.101254](https://doi.org/10.1016/j.diabet.2021.101254)]
- 26 **Bode B**, Garrett V, Messler J, McFarland R, Crowe J, Booth R, Klonoff DC. Glycemic Characteristics and Clinical Outcomes of COVID-19 Patients Hospitalized in the United States. *J Diabetes Sci Technol* 2020; **14**: 813-821 [PMID: [32389027](https://pubmed.ncbi.nlm.nih.gov/32389027/) DOI: [10.1177/1932296820924469](https://doi.org/10.1177/1932296820924469)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

