

World Journal of *Gastrointestinal Endoscopy*

World J Gastrointest Endosc 2022 August 16; 14(8): 474-511



ORIGINAL ARTICLE

Retrospective Cohort Study

- 474 Disparities in colonoscopy utilization for lower gastrointestinal bleeding in rural *vs* urban settings in the United States

Ganta N, Aknouk M, Alnabwani D, Nikiforov I, Bommu VJL, Patel V, Cheriya P, Hollenbeak CS, Hamza A

Retrospective Study

- 487 Percutaneous transluminal angioplasty balloons for endoscopic ultrasound-guided pancreatic duct interventions

AbiMansour JP, Abu Dayyeh BK, Levy MJ, Storm AC, Martin JA, Petersen BT, Law RJ, Topazian MD, Chandrasekhara V

Observational Study

- 495 New application of endocytoscope for histopathological diagnosis of colorectal lesions

Inoue F, Hirata D, Iwatate M, Hattori S, Fujita M, Sano W, Sugai T, Kawachi H, Ichikawa K, Sano Y

CASE REPORT

- 502 Hidden local recurrence of colorectal adenocarcinoma diagnosed by endoscopic ultrasound: A case series

Okasha HH, Wahba M, Fontagnier E, Abdellatef A, Haggag H, AbouElenin S

LETTER TO THE EDITOR

- 508 Laparoscopic and endoscopic cooperative surgery for full-thickness resection and sentinel node dissection for early gastric cancer

Vanella S, Godas M, Pereira JC, Pereira A, Apicella I, Crafa F

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Endoscopy*, Murali Dharan, FASGE, MRCP, Assistant Professor, Department of Gastroenterology and Hepatology, University of Connecticut Health Center, Farmington, CO 06030, United States. dharan@uchc.edu

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Endoscopy* (WJGE, *World J Gastrointest Endosc*) is to provide scholars and readers from various fields of gastrointestinal endoscopy with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGE mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal endoscopy and covering a wide range of topics including capsule endoscopy, colonoscopy, double-balloon enteroscopy, duodenoscopy, endoscopic retrograde cholangiopancreatography, endosonography, esophagoscopy, gastrointestinal endoscopy, gastroscopy, laparoscopy, natural orifice endoscopic surgery, proctoscopy, and sigmoidoscopy.

INDEXING/ABSTRACTING

The WJGE is now abstracted and indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 Journal Citation Indicator (JCI) for WJGE as 0.33.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yi-Xuan Cai; Production Department Director: Xu Guo; Editorial Office Director: Jia-Ping Yan.

NAME OF JOURNAL

World Journal of Gastrointestinal Endoscopy

ISSN

ISSN 1948-5190 (online)

LAUNCH DATE

October 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Anastasios Koulaouzidis, Bing Hu, Sang Chul Lee, Joo Young Cho

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5190/editorialboard.htm>

PUBLICATION DATE

August 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/gerinfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/gerinfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/gerinfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Observational Study

New application of endocytoscope for histopathological diagnosis of colorectal lesions

Fumihiro Inoue, Daizen Hirata, Mineo Iwatate, Santa Hattori, Mikio Fujita, Wataru Sano, Tamotsu Sugai, Hiroshi Kawachi, Kazuhito Ichikawa, Yasushi Sano

Specialty type: Gastroenterology and hepatology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): D
Grade E (Poor): 0

P-Reviewer: Mijwil MM, Iraq;
Tousidonis M, Spain

Received: February 18, 2022

Peer-review started: February 18, 2022

First decision: April 12, 2022

Revised: April 23, 2022

Accepted: July 20, 2022

Article in press: July 20, 2022

Published online: August 16, 2022



Fumihiro Inoue, Daizen Hirata, Mineo Iwatate, Santa Hattori, Mikio Fujita, Wataru Sano, Yasushi Sano, Gastrointestinal Center and Institute of Minimally-invasive Endoscopic Care (i-MEC), Sano Hospital, Kobe 655-0031, Hyogo, Japan

Tamotsu Sugai, Molecular Diagnostic Pathology, Iwate Medical University School of Medicine, Shiwa-gun 028-3694, Iwate, Japan

Hiroshi Kawachi, Department of Pathology, Cancer Institute Hospital of Japanese Foundation for Cancer Research, Koto 135-8550, Tokyo, Japan

Kazuhito Ichikawa, Department of Pathology, Noda Hospital, Noda 270-0237, Chiba, Japan

Corresponding author: Mineo Iwatate, Doctor, Gastrointestinal Center and Institute of Minimally-invasive Endoscopic Care (i-MEC), Sano Hospital, 2-5-1 Shimizugaoka, Tarumi-ku, Kobe 655-0031, Hyogo, Japan. m.iwatate15@gmail.com

Abstract

BACKGROUND

The endocytoscope with ultra-high magnification (x 520) allows us to observe the cellular structure of the colon epithelium during colonoscopy, known as virtual histopathology. We hypothesized that the endocytoscope could directly observe colorectal histopathological specimens and store them as endocyto-pathological images by the endoscopists without a microscope, potentially saving the burden on histopathologists.

AIM

To assess the feasibility of endocyto-pathological images taken by an endoscopist as adequate materials for histopathological diagnosis.

METHODS

Three gastrointestinal pathologists were invited and asked to diagnose 40 cases of endocyto-pathological images of colorectal specimens. Each case contained seven endocyto-pathological images taken by an endoscopist, consisting of one loupe image, three low-magnification images, and three ultra-high magnification images. The participants chose hyperplastic polyp or low-grade adenoma for 20 cases of endocyto-pathological images (10 hyperplastic polyps, and 10 Low-grade adenomas in conventional histopathology) in study 1 and high-grade adenoma/

shallow invasive cancer or deep invasive cancer for 20 cases [10 tumor in situ/T1a and 10 T1b] in study 2. We investigated the agreement between the histopathological diagnosis using the endocyto-pathological images and conventional histopathological diagnosis.

RESULTS

Agreement between the endocyto-pathological and conventional histopathological diagnosis by the three gastrointestinal pathologists was 100% (95%CI: 94.0%–100%) in studies 1 and 2. The interobserver agreement among the three gastrointestinal pathologists was 100%, and the κ coefficient was 1.00 in both studies.

CONCLUSION

Endocyto-pathological images were adequate and reliable materials for histopathological diagnosis.

Key Words: Cancer; Colon; Endocytoscopy; Histopathology; Specimen

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The endocytoscope allows us to observe the histological structure of the colon epithelium, but it is a virtual histopathology. We directly observed pathological specimens by the endocytoscope and evaluated the practical usefulness of endocyto-pathology in this pilot study.

Citation: Inoue F, Hirata D, Iwatate M, Hattori S, Fujita M, Sano W, Sugai T, Kawachi H, Ichikawa K, Sano Y. New application of endocytoscope for histopathological diagnosis of colorectal lesions. *World J Gastrointest Endosc* 2022; 14(8): 495-501

URL: <https://www.wjgnet.com/1948-5190/full/v14/i8/495.htm>

DOI: <https://dx.doi.org/10.4253/wjge.v14.i8.495>

INTRODUCTION

The endocytoscope, which was launched in early 2018 by Olympus Medical Systems Corporation (Tokyo, Japan), can provide ultra-high magnification (x 520) images in real time during colonoscopy. The endocytoscopy allows us to observe the cellular structure of the colorectal lesions, known as virtual histopathology and has provided high diagnostic performance in estimating their histopathology[1-5]. There is growing evidence that the diagnostic accuracy of endocytoscopy with computer-aided diagnosis (CAD) was greater than that of non-expert and comparable to expert endoscopists[6-12].

Based on the background of the shortage of histopathologists, we have explored a new application of endocytoscope for histopathological diagnosis of colorectal lesions[13]. We hypothesized that the endocytoscope could directly observe colorectal histopathological specimens and store them as endocyto-pathological images by the endoscopists themselves without a microscope. The endocyto-pathological images taken by endoscopists can be stored in the same system as the endoscopic images so that both images can be obtained as needed, making it possible to hold clinicopathological conferences efficiently even in countries with a few pathologists. Furthermore, a combination of endocyto-pathological images and the CAD system may lead to saving the burden of histopathologists in the future.

This pilot study aimed to assess the feasibility of endocyto-pathological images taken by an endoscopist as adequate materials for histopathological diagnosis.

MATERIALS AND METHODS


Endocyto-pathological images

First, each specimen was placed horizontally in a white container filled with water to control the diffuse reflection of the scope light. An endoscopist (FI) took the ultra-magnifying images of the specimens (endocyto-pathological images) with the right hand firmly fixed by touching the edge of the container and holding the tip of the scope using a penhold grip (Figure 1). This method helps bring high-quality endocyto-pathological images into focus. Seven endocyto-pathological images were obtained for each case (one loupe image, three low-magnification images, and three ultra-high magnification images) (Figures 2 and 3).




DOI: 10.4253/wjge.v14.i8.495 Copyright ©The Author(s) 2022.

Figure 1 How to take the endocyto-pathological images using an endocytoscope: The right hand was firmly fixed by touching the edge of the container, and the tip of the scope was held in the penhold method.



DOI: 10.4253/wjge.v14.i8.495 Copyright ©The Author(s) 2022.

Figure 2 Endocyto-pathological images of low-grade adenoma. A: Loupe image. B: Low-magnification image. C: Ultra-high magnification image.



DOI: 10.4253/wjge.v14.i8.495 Copyright ©The Author(s) 2022.

Figure 3 Endocyto-pathological images of T1b cancer. A: Loupe image. B: Low-magnification image. C: Ultra-high magnification image.

Selection of colorectal specimens

Candidate colorectal specimens were selected from histopathologically-known material obtained by endoscopic or surgical resection at Sano Hospital between January 2017 and January 2021. Candidates samples with poor preservation, incomplete resection of the lesion, or other candidates deemed inappropriate by the investigators were excluded. Among these candidates samples, 10 specimens for each of the following categories hyperplastic polyps, low-grade adenoma, high-grade adenoma/shallow invasive cancer (10 tumor in situ (Tis)/T1a), and deep invasive cancer (T1b) were randomly selected. The number of specimens in each category was masked to the participants.

Evaluation of endocyto-pathological images by gastrointestinal pathologists

Three gastrointestinal pathologists (TS, HK, KI) were invited and asked to read the endocyto-pathological images for 40 cases (7 images for each case) of colorectal specimens from May to July 2021. The participants were asked to choose hyperplastic polyp or low-grade adenoma for 20 cases of endocyto-pathological images (10 hyperplastic polyps and 10 Low-grade adenomas diagnosed by the conventional method) in study 1 and high-grade adenoma/shallow invasive cancer (Tis/T1a) or deep invasive cancer (T1b) for 20 cases (10 Tis/T1a and 10 Tib cancer) in study 2.

The study protocol was reviewed and approved by the Institutional Review Board at Sano Hospital (202106-02). This study was registered with Japan Registry of Clinical Trials (jRCT1050210046).

Outcome measures

The primary outcome measure was the agreement between the histopathological diagnosis using the endocyto-pathological images and conventional histopathological diagnosis.

The secondary outcome measure was the interobserver agreement rate and Fleiss's Kappa statistics among three pathologists.

Statistical analysis

This study was conducted as an exploratory research investigation without calculating sample size due to the lack of data in previous studies.

RESULTS

Tables 1 and 2 show the agreement between the histopathological diagnosis by three gastrointestinal pathologists using the endocyto-pathological images and conventional histopathological diagnosis in differentiating low-grade adenoma from hyperplastic polyp (study 1) and T1b from Tis/T1a cancer (study 2). The agreement between the endocyto-pathological and conventional histopathological diagnosis was 100% (95%CI: 94.0%-100%) in study 1 and 100% (94.0%-100%) in study 2. The interobserver agreement among the three gastrointestinal pathologists was 100%, and the κ coefficient was 1.00 in both studies.

DISCUSSION

To our knowledge, this is the first report of a new clinical application of the endocytoscope for histopathological specimens. The quality of endocyto-pathological images taken by an endoscopist was sufficiently high to make a histopathological diagnosis. We attempted to take pathological images of histopathological specimens by conventional magnifying endoscopy ($\times 85$ maximum optical magnification with approximately 2mm of a minimum depth of observation); however, cytological findings could not be evaluated owing to a lack of resolution power and focus depth. In contrast, the endocytoscope easily enables the evaluation of cytological findings by taking ultra-high power magnification images with contact on the histological slides. For better quality, the specimens were placed horizontally in a white container filled with water to control the diffuse reflection of the diffuse reflection of the scope light.

Linking endoscopic and histopathological images is a clinically essential step for endoscopists to improve endoscopic diagnosis for estimating the histopathology of gastrointestinal lesions. In situations where pathologists are scarce, it would be better to have endoscopists obtain histopathological images using a microscope. However, most endoscopists do not have microscopes in their institutions or are generally unfamiliar with using them. In this context, we considered it meaningful to have endoscopists obtain histopathological images using endocytoscopes. Additionally, our endocyto-pathological images have the advantage of being stored with endoscopic images in the same endoscopic system, which is helpful when holding clinicopathological conferences. We believe the endocyto-pathological diagnosis will reduce the growing burden on histopathologists, including their time and cost, when especially made with the CAD system. Further studies will be required to prove the hypothesis.

This study has limitations. First, knowledge of histopathology is required for endoscopists to take diagnosable ultra-high magnification images, especially for cancer depth diagnosis. Taking inadequate images would lead to the wrong endocyto-pathological diagnosis. Second, endocytoscopes have not yet been disseminated worldwide. However, the results of this study may encourage the spread of the endocytoscopes, especially in countries with a few pathologists.

Table 1 The agreement between endocyto-pathological and conventional histopathological diagnosis for differentiating low-grade adenoma from hyperplastic polyp by three gastrointestinal pathologists

	Conventional pathological diagnosis	
	Low-grade adenoma (n = 30)	Hyperplastic polyp (n = 30)
Endocyto-pathological diagnosis		
Low-grade adenoma	30	0
Hyperplastic polyp	0	30

Table 2 The agreement between endocyto-pathological and conventional histopathological diagnosis for differentiating T1b from Tis/T1a cancer by three gastrointestinal pathologists

	Conventional pathological diagnosis	
	T1b cancer (n = 30)	Tis/T1a cancer (n = 30)
Endocyto-pathological diagnosis		
T1b cancer	30	0
Tis/T1a cancer	0	30

CONCLUSION

In conclusion, endocyto-pathological images of colorectal lesions were adequate and reliable materials for histopathological diagnosis. Endocytoscopes will be disseminated in the future and have the potential for endocyto-pathology worldwide.

ARTICLE HIGHLIGHTS

Research background

Based on the background of the shortage of histopathologists, we explore the new application of endocytoscope for directly observing histopathological specimens of colorectal lesions and storing them as endocyto-pathological images with their endoscopic images.

Research motivation

Endocyto-pathological images taken by endoscopists potentially reduce the burden of histopathologists and facilitate holding clinicopathological conferences more simply.

Research objectives

To assess the feasibility of endocyto-pathological images taken by an endoscopist as adequate materials for histopathological diagnosis.

Research methods

This was a single-center prospective pilot study. Three gastrointestinal pathologists were asked to diagnose 40 cases of endocyto-pathological images of colorectal specimens (Each case contained seven images: one loupe image, three low-magnification images, and three ultra-high magnification images). The participants chose hyperplastic polyp or low-grade adenoma for 20 cases of endocyto-pathological images (10 hyperplastic polyps, and 10 Low-grade adenomas in conventional histopathology) in study 1 and high-grade adenoma/shallow invasive cancer or deep invasive cancer for 20 cases [10 tumor in situ (Tis)/T1a and 10 T1b] in study 2.

Research results

Agreement between the endocyto-pathological and conventional histopathological diagnosis by the three gastrointestinal pathologists was 100% (95%CI: 94.0%–100%) in studies 1 and 2. The interobserver agreement among the three gastrointestinal pathologists was 100%, and the κ coefficient was 1.00 in both studies.

Research conclusions

Endocytology-pathological images were adequate and reliable materials for histopathological diagnosis.

Research perspectives

Endocytology-pathological images taken by endoscopists will reduce the growing burden on histopathologists, including their time and cost, when especially used with the computer-aided diagnosis system.

FOOTNOTES

Author contributions: Inoue F, Hirata D, Iwatate M, Hattori S, Fujita M, Sano W and Sano Y contributed to the study concept and design; Sugai T, Kawachi H and Ichikawa K contributed to read endocytology-pathological images; Inoue F, Hirata D, Iwatate M and Sano Y contributed to the data analysis and interpretation; Inoue F contributed to draft the manuscript; and Sugai T, Kawachi H, Ichikawa K and Sano Y contributed to the critical revision of the manuscript for intellectual content.

Institutional review board statement: The study protocol was reviewed and approved by the Institutional Review Board at Sano Hospital (No. 202106-02).

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: All authors have no conflicts of interest to report.

Data sharing statement: No additional data are available.

STROBE statement: The authors have read the STROBE Statement—checklist of items, and the manuscript was prepared and revised according to the STROBE Statement—checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Japan

ORCID number: Fumihiro Inoue 0000-0002-1689-2977; Daizen Hirata 0000-0001-7255-6129; Mineo Iwatate 0000-0003-3782-3687; Santa Hattori 0000-0002-5926-3417; Mikio Fujita 0000-0003-4673-7545; Wataru Sano 0000-0002-1401-5591; Tamotsu Sugai 0000-0002-4896-3557; Hiroshi Kawachi 0000-0002-8270-791X; Kazuhito Ichikawa 0000-0003-3462-6012; Yasushi Sano 0000-0002-3352-5757.

S-Editor: Ma YJ

L-Editor: A

P-Editor: Ma YJ

REFERENCES

- 1 **Takamaru H**, Wu SYS, Saito Y. Endocytoscopy: technology and clinical application in the lower GI tract. *Transl Gastroenterol Hepatol* 2020; **5**: 40 [PMID: 32632391 DOI: 10.21037/tgh.2019.12.04]
- 2 **Rotondano G**, Bianco MA, Salerno R, Meucci C, Prisco A, Garofano ML, Sansone S, Cipolletta L. Endocytoscopic classification of preneoplastic lesions in the colorectum. *Int J Colorectal Dis* 2010; **25**: 1111-1116 [PMID: 20532533 DOI: 10.1007/s00384-010-0969-7]
- 3 **Mori Y**, Kudo SE, Ogawa Y, Wakamura K, Kudo T, Misawa M, Hayashi T, Katagiri A, Miyachi H, Inoue H, Oka S, Matsuda T. Diagnosis of sessile serrated adenomas/polyps using endocytoscopy (with videos). *Dig Endosc* 2016; **28** Suppl 1: 43-48 [PMID: 26748690 DOI: 10.1111/den.12601]
- 4 **Sugihara Y**, Kudo SE, Miyachi H, Wakamura K, Mori Y, Misawa M, Hisayuki T, Kudo T, Hayashi T, Hamatani S, Okoshi S, Okada H. *In vivo* detection of desmoplastic reaction using endocytoscopy: A new diagnostic marker of submucosal or more extensive invasion in colorectal carcinoma. *Mol Clin Oncol* 2017; **6**: 291-295 [PMID: 28451401 DOI: 10.3892/mco.2017.1138]
- 5 **Kudo T**, Kudo SE, Mori Y, Wakamura K, Misawa M, Hayashi T, Miyachi H, Katagiri A, Ishida F, Inoue H. Classification of nuclear morphology in endocytoscopy of colorectal neoplasms. *Gastrointest Endosc* 2017; **85**: 628-638 [PMID: 27876633 DOI: 10.1016/j.gie.2016.10.039]
- 6 **Mori Y**, Kudo SE, Wakamura K, Misawa M, Ogawa Y, Kutsukawa M, Kudo T, Hayashi T, Miyachi H, Ishida F, Inoue H. Novel computer-aided diagnostic system for colorectal lesions by using endocytoscopy (with videos). *Gastrointest Endosc*

- 2015; **81**: 621-629 [PMID: [25440671](#) DOI: [10.1016/j.gie.2014.09.008](#)]
- 7 **Mori Y**, Kudo SE, Chiu PW, Singh R, Misawa M, Wakamura K, Kudo T, Hayashi T, Katagiri A, Miyachi H, Ishida F, Maeda Y, Inoue H, Nimura Y, Oda M, Mori K. Impact of an automated system for endocytoscopic diagnosis of small colorectal lesions: an international web-based study. *Endoscopy* 2016; **48**: 1110-1118 [PMID: [27494455](#) DOI: [10.1055/s-0042-113609](#)]
 - 8 **Mori Y**, Kudo SE, Misawa M, Saito Y, Ikematsu H, Hotta K, Ohtsuka K, Urushibara F, Kataoka S, Ogawa Y, Maeda Y, Takeda K, Nakamura H, Ichimasa K, Kudo T, Hayashi T, Wakamura K, Ishida F, Inoue H, Itoh H, Oda M, Mori K. Real-Time Use of Artificial Intelligence in Identification of Diminutive Polyps During Colonoscopy: A Prospective Study. *Ann Intern Med* 2018; **169**: 357-366 [PMID: [30105375](#) DOI: [10.7326/M18-0249](#)]
 - 9 **Takeda K**, Kudo SE, Mori Y, Misawa M, Kudo T, Wakamura K, Katagiri A, Baba T, Hidaka E, Ishida F, Inoue H, Oda M, Mori K. Accuracy of diagnosing invasive colorectal cancer using computer-aided endocytoscopy. *Endoscopy* 2017; **49**: 798-802 [PMID: [28472832](#) DOI: [10.1055/s-0043-105486](#)]
 - 10 **Misawa M**, Kudo SE, Mori Y, Takeda K, Maeda Y, Kataoka S, Nakamura H, Kudo T, Wakamura K, Hayashi T, Katagiri A, Baba T, Ishida F, Inoue H, Nimura Y, Oda M, Mori K. Accuracy of computer-aided diagnosis based on narrow-band imaging endocytoscopy for diagnosing colorectal lesions: comparison with experts. *Int J Comput Assist Radiol Surg* 2017; **12**: 757-766 [PMID: [28247214](#) DOI: [10.1007/s11548-017-1542-4](#)]
 - 11 **Misawa M**, Kudo SE, Mori Y, Nakamura H, Kataoka S, Maeda Y, Kudo T, Hayashi T, Wakamura K, Miyachi H, Katagiri A, Baba T, Ishida F, Inoue H, Nimura Y, Mori K. Characterization of Colorectal Lesions Using a Computer-Aided Diagnostic System for Narrow-Band Imaging Endocytoscopy. *Gastroenterology* 2016; **150**: 1531-1532.e3 [PMID: [27072671](#) DOI: [10.1053/j.gastro.2016.04.004](#)]
 - 12 **Mori Y**, Kudo SE, Misawa M, Hotta K, Kazuo O, Saito S, Ikematsu H, Saito Y, Matsuda T, Kenichi T, Kudo T, Nemoto T, Itoh H, Mori K. Artificial intelligence-assisted colonic endocytoscopy for cancer recognition: a multicenter study. *Endosc Int Open* 2021; **9**: E1004-E1011 [PMID: [34222622](#) DOI: [10.1055/a-1475-3624](#)]
 - 13 **Provenzano E**, Driskell OJ, O'Connor DJ, Rodriguez-Justo M, McDermott J, Wong N, Kendall T, Zhang YZ, Robinson M, Kurian KM, Pell R, Shaaban AM. The important role of the histopathologist in clinical trials: challenges and approaches to tackle them. *Histopathology* 2020; **76**: 942-949 [PMID: [32145084](#) DOI: [10.1111/his.14099](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

