

World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2022 July 27; 14(7): 632-730



Contents

Monthly Volume 14 Number 7 July 27, 2022

REVIEW

- 632 From advanced diagnosis to advanced resection in early neoplastic colorectal lesions: Never-ending and trending topics in the 2020s

Auriemma F, Sferrazza S, Bianchetti M, Savarese MF, Lamonaca L, Paduano D, Piazza N, Giuffrida E, Mete LS, Tucci A, Milluzzo SM, Iannelli C, Repici A, Mangiavillano B

ORIGINAL ARTICLE

Retrospective Study

- 656 Adult patients with allied disorders of Hirschsprung's disease in emergency department: An 11-year retrospective study

Jiang S, Song CY, Feng MX, Lu YQ

- 670 Impact of comorbid renal dysfunction in patients with hepatocellular carcinoma on long-term outcomes after curative resection

Sakamoto Y, Shimada S, Kamiyama T, Sugiyama K, Asahi Y, Nagatsu A, Orimo T, Kakisaka T, Kamachi H, Ito YM, Taketomi A

- 685 Individualized risk estimation for postoperative pulmonary complications after hepatectomy based on perioperative variables

Xu LN, Xu YY, Li GP, Yang B

SYSTEMATIC REVIEWS

- 696 Skeletal muscle metastasis from colorectal adenocarcinoma: A literature review

Kulkarni N, Khalil A, Bodapati S

CASE REPORT

- 706 Percutaneous aspiration and sclerotherapy of a giant simple hepatic cyst causing obstructive jaundice: A case report and review of literature

He XX, Sun MX, Lv K, Cao J, Zhang SY, Li JN

- 714 Indocyanine green fluorescence imaging for spleen preservation in laparoscopic splenic artery aneurysm resection: A case report

Cheng J, Sun LY, Liu J, Zhang CW

LETTER TO THE EDITOR

- 720 Total mesopancreas excision is the better staging tool of the mesopancreas in pancreatic head carcinoma

Peparini N

- 723 Three-dimensional visualization and virtual reality simulation role in hepatic surgery: Further research warranted

Ahmed F, Jahagirdar V, Gudapati S, Mouchli M

- 727** Signs and syndromes in acute appendicitis: A pathophysiologic approach

Yale SH, Tekiner H, Yale ES

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Francesco Saverio Mari, FACS, MD, PhD, Doctor, Lecturer, Postdoctoral Fellow, Research Fellow, Surgeon, Surgical Oncologist, Department of Medical and Surgical Sciences and Translational Medicine, School of Medicine and Psychology, University Sapienza of Rome, Rome 00060, Italy. frasavmari@yahoo.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (WJGS, *World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

INDEXING/ABSTRACTING

The WJGS is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJGS as 2.505; IF without journal self cites: 2.473; 5-year IF: 3.099; Journal Citation Indicator: 0.49; Ranking: 104 among 211 journals in surgery; Quartile category: Q2; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Rui-Rui Wu; Production Department Director: Xiang Li; Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

July 27, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Percutaneous aspiration and sclerotherapy of a giant simple hepatic cyst causing obstructive jaundice: A case report and review of literature

Xu-Xia He, Mei-Xing Sun, Ke Lv, Jian Cao, Sheng-Yu Zhang, Jing-Nan Li

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Ajiki T, Japan; Elshimi E, Egypt

Received: March 2, 2022

Peer-review started: March 2, 2022

First decision: April 25, 2022

Revised: April 30, 2022

Accepted: June 20, 2022

Article in press: June 20, 2022

Published online: July 27, 2022



Xu-Xia He, Mei-Xing Sun, Sheng-Yu Zhang, Jing-Nan Li, Department of Gastroenterology, Peking Union Medical College Hospital, Beijing 10000, China

Ke Lv, Department of Ultrasound, Peking Union Medical College Hospital, Beijing 10000, China

Jian Cao, Department of Radiology, Peking Union Medical College Hospital, Beijing 10000, China

Corresponding author: Sheng-Yu Zhang, MD, Doctor, Department of Gastroenterology, Peking Union Medical College Hospital, No. 9 Dongdan Santiao, Dongcheng District, Beijing 10000, China. hexvxia@163.com

Abstract

BACKGROUND

Giant simple hepatic cysts causing intrahepatic duct dilatation and obstructive jaundice are uncommon. A variety of measures with different clinical efficacies and invasiveness have been developed. Nonsurgical management, such as percutaneous aspiration and sclerotherapy, is often applied.

CASE SUMMARY

The case is a 39-year-old female with a 5-mo history of cutaneous and scleral icterus, loss of appetite, and dark urine. Lab tests showed jaundice and liver function abnormalities. Imaging revealed a giant simple hepatic cyst obstructing the intrahepatic bile ducts. A combination of percutaneous catheter aspiration and lauromacrogol sclerotherapy was successfully performed and the effects were satisfactory with the size of cyst decreasing from 13.7 cm × 13.1 cm to 3.0 cm × 3.0 cm. Further literature review presented the challenges of managing giant simple hepatic cysts that cause obstructive jaundice and compared the safety and efficacy of a combination of percutaneous aspiration and lauromacrogol sclerotherapy with other management strategies.

CONCLUSION

Giant simple hepatic cysts can cause obstructive jaundice, and a combination of percutaneous catheter aspiration and sclerotherapy with lauromacrogol are suggested to treat such cases.

Key Words: Simple hepatic cyst; Obstructive jaundice; Aspiration; Sclerotherapy; Lauromacrogol; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Giant simple hepatic cysts causing obstructive jaundice are uncommon. Here we presented the challenges of managing giant simple hepatic cysts causing obstructive jaundice and compared the safety and efficacy of percutaneous aspiration and lauromacrogol sclerotherapy with other management strategies. The case is a 39-year-old female with jaundice and liver function abnormalities. Images revealed a giant simple hepatic cyst with obstruction of intrahepatic bile ducts. A combination of percutaneous catheter aspiration and lauromacrogol sclerotherapy was conducted successively, achieving satisfactory efficacy. Therefore, a combination of percutaneous aspiration and lauromacrogol sclerotherapy may be suggested to solve such cases.

Citation: He XX, Sun MX, Lv K, Cao J, Zhang SY, Li JN. Percutaneous aspiration and sclerotherapy of a giant simple hepatic cyst causing obstructive jaundice: A case report and review of literature. *World J Gastrointest Surg* 2022; 14(7): 706-713

URL: <https://www.wjgnet.com/1948-9366/full/v14/i7/706.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v14.i7.706>

INTRODUCTION

Hepatic cysts occur in 2.5%-18% of the population[1-3]. They generally include a cluster of diseases with heterogeneous pathogenesis and etiology, including simple hepatic cysts, infectious cysts, cystic neoplasms, biliary duct-related cysts and some congenital polycystic liver diseases[4]. Most simple cysts are asymptomatic and are incidentally identified during imaging examinations, including ultrasonography (US), computed tomography (CT) or magnetic resonance imaging[5,6]. Only 5%-16% of simple hepatic cysts become symptomatic due to mass effects, rupture, hemorrhaging, or infection[5,7,8]. They mainly present as abdominal pain, nausea, vomiting and occasional jaundice[9,10].

The management of simple hepatic cysts widely differs according to clinical manifestations, imaging features, and, sometimes, patient preference. A watch-and-see strategy is acceptable for asymptomatic simple cysts, whereas interventions are required if cysts cause severe symptoms or complications. Various treatment methods with different clinical efficacies and levels of invasiveness have been developed. For nonsurgical management, percutaneous aspiration, sclerotherapy, and internal drainage are often used[8,9]. Surgical treatment mainly includes unroofing, cyst fenestration, hepatectomy, and open or laparoscopic liver transplantation[11]. Treatment selection depends on cyst location, size, surroundings and other factors[12,13].

Here, we report a case of a giant simple hepatic cyst in the hepatic hilum causing intrahepatic duct dilatation and obstructive jaundice. A combination of percutaneous aspiration and lauromacrogol sclerotherapy was performed and achieved satisfactory effects. The related literature was reviewed to better understand management in similar patients.

CASE PRESENTATION

Chief complaints

A 39-year-old female was admitted for cutaneous and scleral icterus, loss of appetite, and dark urine for 5 mo.

History of present illness

A 39-year-old female was admitted for cutaneous and scleral icterus, loss of appetite, and dark urine for 5 mo.

History of past illness

The patient used to be in good health and had no previous medical history.

Personal and family history

The patient's personal habits, customs, and family history were unremarkable.

Physical examination

Physical examination revealed moderate jaundice without abdominal tenderness, hepatomegaly, or Murphy's sign.

Laboratory examinations

Lab tests showed jaundice [total bilirubin (TBil) level was 149.8 $\mu\text{mol/L}$, and direct bilirubin (DBil) level was 118.7 $\mu\text{mol/L}$], liver function abnormalities (liver function test levels included the following: Alanine transaminase (ALT) was 175 U/L, aspartate aminotransferase (AST) was 130 U/L, gamma-glutamyl transpeptidase was 454 U/L, alkaline phosphatase was 314 U/L) and moderate anemia [the hemoglobin (HGB) level was 75 g/L]. Tumor markers were unremarkable except for a slightly elevated carcinoembryonic antigen (CEA) level of 6.1 ng/mL (normal range: 0-5). Antibodies for hepatitis virus, primary biliary cholangitis and autoimmune hepatitis were all within the normal limits.

Imaging examinations

The abdominal US and the endoscopic US showed an enlarged liver (3.7 cm below the xiphoid process) and an anechoic area (increasing from 11.2 cm \times 9.9 cm to 13.7 cm \times 13.1 cm in three months) with a clear boundary and no peripheral blood flow, and the intrahepatic bile duct of the left lateral segment was approximately 0.6 cm wide. Magnetic resonance cholangiopancreatography showed several hepatic cysts. The largest cyst was approximately 9.5 cm \times 11 cm in size, located in the hilum, and obstructed the intrahepatic bile ducts. Three-dimensional reconstruction of the biliary tract showed dilated intrahepatic bile ducts and compressed hepatic vessels and branches of the portal vein (Figure 1).

Notably, esophagogastroduodenoscopy and colonoscopy were performed and excluded gastrointestinal neoplastic diseases.

FINAL DIAGNOSIS

A giant simple hepatic cyst complicated with obstructive jaundice was the diagnosis.

TREATMENT

We successfully performed a combination of percutaneous catheter aspiration and sclerotherapy with lauromacrogol. During percutaneous catheter aspiration under the guidance of US, the giant cyst was punctured with an 18-gauge pig-tail catheter. Postoperative drainage was favorable, and a total of 800 milliliters of clear yellow fluid was drained; bilirubin levels, tumor markers (such as CEA level) and cytology tests were unremarkable. Jaundice (TBil was 66.4 $\mu\text{mol/L}$, DBil was 51.2 $\mu\text{mol/L}$) and liver function anomalies (ALT was 90 U/L, AST was 59 U/L) were significantly relieved soon after drainage.


Then, two sessions of sclerotherapy (lauromacrogol) of the hepatic cyst were performed (30 mL and 20 mL lauromacrogol mixed with triple amounts of air) at one week. Of note, before sclerotherapy, the communications of the cyst with the surrounding bile ducts were ruled out by injecting a diluted contrast medium into the cyst cavity. After sclerotherapy, a small amount of cyst fluid was drained, and the tube was removed. The patient was generally in good condition. He was discharged and experienced further improvement in his liver function (ALT level was 38 U/L, TBil level was 34.9 $\mu\text{mol/L}$, and DBil level was 33.5 $\mu\text{mol/L}$; Figure 2).

OUTCOME AND FOLLOW-UP

During follow-up, the patient reported continued resolution of his symptoms. Three months after treatment, the size of the liver cyst decreased to 6.5 cm \times 5.6 cm, and liver function returned to normal limits. Fourteen months after treatment, the size of the cyst had decreased to 3.0 cm \times 3.0 cm on US.


DISCUSSION

Most simple liver cysts are asymptomatic and stable in size and structure, which allows for observation. However, some of these tumors gradually grow and eventually cause symptoms due to large size, rupture, hemorrhaging, infection, or neoplasm in rare cases[8,14]. Symptoms, including abdominal discomfort or pain, nausea, vomiting, jaundice, early satiety, and even dyspnea[9,10], are largely related to cyst size and location and are more often attributed to larger cysts and right-sided cysts[9,15]. In a recent review, abdominal pain was reported to be the most common symptom of simple hepatic cysts and was reported by 60% (456 of 764) of the patients[16].



DOI: 10.4240/wjgs.v14.i7.706 Copyright ©The Author(s) 2022.

Figure 1 Initial radiographic assessments of the cyst. A: Magnetic resonance cholangiopancreatography displayed a giant hepatic cyst approximately 9.5 cm × 11 cm in size located in the hilum and obstructed intrahepatic bile ducts; B-D: Abdominal contrast-enhanced computed tomography and three-dimensional reconstruction of the biliary tract displayed a hepatic cyst 11.0 cm × 10.6 cm × 12.7 cm in size with compressed hepatic arteries and veins and dilated intrahepatic bile ducts. Multiple portal vein branches were also involved, and tortuous vessels were seen around the gastric fundus and the spleen.



DOI: 10.4240/wjgs.v14.i7.706 Copyright ©The Author(s) 2022.

Figure 2 Changes in hepatic indicators after treatment. The patient successfully underwent percutaneous catheter aspiration on December 17, 2020, and then two lauromacrogol sclerotherapies on December 21 and December 27. After treatment, liver function continuously dropped to normal limits. ALT: Alanine transaminase; HGB: Hemoglobin; TBil: Total bilirubin; DBil: Direct bilirubin.

Obstructive jaundice caused by solitary simple liver cysts is quite rare. A total of 17 cases of simple or benign liver cysts accompanied by obstructive jaundice were reviewed (Table 1)[17-33]. The average age of the patients was 65.2 years old, with a 7:10 female to male ratio. These cysts tended to be large (greater than 10 cm) and centrally located when compression of the main intrahepatic duct or even the hepatic hilum was present. Treatment for these patients varied from aspiration to resection. In recent years, a combination of drainage, sclerosing agent injection, and deroofing seem to be the most common treatment methods. Choledochoscopy was also proven to effectively treat these patients[33]. In our

Table 1 Published cases with simple or benign hepatic cysts causing obstructive jaundice

| No. | Ref. | Age/sex | Cyst (cm) | Location (segments) | Total bilirubin (mg/dL) | Treatment | Prognosis | Follow-up period |
|-----|-------------------------------------|---------|-----------|---------------------|-------------------------|--|-----------|------------------|
| 1 | Caravati <i>et al</i> [17], 1950 | 33/M | NA | IV, V | NA | Aspiration + marsupialization | Improved | 7 mo |
| 2 | Hudson[18], 1963 | 55/F | 25 | III, IV, V | 14 | Cystenterostomy | Improved | 1 mo |
| 3 | Dardik <i>et al</i> [19], 1964 | 69/F | 15 | V | 9 | Cystectomy | Improved | 1 mo |
| 4 | Sacks <i>et al</i> [20], 1967 | 81/M | 20 | IV | 19 | Aspiration | Improved | 2 mo |
| 5 | Santman <i>et al</i> [21], 1977 | 61/M | 15 | IV | 29 | Partial resection | Improved | NA |
| 6 | Machell <i>et al</i> [22], 1978 | 67/F | NA | III, IV, V | NA | Drainage + transhepatic T-tube | Improved | 7 mo |
| 7 | Morin <i>et al</i> [23], 1980 | 80/M | 17 | IV, V | 15 | Aspiration only | Improved | 10 mo |
| 8 | Fernandez <i>et al</i> [24], 1984 | 61/F | 30 | III, IV, V | 22 | Partial resection | Improved | 24 mo |
| 9 | Clinkscales <i>et al</i> [25], 1985 | 80/M | 8 | IV | 8 | Aspiration only | Improved | 1 mo |
| 10 | Cappel <i>et al</i> [26], 1988 | 44/F | 12 | IV, V | 5 | Aspiration | Improved | 3 mo |
| 11 | Spivey <i>et al</i> [27], 1990 | 73/M | 11 | IV, V | 10 | Drainage + deroofing | Improved | NA |
| 12 | Terada <i>et al</i> [28], 1993 | 71/F | 12 | III, IV, V | 9 | Drainage + cystectomy | Improved | 1 mo |
| 13 | Yoshihara <i>et al</i> [29], 1996 | 88/M | 16 | IV, V | 8 | Drainage + minocycline injection | Improved | 9 mo |
| 14 | Kanai <i>et al</i> [30], 1999 | 71/M | 15 | IV, V, VIII | 5 | Drainage + deroofing | Improved | 15 mo |
| 15 | Ishikawa <i>et al</i> [31], 2002 | 70/M | 18 | IV, V, VIII | 9 | Drainage + minocycline injection | Improved | 20 mo |
| 16 | Ogawa <i>et al</i> [32], 2004 | 64/M | 9 | NA | NA | Drainage + minocycline injection | Improved | NA |
| 17 | Zhang <i>et al</i> [33], 2018 | 41/F | 5 | IV | 24 | Choledochoscopic high-frequency needle-knife electrotomy | Improved | 36 mo |

NA: Not available.

patients, the giant liver cyst caused obstructive jaundice and dilatation of the intrahepatic bile duct of the left lateral segment of the liver, which largely accounted for the patient's symptoms.

Aspiration is generally associated with high recurrence rates[34]. In recent years, percutaneous aspiration combined with sclerotherapy has been widely used as a minimally invasive procedure for simple hepatic cysts with satisfactory results[35-39]. During percutaneous aspiration and sclerotherapy, US- or CT-guided aspiration and drainage are combined with the injection of a sclerosing agent[7,40,41]. Sclerosing agents with good efficacy include ethanol, iophendylate, tetracycline chloride, doxycycline, minocycline chloride, and hypertonic saline solution[42].

While liquid sclerosing agents may mix with cyst contents and reduce sclerosing effects, foam sclerotherapy was initially used for vascular malformations and has evolved as an alternative for treating simple hepatic cysts[43]. The agents in a foam vehicle can completely destroy the intimal barrier after 2 min of exposure, causing endothelial edema, exfoliation from the tunica media, and thrombogenesis in the tunica media in 30 min[44]. Sclerotherapy using lauromacrogol foam is rarely reported for treating hepatic cysts. In one case report, laparoscopic lauromacrogol sclerotherapy surgery was reported to be safe and effective in patients with IVa, VII and VIII segment simple hepatic cysts, but more studies are needed to confirm their conclusion[45]. Our case report is the first to combine percutaneous aspiration with sclerotherapy using lauromacrogol in treating a giant simple hepatic cyst, thus proving the safety and efficacy of the therapy. Single or multiple sessions of percutaneous aspiration and sclerotherapy for persistent or recurrent symptoms are adaptable based on cyst features, efficacy and doctor or patient

preference[7]. In our patients, sclerotherapy with lauromacrogol was planned and administered twice to achieve a better sclerosing effect.

Surgical treatment of simple hepatic cysts, such as open or laparoscopic cyst deroofing or hepatectomy, can be effective but may contribute to recurrence and complications[46,47]. Generally, percutaneous aspiration combined with sclerotherapy and laparoscopic deroofing is reasonable for most symptomatic simple hepatic cysts. A systematic review showed that the outcome of percutaneous aspiration and sclerotherapy was excellent, with symptoms that persisted in less than 4% of patients, and both complication and recurrence rates were < 1%[16]. Major complications were reported in 2/265 (0.8%), 6/348 (1.7%) and 3/123 (2.4%), and cyst recurrence rates were 0.0%, 5.6% and 7.7% in patients treated with percutaneous aspiration and sclerotherapy and laparoscopic and open surgery, respectively[16]. Other studies on the advantage of percutaneous aspiration and sclerotherapy compared to surgical techniques reported similar results[13]. These results supported the safety and efficacy of percutaneous aspiration and sclerotherapy in treating symptomatic simple hepatic cysts prior to surgical procedures. Our patient's outcome suggested that percutaneous aspiration and sclerotherapy could effectively treat simple giant hepatic cysts. Studies concerning cost, hospitalization time, and quality of life are needed to further compare these measures.

CONCLUSION

Giant simple hepatic cysts can obstruct the intrahepatic bile ducts and cause obstructive jaundice. A combination of percutaneous catheter aspiration and sclerotherapy using lauromacrogol can achieve satisfactory results without evident complications compared to surgical interventions.

FOOTNOTES

Author contributions: He XX and Sun MX compiled all relevant information concerning that case and did the literature research; He XX did the drafting and review of the manuscript; Lv K and Cao J did the radiological analysis; Zhang SY did the study concept and design; Zhang SY and Li JN evaluated the whole treatment of the patient and supervised the study; all authors had reviewed and approved the final version of this manuscript.

Supported by the Peking Union Medical College Education Reform Program, No. 2019zlgc0116; the 13th Five-Year Plan for National Key R&D Program of China, No. 2018YFC1705402; National Natural Science Foundation of China, No. 81770559 and No. 81370500; and Medical and Health Technology Innovation Project of the Chinese Academy of Medical Sciences, No. 2020-I2M-2-013.

Informed consent statement: Informed consent was obtained from the patient for publication of this case report.

Conflict-of-interest statement: The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Sheng-Yu Zhang 0000-0002-9218-6409.

S-Editor: Yan JP

L-Editor: A

P-Editor: Yan JP

REFERENCES

- 1 Caremani M, Vincenti A, Benci A, Sassoli S, Tacconi D. Ecographic epidemiology of non-parasitic hepatic cysts. *J Clin Ultrasound* 1993; **21**: 115-118 [PMID: 8381130 DOI: 10.1002/jcu.1870210207]
- 2 Carrim ZI, Murchison JT. The prevalence of simple renal and hepatic cysts detected by spiral computed tomography. *Clin*

- Radiol* 2003; **58**: 626-629 [PMID: [12887956](#) DOI: [10.1016/s0009-9260\(03\)00165-x](#)]
- 3 **European Association for the Study of the Liver (EASL)**. EASL Clinical Practice Guidelines on the management of benign liver tumours. *J Hepatol* 2016; **65**: 386-398 [PMID: [27085809](#) DOI: [10.1016/j.jhep.2016.04.001](#)]
- 4 **Taylor BR**, Langer B. Current surgical management of hepatic cyst disease. *Adv Surg* 1997; **31**: 127-148 [PMID: [9408491](#)]
- 5 **Cowles RA**, Mulholland MW. Solitary hepatic cysts. *J Am Coll Surg* 2000; **191**: 311-321 [PMID: [10989905](#) DOI: [10.1016/s1072-7515\(00\)00345-8](#)]
- 6 **Lantinga MA**, Gevers TJ, Drenth JP. Evaluation of hepatic cystic lesions. *World J Gastroenterol* 2013; **19**: 3543-3554 [PMID: [23801855](#) DOI: [10.3748/wjg.v19.i23.3543](#)]
- 7 **Karam AR**, Connolly C, Fulwadhva U, Hussain S. Alcohol sclerosis of a giant liver cyst following failed deroofings. *J Radiol Case Rep* 2011; **5**: 19-22 [PMID: [22470778](#) DOI: [10.3941/jrcr.v5i2.634](#)]
- 8 **Nisenbaum HL**, Rowling SE. Ultrasound of focal hepatic lesions. *Semin Roentgenol* 1995; **30**: 324-346 [PMID: [8539643](#) DOI: [10.1016/s0037-198x\(05\)80021-5](#)]
- 9 **Lai EC**, Wong J. Symptomatic nonparasitic cysts of the liver. *World J Surg* 1990; **14**: 452-456 [PMID: [2382451](#) DOI: [10.1007/BF01658666](#)]
- 10 **Karavias DD**, Tsamandas AC, Payatakes AH, Solomou E, Salakou S, Felekouras ES, Tepetes KN. Simple (non-parasitic) liver cysts: clinical presentation and outcome. *Hepatogastroenterology* 2000; **47**: 1439-1443 [PMID: [11100371](#)]
- 11 **Gomez A**, Wisneski AD, Luu HY, Hirose K, Roberts JP, Hirose R, Freise CE, Nakakura EK, Corvera CU. Contemporary Management of Hepatic Cyst Disease: Techniques and Outcomes at a Tertiary Hepatobiliary Center. *J Gastrointest Surg* 2021; **25**: 77-84 [PMID: [33083858](#) DOI: [10.1007/s11605-020-04821-1](#)]
- 12 **Macutkiewicz C**, Plastow R, Chrispijn M, Filobos R, Ammori BA, Sherlock DJ, Drenth JP, O'Reilly DA. Complications arising in simple and polycystic liver cysts. *World J Hepatol* 2012; **4**: 406-411 [PMID: [23355921](#) DOI: [10.4254/wjh.v4.i12.406](#)]
- 13 **Erdogan D**, van Delden OM, Rauws EA, Busch OR, Lameris JS, Gouma DJ, van Gulik TM. Results of percutaneous sclerotherapy and surgical treatment in patients with symptomatic simple liver cysts and polycystic liver disease. *World J Gastroenterol* 2007; **13**: 3095-3100 [PMID: [17589926](#) DOI: [10.3748/wjg.v13.i22.3095](#)]
- 14 **Monteagudo M**, Vidal G, Moreno M, Bella R, Diaz MJ, Colomer O, Santesmasses A. Squamous cell carcinoma and infection in a solitary hepatic cyst. *Eur J Gastroenterol Hepatol* 1998; **10**: 1051-1053 [PMID: [9895053](#) DOI: [10.1097/00042737-199812000-00012](#)]
- 15 **Sanchez H**, Gagner M, Rossi RL, Jenkins RL, Lewis WD, Munson JL, Braasch JW. Surgical management of nonparasitic cystic liver disease. *Am J Surg* 1991; **161**: 113-8; discussion 118 [PMID: [1987844](#) DOI: [10.1016/0002-9610\(91\)90370-s](#)]
- 16 **Furumaya A**, van Rosmalen BV, de Graeff JJ, Haring MPD, de Meijer VE, van Gulik TM, Verheij J, Besselink MG, van Delden OM, Erdmann JJ; Dutch Benign Liver Tumor Group. Systematic review on percutaneous aspiration and sclerotherapy versus surgery in symptomatic simple hepatic cysts. *HPB (Oxford)* 2021; **23**: 11-24 [PMID: [32830070](#) DOI: [10.1016/j.hpb.2020.07.005](#)]
- 17 **Caravati CM**, Watts TD. Benign solitary non-parasitic cyst of the liver. *Gastroenterology* 1950; **14**: 317-320 [PMID: [15404887](#)]
- 18 **Hudson EK**. Obstructive jaundice from solitary hepatic cyst. *Am J Gastroenterol* 1963; **39**: 161-164 [PMID: [13955453](#)]
- 19 **Dardik H**, Glotzer P, Silver C. Congenital Hepatic Cyst Causing Jaundice: Report Of A Case And Analogies With Respiratory Malformations. *Ann Surg* 1964; **159**: 585-592 [PMID: [14138206](#) DOI: [10.1097/00000658-196404000-00018](#)]
- 20 **Sacks HJ**, Robbins LS. Fistulization of a solitary hepatic cyst. *JAMA* 1967; **200**: 415-417 [PMID: [6071519](#)]
- 21 **Santman FW**, Thijs LG, Van Der Veen EA, Den Otter G, Blok P. Intermittent jaundice: a rare complication of a solitary nonparasitic liver cyst. *Gastroenterology* 1977; **72**: 325-328 [PMID: [830585](#)]
- 22 **Machell RJ**, Calne RY. Solitary non-parasitic hepatic cyst presenting with jaundice. *Br J Radiol* 1978; **51**: 631-632 [PMID: [678759](#) DOI: [10.1259/0007-1285-51-608-631](#)]
- 23 **Morin ME**, Baker DA, Vanagunas A, Tan A, Sue HK. Solitary nonparasitic hepatic cyst causing obstructive jaundice. *Am J Gastroenterol* 1980; **73**: 434-436 [PMID: [7416140](#)]
- 24 **Fernandez M**, Cacioppo JC, Davis RP, Nora PF. Management of solitary nonparasitic liver cyst. *Am Surg* 1984; **50**: 205-208 [PMID: [6712014](#)]
- 25 **Clinkscales NB**, Trigg LP, Poklepovic J. Obstructive jaundice secondary to benign hepatic cyst. *Radiology* 1985; **154**: 643-644 [PMID: [3969466](#) DOI: [10.1148/radiology.154.3.3969466](#)]
- 26 **Cappell MS**. Obstructive jaundice from benign, nonparasitic hepatic cysts: identification of risk factors and percutaneous aspiration for diagnosis and treatment. *Am J Gastroenterol* 1988; **83**: 93-96 [PMID: [3276152](#)]
- 27 **Spivey JR**, Garrido JA, Reddy KR, Jeffers LJ, Schiff ER. ERCP documentation of obstructive jaundice caused by a solitary, centrally located, benign hepatic cyst. *Gastrointest Endosc* 1990; **36**: 521-523 [PMID: [2227332](#) DOI: [10.1016/s0016-5107\(90\)71133-9](#)]
- 28 **Terada N**, Shimizu T, Imai Y, Kobayashi T, Terashima M, Furukawa K, Kumazawa S, Kiyosawa K. Benign, non-parasitic hepatic cyst causing obstructive jaundice. *Intern Med* 1993; **32**: 857-860 [PMID: [8012087](#) DOI: [10.2169/internalmedicine.32.857](#)]
- 29 **Yoshihara K**, Yamashiro S, Koizumi S, Matsuo Y, Shigeru J, Kanegae S, Oda Y. Obstructive jaundice caused by non-parasitic hepatic cyst treated with percutaneous drainage and instillation of minocycline hydrochloride as a sclerosing agent. *Intern Med* 1996; **35**: 373-375 [PMID: [8797049](#) DOI: [10.2169/internalmedicine.35.373](#)]
- 30 **Kanai T**, Kenmochi T, Takabayashi T, Hangai N, Kawano Y, Suwa T, Yonekawa H, Miyazawa N. Obstructive jaundice caused by a huge liver cyst riding on the hilum: report of a case. *Surg Today* 1999; **29**: 791-794 [PMID: [10483760](#) DOI: [10.1007/BF02482330](#)]
- 31 **Ishikawa H**, Uchida S, Yokokura Y, Iwasaki Y, Horiuchi H, Hiraki M, Kinoshita H, Shirouzu K. Nonparasitic solitary huge liver cysts causing intracystic hemorrhage or obstructive jaundice. *J Hepatobiliary Pancreat Surg* 2002; **9**: 764-768 [PMID: [12658414](#) DOI: [10.1007/s005340200107](#)]
- 32 **Ogawa M**, Kubo S, Uenishi T, Hirohashi K, Tanaka H, Shuto T, Yamamoto T, Takemura S. Nonoperative management of obstructive jaundice caused by a benign hepatic cyst. *Osaka City Med J* 2004; **50**: 95-99 [PMID: [15819304](#)]

- 33 **Zhang C**, Ma YF, Yang YL. Jaundice caused by protrusion of a hepatic cyst into common bile duct that was resolved by choledochoscopic needle-knife electrotomy: a case report. *BMC Gastroenterol* 2018; **18**: 90 [PMID: 29921238 DOI: 10.1186/s12876-018-0815-x]
- 34 **Koperna T**, Vogl S, Satzinger U, Schulz F. Nonparasitic cysts of the liver: results and options of surgical treatment. *World J Surg* 1997; **21**: 850-4; discussion 854 [PMID: 9327677 DOI: 10.1007/s002689900316]
- 35 **Vardakostas D**, Damaskos C, Garmpis N, Antoniou EA, Kontzoglou K, Kouraklis G, Dimitroulis D. Minimally invasive management of hepatic cysts: indications and complications. *Eur Rev Med Pharmacol Sci* 2018; **22**: 1387-1396 [PMID: 29565498 DOI: 10.26355/eurrev_201803_14484]
- 36 **Wijnands TF**, Görtjes AP, Gevers TJ, Jenniskens SF, Kool LJ, Potthoff A, Ronot M, Drenth JP. Efficacy and Safety of Aspiration Sclerotherapy of Simple Hepatic Cysts: A Systematic Review. *AJR Am J Roentgenol* 2017; **208**: 201-207 [PMID: 27824501 DOI: 10.2214/AJR.16.16130]
- 37 **Wijnands TF**, Ronot M, Gevers TJ, Benzimra J, Kool LJ, Vilgrain V, Drenth JP. Predictors of treatment response following aspiration sclerotherapy of hepatic cysts: an international pooled analysis of individual patient data. *Eur Radiol* 2017; **27**: 741-748 [PMID: 27180184 DOI: 10.1007/s00330-016-4363-x]
- 38 **Yang CF**, Liang HL, Pan HB, Lin YH, Mok KT, Lo GH, Lai KH. Single-session prolonged alcohol-retention sclerotherapy for large hepatic cysts. *AJR Am J Roentgenol* 2006; **187**: 940-943 [PMID: 16985138 DOI: 10.2214/AJR.05.0621]
- 39 **Yu JH**, Du Y, Li Y, Yang HF, Xu XX, Zheng HJ, Li B. Effectiveness of CT-guided sclerotherapy with estimated ethanol concentration for treatment of symptomatic simple hepatic cysts. *Clin Res Hepatol Gastroenterol* 2014; **38**: 190-194 [PMID: 24210773 DOI: 10.1016/j.clinre.2013.09.008]
- 40 **Debs T**, Kassir R, Reccia I, Elias B, Ben Amor I, Iannelli A, Gugenheim J, Johann M. Technical challenges in treating recurrent non-parasitic hepatic cysts. *Int J Surg* 2016; **25**: 44-48 [PMID: 26654896 DOI: 10.1016/j.ijssu.2015.11.051]
- 41 **Wijnands TF**, Lantinga MA, Drenth JP. Hepatic cyst infection following aspiration sclerotherapy: a case series. *J Gastrointest Liver Dis* 2014; **23**: 441-444 [PMID: 25532005 DOI: 10.15403/jgld.2014.1121.234.hcy]
- 42 **Blonski WC**, Campbell MS, Faust T, Metz DC. Successful aspiration and ethanol sclerosis of a large, symptomatic, simple liver cyst: case presentation and review of the literature. *World J Gastroenterol* 2006; **12**: 2949-2954 [PMID: 16718826 DOI: 10.3748/wjg.v12.i18.2949]
- 43 **Itou C**, Koizumi J, Hashimoto T, Myojin K, Kagawa T, Mine T, Imai Y. Foam sclerotherapy for a symptomatic hepatic cyst: a preliminary report. *Cardiovasc Intervent Radiol* 2014; **37**: 800-804 [PMID: 24170168 DOI: 10.1007/s00270-013-0761-5]
- 44 **Orsini C**, Brotto M. Immediate pathologic effects on the vein wall of foam sclerotherapy. *Dermatol Surg* 2007; **33**: 1250-1254 [PMID: 17903159 DOI: 10.1111/j.1524-4725.2007.33261.x]
- 45 **Xu S**, Rao M, Pu Y, Zhou J, Zhang Y. The efficacy of laparoscopic lauromacrogol sclerotherapy in the treatment of simple hepatic cysts located in posterior segments: a refined surgical approach. *Ann Palliat Med* 2020; **9**: 3462-3471 [PMID: 33065797 DOI: 10.21037/apm-20-1723]
- 46 **Moorthy K**, Mihssin N, Houghton PW. The management of simple hepatic cysts: sclerotherapy or laparoscopic fenestration. *Ann R Coll Surg Engl* 2001; **83**: 409-414 [PMID: 11777137]
- 47 **Katkhouda N**, Mavor E. Laparoscopic management of benign liver disease. *Surg Clin North Am* 2000; **80**: 1203-1211 [PMID: 10987031 DOI: 10.1016/s0039-6109(05)70220-6]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

