

ANSWERING REVIEWERS

March 12, 2014

Dear Editor,

Please find enclosed the edited manuscript in Word format.

Title: Advancing Frontiers in Anaesthesiology with Laparoscopy

Author: Jayashree Sood

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 7716

The manuscript has been improved according to the suggestions of reviewers:

1 Format has been updated

2 Revision has been made according to the suggestions of the reviewer

Reviewed by 02549484

- 1) The pneumoperitoneum should always be established slowly and gradually at the rate of 1L/min with the Veress needle till 1 litre has been insufflated. Then the rate is increased to 2L/min till an IAP of 15 mm Hg is achieved.
- 2) Usually the IAP does not change when the instruments are changed, since the instruments and trocars are leak proof.
- 3) The volume of CO₂ required for achieving an adequate IAP (13-15 mm Hg) varies in different patients. This does not have any limitation since it is the IAP which is the most important factor and it has to be reached for successful surgery.

Reviewed by 00506142

- 1) Future direction of NOTES has been added
- 2) Pregnancy and patients with limited cardiopulmonary function has been added
- 3) The wrongly inserted paragraph has been corrected

Reviewed by 02549363

- 1) The article has been modified with a clearer aim

3 References and typesetting were corrected

Thank you for publishing my manuscript in the *World Journal of Gastroenterology*.

Sincerely yours,



Dr. Jayashree Sood

Chairperson

Department of Anaesthesiology, Pain & Perioperative Medicine

Sir Ganga Ram Hospital, New Delhi 110060, INDIA

Mobile: +91-98112 94608

Phone: +91-11-42252502

Email: jayashreesood@hotmail.com