

World Journal of *Clinical Cases*

World J Clin Cases 2022 August 16; 10(23): 8057-8431



Contents

Thrice Monthly Volume 10 Number 23 August 16, 2022

OPINION REVIEW

- 8057** Invasive intervention timing for infected necrotizing pancreatitis: Late invasive intervention is not late for collection

Xiao NJ, Cui TT, Liu F, Li W

- 8063** Clinical utility of left atrial strain in predicting atrial fibrillation recurrence after catheter ablation: An up-to-date review

Yu ZX, Yang W, Yin WS, Peng KX, Pan YL, Chen WW, Du BB, He YQ, Yang P

MINIREVIEWS

- 8076** Gut microbiota and COVID-19: An intriguing pediatric perspective

Valentino MS, Esposito C, Colosimo S, Caprio AM, Puzone S, Guarino S, Marzuillo P, Miraglia del Giudice E, Di Sessa A

- 8088** Beta receptor blocker therapy for the elderly in the COVID-19 era

Santillo E, Migale M

ORIGINAL ARTICLE

Retrospective Cohort Study

- 8097** Nonselective beta-blocker use is associated with increased hepatic encephalopathy-related readmissions in cirrhosis

Fallahzadeh MA, Asrani SK, Tapper EB, Saracino G, Rahimi RS

Retrospective Study

- 8107** Different squatting positions after total knee arthroplasty: A retrospective study

Li TJ, Sun JY, Du YQ, Shen JM, Zhang BH, Zhou YG

- 8115** Outcomes of seromuscular bladder augmentation compared with standard bladder augmentation in the treatment of children with neurogenic bladder

Sun XG, Li YX, Ji LF, Xu JL, Chen WX, Wang RY

- 8124** Distinctive clinical features of spontaneous pneumoperitoneum in neonates: A retrospective analysis

Kim SH, Cho YH, Kim HY

- 8133** Cognitive training for elderly patients with early Alzheimer's disease in the Qinghai-Tibet Plateau: A pilot study

Wang XH, Luo MQ

- 8141** Diagnostic value of elevated serum carbohydrate antigen 125 level in sarcoidosis

Zhang Q, Jing XY, Yang XY, Xu ZJ

- 8152** Evaluation of progressive early rehabilitation training mode in intensive care unit patients with mechanical ventilation

Qie XJ, Liu ZH, Guo LM

- 8161** Comparison of demographic features and laboratory parameters between COVID-19 deceased patients and surviving severe and critically ill cases

Wang L, Gao Y, Zhang ZJ, Pan CK, Wang Y, Zhu YC, Qi YP, Xie FJ, Du X, Li NN, Chen PF, Yue CS, Wu JH, Wang XT, Tang YJ, Lai QQ, Kang K

Clinical Trials Study

- 8170** Role of H₂receptor blocker famotidine over the clinical recovery of COVID-19 patients: A randomized controlled trial

Mohiuddin Chowdhury ATM, Kamal A, Abbas MKU, Karim MR, Ali MA, Talukder S, Hamidullah Mehedi H, Hassan H, Shahin AH, Li Y, He S

Observational Study

- 8186** Short-term prognostic factors for hepatitis B virus-related acute-on-chronic liver failure

Ye QX, Huang JF, Xu ZJ, Yan YY, Yan Y, Liu LG

- 8196** Three-dimensional psychological guidance combined with evidence-based health intervention in patients with liver abscess treated with ultrasound

Shan YN, Yu Y, Zhao YH, Tang LL, Chen XM

- 8205** Role of serum β 2-microglobulin, glycosylated hemoglobin, and vascular endothelial growth factor levels in diabetic nephropathy

Yang B, Zhao XH, Ma GB

SYSTEMATIC REVIEWS

- 8212** Gallbladder neuroendocrine carcinoma diagnosis, treatment and prognosis based on the SEER database: A literature review

Cai XC, Wu SD

CASE REPORT

- 8224** Sepsis complicated with secondary hemophagocytic syndrome induced by giant gouty tophi rupture: A case report

Lai B, Pang ZH

- 8232** Spontaneous remission of autoimmune pancreatitis: Four case reports

Zhang BB, Huo JW, Yang ZH, Wang ZC, Jin EH

- 8242** Epstein-Barr-virus-associated hepatitis with aplastic anemia: A case report

Zhang WJ, Wu LQ, Wang J, Lin SY, Wang B

- 8249** Aspiration as the first-choice procedure for airway management in an infant with large epiglottic cysts: A case report

Zheng JQ, Du L, Zhang WY

- 8255** Sequential multidisciplinary minimally invasive therapeutic strategy for heart failure caused by four diseases: A case report
Zhao CZ, Yan Y, Cui Y, Zhu N, Ding XY
- 8262** Primary ascending colon cancer accompanying skip metastases in left shoulder skin and left neck lymph node: A case report
Zhou JC, Wang JJ, Liu T, Tong Q, Fang YJ, Wu ZQ, Hong Q
- 8271** Clinical and genetic study of ataxia with vitamin E deficiency: A case report
Zhang LW, Liu B, Peng DT
- 8277** Complete resection of large-cell neuroendocrine and hepatocellular carcinoma of the liver: A case report
Noh BG, Seo HI, Park YM, Kim S, Hong SB, Lee SJ
- 8284** Immunotherapy combined with antiangiogenic agents in patients with advanced malignant pleural mesothelioma: A case report
Xuan TT, Li GY, Meng SB, Wang ZM, Qu LL
- 8291** Bladder malacoplakia: A case report
Wang HK, Hang G, Wang YY, Wen Q, Chen B
- 8298** Delayed inflammatory response evoked in nasal alloplastic implants after COVID-19 vaccination: A case report
Seo MG, Choi EK, Chung KJ
- 8304** Phosphoglyceride crystal deposition disease requiring differential diagnosis from malignant tumors and confirmed by Raman spectroscopy: A case report
Ohkura Y, Uruga H, Shiiba M, Ito S, Shimoyama H, Ishihara M, Ueno M, Udagawa H
- 8312** Vulvovaginal myeloid sarcoma with massive pelvic floor infiltration: A case report and review of literature
Wang JX, Zhang H, Ning G, Bao L
- 8323** Femoral neck stress fracture and medial tibial stress syndrome following high intensity interval training: A case report and review of literature
Tan DS, Cheung FM, Ng D, Cheung TLA
- 8330** Periosteal chondroma of the rib: A case report
Gao Y, Wang JG, Liu H, Gao CP
- 8336** Papillary thyroid carcinoma occurring with undifferentiated pleomorphic sarcoma: A case report
Lee YL, Cheng YQ, Zhu CF, Huo HZ
- 8344** Laparoscopic treatment of bilateral duplex kidney and ectopic ureter: A case report
Wang SB, Wan L, Wang Y, Yi ZJ, Xiao C, Cao JZ, Liu XY, Tang RP, Luo Y
- 8352** Incontinentia pigmenti with intracranial arachnoid cyst: A case report
Li WC, Li ML, Ding JW, Wang L, Wang SR, Wang YY, Xiao LF, Sun T

- 8360** Relapsing polychondritis causing breathlessness: Two case reports
Zhai SY, Zhang YH, Guo RY, Hao JW, Wen SX
- 8367** Endodontic management of a fused left maxillary second molar and two paramolars using cone beam computed tomography: A case report
Mei XH, Liu J, Wang W, Zhang QX, Hong T, Bai SZ, Cheng XG, Tian Y, Jiang WK
- 8375** Infant biliary cirrhosis secondary to a biliary inflammatory myofibroblastic tumor: A case report and review of literature
Huang Y, Shu SN, Zhou H, Liu LL, Fang F
- 8384** Metastatic low-grade endometrial stromal sarcoma with variable morphologies in the ovaries and mesentery: A case report
Yu HY, Jin YL
- 8392** Bronchogenic cysts with infection in the chest wall skin of a 64-year-old asymptomatic patient: A case report
Ma B, Fu KW, Xie XD, Cheng Y, Wang SQ
- 8400** Incidental accumulation of Technetium-99m pertechnetate in subacute cerebral infarction: A case report
Han YH, Jeong HJ, Kang HG, Lim ST
- 8406** Metal stent combined with ileus drainage tube for the treatment of delayed rectal perforation: A case report
Cheng SL, Xie L, Wu HW, Zhang XF, Lou LL, Shen HZ
- 8417** Using ketamine in a patient with a near-occlusion tracheal tumor undergoing tracheal resection and reconstruction: A case report
Xu XH, Gao H, Chen XM, Ma HB, Huang YG

LETTER TO THE EDITOR

- 8422** Reflections on the prevalence of human leukocyte antigen-B27 and human leukocyte antigen-B51 co-occurrence in patients with spondylarthritis
Gonçalves Júnior J, Sampaio-Barros PD, Shinjo SK
- 8425** Comment on "Disease exacerbation is common in inflammatory bowel disease patients treated with immune checkpoint inhibitors for malignancy"
Argyriou K, Kotsakis A
- 8428** Intranasal sufentanil combined with intranasal dexmedetomidine: A promising method for non-anesthesiologist sedation during endoscopic ultrasonography
Wang Y, Ge ZJ, Han C

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Peng Liang, MD, Associate Professor, Day Surgery Center, Department of Anesthesiology, West China Hospital of Sichuan University, Chengdu 610041, Sichuan Province, China. 39485572@qq.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yin; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

August 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Observational Study

Role of serum β 2-microglobulin, glycosylated hemoglobin, and vascular endothelial growth factor levels in diabetic nephropathy

Bing Yang, Xiao-Hong Zhao, Guo-Bin Ma

Specialty type: Endocrinology and metabolism

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Dascalu AM, Romania; Kilanowska A, Poland

Received: May 6, 2022

Peer-review started: May 6, 2022

First decision: June 7, 2022

Revised: June 18, 2022

Accepted: July 6, 2022

Article in press: July 6, 2022

Published online: August 16, 2022



Bing Yang, Xiao-Hong Zhao, Guo-Bin Ma, Department of Endocrinology and Metabolism, 3201 Hospital, Xi'an Jiaotong University Health Science Center, Hanzhong 723099, Shaanxi Province, China

Corresponding author: Bing Yang, MM, Associate Chief Physician, Department of Endocrinology and Metabolism, 3201 Hospital, Xi'an Jiaotong University Health Science Center, No. 783 Tianhan Avenue, Hanzhong 723099, Shaanxi Province, China.

ybing3201@163.com

Abstract

BACKGROUND

Diabetic nephropathy (DN) is a common complication of type 1 and type 2 diabetes that can lead to kidney damage and high blood pressure. Increasing evidence support the important roles of microproteins and cytokines, such as β 2-microglobulin (β 2-MG), glycosylated hemoglobin (HbA1c), and vascular endothelial growth factor (VEGF), in the pathogenesis of this disease. In this study, we identified novel therapeutic options for this disease.

AIM

To analyze the guiding significance of β 2-MG, HbA1c, and VEGF levels in patients with DN.

METHODS

A total of 107 patients with type 2 diabetes mellitus complicated with nephropathy and treated in our hospital from May 2018 to February 2021 were included in the study. Additionally, 107 healthy individuals and 107 patients with simple diabetes mellitus were selected as the control groups. Changes in β 2-MG, HbA1c, and VEGF levels in the three groups as well as the different proteinuria exhibited by the three groups were examined.

RESULTS

Changes in β 2-MG, HbA1c, and VEGF levels in the disease, healthy, and simple diabetes groups were significantly different ($P < 0.05$). The expression of these factors from high to low were evaluated in different groups by pairwise comparison. In the disease group, high to low changes in β 2-MG, HbA1c, and VEGF levels were noted in the massive proteinuria, microproteinuria, and normal urinary protein groups, respectively. Changes in these factors were positively correlated with disease progression.

CONCLUSION

The expression of serum β 2-MG, HbA1c, and VEGF was closely correlated with DN progression, and disease progression could be evaluated by these factors.

Key Words: Type 2 diabetic nephropathy; β 2-microglobulin; Glycosylated hemoglobin; Vascular endothelial growth factor; Disease progression

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This study investigated the relationship between diabetic nephropathy (DN) and the expression of serum β 2-microglobulin (β 2-MG), glycosylated hemoglobin (HbA1c), and vascular endothelial growth factor (VEGF). In total, 107 patients with type 2 diabetes mellitus complicated by nephropathy were included in this study. Additionally, 107 healthy individuals were included in the control group. The expression levels of these factors, from high to low, were evaluated in the different groups by pairwise comparison. Serum β 2-MG, HbA1c, and VEGF were all closely correlated with DN progression based on all indicators.

Citation: Yang B, Zhao XH, Ma GB. Role of serum β 2-microglobulin, glycosylated hemoglobin, and vascular endothelial growth factor levels in diabetic nephropathy. *World J Clin Cases* 2022; 10(23): 8205-8211

URL: <https://www.wjgnet.com/2307-8960/full/v10/i23/8205.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i23.8205>

INTRODUCTION

Diabetic nephropathy (DN) is a common clinical diabetic microangiopathy that is known to be an important cause of death in patients with end-stage renal disease[1]. Studies have reported that inflammatory reactions and vascular endothelial cell damage are important factors in the pathogenesis of DN [2]. β 2-microglobulin (β 2-MG) is a microprotein formed by lymphocytes, polymorphonuclear leukocytes, and platelets, which has a positive effect on the inflammatory response[3]. Glycosylated hemoglobin (HbA1c) levels can reflect the specific control of blood glucose levels in patients in recent months. Excessively elevated HbA1c levels indicate the worsening of hyperglycemic injury in patients, increasing the influence of hyperglycemia on the development of microvascular lesions[4]. Vascular endothelial growth factor (VEGF) reflects the development of capillary malformations caused by pathological changes to a certain extent, and the degree of kidney disease can be determined by analyzing the development of renal microvascular malformations[5]. Considering all of these, this study examined the relationship between the expression of serum β 2-MG, HbA1c, and VEGF and the evaluation of DN patients, providing a scientific basis for clinical treatment and analysis of therapeutic effects.

MATERIALS AND METHODS

Study population

A total of 107 patients with type 2 diabetes complicated with nephropathy and treated in our hospital from May 2018 to February 2021 were included in the study. Among them, 59 were male and 48 were female, with a mean age of 49.27 ± 4.26 years old and mean body mass index (BMI) of 24.39 ± 1.54 kg/m². These patients were divided into three groups based on urine protein content: normal urinary protein group, < 30 mg/g, 32 patients; microproteinuria group, 30–300 mg/g, 35 patients; and massive proteinuria group, > 300 mg/g, 40 patients. 107 healthy individuals and 107 patients with simple diabetes were recruited, and were established as the two control groups. There were no obvious differences in sex, age, or BMI among the five groups ($P > 0.05$) (Table 1). All recruited patients provided written informed consent, which was approved by the Ethics Committee.

The inclusion criteria were as follows: (1) diagnosis of DN[6]; (2) blood glomerular filtration rate < 15 mL/min; (3) serum creatinine level > 177 μ mol/L; and (4) exhibited normal level of consciousness and ability to communicate.

The exclusion criteria were as follows: (1) tumors; (2) communication disorders; (3) any hormone therapy; (4) abnormal routine blood laboratory findings; and (5) refusal to cooperate with the treatment plan of this study.

Table 1 Comparison of baseline data among the five groups (mean $\pm$ SE)

Group	n	Age (yr)	BMI (kg/m ²)	Sex (male/female)
Normal urinary protein group	32	49.34 $\pm$ 4.28	24.33 $\pm$ 1.65	15/17
Microproteinuria group	35	49.22 $\pm$ 4.34	24.31 $\pm$ 1.51	15/20
Massive proteinuria group	40	49.25 $\pm$ 4.29	24.52 $\pm$ 1.52	29/11
Healthy group	107	49.72 $\pm$ 4.11	24.49 $\pm$ 1.62	55/52
Simple diabetes group	107	49.88 $\pm$ 4.69	24.59 $\pm$ 1.97	57/50
χ^2 /F value	-	0.281	0.259	8.142
P value	-	0.890	0.904	0.087

BMI: Body mass index.

Enzyme-linked immunosorbent assay

Four milliliters of fasting blood were extracted from all patients and centrifuged at 3500 r/min for 15 min. The supernatant of each sample was then collected, which were used to determine serum β 2-MG, HbA1c, and VEGF levels by Enzyme-linked immunosorbent assay. All detection reagents were obtained from Roche Shanghai, and the operating procedures were strictly followed according to the manufacturer's instructions.

Observational index

Changes in β 2-MG, HbA1c, and VEGF levels in the disease, healthy, and simple diabetes groups were compared. In the disease group, the same three factors were compared between the normal, microproteinuria, and massive proteinuria groups. Moreover, the correlation between disease progression and β 2-MG, HbA1c, and VEGF levels were also analyzed.

Statistical analysis

SPSS ver. 22.0 was used in the data analysis, and data were expressed as mean $\pm$ SE. Analysis of variance was used for multi-group comparisons and the least significant difference-*t* test was used for pairwise comparisons. The count data were expressed as (*n*) %, and the χ^2 test was used. A *P* value < 0.05 indicated a statistically significant difference.

RESULTS

Comparison of changes in β 2-MG, HbA1c, and VEGF levels in the disease, healthy, and simple diabetes groups

The changes in β 2-MG, HbA1c, and VEGF levels in the disease, healthy, and simple diabetes groups were significantly different (*P* < 0.05). Changes in β 2-MG, HbA1c, and VEGF levels, from high to low, were noted in the disease, simple diabetes, and healthy groups, and the differences were statistically significant (*P* < 0.05, Table 2).

Comparison of changes in β 2-MG, HbA1c, and VEGF levels among the different disease groups

There were statistically significant differences observed in terms of changes in β 2-MG, HbA1c, and VEGF levels among the different disease groups (*P* < 0.05). β 2-MG, HbA1c, and VEGF levels, from high to low, were noted in the massive proteinuria, microproteinuria, and normal urinary protein groups (Table 3).

Correlation analysis

Correlation analysis indicated that changes in serum β 2-MG, HbA1c, and VEGF levels were positively correlated with disease progression in patients with DN, showing statistically significant differences for all three factors (*P* < 0.05, Table 4).

DISCUSSION

With an aging population in China, the lifestyle of patients with diabetes has changed significantly, resulting an increase in the incidence of diabetes annually[7]. Some studies have reported that diabetes

Table 2 Comparison of changes in β 2-microglobulin, glycosylated hemoglobin, and Vascular endothelial growth factor levels in the disease, healthy, and simple diabetes groups (mean $\pm$ SE)

Group	<i>n</i>	β 2-MG (mg/L)	HbA1c (%)	VEGF (ng/L)
Disease group	107	4.34 $\pm$ 1.53	10.84 $\pm$ 1.81	176.69 $\pm$ 11.84
Simple diabetes group	107	2.56 $\pm$ 1.14	8.40 $\pm$ 1.24	137.84 $\pm$ 10.36
Healthy groups	107	1.11 $\pm$ 0.82	4.58 $\pm$ 0.95	109.32 $\pm$ 12.44
<i>F</i> value		195.518	558.284	912.281
<i>P</i> value		0.000	0.000	0.000
LSD- <i>t</i> (disease <i>vs</i> simple diabetes)		10.862	12.921	24.535
<i>P</i> value		0.000	0.000	0.000
LSD- <i>t</i> (disease <i>vs</i> healthy)		19.742	33.148	42.548
<i>P</i> value		0.000	0.000	0.000
LSD- <i>t</i> (healthy <i>vs</i> simple diabetes)		8.879	20.228	18.013
<i>P</i> value		0.000	0.000	0.000

LSD-*t*: Least significant difference; β 2-MG: β 2-microglobulin; HbA1c: Glycosylated hemoglobin; VEGF: Vascular endothelial growth factor.

Table 3 Comparison of β 2-microglobulin, glycosylated hemoglobin, and vascular endothelial growth factor levels among the different disease groups (mean $\pm$ SE)

Group	<i>n</i>	β 2-MG (mg/L)	HbA1c (%)	VEGF (ng/L)
Normal urinary protein group	32	3.37 $\pm$ 1.32	9.26 $\pm$ 1.33	170.16 $\pm$ 10.62
Microproteinuria group	35	4.19 $\pm$ 1.39	10.57 $\pm$ 1.21	175.45 $\pm$ 10.39
Massive proteinuria group	40	5.26 $\pm$ 1.29	12.34 $\pm$ 1.35	182.99 $\pm$ 10.97
<i>F</i> value		18.250	51.067	13.179
<i>P</i> value		0.000	0.000	0.000
LSD- <i>t</i> (normal urinary protein <i>vs</i> Microproteinuria)		2.516	4.119	2.025
<i>P</i> value		0.013	0.000	0.045
LSD- <i>t</i> (normal urinary protein <i>vs</i> massive proteinuria)		5.981	9.994	5.066
<i>P</i> value		0.000	0.000	0.000
LSD- <i>t</i> (microproteinuria <i>vs</i> massive proteinuria)		3.471	5.887	3.051
<i>P</i> value		0.001	0.000	0.003

LSD-*t*: Least significant difference; β 2-MG: β 2-microglobulin; HbA1c: Glycosylated hemoglobin; VEGF: Vascular endothelial growth factor.

Table 4 Correlation analysis

Index	β 2-MG (mg/L)		HbA1c (%)		VEGF (ng/L)	
	<i>r</i> value	<i>P</i> value	<i>r</i> value	<i>P</i> value	<i>r</i> value	<i>P</i> value
Disease progression	0.705	0.000	0.707	0.000	0.518	0.000

β 2-MG: β 2-microglobulin; HbA1c: Glycosylated hemoglobin; VEGF: Vascular endothelial growth factor.

is the third most common chronic non-communicable disease. In the disease progression of diabetes mellitus, the risk of DN gradually increases with continuous changes in the patient's microvasculature [8]. Epidemiological investigations show that nearly 37.4% of patients with diabetes mellitus in China have different severities of DN[9]. Currently, the pathogenesis of DN is unclear, but most studies

believe that carbohydrate and lipid metabolism disorders, hemodynamic changes, and abnormal secretion of cytokines lead to the often insidious onset of DN[10]. In most patients, DN manifests as massive proteinuria, which significantly affects patient safety. Considering these, thorough analysis of serological indicators, timely and effective assessment of the patient's condition, and early preventive intervention should therefore have a positive effect on the prognosis of patients.

VEGF has a positive promoting effect on the proliferation of vascular endothelial cells, and is currently clinically considered an important indicator of diabetic retinopathy. It plays a critical role in the effective maintenance of the functional integrity of patients' new blood vessels and endothelial cells [11]. In this study, VEGF levels increased significantly with the progression of the disease, suggesting that with the increase in VEGF levels, the risks of renal malformed vessels as well as vascular permeability changes significantly increase, ultimately resulting in the progression of DN. Through the comparison of VEGF levels in patients with diabetes mellitus, Wang *et al*[12] found that VEGF levels in patients with DN were distinctly higher than those in patients with simple diabetes, which was consistent with this study.

HbA1c has been a common indicator used to evaluate patients' blood glucose control in recent months. The more obvious the increase in HbA1c level, the worse the blood glucose control[13,14]. However, excessive blood glucose levels can damage vascular endothelial cells in patients with DN and easily induce spasms of renal afferent arterioles, thus causing damage to the renal units due to ischemia and hypoxia[15,16]. In the early stages of the disease, kidney injury remains reversible to some extent. However, with gradual enhancement of kidney cell injury, the patient's kidney self-repair ability is eventually lost. By this time, kidney transplantation or dialysis treatment is usually required, which seriously threatens a patient's life[17,18]. Guo *et al*[19] analyzed the HbA1c level of patients with DN, and showed that the determination of HbA1c level in patients has a positive effect on early kidney injury. Yuan *et al*[20] obtained similar results.

β 2-MG is a microglobulin protein that can be used as a clinical indicator of chronic kidney disease. During the progression of chronic kidney disease, the glomerular rate of patients significantly decreases as the severity of the disease increases, resulting in a large amount of microglobulin extravasation. In clinical treatment, the therapeutic effect and condition of patients can be determined according to serum β 2-MG levels. In this study, β 2-MG levels significantly increased as the degree of kidney injury increased, which is consistent with the results of Guo *et al*[19].

There are limitations to this study: (1) The sample size was limited to retrospective studies from a single center; these findings need further multi-institutional validation with a larger sample size[21]; and (2) This was a retrospective study that could not completely avoid missing data and measurement biases. Therefore, further studies must include more candidate biomarkers to develop predictive models in the future[22].

CONCLUSION

Our research shows that the expression of serum β 2-MG, HbA1c, and VEGF are closely correlated with DN progression, and the progression of the disease can be evaluated by the expression of these factors in patients in the future.

ARTICLE HIGHLIGHTS

Research background

Diabetic nephropathy (DN) is a common complication of diabetes and is the leading cause of chronic kidney disease. Many identified biomarkers related to DN have been reported in previous studies, but none have been tested at bedside or in clinical trials. Moreover, their validation in larger patient cohorts and longitudinal studies are lacking.

Research motivation

Recent studies have demonstrated that inflammatory reactions and vascular endothelial cell damage are important factors in the pathogenesis of DN. The early detection of changes in renal function is of great benefit for the treatment of DN. Therefore, we aimed to analyze the significance of serum β 2 microglobulin (β 2-MG), glycosylated hemoglobin (HbA1c), and vascular endothelial growth factor (VEGF) levels in DN.

Research objectives

The objective of this study was to determine whether the expression levels of serum β 2-MG, HbA1c, and VEGF are associated with DN.

Research methods

A total of 107 patients with type 2 diabetes mellitus complicated by nephropathy were included in the study, and 107 healthy individuals as well as 107 patients with simple diabetes mellitus were selected as the control groups. Changes in β 2-MG, HbA1c, and VEGF levels in the three groups and the different proteinuria groups of the disease group were examined.

Research results

The expression levels of these factors were evaluated in different groups by pairwise comparison. Changes in β 2-MG, HbA1c, and VEGF levels in the different disease groups were significantly different. By pairwise comparison, changes in β 2-MG, HbA1c, and VEGF levels, from high to low, were noted in the massive proteinuria, microproteinuria, and normal urinary protein groups. Changes in these factors were positively correlated with disease progression.

Research conclusions

The expression of serum β 2-MG, HbA1c, and VEGF are closely correlated with DN progression, and the progression of the disease can be evaluated by the expression of these factors in patients in the future.

Research perspectives

Diabetic kidney disease is a major health care challenge that complicates the course of many people living with diabetes. The current study showed that the 22 studied biomarkers had different levels of diagnostic accuracy, ranging from excellent to very good to good, and specificity values. The combined diagnosis of multiple biomarkers may improve the accuracy of early diagnosis of DN. In the future, optimization of biomarkers for clinical situations requires prospective validation in many patients with diabetic nephropathy, and needs to be performed in different critically ill populations.

FOOTNOTES

Author contributions: Yang B contributed to conceptualization, data analysis, and writing; Zhao XH contributed to data analysis and writing; Ma GB revised the manuscript; and all authors have read and approved the manuscript.

Institutional review board statement: The study was reviewed and approved by the 3201 Hospital Institutional Review Board.

Informed consent statement: Informed consent was obtained from all study participants or their legal guardians prior to enrolment.

Conflict-of-interest statement: The authors have no conflicts of interest to declare.

Data sharing statement: Data supporting the findings of this study can be found within the article.

STROBE statement: The manuscript was prepared and revised according to the STROBE statement.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Bing Yang 0000-0003-1305-4211; Xiao-Hong Zhao 0000-0001-9752-5783; Guo-Bin Ma 0000-0002-7896-5000.

S-Editor: Wang JL

L-Editor: A

P-Editor: Wang JL

REFERENCES

- 1 Srivastava SP, Zhou H, Setia O, Liu B, Kanasaki K, Koya D, Dardik A, Fernandez-Hernando C, Goodwin J. Loss of endothelial glucocorticoid receptor accelerates diabetic nephropathy. *Nat Commun* 2021; **12**: 2368 [PMID: 33888696 DOI: 10.1038/s41467-021-22617-y]
- 2 Alomari G, Al-Trad B, Hamdan S, Aljabali AAA, Al Zoubi MS, Al-Batanyeh K, Qar J, Eaton GJ, Alkaraki AK, Alshaer

- W, Haifawi S, Jemon K, Chellappan DK, Dua K, Tambuwala MM. Alleviation of diabetic nephropathy by zinc oxide nanoparticles in streptozotocin-induced type 1 diabetes in rats. *IET Nanobiotechnol* 2021; **15**: 473-483 [PMID: [34694755](#) DOI: [10.1049/nbt2.12026](#)]
- 3 **Nongnuch A**, Kitiyakara C, Sappadungsuk S, Sathirapongsasuti N, Vipattawat K, Zhang P, Davies N, Davenport A. Pilot study to investigate differences in middle molecules, oxidative stress and markers of peripheral vascular disease in patients treated by high flux haemodialysis and haemodiafiltration. *PLoS One* 2021; **16**: e0258223 [PMID: [34614018](#) DOI: [10.1371/journal.pone.0258223](#)]
 - 4 **Ratra D**, Dalan D, Prakash N, Kaviarasan K, Thanikachalam S, Das UN, Angayarkanni N. Quantitative analysis of retinal microvascular changes in prediabetic and diabetic patients. *Indian J Ophthalmol* 2021; **69**: 3226-3234 [PMID: [34708778](#) DOI: [10.4103/ijo.IJO_1254_21](#)]
 - 5 **Zheng Q**, Wang Y, Yang H, Sun L, Fu X, Wei R, Liu YN, Liu WJ. Efficacy and Safety of Daprodustat for Anemia Therapy in Chronic Kidney Disease Patients: A Systematic Review and Meta-Analysis. *Front Pharmacol* 2020; **11**: 573645 [PMID: [33597868](#) DOI: [10.3389/fphar.2020.573645](#)]
 - 6 **Feng XF**, Li AM, Xu SL, Jia P, Shen SH. [Clinical value of glomerular filtration rate, urinary microalbumin, blood β 2-microglobulin and cystatin C in the diagnosis of diabetic nephropathy]. *Zhonghua Heyixue Yu Fenzixiangxue Zazhi* 2017; **37**: 331-336 [DOI: [10.3760/cma.j.issn.2095-2848.2017.06.003](#)]
 - 7 **Li Y**, Guo C, Cao Y. Secular incidence trends and effect of population aging on mortality due to type 1 and type 2 diabetes mellitus in China from 1990 to 2019: findings from the Global Burden of Disease Study 2019. *BMJ Open Diabetes Res Care* 2021; **9** [PMID: [34732399](#) DOI: [10.1136/bmjdr-2021-002529](#)]
 - 8 **Huang Y**, Huang W, Zhang S, Jiang Y, Zhu JH. [Preliminary exploration on value of serum neutrophil gelatinase-associated lipocalin, high-sensitivity C-reaction protein, cystatin C and urine-microalbumin in early diagnosis of diabetic nephropathy]. *Shiyong Yufang Yixue* 2017; **24**: 1168-1171 [DOI: [10.3969/j.issn.1006-3110.2017.10.005](#)]
 - 9 **Bramham K**. Diabetic Nephropathy and Pregnancy. *Semin Nephrol* 2017; **37**: 362-369 [PMID: [28711075](#) DOI: [10.1016/j.semnephrol.2017.05.008](#)]
 - 10 **Hu M**, Wang R, Li X, Fan M, Lin J, Zhen J, Chen L, Lv Z. LncRNA MALAT1 is dysregulated in diabetic nephropathy and involved in high glucose-induced podocyte injury via its interplay with β -catenin. *J Cell Mol Med* 2017; **21**: 2732-2747 [PMID: [28444861](#) DOI: [10.1111/jcmm.13189](#)]
 - 11 **Zhang XG**, Zhang YQ, Cheng QP, Cao Y, Sun JM, Lv XF. The impact of insulin pump therapy to oxidative stress in patients with diabetic nephropathy. *Eur J Med Res* 2018; **23**: 7 [PMID: [29433562](#) DOI: [10.1186/s40001-018-0304-2](#)]
 - 12 **Wang F**, Gao XY, Zhang R. [Changes and detection significance of serum inflammatory factors, vWf, VEGF and adhesion molecules in patients with diabetic nephropathy]. *Hainan Yixueyuan Xuebao* 2017; **23**: 2207-2210 [DOI: [10.13210/j.cnki.jhmu.20170816.004](#)]
 - 13 **Nally LM**, Wagner J, Sherr J, Tichy E, Weyman K, Ginley MK, Zajac K, Desousa M, Shabanova V, Petry NM, Tamborlane WV, Van Name M. A Pilot Study of Youth With Type 1 Diabetes Initiating Use of a Hybrid Closed-Loop System While Receiving a Behavioral Economics Intervention. *Endocr Pract* 2021; **27**: 545-551 [PMID: [34120699](#) DOI: [10.1016/j.epr.2020.11.017](#)]
 - 14 **Levin P**, Hoogwerf BJ, Snell-Bergeon J, Vigers T, Pyle L, Bromberger L. Ultra Rapid-Acting Inhaled Insulin Improves Glucose Control in Patients With Type 2 Diabetes Mellitus. *Endocr Pract* 2021; **27**: 449-454 [PMID: [33853718](#) DOI: [10.1016/j.epr.2020.10.004](#)]
 - 15 **Liu S**, Yang Y, Gao H, Zhou N, Wang P, Zhang Y, Zhang A, Jia Z, Huang S. Trehalose attenuates renal ischemia-reperfusion injury by enhancing autophagy and inhibiting oxidative stress and inflammation. *Am J Physiol Renal Physiol* 2020; **318**: F994-F1005 [PMID: [32068461](#) DOI: [10.1152/ajprenal.00568.2019](#)]
 - 16 **Nespoux J**, Patel R, Hudkins KL, Huang W, Freeman B, Kim YC, Koepsell H, Alpers CE, Vallon V. Gene deletion of the Na⁺-glucose cotransporter SGLT1 ameliorates kidney recovery in a murine model of acute kidney injury induced by ischemia-reperfusion. *Am J Physiol Renal Physiol* 2019; **316**: F1201-F1210 [PMID: [30995111](#) DOI: [10.1152/ajprenal.00111.2019](#)]
 - 17 **Campos CS**, Ferreira GF, Bastos KV, da Costa Carbogim F, de Brito Poveda V, de Araújo Püschel VA, Dos Santos KB. Immunosuppressive adherence after kidney transplantation in adult patients treated at an outpatient clinic: a better practice implementation project. *JBIM Evid Implement* 2020; **19**: 306-314 [PMID: [34491926](#) DOI: [10.1097/XEB.0000000000000262](#)]
 - 18 **Yu H**, Liu D, Shu G, Jin F, Du Y. Recent advances in nanotherapeutics for the treatment and prevention of acute kidney injury. *Asian J Pharm Sci* 2021; **16**: 432-443 [PMID: [34703493](#) DOI: [10.1016/j.ajps.2020.11.001](#)]
 - 19 **Guo WH**, Zhang J. [The value of serum glycosylated albumin in early diagnosis of diabetic nephropathy]. *Jiannan Yixue Yu Linchuang* 2017; **14**: 27-29 [DOI: [10.3969/j.issn.1672-9455.2017.25.011](#)]
 - 20 **Yuan TF**, Li Y, Peng R. [Change of serum SCC and β 2-microglobulin in patients with chronic kidney disease and their correlation]. *Jiannan Yixue Yu Linchuang* 2017; **14**: 756-758 [DOI: [10.3969/j.issn.1672-9455.2017.06.005](#)]
 - 21 **Sen S**, Cohen AS. The Impact of Sample Size and Various Other Factors on Estimation of Dichotomous Mixture IRT Models. *Educ Psychol Meas* 2022 [DOI: [10.1177/00131644221094325](#)]
 - 22 **Sewell K**, Halanach JH, Russell LB, Andreae SJ, Cherrington AL, Martin MY, Pisu M, Safford MM. Blood Pressure Measurement Biases in Clinical Settings, Alabama, 2010-2011. *Prev Chronic Dis* 2016; **13**: E01 [PMID: [26741995](#) DOI: [10.5888/pcd13.150348](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

