

World Journal of *Clinical Cases*

World J Clin Cases 2022 October 16; 10(29): 10391-10822



STANDARD AND CONSENSUS

- 10391** Baishideng's Reference Citation Analysis database announces the first Article Influence Index of multidisciplinary scholars
Wang JL, Ma YJ, Ma L, Ma N, Guo DM, Ma LS

REVIEW

- 10399** Cholecystectomy for asymptomatic gallstones: Markov decision tree analysis
Lee BJH, Yap QV, Low JK, Chan YH, Shelat VG
- 10413** Liver transplantation for hepatocellular carcinoma: Historical evolution of transplantation criteria
Ince V, Sahin TT, Akbulut S, Yilmaz S

MINIREVIEWS

- 10428** Prostate only radiotherapy using external beam radiotherapy: A clinician's perspective
Lee JW, Chung MJ

ORIGINAL ARTICLE**Retrospective Study**

- 10435** Age-adjusted NT-proBNP could help in the early identification and follow-up of children at risk for severe multisystem inflammatory syndrome associated with COVID-19 (MIS-C)
Rodriguez-Gonzalez M, Castellano-Martinez A
- 10451** Clinicopathological characteristics and prognosis of gastric signet ring cell carcinoma
Tian HK, Zhang Z, Ning ZK, Liu J, Liu ZT, Huang HY, Zong Z, Li H

- 10467** Development and validation of a prognostic nomogram for decompensated liver cirrhosis
Zhang W, Zhang Y, Liu Q, Nie Y, Zhu X

Observational Study

- 10478** Effect of medical care linkage-continuous management mode in patients with posterior circulation cerebral infarction undergoing endovascular interventional therapy
Zhu FX, Ye Q
- 10487** Effect of the COVID-19 pandemic on patients with presumed diagnosis of acute appendicitis
Akbulut S, Tuncer A, Ogut Z, Sahin TT, Koc C, Guldogan E, Karabulut E, Tanriverdi ES, Ozer A

EVIDENCE-BASED MEDICINE

- 10501** Delineation of a SMARCA4-specific competing endogenous RNA network and its function in hepatocellular carcinoma

Zhang L, Sun T, Wu XY, Fei FM, Gao ZZ

SYSTEMATIC REVIEWS

- 10516** Comparison of laboratory parameters, clinical symptoms and clinical outcomes of COVID-19 and influenza in pediatric patients: A systematic review and meta-analysis

Yu B, Chen HH, Hu XF, Mai RZ, He HY

CASE REPORT

- 10529** Surgical treatment of bipolar segmental clavicle fracture: A case report

Liang L, Chen XL, Chen Y, Zhang NN

- 10535** Multiple disciplinary team management of rare primary splenic malignancy: Two case reports

Luo H, Wang T, Xiao L, Wang C, Yi H

- 10543** Klippel-Trenaunay-Weber syndrome with ischemic stroke: A case report

Lee G, Choi T

- 10550** Vedolizumab in the treatment of immune checkpoint inhibitor-induced colitis: Two case reports

Zhang Z, Zheng CQ

- 10559** Novel way of patent foramen ovale detection and percutaneous closure by intracardiac echocardiography: A case report

Han KN, Yang SW, Zhou YJ

- 10565** Treatment failure in a patient infected with *Listeria* sepsis combined with latent meningitis: A case report

Wu GX, Zhou JY, Hong WJ, Huang J, Yan SQ

- 10575** Three-in-one incidence of hepatocellular carcinoma, cholangiocellular carcinoma, and neuroendocrine carcinoma: A case report

Wu Y, Xie CB, He YH, Ke D, Huang Q, Zhao KF, Shi RS

- 10583** Intestinal microbiome changes in an infant with right atrial isomerism and recurrent necrotizing enterocolitis: A case report and review of literature

Kaplina A, Zaikova E, Ivanov A, Volkova Y, Alkhova T, Nikiforov V, Latypov A, Khavkina M, Fedoseeva T, Pervunina T, Skorobogatova Y, Volkova S, Ulyantsev V, Kalinina O, Sitkin S, Petrova N

- 10600** *Serratia fonticola* and its role as a single pathogen causing emphysematous pyelonephritis in a non-diabetic patient: A case report

Villasuso-Alcocer V, Flores-Tapia JP, Perez-Garfias F, Rochel-Perez A, Mendez-Dominguez N

- 10606** Cardiac myxoma shedding leads to lower extremity arterial embolism: A case report

Meng XH, Xie LS, Xie XP, Liu YC, Huang CP, Wang LJ, Zhang GH, Xu D, Cai XC, Fang X

- 10614** Extracorporeal membrane oxygenation in curing a young man after modified Fontan operation: A case report
Guo HB, Tan JB, Cui YC, Xiong HF, Li CS, Liu YF, Sun Y, Pu L, Xiang P, Zhang M, Hao JJ, Yin NN, Hou XT, Liu JY
- 10622** Wandering small intestinal stromal tumor: A case report
Su JZ, Fan SF, Song X, Cao LJ, Su DY
- 10629** Acute mesenteric ischemia secondary to oral contraceptive-induced portomesenteric and splenic vein thrombosis: A case report
Zhao JW, Cui XH, Zhao WY, Wang L, Xing L, Jiang XY, Gong X, Yu L
- 10638** Perioperative anesthesia management in pediatric liver transplant recipient with atrial septal defect: A case report
Liu L, Chen P, Fang LL, Yu LN
- 10647** Multiple tophi deposits in the spine: A case report
Chen HJ, Chen DY, Zhou SZ, Chi KD, Wu JZ, Huang FL
- 10655** Myeloproliferative neoplasms complicated with β -thalassemia: Two case report
Xu NW, Li LJ
- 10663** Synchronous renal pelvis carcinoma associated with small lymphocytic lymphoma: A case report
Yang HJ, Huang X
- 10670** *Leclercia adecarboxylata* infective endocarditis in a man with mitral stenosis: A case report and review of the literature
Tan R, Yu JQ, Wang J, Zheng RQ
- 10681** Progressive ataxia of cerebrotendinous xanthomatosis with a rare c.255+1G>T splice site mutation: A case report
Chang YY, Yu CQ, Zhu L
- 10689** Intravesical explosion during transurethral resection of bladder tumor: A case report
Xu CB, Jia DS, Pan ZS
- 10695** Submucosal esophageal abscess evolving into intramural submucosal dissection: A case report
Jiao Y, Sikong YH, Zhang AJ, Zuo XL, Gao PY, Ren QG, Li RY
- 10701** Immune checkpoint inhibitor-associated arthritis in advanced pulmonary adenocarcinoma: A case report
Yang Y, Huang XJ
- 10708** Chondroid syringoma of the lower back simulating lipoma: A case report
Huang QF, Shao Y, Yu B, Hu XP
- 10713** Tension-reduced closure of large abdominal wall defect caused by shotgun wound: A case report
Li Y, Xing JH, Yang Z, Xu YJ, Yin XY, Chi Y, Xu YC, Han YD, Chen YB, Han Y

- 10721** Myocardial bridging phenomenon is not invariable: A case report
Li HH, Liu MW, Zhang YF, Song BC, Zhu ZC, Zhao FH
- 10728** Recurrent atypical leiomyoma in bladder trigone, confused with uterine fibroids: A case report
Song J, Song H, Kim YW
- 10735** Eczema herpeticum vs dermatitis herpetiformis as a clue of dedicator of cytokinesis 8 deficiency diagnosis: A case report
Alshengeti A
- 10742** Cutaneous allergic reaction to subcutaneous vitamin K₁: A case report and review of literature
Zhang M, Chen J, Wang CX, Lin NX, Li X
- 10755** Perithyroidal hemorrhage caused by hydrodissection during radiofrequency ablation for benign thyroid nodules: Two case reports
Zheng BW, Wu T, Yao ZC, Ma YP, Ren J
- 10763** Malignant giant cell tumors of the tendon sheath of the right hip: A case report
Huang WP, Gao G, Yang Q, Chen Z, Qiu YK, Gao JB, Kang L
- 10772** Atypical Takotsubo cardiomyopathy presenting as acute coronary syndrome: A case report
Wang ZH, Fan JR, Zhang GY, Li XL, Li L
- 10779** Secondary light chain amyloidosis with Waldenström's macroglobulinemia and internodal marginal zone lymphoma: A case report
Zhao ZY, Tang N, Fu XJ, Lin LE
- 10787** Bilateral occurrence of sperm granulomas in the left spermatic cord and on the right epididymis: A case report
Ly DY, Xie HJ, Cui F, Zhou HY, Shuang WB
- 10794** Glucocorticoids combined with tofacitinib in the treatment of Castleman's disease: A case report
Liu XR, Tian M
- 10803** Giant bilateral scrotal lipoma with abnormal somatic fat distribution: A case report
Chen Y, Li XN, Yi XL, Tang Y
- 10811** Elevated procalcitonin levels in the absence of infection in procalcitonin-secreting hepatocellular carcinoma: A case report
Zeng JT, Wang Y, Wang Y, Luo ZH, Qing Z, Zhang Y, Zhang YL, Zhang JF, Li DW, Luo XZ

LETTER TO THE EDITOR

- 10817** "Helicobacter pylori treatment guideline: An Indian perspective": Letter to the editor
Swarnakar R, Yadav SL
- 10820** Effect of gender on the reliability of COVID-19 rapid antigen test among elderly
Nori W, Akram W

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Natalia Stepanova, DSc, MD, PhD, Academic Research, Chief Doctor, Full Professor, Department of Nephrology and Dialysis, State Institution "Institute of Nephrology of the National Academy of Medical Sciences of Ukraine", Kyiv 04050, Ukraine. nmstep88@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for *WJCC* as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The *WJCC*'s CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Hua-Ge Yu*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

October 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Intravesical explosion during transurethral resection of bladder tumor: A case report

Chuan-Bing Xu, Dong-Sheng Jia, Zheng-Sheng Pan

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Hakimi T, Afghanistan; Markic D, Croatia

Received: May 11, 2022

Peer-review started: May 11, 2022

First decision: July 12, 2022

Revised: July 15, 2022

Accepted: August 30, 2022

Article in press: August 30, 2022

Published online: October 16, 2022



Chuan-Bing Xu, Dong-Sheng Jia, Zheng-Sheng Pan, Department of Urology, Zibo Central Hospital, Zibo 255036, Shandong Province, China

Corresponding author: Zheng-Sheng Pan, MD, Attending Doctor, Department of Urology, Zibo Central Hospital, No. 54 Gongqingtuan West Road, Zibo 255036, Shandong Province, China. panzhengsheng@aliyun.com

Abstract

BACKGROUND

Intravesical explosion during transurethral resection of bladder tumor (TUR-BT) is a very rare complication, and it may result in rupture of the bladder, which usually requires surgical correction and causes a potential threat to the patient's life.

CASE SUMMARY

This paper reports a case of intravesical explosion during TUR-BT. Combined with the literature review, the risk factors are analyzed and measures of prevention and treatment are discussed.

CONCLUSION

Although rare, intravesical explosions can cause serious consequences, and the loud explosion can also lead to a profound psychological shadow on the patient. Urologists must be aware of this potential complication. Careful operative techniques and special precautions can reduce the risk of this complication.

Key Words: Transurethral resection of bladder tumor; Intravesical explosion; Vesical rupture; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Transurethral resection of bladder tumor (TUR-BT) is the most common operation performed for non-muscle invasive bladder cancer. Although it is considered as a safe and minimally invasive procedure, occasionally life-threatening complications may be seen. Intravesical explosion is a very rare complication, and it may result in rupture of the bladder and has received special attention in consensus on safety of TUR-BT. Here, we present a case of an intravesical explosion during TUR-BT leading to bladder rupture to remind urologists of this rare complication with suggestions on how to manage and prevent this complication.

Citation: Xu CB, Jia DS, Pan ZS. Intravesical explosion during transurethral resection of bladder tumor: A case report. *World J Clin Cases* 2022; 10(29): 10689-10694

URL: <https://www.wjgnet.com/2307-8960/full/v10/i29/10689.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i29.10689>

INTRODUCTION

Transurethral resection of bladder tumor (TUR-BT) is the most common operation performed for non-muscle invasive bladder cancer. Although it is considered as a safe and minimally invasive procedure, occasionally life-threatening complications may be seen. Intravesical explosion, which was first reported in 1926 by Cassuto[1], is a very rare complication that can cause serious consequences, and the loud explosion can also lead to a profound psychological shadow on the patient. It may result in rupture of the bladder and has received special attention in consensus on safety of TUR-BT[2]. If bladder explosion happens and is not diagnosed and treated timely, it will cause serious consequences. Here, we present a case of intravesical explosion during TUR-BT leading to bladder rupture to remind urologists of this rare complication with suggestions on how to manage and prevent this complication.

CASE PRESENTATION

Chief complaints

A 44-year-old man was admitted to the hospital with painless gross haematuria that had persisted for 1 wk.

History of present illness

The patient experienced haematuria throughout urination, with blood clots, but without dysuria, frequent or urgent urination, or lower back pain.

History of past illness

The patient had a history of hypertension.

Personal and family history

The patient had no personal and family history.

Physical examination


There was no specific physical examination.

Laboratory examinations

Routine urine test and microscopic examination showed plenty of red cells.

Imaging examinations

Ultrasound showed multiple irregular, heterogeneous, slightly high echoes in the bladder, of which the smallest was located in the left bladder wall (approximately 7 mm × 5 mm in size) and the largest one in the right posterior wall (approximately 61 mm × 47 mm in size). Colour Doppler flow imaging detected obvious blood flow signals within the region. Cystoscopy identified many cauliflower-like tumors in the left apical and posterior walls of the bladder (Figure 1A), the largest of which was in the posterior wall of the bladder, with local necrosis and bleeding. Computed tomography (CT) examination showed a tumor in the right lateral wall of the bladder (Figure 1B). Despite its size, this could not be visualised entirely because of its location. There was no obvious abnormality in the ureteral orifice.



DOI: 10.12998/wjcc.v10.i29.10689 Copyright ©The Author(s) 2022.

Figure 1 Clinical examination. A: Cystoscopy examination revealed a large cauliflower tumor in the bladder; B: Computed tomography examination showed a tumor in the right lateral wall of the bladder.

FINAL DIAGNOSIS

Pathological examination indicated possible high-grade urothelial carcinoma. The recommended surgical procedure was radical cystectomy combined with urinary diversion. However, the patient was still young and was unwilling to undergo cystectomy. Therefore, after some discussion, we performed a TUR-BT.

TREATMENT

The patient was placed under general anaesthesia. An Olympus F24 plasma resectoscope was used intraoperatively. We used 0.9% warm normal saline as a flushing solution, and the monopolar electrocautery current was set at 120 W for coagulation and 280 W for cutting. The tumor was located and found to have a wide base and an extensive range. Resection was performed using step-by-step electroexcision. However, the surgical field of vision was poor because of the large tumor size and its abundant blood supply, which resulted in substantial haemorrhage. There was also an accumulation of resected tumor tissue in the bladder. The excised tissue was repeatedly rinsed using Ellikrinser. After approximately 120 min, we encountered difficulty accessing the tumor tissue in the apical wall of the bladder with the electric cutting ring. The assistant pressed down on the abdomen for assistance, and an explosive bang sound was emitted from the lower abdomen, leading to the blurring of the field of vision of the resectoscope and failure of bladder filling. We surmised this to be a bladder rupture, and an exploratory laparotomy was performed immediately. During this procedure, the bladder was broken into several irregular pieces, with irregular wound margins, a lacerated appearance, and active bleeding. The rupture was traced to the abdominal cavity. Approximately 300 mL of light red liquid was sucked out of the abdominal cavity. No obvious damage was found to the intestines or other abdominal structures, but a large number of blood clots were found in the bladder. Following careful exploration of the bleeding sites and complete haemostasis, the residual tumor tissue was completely resected, after which the bladder was checked carefully. No other lesions were found, and the urine spraying at the bilateral ureteral orifices showed no abnormality. After trimming the wound margin of the bladder and confirming that there was no further bleeding, the muscular layer and serosa layer were continuously sutured with 2.0 absorbable sutures. A test of bladder infusion showed no suture leakage. The pelvic and abdominal drainage tubes were placed, and the incision closed.

OUTCOME AND FOLLOW-UP

The drainage tubes were removed after 1 wk, and the urinary tube after 2 wk. There were no abnormalities in urinary function. The patient received two cycles of gemcitabine and cisplatin chemotherapy combined with regular intravesical gemcitabine. Both CT and cystoscopy at the 3-mo postoperative review revealed no abnormalities of the bladder or abdominal cavity. The patient had three recurrences of bladder tumors in the next 3 years and was treated in each instance by TUR-BT because of the smaller size of the tumors. The patient has received regular follow-ups up to the present time without further tumor recurrence.

DISCUSSION

Intravesical explosion is rare in TUR of the prostate (TURP) with a low incidence during TUR-BT. We conducted a Chinese and English literature review through the PubMed database and the Wanfang database for research on bladder explosions up to December 2021. Only three papers related to TUR-BT, and the rest related to TURP. TUR-BT and TURP are both transurethral procedures for conditions with similar causes, clinical manifestations, treatments, and preventative measures. Martov *et al*[3] investigated 5401 cases of intraoperative and postoperative complications of TURP, and one case of bladder gas explosion was found. This case developed into bladder rupture. Liu *et al*[4] conducted a retrospective analysis of the surgical data from 1950 cases of benign prostatic hyperplasia and found four cases of bladder gas explosion followed by bladder repair, amounting to an incidence of < 0.2%. In most cases, bladder gas explosion is characterised by a dull explosive sound in the middle to late stage of the operation, without obvious damage to the bladder and with no special treatment required. In instances where a mild injury occurs, it is in the form of congestion and slight tearing and bleeding of the bladder mucosa, which can be treated by electrosurgical resection and haemostasis. When a severe injury occurs, it is predominantly bladder rupture. Intraperitoneal injury is more common than extraperitoneal[5,6]. At this time, there are no reports in the literature of abdominal involvement or major vascular injury induced by bladder gas explosion. However, Seitz *et al*[7] have reported a case requiring expanded excision of the bladder due to poor blood supply at the bladder wound margin. In another case, the bladder repair was further complicated by bladder dehiscence due to the severe damage and this had required a second repair[8]. Additionally, improper treatment of fluid and electrolyte disorders can lead to patient mortality[9]. Therefore, in cases of bladder rupture, open or laparoscopic bladder repair must be performed on time, with additional attention paid to any injury of the intestinal tract or surrounding structures. As suggested by Georgios *et al*[10], laparoscopic repair of the bladder has the advantages of reducing surgical trauma, enabling easy aspiration of fluid in the abdominal cavity, and facilitating more comprehensive observation of the abdominal organs. However, open surgery is more advantageous for cases with large, irregular, and numerous bladder lacerations. Another serious problem of intravesical explosion is seeding of the tumor cells caused by intraperitoneal bladder rupture. Bus *et al*[11] reported the first case of tumor seeding to both adnexa in a patient with low grade urothelial cancer conservatively treated by TUR-BT that had intraperitoneal perforation.

An intravesical explosion is a violent chemical reaction of combustible gas under specific conditions. Bladder rupture may occur when the volume of gas and liquid that the bladder contains exceeds its maximum capacity. A gas explosion requires certain physical and chemical conditions, including a combustion agent (*e.g.*, hydrogen), an oxidant (*e.g.*, oxygen), and an ignition source (*e.g.*, an electric spark). During transurethral surgery, the heat of electroexcision may induce cellular inflammation, rupture, and gasification. The water content of the cells can then evaporate, leading surgical electrocoagulation to deform proteins and destroy cell walls[12]. Ning *et al*[13] analysed the gas produced by electrocautery of tissues through *in vitro* experiments and found its composition to include 40%-50% hydrogen and less than 3% oxygen. Similar results have been reported by Davis[14]. These concentrations of gases are not sufficient to produce an explosion, suggesting that electroexcision or electric coagulation alone does not cause intravesical explosions. It has been proposed that with a 40% volume ratio of hydrogen in the combustible gas generated by electroexcision, 11.4%-90.6% oxygen is required to induce a hydrogen explosion. Therefore, the accidental intraoperative introduction of air *via* a poorly sealed endoscopic sheath, through the flushing tube during the continuous bladder irrigation, or during repeated rinsing with Ellik, can provide sufficient oxidant for the gas to become explosive. The gas in the bladder accumulates when the operation is lengthy. In the present case, surgery was prolonged because of the large tumor size, massive haemorrhage, quantity of resected tissue, and repeated rinsing to clear the visual field. Air was introduced into the bladder during rinsing, resulting in the internal oxygen concentration required for a hydrogen explosion. Hydrogen has an extremely low minimum ignition energy of 0.019 mJ[15]. The electric spark produced by electroexcision using an electric cutting ring or electric coagulation may ignite the mixed gases. The energy released by an explosion will determine the extent of the damage. Bladder rupture occurs when the pressure generated by the explosion exceeds that which can be borne by the bladder wall. Multiple trabeculations and bladder diverticulum have been identified as risk factors for bladder explosion and rupture[16]. Presently, most scholars believe that the flushing solution does not affect bladder explosion, with no reported cases being caused by mannitol, glucose, or normal saline. Additionally, neither intraspinal anaesthesia nor general anaesthesia has any obvious relationship with bladder explosion. However, Hirai *et al*[17] have reported a case of bladder explosion caused by anaesthesia by nitrous oxide (N₂O) inhalation. This was due to the flammable and explosive nature of N₂O in confined spaces (such as the bladder).

In addition to the aforementioned objective contributors, there may also be human factors involved in bladder explosion. These can include the habits, safety awareness, and experience and personality of the surgeon. For instance, some surgeons are used to performing cystostomies before complex TURs. This can certainly increase the risk of bladder tumor dissemination but can avoid the accumulation of gas in the bladder[18]. Some surgeons adhere to the principle that safety is the top priority and adhere rigidly to the operation specifications. They may avoid the introduction of air and carefully observe any gas accumulation in the bladder intraoperatively, ensuring that it is discharged before there is excessive

build-up. Other surgeons may have insufficient surgical experience or poor safety awareness. They may be impatient, make rash decisions during operations, or ignore surgical specifications.

Our literature review suggests a lower incidence of bladder explosions in TUR-BT than in TURP. When removing tumors in the trigone, lateral, and posterior walls of the bladder, keeping an adequate distance from the gas accumulation area of the anterior wall may reduce the risk of bladder explosion. Operations may be disturbed by bubbles when removing tumors from the anterior and apical walls. When this occurs, surgeons usually discharge the gas or change the body position to avoid bubble interference during TUR-BT. However, the prostate has a relatively fixed position in TURP, and removal of the tissue at the 12 o'clock position carries a high risk of disturbing gas accumulated in the anterior wall of the bladder and causing an explosion. Hence, the conditions necessary for bladder explosion are less frequently met during TUR-BT, resulting in a lower incidence.

Special precautions should be taken to avoid bladder explosion during transurethral surgery. First, the risk of explosion can be reduced with the use of low or medium power as this reduces the energy of the ignition source and lowers the local temperature during the electroexcision and electrocoagulation processes. Second, to carefully check for gas mixing in the flushing solution and to minimise the flushing times, maintaining a tight connection on the endoscopic sheath is important. The Ellik should be filled with normal saline during each flushing. Third, despite the inevitable generation of combustible gas, bladder explosion can be avoided by discharging the build-up of gas within the bladder promptly. An appropriate method for achieving this has been proposed in previous research [19]. The procedure is to tilt the resectoscope towards the top of the bladder and align it with the bubbles, close the exhalant canal, unplug the water inlet, and open the water inlet control valve to discharge the bubbles. Fourth, the duration of electroexcision should be kept to a minimum to reduce gas production, and this is best achieved through the adoption of punctate electroexcision and electrocoagulation to avoid blind inch-by-inch electrocoagulation. Fifth, the surgical position can be adjusted according to the intraoperative situation to change the bubble position. In this way, the surgeon can avoid contact between the gas bubbles and the electric spark produced by the electric cutting ring. Lastly, it is critical to be fully conversant with the operation indications and to conduct a thorough assessment of the surgical difficulty and risks. In our case, bladder explosion occurred because TUR of the bladder was an inappropriate treatment given the large tumor size. This led to the generation of a considerable amount of flammable gas because of the duration of the operation and the introduction of a great deal of air during repeated flushing. After this bladder explosion incident, our team conducted a systematic analysis and evaluation to improve operational details and raise surgeons' awareness of the risk of such events. There have been no further intravesical explosions at our institution since this case.

CONCLUSION

Although rare, intravesical explosions can cause serious consequences, and the loud explosion can also lead to a profound psychological shadow on the patient. Urologists must be aware of this potential complication. Careful operative techniques and special precautions can reduce the risk of this complication.

FOOTNOTES

Author contributions: Pan ZS conceived and wrote the manuscript; Xu CB reviewed the manuscript and reviewed the literature; Jia DS supervised the treatment of the patient; and all authors read and approved the final manuscript.

Informed consent statement: Informed written consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Zheng-Sheng Pan [0000-0002-4395-7094](https://orcid.org/0000-0002-4395-7094).

S-Editor: Wang JJ

L-Editor: Wang TQ

P-Editor: Wang JJ

REFERENCES

- 1 **Cassuto A.** Explosion dans la vessie au cours d'une electrocoagulation. *J Urol* 1926; **22**: 263
- 2 **Wang XH,** Qi L, He DL. Consensus on safety of transurethral plasma resection of bladder tumors. *J Modern Urol* 2018; **23**: 895-901 [DOI: [10.3969/j.issn.10098291.2018.12.003](https://doi.org/10.3969/j.issn.10098291.2018.12.003)]
- 3 **Martov AG,** Kornienko SI, Gushchin BL, Ergakov DV, Sazonov OA. [Intraoperative urological complications in transurethral surgical interventions on the prostate for benign hyperplasia]. *Urologiia* 2005; 3-8 [PMID: [16158736](https://pubmed.ncbi.nlm.nih.gov/16158736/)]
- 4 **Liu HQ,** Chen YS, Zou B, Kong J, Tao LS, Zhu GB. Serious complications of transurethral resection of the prostate. *Chin J Urol* 2016; **37**: 515-518 [DOI: [10.3760/cma.j.issn.1000-6702.2016.07.010](https://doi.org/10.3760/cma.j.issn.1000-6702.2016.07.010)]
- 5 **Di Tonno F,** Fusaro V, Bertoldin R, Lavelli D. Bladder explosion during transurethral resection of the prostate. *Urol Int* 2003; **71**: 108-109 [PMID: [12845273](https://pubmed.ncbi.nlm.nih.gov/12845273/) DOI: [10.1159/000071106](https://doi.org/10.1159/000071106)]
- 6 **Dublin N,** Razack AH, Loh CS. Intravesical explosion during transurethral resection of the prostate. *ANZ J Surg* 2001; **71**: 384-385 [PMID: [11409027](https://pubmed.ncbi.nlm.nih.gov/11409027/) DOI: [10.1046/j.1440-1622.2001.02134.x](https://doi.org/10.1046/j.1440-1622.2001.02134.x)]
- 7 **Seitz M,** Soljanik I, Stanislaus P, Sroka R, Stief C. Explosive gas formation during transurethral resection of the prostate (TURP). *Eur J Med Res* 2008; **13**: 399-400 [PMID: [18952523](https://pubmed.ncbi.nlm.nih.gov/18952523/)]
- 8 **Ribeiro da Silva MN,** Lopes Neto AC, Zamboni JP, Verotti MJ, Wroclawski ER. Vesical explosion during transurethral resection of the prostate: report of a case. *Arch Esp Urol* 2006; **59**: 651-652 [PMID: [16933499](https://pubmed.ncbi.nlm.nih.gov/16933499/) DOI: [10.4321/S0004-06142006000600017](https://doi.org/10.4321/S0004-06142006000600017)]
- 9 **Vacchiano G,** Rocca A, Compagna R, Zamboli AGI, Cirillo V, Di Domenico L, Di Nardo V, Servillo G, Amato B. Transurethral Resection of the Prostate, Bladder Explosion and Hyponatremic Encephalopathy: A Rare Case Report of Malpractice. *Open Med (Wars)* 2017; **12**: 50-57 [PMID: [28435905](https://pubmed.ncbi.nlm.nih.gov/28435905/) DOI: [10.1515/med-2017-0010](https://doi.org/10.1515/med-2017-0010)]
- 10 **Georgios K,** Evangelos B, Helai H, Ioannis G. Intravesical explosion during transurethral electrocautery. *Scott Med J* 2015; **60**: e25-e26 [PMID: [25680867](https://pubmed.ncbi.nlm.nih.gov/25680867/) DOI: [10.1177/0036933015572091](https://doi.org/10.1177/0036933015572091)]
- 11 **Bus MT,** Cordeiro ER, Anastasiadis A, Klioueva NM, de la Rosette JJ, de Reijke TM. Urothelial carcinoma in both adnexa following perforation during transurethral resection of a non-muscle-invasive bladder tumor: a case report and literature review. *Expert Rev Anticancer Ther* 2012; **12**: 1529-1536 [PMID: [23253219](https://pubmed.ncbi.nlm.nih.gov/23253219/) DOI: [10.1586/era.12.136](https://doi.org/10.1586/era.12.136)]
- 12 **Liu S,** Liu YH. High frequency electrotome and safe clinical use. *Chin J Pract Sur* 2005; **25**: 383-384
- 13 **Ning TC Jr,** Atkins DM, Murphy RC. Bladder explosions during transurethral surgery. *J Urol* 1975; **114**: 536-539 [PMID: [1242014](https://pubmed.ncbi.nlm.nih.gov/1242014/) DOI: [10.1016/s0022-5347\(17\)67077-0](https://doi.org/10.1016/s0022-5347(17)67077-0)]
- 14 **Davis TR.** The composition and origin of the gas produced during urological endoscopic resections. *Br J Urol* 1983; **55**: 294-297 [PMID: [6850245](https://pubmed.ncbi.nlm.nih.gov/6850245/) DOI: [10.1111/j.1464-410X.1983.tb03301.x](https://doi.org/10.1111/j.1464-410X.1983.tb03301.x)]
- 15 **He LQ.** Gas combustion. In: *Fire combustion science*. Beijing: China Machine Press, 2014: 129-143
- 16 **Vincent DP.** Bladder explosion during transurethral resection of prostate: Bladder diverticula as an additional risk factor. *Urol Ann* 2017; **9**: 68-70 [PMID: [28216933](https://pubmed.ncbi.nlm.nih.gov/28216933/) DOI: [10.4103/0974-7796.198887](https://doi.org/10.4103/0974-7796.198887)]
- 17 **Hirai E,** Tokumine J, Lefor AK, Ogura S, Kawamata M. Bladder Explosion during Transurethral Resection of the Prostate with Nitrous Oxide Inhalation. *Case Rep Anesthesiol* 2015; **2015**: 464562 [PMID: [26294981](https://pubmed.ncbi.nlm.nih.gov/26294981/) DOI: [10.1155/2015/464562](https://doi.org/10.1155/2015/464562)]
- 18 **Chang SS,** Boorjian SA, Chou R, Clark PE, Daneshmand S, Konety BR, Pruthi R, Quale DZ, Ritch CR, Seigne JD, Skinner EC, Smith ND, McKiernan JM. Diagnosis and Treatment of Non-Muscle Invasive Bladder Cancer: AUA/SUO Guideline. *J Urol* 2016; **196**: 1021-1029 [PMID: [27317986](https://pubmed.ncbi.nlm.nih.gov/27317986/) DOI: [10.1016/j.juro.2016.06.049](https://doi.org/10.1016/j.juro.2016.06.049)]
- 19 **Takeshita H,** Moriyama S, Chiba K, Noro A. A simple technique for evacuating air bubbles with scum from the bladder dome during transurethral resection of bladder tumor. *Wideochir Inne Tech Maloinwazyjne* 2014; **9**: 619-622 [PMID: [25562002](https://pubmed.ncbi.nlm.nih.gov/25562002/) DOI: [10.5114/wiitm.2014.47092](https://doi.org/10.5114/wiitm.2014.47092)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

