

World Journal of *Clinical Cases*

World J Clin Cases 2022 November 6; 10(31): 11214-11664



REVIEW

- 11214** Diabetes and skin cancers: Risk factors, molecular mechanisms and impact on prognosis
Dobrică EC, Banciu ML, Kipkorir V, Khazeei Tabari MA, Cox MJ, Simhachalam Kutikuppala LV, Găman MA
- 11226** Endocrine disruptor chemicals as obesogen and diabetogen: Clinical and mechanistic evidence
Kurşunoğlu NE, Sarer Yurekli BP
- 11240** Intestinal microbiota in the treatment of metabolically associated fatty liver disease
Wang JS, Liu JC

MINIREVIEWS

- 11252** Lactation mastitis: Promising alternative indicators for early diagnosis
Huang Q, Zheng XM, Zhang ML, Ning P, Wu MJ
- 11260** Clinical challenges of glycemic control in the intensive care unit: A narrative review
Sreedharan R, Martini A, Das G, Aftab N, Khanna S, Ruetzler K
- 11273** Concise review on short bowel syndrome: Etiology, pathophysiology, and management
Lakkasani S, Seth D, Khokhar I, Touza M, Dacosta TJ
- 11283** Role of nickel-regulated small RNA in modulation of *Helicobacter pylori* virulence factors
Freire de Melo F, Marques HS, Fellipe Bueno Lemos F, Silva Luz M, Rocha Pinheiro SL, de Carvalho LS, Souza CL, Oliveira MV
- 11292** Surgical intervention for acute pancreatitis in the COVID-19 era
Su YJ, Chen TH

ORIGINAL ARTICLE**Clinical and Translational Research**

- 11299** Screening of traditional Chinese medicine monomers as ribonucleotide reductase M2 inhibitors for tumor treatment
Qin YY, Feng S, Zhang XD, Peng B

Case Control Study

- 11313** Covered transjugular intrahepatic portosystemic stent-shunt *vs* large volume paracentesis in patients with cirrhosis: A real-world propensity score-matched study
Dhaliwal A, Merhzad H, Karkhanis S, Tripathi D

Retrospective Cohort Study

- 11325** Endoscopic submucosal tunnel dissection for early esophageal squamous cell carcinoma in patients with cirrhosis: A propensity score analysis
Zhu LL, Liu LX, Wu JC, Gan T, Yang JL

Retrospective Study

- 11338** Nomogram for predicting overall survival in Chinese triple-negative breast cancer patients after surgery
Lin WX, Xie YN, Chen YK, Cai JH, Zou J, Zheng JH, Liu YY, Li ZY, Chen YX
- 11349** Early patellar tendon rupture after total knee arthroplasty: A direct repair method
Li TJ, Sun JY, Du YQ, Shen JM, Zhang BH, Zhou YG
- 11358** Coxsackievirus A6 was the most common enterovirus serotype causing hand, foot, and mouth disease in Shiyan City, central China
Li JF, Zhang CJ, Li YW, Li C, Zhang SC, Wang SS, Jiang Y, Luo XB, Liao XJ, Wu SX, Lin L
- 11371** Dynamic changes of estimated glomerular filtration rate are conversely related to triglyceride in non-overweight patients
Liu SQ, Zhang XJ, Xue Y, Huang R, Wang J, Wu C, He YS, Pan YR, Liu LG
- 11381** C-reactive protein as a non-linear predictor of prolonged length of intensive care unit stay after gastrointestinal cancer surgery
Yan YM, Gao J, Jin PL, Lu JJ, Yu ZH, Hu Y

Clinical Trials Study

- 11391** Dan Bai Xiao Formula combined with glucocorticoids and cyclophosphamide for pediatric lupus nephritis: A pilot prospective study
Cao TT, Chen L, Zhen XF, Zhao GJ, Zhang HF, Hu Y

Observational Study

- 11403** Relationship between lipids and sleep apnea: Mendelian randomization analysis
Zhang LP, Zhang XX
- 11411** Efficacy and safety profile of two-dose SARS-CoV-2 vaccines in cancer patients: An observational study in China
Cai SW, Chen JY, Wan R, Pan DJ, Yang WL, Zhou RG

Prospective Study

- 11419** Pressure changes in tapered and cylindrical shaped cuff after extension of head and neck: A randomized controlled trial
Seol G, Jin J, Oh J, Byun SH, Jeon Y

Randomized Controlled Trial

- 11427** Effect of intradermal needle therapy at combined acupoints on patients' gastrointestinal function following surgery for gastrointestinal tumors
Guo M, Wang M, Chen LL, Wei FJ, Li JE, Lu QX, Zhang L, Yang HX

SYSTEMATIC REVIEWS

- 11442** Video-assisted bystander cardiopulmonary resuscitation improves the quality of chest compressions during simulated cardiac arrests: A systemic review and meta-analysis
Pan DF, Li ZJ, Ji XZ, Yang LT, Liang PF

META-ANALYSIS

- 11454** Efficacy of the femoral neck system in femoral neck fracture treatment in adults: A systematic review and meta-analysis
Wu ZF, Luo ZH, Hu LC, Luo YW
- 11466** Prevalence of polymyxin-induced nephrotoxicity and its predictors in critically ill adult patients: A meta-analysis
Wang JL, Xiang BX, Song XL, Que RM, Zuo XC, Xie YL

CASE REPORT

- 11486** Novel compound heterozygous variants in the LHX3 gene caused combined pituitary hormone deficiency: A case report
Lin SZ, Ma QJ, Pang QM, Chen QD, Wang WQ, Li JY, Zhang SL
- 11493** Fatal bleeding due to an aorto-esophageal fistula: A case report and literature review
Ćeranić D, Nikolić S, Lučev J, Slanić A, Bujas T, Ocepek A, Skok P
- 11500** Tolvaptan ameliorated kidney function for one elderly autosomal dominant polycystic kidney disease patient: A case report
Zhou L, Tian Y, Ma L, Li WG
- 11508** Extensive right coronary artery thrombosis in a patient with COVID-19: A case report
Dall'Orto CC, Lopes RPF, Cancela MT, de Sales Padilha C, Pinto Filho GV, da Silva MR
- 11517** Yokoyama procedure for a woman with heavy eye syndrome who underwent multiple recession-resection operations: A case report
Yao Z, Jiang WL, Yang X
- 11523** Rectal cancer combined with abdominal tuberculosis: A case report
Liu PG, Chen XF, Feng PF
- 11529** Malignant obstruction in the ileocecal region treated by self-expandable stent placement under the fluoroscopic guidance: A case report
Wu Y, Li X, Xiong F, Bao WD, Dai YZ, Yue LJ, Liu Y
- 11536** Granulocytic sarcoma with long spinal cord compression: A case report
Shao YD, Wang XH, Sun L, Cui XG
- 11542** Aortic dissection with epileptic seizure: A case report
Zheng B, Huang XQ, Chen Z, Wang J, Gu GF, Luo XJ

- 11549** Multiple bilateral and symmetric C1-2 ganglioneuromas: A case report
Wang S, Ma JX, Zheng L, Sun ST, Xiang LB, Chen Y
- 11555** Acute myocardial infarction due to Kounis syndrome: A case report
Xu GZ, Wang G
- 11561** Surgical excision of a large retroperitoneal lymphangioma: A case report
Park JH, Lee D, Maeng YH, Chang WB
- 11567** Mass-like extragonadal endometriosis associated malignant transformation in the pelvis: A rare case report
Chen P, Deng Y, Wang QQ, Xu HW
- 11574** Gastric ulcer treated using an elastic traction ring combined with clip: A case report
Pang F, Song YJ, Sikong YH, Zhang AJ, Zuo XL, Li RY
- 11579** Novel liver vein deprivation technique that promotes increased residual liver volume (with video): A case report
Wu G, Jiang JP, Cheng DH, Yang C, Liao DX, Liao YB, Lau WY, Zhang Y
- 11585** Linear porokeratosis of the foot with dermoscopic manifestations: A case report
Yang J, Du YQ, Fang XY, Li B, Xi ZQ, Feng WL
- 11590** Primary hepatic angiosarcoma: A case report
Wang J, Sun LT
- 11597** Hemorrhagic shock due to ruptured lower limb vascular malformation in a neurofibromatosis type 1 patient: A case report
Shen LP, Jin G, Zhu RT, Jiang HT
- 11607** Gastric linitis plastica with autoimmune pancreatitis diagnosed by an endoscopic ultrasonography-guided fine-needle biopsy: A case report
Sato R, Matsumoto K, Kanzaki H, Matsumi A, Miyamoto K, Morimoto K, Terasawa H, Fujii Y, Yamazaki T, Uchida D, Tsutsumi K, Horiguchi S, Kato H
- 11617** Favorable response of primary pulmonary lymphoepithelioma-like carcinoma to sintilimab combined with chemotherapy: A case report
Zeng SY, Yuan J, Lv M
- 11625** Benign paroxysmal positional vertigo with congenital nystagmus: A case report
Li GF, Wang YT, Lu XG, Liu M, Liu CB, Wang CH
- 11630** Secondary craniofacial necrotizing fasciitis from a distant septic emboli: A case report
Lee DW, Kwak SH, Choi HJ
- 11638** Pancreatic paraganglioma with multiple lymph node metastases found by spectral computed tomography: A case report and review of the literature
Li T, Yi RQ, Xie G, Wang DN, Ren YT, Li K

- 11646** Apnea caused by retrobulbar anesthesia: A case report
Wang YL, Lan GR, Zou X, Wang EQ, Dai RP, Chen YX
- 11652** Unexplained septic shock after colonoscopy with polyethylene glycol preparation in a young adult: A case report
Song JJ, Wu CJ, Dong YY, Ma C, Gu Q
- 11658** Metachronous isolated penile metastasis from sigmoid colon adenocarcinoma: A case report
Yin GL, Zhu JB, Fu CL, Ding RL, Zhang JM, Lin Q

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Muhammad Hamdan Gul, MD, Assistant Professor, Department of Internal Medicine, University of Kentucky, Chicago, IL 60657, United States.
hamdan3802@hotmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for *WJCC* as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The *WJCC*'s CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Xu Guo*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

November 6, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Lactation mastitis: Promising alternative indicators for early diagnosis

Qian Huang, Xue-Mei Zheng, Mao-Lin Zhang, Ping Ning, Meng-Jun Wu

Specialty type: Medicine, general and internal

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Bairwa DBL, India; Mahmoud MZ, Saudi Arabia

Received: May 9, 2022

Peer-review started: May 9, 2022

First decision: August 21, 2022

Revised: August 31, 2022

Accepted: September 22, 2022

Article in press: September 22, 2022

Published online: November 6, 2022



Qian Huang, Xue-Mei Zheng, School of Medicine, University of Electronic Science and Technology of China, Chengdu 610000, Sichuan Province, China

Mao-Lin Zhang, Department of Anesthesiology, Chongqing Medical University, Chongqing 400016, Sichuan Province, China

Ping Ning, Department of Breast, Chengdu Women's and Children's Central Hospital, Chengdu 610000, Sichuan Province, China

Meng-Jun Wu, Department of Anesthesiology, Chengdu Women's and Children's Central Hospital, Chengdu 610000, Sichuan Province, China

Corresponding author: Ping Ning, MD, Doctor, Department of Breast, Chengdu Women's and Children's Central Hospital, No. 1617 Riyue Avenue, Qingyang District, Chengdu 610000, Sichuan Province, China. nping11@163.com

Abstract

Although lactation mastitis (LM) has been extensively researched, the incidence rate of LM remains a salient clinical problem. To reduce this incidence rate and achieve a better prognosis, early and specific quantitative indicators are particularly important. It has been found that milk electrolyte concentrations (chloride, potassium, and sodium) and electrical conductivity (EC) significantly change in the early stages of LM in an animal model. Several studies have evaluated EC for the detection of subclinical mastitis in cows. EC, chloride, and sodium content of milk were more accurate for predicting infection status than were other variables. In the early stages of LM, lactic sodium, chloride, and EC increase, but potassium decreases. However, these indicators have not been reported in the diagnosis of LM in humans. This review summarizes the pathogenesis and the mechanism of LM in terms of milk electrolyte concentration and EC, and aim to provide new ideas for the detection of sub-clinical mastitis in humans.

Key Words: Lactation mastitis; Milk electrolyte concentration; Milk electrical conductivity; Pathogenesis; Mechanism; Early diagnosis

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: It has been found that milk electrolyte concentrations and electrical conductivity (EC) significantly change in the early stages of lactation mastitis (LM) in an animal model, allowing the early and specific diagnosis of LM. These indicators have not been reported in the diagnosis of LM in humans. We summarize the pathogenesis and the mechanism of LM in terms of milk electrolyte concentrations and EC and aim to provide new ideas for the early diagnosis of LM in humans.

Citation: Huang Q, Zheng XM, Zhang ML, Ning P, Wu MJ. Lactation mastitis: Promising alternative indicators for early diagnosis. *World J Clin Cases* 2022; 10(31): 11252-11259

URL: <https://www.wjgnet.com/2307-8960/full/v10/i31/11252.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i31.11252>


INTRODUCTION

Lactation mastitis (LM), which is universally described as a suite of breast conditions that present with local, and often systemic, inflammatory symptoms and signs during lactation[1]. It is an enervating and common disease that affects up to 33% of lactating women[2-4]. LM has a negative impact on both the mother and the baby. Women with LM may develop pain, localized skin redness, and can have associated systemic symptoms, including fever. However, breast pain is the most common and the most distressing symptom for mothers[5]. The symptoms of LM can lead to a compromised maternal psychological state. In addition, approximately 3% of women with LM will develop a breast abscess[6], and an incidence rate of up to 11% has been reported[7], which may cause permanent damage to the shape of the breast. In order to minimize the detrimental effects of LM, researchers have explored various strategies, including the associated risk factors and etiology[8,9]. However, none of these are known to improve clinical prognosis.

To date, the clinical diagnosis of LM has mostly relied on empirical diagnoses, such as a tender, hot, swollen, wedge-shaped area of redness on the affected breast which is associated with an elevated temperature and systemic symptoms, such as a temperature of 38.5 °C or higher, chills, flu-like aching, and systemic illness[10,11]. Due to the lack of early standardized diagnostic criteria, this causes repeated outbreaks and even progresses to a breast abscess, causing great physical and mental pain in mothers with LM. In some Western countries, for example, 73% of children born in Sweden in 1996 were breastfed for 6 mo[12], but LM frequently results in the cessation of exclusive breastfeeding in the absence of appropriate treatment. As a result, it frequently reduces the protective effect of breastfeeding in mothers. For example, related data show that breastfeeding is associated with a 24% lower risk of invasive ovarian cancer[13], and aggregate results indicate that breastfeeding is inversely associated with the risk of breast cancer[14] and reduces the incidence of osteoporosis and type 2 diabetes[15]. Furthermore, LM is linked to lower levels of fat, carbohydrate, and energy in breast milk[16], which may lead to nutritional deficiencies, lowered immunity, and mental effects in infants.

Several studies have evaluated electrical conductivity (EC) for the detection of subclinical mastitis in cows, some have even identified mastitis causing pathogens using EC[17,18]. EC, chloride (Cl⁻), and sodium (Na⁺) content in milk were more accurate in predicting infection status than other variables. The electrolyte concentration and EC of milk are the physicochemical properties of milk, and EC measurements were used as an experimental screening indicator for LM in animals as early as the 1990s [19]. Paudyal *et al*[18] showed that absolute changes in cow's milk electrolyte concentrations and EC can be used to screen breast milk samples for LM. A previous study provided evidence that when an inflammatory response occurs in the breast tissue of animals, breast permeability increases, and as a result, the potassium (K⁺) concentration reduces, but the Na⁺, Cl⁻ concentrations, and EC increase[20-22]. Singh *et al*[23] recently conducted an observational animal study and concluded that the magnitude of changes in the milk electrolyte concentration and EC may have diagnostic and prognostic values. Due to a lack of research, more data are needed to identify the variation in human milk electrolyte concentration and EC during LM.

However, it is uncertain whether human milk electrolyte concentrations and EC have a positive effect on the early diagnosis of LM. To enable more patients with LM to be effectively diagnosed and treated at an early stage of the disease, the pathogenesis of LM and the relationship between LM and electrolyte concentrations and EC (Figure 1) are discussed in this review. We investigate the changes in electrolyte concentrations and EC from two perspectives: Altered cell membrane permeability and osmotic pressure level. This will provide new ideas for the detection of subclinical mastitis in humans.



DOI: 10.12998/wjcc.v10.i31.11252 Copyright ©The Author(s) 2022.

Figure 1 The mechanism of lactation mastitis about milk electrolyte concentration and electrical conductivity in animals. DAMPs: Danger-associated molecular patterns; LPS: Lipopolysaccharide; MAMPs: Microbe-associated molecular patterns; TLR: Toll-like receptors; EC: Electrical conductivity.

LITERATURE SEARCH

A descriptive review was conducted on the mechanism of changes in milk electrolyte concentration and EC during LM. Pub-Med was searched for articles published between July 1966 and February 2022. The following Medical Subject Headings or free-text terms were used in the search: LM, milk electrolyte concentration, milk EC, early diagnosis, pathogenesis, mechanism, inflammatory injury, and altered cell membrane permeability. The search was limited to papers written in English, with no restrictions on the type of article.

PATHOGENESIS OF LM

The susceptibility and severity of LM are positively correlated with inflammatory factors. Some researchers consider LM to be an infectious disorder[19,24,25], but the etiology of LM has now shifted from infection to inflammation.

Infectious mechanism

Due to the characteristics of LM, it was previously considered to be an infectious disease. Based on the maturation of milk culture technology, some infectious pathogenic bacteria, such as *Staphylococcus aureus*, *Escherichia coli*, and *Streptococcus*, have all been isolated from breast milk, confirming the importance of infective pathogenic bacteria in the pathogenesis of LM[26,27]. In recent years, *Staphylococcus epidermidis*, the most common species on human skin and mucosa, has also been isolated from the milk of patients with LM and may be another causative agent of LM[28-30]. Based on the fact that a variety of infectious pathogens are present in the milk of patients with LM, researchers believed that the occurrence of LM was closely related to infectious pathogens in the milk[31]. Milk stagnation, breast trauma, excessive emptying of the breast, nipple cracking, and dysbiosis of the breast flora can all contribute to the development of infectious LM.

Inflammatory mechanism

Recently, in-depth studies on LM have revealed that the presence of infectious agents is not positively associated with the occurrence of LM[26]. In addition, there is no positive correlation between the severity of LM and the bacterial count in milk[32]. Several studies showed that the susceptibility and severity of LM are positively correlated with inflammatory factors, including C-reactive protein, interleukin (IL)-1, IL-6, IL-8, and tumor necrosis factor- α (TNF- α)[33-35]. Elevated serum cytokines IL-1, IL-6, IL-8, and TNF- α reveal the activation of transcription factor nuclear factor-kappa B (NF- κ B) in the host[36]. Furthermore, activation of the NF- κ B pathway in LM hosts has been demonstrated in numerous animal studies, *in vitro* experiments, and genetic studies[37-40].

As upstream candidate receptors of NF- κ B, Toll-like receptors (TLRs) are important inflammatory mediators and their isoforms include TLR1-TLR11[41,42], which are recognized as pattern recognition receptors (PRRs) by microbe-associated molecular patterns (MAMPs) or danger-associated molecular patterns (DAMPs) and activate downstream NF- κ B signaling pathways[34].

MAMP-mediated TLR signaling pathway

Of all the TLRs, TLR2, TLR4, and TLR5 bind to bacterial byproducts in MAMPs, such as lipopolysaccharide, phospholipid wall acids, bacterial lipopeptides, and flagellin, activate the corresponding TLRs, which activate transcription, translation, and the release of a series of inflammatory factors, chemokines, and adhesion molecules, and recruit other molecules involved in the innate immune response (neutrophils, *etc.*) to the site of infection[33,34]. Pathogenic byproducts bind to TLRs on the surface of PRRs, and the signal is transmitted from extracellular to intracellular, activating TLRs, followed by signal transduction, and activation of the transcription factor NF- κ B. This is the pathogenesis of infectious LM *via* the inflammatory mechanism.

DAMP-mediated TLR signaling pathway

TLR activation *via* the DAMP pathway explains how LM can lead to disease in the absence of infectious agents. DAMPs are endogenous proteins and can activate TLRs in a sterile environment[43,44]. DAMPs can activate TLRs in two ways: first, some inflammatory mediators activated by DAMPs activate TLRs and downstream NF- κ B[45]; second, DAMPs can enhance the susceptibility of the TLR immune response in a sterile environment, thus activating TLRs and downstream NF- κ B leading to the development of noninfectious LM[46].

MECHANISM OF LM-INDUCED CHANGES IN MILK ELECTROLYTE CONCENTRATION AND EC

Mechanism of cell membrane permeability alteration

Changes in milk electrolyte concentration and EC are linked to the inflammatory response during LM, and pathological changes in breast tissues caused by inflammatory factors result in changes in milk composition. Our understanding of the changes in milk electrolyte concentration and EC during LM is linked to the research by Smith *et al*[47] in the 1960s. The essence of LM is the inflammatory response of the breast tissue[48], and inflammatory factors cause increased epithelial permeability and both vascular and parenchyma damage[20].

The sodium pump on the basolateral membrane, which keeps intracellular K⁺ high and Na⁺ low, and the distribution of these ions according to the gradient of electric potential across the luminal membranes are the most important characteristics in terms of Na⁺ and K⁺[49]. The Na⁺/K⁺ ratio of the intracellular fluid is maintained at approximately 1:3 and the ratio of the extracellular fluid is maintained at approximately 3:1 by active transport of the sodium-potassium ion pump. As milk is electrically positive compared to the interior of the cell, the concentrations of these ions are lower in milk. However, because milk is nearly isosmotic to plasma[50], the ratio between them is similar, with a Na⁺/K⁺ ratio of about 1:3[51,52].

When an inflammatory response occurs in the breast tissue, all of the inflammatory factors produced damage the ducts and secretory epithelial cells, disrupt the tight junctions between secretory cells, and increase capillary permeability. As a result, higher levels of Na⁺ and Cl⁻ in the extracellular fluid enter the mammary gland alveolar lumen through the tight junctions that are opened, while the K⁺ concentration decreases in order to maintain the osmotic pressure of milk in the alveolar lumen[53]. The increase in ionic concentrations in breast milk in the presence of mastitis, such as Na⁺ and Cl⁻, is consistent with other animal studies[54-56].

Mechanism of the milk osmolality level

Milk is rich in lactose. Lactose excretion may provide a reliable basis for fateful changes in breast permeability as lactose in food contributes minimally to the circulation[54]. Fetherston *et al*[57] conducted relevant studies, and the 24-h urinary excretion of lactose during LM has been extensively discussed. The correspondingly low steady-state urinary excretion of lactose demonstrates that these variations are not the result of increased paracellular pathway permeability. It indicates that the higher than normal concentrations of milk Na⁺ and Cl⁻ observed in "normal breasts" could be a normal physiologic response to a lower concentration of lactose, ensuring that the osmolality of milk remains isotonic with plasma[20]. This has significant implications for the supposition that a raised Na⁺ concentration is an outcome of subclinical LM.

ADDITIONAL DIAGNOSTIC PROSPECTS: MILK ELECTROLYTE CONCENTRATION AND EC

The electrolyte concentration and EC are physical properties of body fluids that are used to assess human health status and disease severity[58,59]. Previous animal studies have indicated that they can be used as an alternative method for the early diagnosis of LM[60]. In addition, Kitchen *et al*[61] demonstrated that the measurements of electrolyte concentration and EC in milk samples for the diagnosis of LM in animals were comparable to other diagnostic methods.

Similar to the pathological response in animals with LM[62,63], when a woman is diagnosed with LM, inflammatory factors, such as procalcitonin, IL-1, IL-6, IL-8, and TNF- α [36], cause disruption of the tight junctions between cells. Furthermore, as a result of both the decrease in available glucose, particularly during severe symptoms, and the damage or death of lactocytes due to inflammation, lactose synthesis decreases[20]. Both of these factors can lead to changes in the electrolyte concentration and EC of milk. As mentioned earlier, numerous animal studies have indicated that milk electrolyte concentration and EC can be used in the early diagnosis of LM; therefore, based on the similar pathological responses, it is feasible to use milk electrolyte concentration and EC to diagnose early LM in humans.

In a recent study, we collected milk specimens from approximately 30 healthy women and 15 patients diagnosed with LM, and measured the electrolyte concentration (including Na⁺, K⁺, and Cl⁻) and EC of bilateral breast milk in all patients. The final test results revealed that the concentrations of Na⁺, Cl⁻ and EC in breast milk were significantly higher in women with LM than in healthy women, and in some cases several times higher. In addition, in women with LM, the concentrations of Na⁺, Cl⁻ and EC in breast milk were markedly higher on the affected side than on the healthy side. Furthermore, we also found that the concentrations of Na⁺, Cl⁻ and EC in milk were increased to varying degrees in patients who had symptoms but were not diagnosed with LM.

CONCLUSION

The early diagnosis and prevention of LM still face many challenges. We summarized the pathogenesis of LM, the mechanism of LM-induced changes in milk electrolyte concentration and EC, and found that changes in milk electrolyte concentration and EC in humans were primarily correlated with LM. It is clear from animal studies that there is a correlation between LM and milk electrolyte concentration and EC, and from the significant changes in these indicators, LM can be diagnosed at an early stage and thus achieve a better prognosis. However, there have been few studies carried out on this topic. As a result, more data are required to verify these findings. If the changes in milk electrolyte concentration and EC are beneficial, these will have an enormous impact on clinical practice.

FOOTNOTES

Author contributions: Huang Q, Zheng XM and Zhang ML contributed equally to this work; Ning P, Wu MJ designed the research study; Huang Q, Zheng XM collected data from the literature; Huang Q, Zheng XM and Zhang ML wrote the manuscript; Ning P, Wu MJ conducted a review of the manuscript; all authors have read and approve the final manuscript.

Conflict-of-interest statement: The authors have no potential conflicts of interest to disclose.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Qian Huang 0000-0002-3064-8604; Xue-Mei Zheng 0000-0002-5380-5457; Mao-Lin Zhang 0000-0003-3366-4156; Ping Ning 0000-0001-5799-7030; Meng-Jun Wu 0000-0002-6899-4318.

S-Editor: Chang KL

L-Editor: A

P-Editor: Chang KL

REFERENCES

- 1 **Cooper M**, Lowe H, McArdle A. Development of a novel patient focussed symptom severity index for use in assessing and treating inflammatory conditions of the lactating breast: a Delphi study. *Int J Evid Based Healthc* 2020; **18**: 231-240 [PMID: 32487966 DOI: 10.1097/XEB.000000000000225]
- 2 **Angelopoulou A**, Field D, Ryan CA, Stanton C, Hill C, Ross RP. The microbiology and treatment of human mastitis. *Med Microbiol Immunol* 2018; **207**: 83-94 [PMID: 29350290 DOI: 10.1007/s00430-017-0532-z]
- 3 **Lai BY**, Yu BW, Chu AJ, Liang SB, Jia LY, Liu JP, Fan YY, Pei XH. Risk factors for lactation mastitis in China: A systematic review and meta-analysis. *PLoS One* 2021; **16**: e0251182 [PMID: 33983987 DOI: 10.1371/journal.pone.0251182]
- 4 **Barker M**, Adelson P, Peters MDJ, Steen M. Probiotics and human lactational mastitis: A scoping review. *Women Birth* 2020; **33**: e483-e491 [PMID: 32146088 DOI: 10.1016/j.wombi.2020.01.001]
- 5 **Altintoprak F**, Kivilcim T, Ozkan OV. Aetiology of idiopathic granulomatous mastitis. *World J Clin Cases* 2014; **2**: 852-858 [PMID: 25516860 DOI: 10.12998/wjcc.v2.i12.852]
- 6 **O'Brien C**, Quinn E, Murphy M, Lehane E, O'Leary DP, Livingstone V, Paul Redmond H, Corrigan MA. Breast abscess: Not just a puerperal problem. *Breast J* 2020; **26**: 339-342 [PMID: 31544305 DOI: 10.1111/tbj.13586]
- 7 **Amir LH**, Forster D, McLachlan H, Lumley J. Incidence of breast abscess in lactating women: report from an Australian cohort. *BJOG* 2004; **111**: 1378-1381 [PMID: 15663122 DOI: 10.1111/j.1471-0528.2004.00272.x]
- 8 **Kvist LJ**, Rydhstroem H. Factors related to breast abscess after delivery: a population-based study. *BJOG* 2005; **112**: 1070-1074 [PMID: 16045520 DOI: 10.1111/j.1471-0528.2005.00659.x]
- 9 **Branch-Elliman W**, Golen TH, Gold HS, Yassa DS, Baldini LM, Wright SB. Risk factors for Staphylococcus aureus postpartum breast abscess. *Clin Infect Dis* 2012; **54**: 71-77 [PMID: 22052894 DOI: 10.1093/cid/cir751]
- 10 **Cooney F**, Petty-Saphon N. The Burden of Severe Lactational Mastitis in Ireland from 2006 to 2015. *Ir Med J* 2019; **112**: 855 [PMID: 30719896]
- 11 **Amir LH**; Academy of Breastfeeding Medicine Protocol Committee. ABM clinical protocol #4: Mastitis, revised March 2014. *Breastfeed Med* 2014; **9**: 239-243 [PMID: 24911394 DOI: 10.1089/bfm.2014.9984]
- 12 **Brown CR**, Dodds L, Legge A, Bryanton J, Semenic S. Factors influencing the reasons why mothers stop breastfeeding. *Can J Public Health* 2014; **105**: e179-e185 [PMID: 25165836 DOI: 10.17269/cjph.105.4244]
- 13 **Babic A**, Sasamoto N, Rosner BA, Tworoger SS, Jordan SJ, Risch HA, Harris HR, Rossing MA, Doherty JA, Fortner RT, Chang-Claude J, Goodman MT, Thompson PJ, Moysich KB, Ness RB, Kjaer SK, Jensen A, Schildkraut JM, Titus LJ, Cramer DW, Bandera EV, Qin B, Sieh W, McGuire V, Sutphen R, Pearce CL, Wu AH, Pike M, Webb PM, Modugno F, Terry KL. Association Between Breastfeeding and Ovarian Cancer Risk. *JAMA Oncol* 2020; **6**: e200421 [PMID: 32239218 DOI: 10.1001/jamaoncol.2020.0421]
- 14 **Zhou Y**, Chen J, Li Q, Huang W, Lan H, Jiang H. Association between breastfeeding and breast cancer risk: evidence from a meta-analysis. *Breastfeed Med* 2015; **10**: 175-182 [PMID: 25785349 DOI: 10.1089/bfm.2014.0141]
- 15 **Aune D**, Norat T, Romundstad P, Vatten LJ. Breastfeeding and the maternal risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies. *Nutr Metab Cardiovasc Dis* 2014; **24**: 107-115 [PMID: 24439841 DOI: 10.1016/j.numecd.2013.10.028]
- 16 **Say B**, Dizdar EA, Degirmencioglu H, Uras N, Sari FN, Oguz S, Canpolat FE. The effect of lactational mastitis on the macronutrient content of breast milk. *Early Hum Dev* 2016; **98**: 7-9 [PMID: 27341630 DOI: 10.1016/j.earlhumdev.2016.03.009]
- 17 **Musser JM**, Anderson KL, Caballero M, Amaya D, Maroto-Puga J. Evaluation of a hand-held electrical conductivity meter for detection of subclinical mastitis in cattle. *Am J Vet Res* 1998; **59**: 1087-1091 [PMID: 9736381]
- 18 **Paudyal S**, Melendez P, Manriquez D, Velasquez-Munoz A, Pena G, Roman-Muniz IN, Pinedo PJ. Use of milk electrical conductivity for the differentiation of mastitis causing pathogens in Holstein cows. *Animal* 2020; **14**: 588-596 [PMID: 31583992 DOI: 10.1017/S1751731119002210]
- 19 **Osterman KL**, Rahm VA. Lactation mastitis: bacterial cultivation of breast milk, symptoms, treatment, and outcome. *J Hum Lact* 2000; **16**: 297-302 [PMID: 11155607 DOI: 10.1177/089033440001600405]
- 20 **Fetherston CM**, Lai CT, Hartmann PE. Relationships between symptoms and changes in breast physiology during lactation mastitis. *Breastfeed Med* 2006; **1**: 136-145 [PMID: 17661590 DOI: 10.1089/bfm.2006.1.136]
- 21 **Bruckmaier RM**, Weiss D, Wiedemann M, Schmitz S, Wendl G. Changes of physicochemical indicators during mastitis and the effects of milk ejection on their sensitivity. *J Dairy Res* 2004; **71**: 316-321 [PMID: 15354578 DOI: 10.1017/s0022029904000366]
- 22 **Pyörälä S**. Indicators of inflammation in the diagnosis of mastitis. *Vet Res* 2003; **34**: 565-578 [PMID: 14556695 DOI: 10.1051/vetres:2003026]
- 23 **Singh M**, Yadav P, Sharma A, Garg VK, Mittal D. Estimation of Mineral and Trace Element Profile in Bubaline Milk Affected with Subclinical Mastitis. *Biol Trace Elem Res* 2017; **176**: 305-310 [PMID: 27631328 DOI: 10.1007/s12011-016-0842-9]
- 24 **Jonsson S**, Pulkkinen MO. Mastitis today: incidence, prevention and treatment. *Ann Chir Gynaecol Suppl* 1994; **208**: 84-87 [PMID: 8092782]
- 25 **Riordan JM**, Nichols FH. A descriptive study of lactation mastitis in long-term breastfeeding women. *J Hum Lact* 1990; **6**: 53-58 [PMID: 2346600 DOI: 10.1177/089033449000600213]
- 26 **Kvist LJ**, Larsson BW, Hall-Lord ML, Steen A, Schalén C. The role of bacteria in lactational mastitis and some considerations of the use of antibiotic treatment. *Int Breastfeed J* 2008; **3**: 6 [PMID: 18394188 DOI: 10.1186/1746-4358-3-6]
- 27 **Rimoldi SG**, Pileri P, Mazzocco MI, Romeri F, Bestetti G, Calvagna N, Tonielli C, Fiori L, Gigantiello A, Pagani C, Magistrelli P, Sartani A, De Silvestri A, Gismondo MR, Cetin I. The Role of *Staphylococcus aureus* in Mastitis : A Multidisciplinary Working Group Experience. *J Hum Lact* 2020; **36**: 503-509 [PMID: 31593644 DOI: 10.1177/0890334419876272]

- 28 **Berens P**, Swaim L, Peterson B. Incidence of methicillin-resistant *Staphylococcus aureus* in postpartum breast abscesses. *Breastfeed Med* 2010; **5**: 113-115 [PMID: 20113200 DOI: 10.1089/bfm.2009.0030]
- 29 **Lasagno M**, Ortiz M, Vissio C, Yaciuk R, Bonetto C, Pellegrino M, Bogni C, Odierno L, Raspanti C. Pathogenesis and inflammatory response in experimental caprine mastitis due to *Staphylococcus chromogenes*. *Microb Pathog* 2018; **116**: 146-152 [PMID: 29360565 DOI: 10.1016/j.micpath.2018.01.031]
- 30 **Rodvold KA**, McConeghy KW. Methicillin-resistant *Staphylococcus aureus* therapy: past, present, and future. *Clin Infect Dis* 2014; **58** Suppl 1: S20-S27 [PMID: 24343828 DOI: 10.1093/cid/cit614]
- 31 **Shuyang X**, Qiang Y. Bacterial factors of mastitis in lactating women and its effect on the physical properties and chemical composition of breast milk. *Cell Mol Biol (Noisy-le-grand)* 2021; **67**: 172-177 [PMID: 34933713 DOI: 10.14715/cmb/2021.67.3.27]
- 32 **Fetherston C**. Mastitis in lactating women: physiology or pathology? *Breastfeed Rev* 2001; **9**: 5-12 [PMID: 11424519]
- 33 **Glynn DJ**, Hutchinson MR, Ingman WV. Toll-like receptor 4 regulates lipopolysaccharide-induced inflammation and lactation insufficiency in a mouse model of mastitis. *Biol Reprod* 2014; **90**: 91 [PMID: 24671877 DOI: 10.1095/biolreprod.114.117663]
- 34 **Kawai T**, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol* 2010; **11**: 373-384 [PMID: 20404851 DOI: 10.1038/ni.1863]
- 35 **Wambach KA**. Lactation mastitis: a descriptive study of the experience. *J Hum Lact* 2003; **19**: 24-34 [PMID: 12587642 DOI: 10.1177/0890334402239731]
- 36 **Ingman WV**, Glynn DJ, Hutchinson MR. Inflammatory mediators in mastitis and lactation insufficiency. *J Mammary Gland Biol Neoplasia* 2014; **19**: 161-167 [PMID: 24961655 DOI: 10.1007/s10911-014-9325-9]
- 37 **Xiao HB**, Wang CR, Liu ZK, Wang JY. LPS induces pro-inflammatory response in mastitis mice and mammary epithelial cells: Possible involvement of NF- κ B signaling and OPN. *Pathol Biol (Paris)* 2015; **63**: 11-16 [PMID: 25468491 DOI: 10.1016/j.patbio.2014.10.005]
- 38 **Cao D**, Luo J, Chen D, Xu H, Shi H, Jing X, Zang W. CD36 regulates lipopolysaccharide-induced signaling pathways and mediates the internalization of *Escherichia coli* in cooperation with TLR4 in goat mammary gland epithelial cells. *Sci Rep* 2016; **6**: 23132 [PMID: 26976286 DOI: 10.1038/srep23132]
- 39 **Liu S**, Shi X, Bauer I, Günther J, Seyfert HM. Lingual antimicrobial peptide and IL-8 expression are oppositely regulated by the antagonistic effects of NF- κ B p65 and C/EBP β in mammary epithelial cells. *Mol Immunol* 2011; **48**: 895-908 [PMID: 21255844 DOI: 10.1016/j.molimm.2010.12.018]
- 40 **Wu J**, Li L, Sun Y, Huang S, Tang J, Yu P, Wang G. Altered molecular expression of the TLR4/NF- κ B signaling pathway in mammary tissue of Chinese Holstein cattle with mastitis. *PLoS One* 2015; **10**: e0118458 [PMID: 25706977 DOI: 10.1371/journal.pone.0118458]
- 41 **Sasai M**, Yamamoto M. Pathogen recognition receptors: ligands and signaling pathways by Toll-like receptors. *Int Rev Immunol* 2013; **32**: 116-133 [PMID: 23570313 DOI: 10.3109/08830185.2013.774391]
- 42 **Lebda MA**, Elmassy IH, Taha NM, Elfeky MS. Nanocurcumin alleviates inflammation and oxidative stress in LPS-induced mastitis via activation of Nrf2 and suppressing TLR4-mediated NF- κ B and HMGB1 signaling pathways in rats. *Environ Sci Pollut Res Int* 2022; **29**: 8294-8305 [PMID: 34482462 DOI: 10.1007/s11356-021-16309-9]
- 43 **McDermott MF**, Tschopp J. From inflammasomes to fevers, crystals and hypertension: how basic research explains inflammatory diseases. *Trends Mol Med* 2007; **13**: 381-388 [PMID: 17822957 DOI: 10.1016/j.molmed.2007.07.005]
- 44 **Buchanan MM**, Hutchinson M, Watkins LR, Yin H. Toll-like receptor 4 in CNS pathologies. *J Neurochem* 2010; **114**: 13-27 [PMID: 20402965 DOI: 10.1111/j.1471-4159.2010.06736.x]
- 45 **El Mezayen R**, El Gazzar M, Seeds MC, McCall CE, Dreskin SC, Nicolls MR. Endogenous signals released from necrotic cells augment inflammatory responses to bacterial endotoxin. *Immunol Lett* 2007; **111**: 36-44 [PMID: 17568691 DOI: 10.1016/j.imlet.2007.04.011]
- 46 **Samuvel DJ**, Sundararaj KP, Nareika A, Lopes-Virella MF, Huang Y. Lactate boosts TLR4 signaling and NF- κ B pathway-mediated gene transcription in macrophages via monocarboxylate transporters and MD-2 up-regulation. *J Immunol* 2009; **182**: 2476-2484 [PMID: 19201903 DOI: 10.4049/jimmunol.0802059]
- 47 **Smith A**, Wheelock JV, Dodd FH. Effect of milking throughout pregnancy on milk yield in the succeeding lactation. *J Dairy Sci* 1966; **49**: 895-896 [PMID: 5967711 DOI: 10.3168/jds.S0022-0302(66)87966-3]
- 48 **Omranipour R**, Vasigh M. Mastitis, Breast Abscess, and Granulomatous Mastitis. *Adv Exp Med Biol* 2020; **1252**: 53-61 [PMID: 32816262 DOI: 10.1007/978-3-030-41596-9_7]
- 49 **Peaker M**. Recent advances in the study of monovalent ions movements across the mammary epithelium: relation to onset of lactation. *J Dairy Sci* 1975; **58**: 1042-1047 [PMID: 1095619 DOI: 10.3168/jds.s0022-0302(75)84677-7]
- 50 **Peaker M**. Mechanism of milk secretion: milk composition in relation to potential difference across the mammary epithelium. *J Physiol* 1977; **270**: 489-505 [PMID: 903903 DOI: 10.1113/jphysiol.1977.sp011964]
- 51 **Linzell JL**, Peaker M. The distribution and movements of carbon dioxide, carbonic acid and bicarbonate between blood and milk in the goat. *J Physiol* 1975; **244**: 771-782 [PMID: 237118 DOI: 10.1113/jphysiol.1975.sp010825]
- 52 **Shennan DB**, McNeillie SA. High affinity (Na⁺ + Cl⁻)-dependent taurine transport by lactating mammary tissue. *J Dairy Res* 1994; **61**: 335-343 [PMID: 7962835 DOI: 10.1017/s0022029900030752]
- 53 **Prentice A**, Prentice AM, Lamb WH. Mastitis in rural Gambian mothers and the protection of the breast by milk antimicrobial factors. *Trans R Soc Trop Med Hyg* 1985; **79**: 90-95 [PMID: 4039482 DOI: 10.1016/0035-9203(85)90245-7]
- 54 **Prosser CG**, Hartmann PE. Comparison of mammary gland function during the ovulatory menstrual cycle and acute breast inflammation in women. *Aust J Exp Biol Med Sci* 1983; **61** (Pt 3): 277-286 [PMID: 6626049 DOI: 10.1038/icb.1983.27]
- 55 **Semba RD**, Kumwenda N, Taha TE, Hoover DR, Lan Y, Eisinger W, Mtimavalye L, Broadhead R, Miotti PG, Van Der Hoeven L, Chiphangwi JD. Mastitis and immunological factors in breast milk of lactating women in Malawi. *Clin Diagn Lab Immunol* 1999; **6**: 671-674 [PMID: 10473515 DOI: 10.1128/CDLI.6.5.671-674.1999]
- 56 **Brew K**, Hill RL. Lactose biosynthesis. *Rev Physiol Biochem Pharmacol* 1975; **72**: 105-158 [PMID: 806951 DOI: 10.1007/BFb0031548]
- 57 **Fetherston CM**, Lai CT, Mitoulas LR, Hartmann PE. Excretion of lactose in urine as a measure of increased permeability

- of the lactating breast during inflammation. *Acta Obstet Gynecol Scand* 2006; **85**: 20-25 [PMID: 16521675 DOI: 10.1080/00016340500324514]
- 58 **Fazil Marickar YM.** Electrical conductivity and total dissolved solids in urine. *Urol Res* 2010; **38**: 233-235 [PMID: 19921168 DOI: 10.1007/s00240-009-0228-y]
- 59 **Seifter JL.** Body Fluid Compartments, Cell Membrane Ion Transport, Electrolyte Concentrations, and Acid-Base Balance. *Semin Nephrol* 2019; **39**: 368-379 [PMID: 31300092 DOI: 10.1016/j.semnephrol.2019.04.006]
- 60 **de Oliveira Moura E,** do Nascimento Rangel AH, de Melo MCN, Borba LHF, de Lima Júnior DM, Novaes LP, Urbano SA, de Andrade Neto JC. Evaluation of microbiological, cellular and risk factors associated with subclinical mastitis in female buffaloes. *Asian-Australas J Anim Sci* 2017; **30**: 1340-1349 [PMID: 28183165 DOI: 10.5713/ajas.16.0744]
- 61 **Kitchen BJ,** Middleton G, Salmon M. Bovine milk N-acetyl-beta-D-glucosaminidase and its significance in the detection of abnormal udder secretions. *J Dairy Res* 1978; **45**: 15-20 [PMID: 621309 DOI: 10.1017/s0022029900016149]
- 62 **Kermack WO,** Miller RA. Electrical conductivity and chloride content of women's milk. Part 2. The effect of factors relating to lactation. *Arch Dis Child* 1951; **26**: 320-324 [PMID: 14878472 DOI: 10.1136/adc.26.128.320]
- 63 **Miller RA,** Jackson II. Electrical conductivity and chloride content of women's milk. Part 4. Results and their relationship to milk yield and to duration of lactation. *Arch Dis Child* 1951; **26**: 329-334 [PMID: 14878474 DOI: 10.1136/adc.26.128.329]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

