

World Journal of *Clinical Cases*

World J Clin Cases 2022 October 16; 10(29): 10391-10822



Contents

Thrice Monthly Volume 10 Number 29 October 16, 2022

STANDARD AND CONSENSUS

- 10391** Baishideng's *Reference Citation Analysis* database announces the first *Article Influence Index* of multidisciplinary scholars
Wang JL, Ma YJ, Ma L, Ma N, Guo DM, Ma LS

REVIEW

- 10399** Cholecystectomy for asymptomatic gallstones: Markov decision tree analysis
Lee BJH, Yap QV, Low JK, Chan YH, Shelat VG
- 10413** Liver transplantation for hepatocellular carcinoma: Historical evolution of transplantation criteria
Ince V, Sahin TT, Akbulut S, Yilmaz S

MINIREVIEWS

- 10428** Prostate only radiotherapy using external beam radiotherapy: A clinician's perspective
Lee JW, Chung MJ

ORIGINAL ARTICLE

Retrospective Study

- 10435** Age-adjusted NT-proBNP could help in the early identification and follow-up of children at risk for severe multisystem inflammatory syndrome associated with COVID-19 (MIS-C)
Rodriguez-Gonzalez M, Castellano-Martinez A
- 10451** Clinicopathological characteristics and prognosis of gastric signet ring cell carcinoma
Tian HK, Zhang Z, Ning ZK, Liu J, Liu ZT, Huang HY, Zong Z, Li H
- 10467** Development and validation of a prognostic nomogram for decompensated liver cirrhosis
Zhang W, Zhang Y, Liu Q, Nie Y, Zhu X

Observational Study

- 10478** Effect of medical care linkage-continuous management mode in patients with posterior circulation cerebral infarction undergoing endovascular interventional therapy
Zhu FX, Ye Q
- 10487** Effect of the COVID-19 pandemic on patients with presumed diagnosis of acute appendicitis
Akbulut S, Tuncer A, Ogut Z, Sahin TT, Koc C, Guldogan E, Karabulut E, Tanriverdi ES, Ozer A

EVIDENCE-BASED MEDICINE

- 10501** Delineation of a SMARCA4-specific competing endogenous RNA network and its function in hepatocellular carcinoma

Zhang L, Sun T, Wu XY, Fei FM, Gao ZZ

SYSTEMATIC REVIEWS

- 10516** Comparison of laboratory parameters, clinical symptoms and clinical outcomes of COVID-19 and influenza in pediatric patients: A systematic review and meta-analysis

Yu B, Chen HH, Hu XF, Mai RZ, He HY

CASE REPORT

- 10529** Surgical treatment of bipolar segmental clavicle fracture: A case report

Liang L, Chen XL, Chen Y, Zhang NN

- 10535** Multiple disciplinary team management of rare primary splenic malignancy: Two case reports

Luo H, Wang T, Xiao L, Wang C, Yi H

- 10543** Klippel-Trenaunay-Weber syndrome with ischemic stroke: A case report

Lee G, Choi T

- 10550** Vedolizumab in the treatment of immune checkpoint inhibitor-induced colitis: Two case reports

Zhang Z, Zheng CQ

- 10559** Novel way of patent foramen ovale detection and percutaneous closure by intracardiac echocardiography: A case report

Han KN, Yang SW, Zhou YJ

- 10565** Treatment failure in a patient infected with *Listeria* sepsis combined with latent meningitis: A case report

Wu GX, Zhou JY, Hong WJ, Huang J, Yan SQ

- 10575** Three-in-one incidence of hepatocellular carcinoma, cholangiocellular carcinoma, and neuroendocrine carcinoma: A case report

Wu Y, Xie CB, He YH, Ke D, Huang Q, Zhao KF, Shi RS

- 10583** Intestinal microbiome changes in an infant with right atrial isomerism and recurrent necrotizing enterocolitis: A case report and review of literature

Kaplina A, Zaikova E, Ivanov A, Volkova Y, Alkhova T, Nikiforov V, Latypov A, Khavkina M, Fedoseeva T, Pervunina T, Skorobogatova Y, Volkova S, Ulyantsev V, Kalinina O, Sitkin S, Petrova N

- 10600** *Serratia fonticola* and its role as a single pathogen causing emphysematous pyelonephritis in a non-diabetic patient: A case report

Villasuso-Alcocer V, Flores-Tapia JP, Perez-Garfias F, Rochel-Perez A, Mendez-Dominguez N

- 10606** Cardiac myxoma shedding leads to lower extremity arterial embolism: A case report

Meng XH, Xie LS, Xie XP, Liu YC, Huang CP, Wang LJ, Zhang GH, Xu D, Cai XC, Fang X

- 10614** Extracorporeal membrane oxygenation in curing a young man after modified Fontan operation: A case report
Guo HB, Tan JB, Cui YC, Xiong HF, Li CS, Liu YF, Sun Y, Pu L, Xiang P, Zhang M, Hao JJ, Yin NN, Hou XT, Liu JY
- 10622** Wandering small intestinal stromal tumor: A case report
Su JZ, Fan SF, Song X, Cao LJ, Su DY
- 10629** Acute mesenteric ischemia secondary to oral contraceptive-induced portomesenteric and splenic vein thrombosis: A case report
Zhao JW, Cui XH, Zhao WY, Wang L, Xing L, Jiang XY, Gong X, Yu L
- 10638** Perioperative anesthesia management in pediatric liver transplant recipient with atrial septal defect: A case report
Liu L, Chen P, Fang LL, Yu LN
- 10647** Multiple tophi deposits in the spine: A case report
Chen HJ, Chen DY, Zhou SZ, Chi KD, Wu JZ, Huang FL
- 10655** Myeloproliferative neoplasms complicated with β -thalassemia: Two case report
Xu NW, Li LJ
- 10663** Synchronous renal pelvis carcinoma associated with small lymphocytic lymphoma: A case report
Yang HJ, Huang X
- 10670** *Leclercia adecarboxylata* infective endocarditis in a man with mitral stenosis: A case report and review of the literature
Tan R, Yu JQ, Wang J, Zheng RQ
- 10681** Progressive ataxia of cerebrotendinous xanthomatosis with a rare c.255+1G>T splice site mutation: A case report
Chang YY, Yu CQ, Zhu L
- 10689** Intravesical explosion during transurethral resection of bladder tumor: A case report
Xu CB, Jia DS, Pan ZS
- 10695** Submucosal esophageal abscess evolving into intramural submucosal dissection: A case report
Jiao Y, Sikong YH, Zhang AJ, Zuo XL, Gao PY, Ren QG, Li RY
- 10701** Immune checkpoint inhibitor-associated arthritis in advanced pulmonary adenocarcinoma: A case report
Yang Y, Huang XJ
- 10708** Chondroid syringoma of the lower back simulating lipoma: A case report
Huang QF, Shao Y, Yu B, Hu XP
- 10713** Tension-reduced closure of large abdominal wall defect caused by shotgun wound: A case report
Li Y, Xing JH, Yang Z, Xu YJ, Yin XY, Chi Y, Xu YC, Han YD, Chen YB, Han Y

- 10721** Myocardial bridging phenomenon is not invariable: A case report
Li HH, Liu MW, Zhang YF, Song BC, Zhu ZC, Zhao FH
- 10728** Recurrent atypical leiomyoma in bladder trigone, confused with uterine fibroids: A case report
Song J, Song H, Kim YW
- 10735** Eczema herpeticum *vs* dermatitis herpetiformis as a clue of dedicator of cytokinesis 8 deficiency diagnosis: A case report
Alshengeti A
- 10742** Cutaneous allergic reaction to subcutaneous vitamin K₁: A case report and review of literature
Zhang M, Chen J, Wang CX, Lin NX, Li X
- 10755** Perithyroidal hemorrhage caused by hydrodissection during radiofrequency ablation for benign thyroid nodules: Two case reports
Zheng BW, Wu T, Yao ZC, Ma YP, Ren J
- 10763** Malignant giant cell tumors of the tendon sheath of the right hip: A case report
Huang WP, Gao G, Yang Q, Chen Z, Qiu YK, Gao JB, Kang L
- 10772** Atypical Takotsubo cardiomyopathy presenting as acute coronary syndrome: A case report
Wang ZH, Fan JR, Zhang GY, Li XL, Li L
- 10779** Secondary light chain amyloidosis with Waldenström's macroglobulinemia and internodal marginal zone lymphoma: A case report
Zhao ZY, Tang N, Fu XJ, Lin LE
- 10787** Bilateral occurrence of sperm granulomas in the left spermatic cord and on the right epididymis: A case report
Ly DY, Xie HJ, Cui F, Zhou HY, Shuang WB
- 10794** Glucocorticoids combined with tofacitinib in the treatment of Castleman's disease: A case report
Liu XR, Tian M
- 10803** Giant bilateral scrotal lipoma with abnormal somatic fat distribution: A case report
Chen Y, Li XN, Yi XL, Tang Y
- 10811** Elevated procalcitonin levels in the absence of infection in procalcitonin-secreting hepatocellular carcinoma: A case report
Zeng JT, Wang Y, Wang Y, Luo ZH, Qing Z, Zhang Y, Zhang YL, Zhang JF, Li DW, Luo XZ

LETTER TO THE EDITOR

- 10817** "Helicobacter pylori treatment guideline: An Indian perspective": Letter to the editor
Swarnakar R, Yadav SL
- 10820** Effect of gender on the reliability of COVID-19 rapid antigen test among elderly
Nori W, Akram W

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Natalia Stepanova, DSc, MD, PhD, Academic Research, Chief Doctor, Full Professor, Department of Nephrology and Dialysis, State Institution "Institute of Nephrology of the National Academy of Medical Sciences of Ukraine", Kyiv 04050, Ukraine. nmstep88@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yin; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

October 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Clinicopathological characteristics and prognosis of gastric signet ring cell carcinoma

Hua-Kai Tian, Zuo Zhang, Zhi-Kun Ning, Jiang Liu, Zi-Tao Liu, Hao-Yu Huang, Zhen Zong, Hui Li

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): D
Grade E (Poor): E

P-Reviewer: Kao JT, Taiwan; Saglam S, Turkey; Silano F, Brazil; Yu Y, China

Received: May 22, 2022

Peer-review started: May 22, 2022

First decision: July 14, 2022

Revised: July 21, 2022

Accepted: September 1, 2022

Article in press: September 1, 2022

Published online: October 16, 2022



Hua-Kai Tian, Department of General Surgery, First Affiliated Hospital of Nanchang University, Nanchang 330006, Jiangxi Province, China

Zuo Zhang, Department of Obstetrics and Gynecology, Second Affiliated Hospital of Nanchang University, Nanchang 330006, Jiangxi Province, China

Zhi-Kun Ning, Department of Day Ward, First Affiliated Hospital of Nanchang University, Nanchang 330006, Jiangxi Province, China

Jiang Liu, Zi-Tao Liu, Hao-Yu Huang, Zhen Zong, Department of Gastrointestinal Surgery, Second Affiliated Hospital of Nanchang University, Nanchang 330006, Jiangxi Province, China

Hui Li, Department of Rheumatology and Immunology, First Affiliated Hospital of Nanchang University, Nanchang 330006, Jiangxi Province, China

Corresponding author: Hui Li, MD, Doctor, Department of Rheumatology and Immunology, First Affiliated Hospital of Nanchang University, No. 17 Yongwaizheng Street, Donghu District, Nanchang 330006, Jiangxi Province, China. lihui0791nc@126.com

Abstract

BACKGROUND

The clinicopathological features and prognosis of gastric signet ring cell carcinoma (GSRC) remain controversial, particularly with regard to sensitivity to postoperative adjuvant therapy.

AIM

To compare the pathological features of GSRC with those of gastric adenocarcinoma of different degrees of differentiation and the differences in survival prognosis between the different disease processes.

METHODS

By screening gastric cancer patients from 2010 to 2015 in the database of Surveillance, Epidemiology and End Results, and collecting the clinicopathological and prognostic data of gastric cancer patients who underwent surgery from January 2014 to December 2016 in the Second Affiliated Hospital of Nanchang University, we analyzed the general pathological characteristics of GSRC by the chi-square test. Univariate and multivariate analyses were conducted to compare the factors affecting the survival and prognosis of early and

advanced gastric adenocarcinoma. The Kaplan-Meier curves were plotted to reveal the survival difference between early and advanced GSRC and different differentiated types of gastric adenocarcinoma. The prognosis model of advanced GSRC was established with R software, and the area under curve (AUC) and C-index were used to assess the accuracy of the model.

RESULTS

Analysis of pathological features revealed that signet ring-cell carcinoma (SRC) was more frequently seen in younger (< 60 years), female, and White patients compared to non-SRC patients. SRC was less commonly associated with early gastric cancer (EGC) (23.60% *vs* 39.10%), lower N0 (38.61% *vs* 61.03%), and larger tumour sizes > 5 cm (31.15% *vs* 27.10%) compared to the differentiated type, while the opposite was true compared to the undifferentiated type. Survival prognostic analysis found no significant difference in the prognosis of SRC patients among EGC patients. In contrast, among advanced gastric cancer (AGC) patients, the prognosis of SRC patients was correlated with age, race, tumour size, AJCC stage, T-stage, and postoperative adjuvant therapy. The predictive model showed that the 3-year AUC was 0.787, 5-year AUC was 0.806, and C-index was 0.766. Compared to non-SRC patients, patients with SRC had a better prognosis in EGC [hazard ratio (HR): 0.626, 95% confidence interval (CI): 0.427-0.919, $P < 0.05$] and a worse prognosis in AGC (HR: 1.139, 95%CI: 1.030-1.258, $P < 0.05$). When non-SRC was divided into differentiated and undifferentiated types for comparison, it was found that in EGC, SRC had a better prognosis than differentiated and undifferentiated types, while there was no significant difference between differentiated and undifferentiated types. In AGC, there was no significant difference in prognosis between SRC and undifferentiated types, both of which were worse than differentiated types. A prognostic analysis of postoperative adjuvant therapy for SRC in patients with AGC revealed that adjuvant postoperative radiotherapy or chemotherapy significantly improved patient survival (34.6% and 36.2% *vs* 18.6%, $P < 0.05$).

CONCLUSION

The prognosis of SRC is better than that of undifferentiated type, especially in EGC, and its prognosis is even better than that of differentiated type. SRC patients can benefit from early detection, surgical resection, and aggressive adjuvant therapy.

Key Words: Gastric adenocarcinoma; Gastric signet ring cell carcinoma; Pathological features; Survival prognosis; Prognostic model; Adjuvant therapy

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This observational study analysed the clinicopathological features and prognosis of gastric signet ring cell carcinoma (GSRC). We compared GSRC with differentiated gastric adenocarcinoma and found that GSRC had unique clinicopathological features, was more common in younger female patients, and was more aggressive, showing higher lymph node metastasis and tumour size. However, the prognosis of early GSRC was relatively good, even better than that of differentiated adenocarcinoma. GSRC should be diagnosed early, and radical surgical resection with adjuvant radiotherapy and chemotherapy can significantly improve the survival rate of patients, though it still needs more clinical data to verify.

Citation: Tian HK, Zhang Z, Ning ZK, Liu J, Liu ZT, Huang HY, Zong Z, Li H. Clinicopathological characteristics and prognosis of gastric signet ring cell carcinoma. *World J Clin Cases* 2022; 10(29): 10451-10466

URL: <https://www.wjgnet.com/2307-8960/full/v10/i29/10451.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i29.10451>

INTRODUCTION

Gastric cancer, a kind of extremely common malignancy, ranked fifth in morbidity and fourth in mortality worldwide[1,2]. In accordance with the classification of the World Health Organization, gastric adenocarcinoma could be roughly divided into four histological types, mucinous adenocarcinoma, tubular adenocarcinoma, papillary adenocarcinoma, and signet ring cell carcinoma (SRC)[3]. In addition, it is divided into undifferentiated, poorly-differentiated, moderately-differentiated, and well-differentiated types depending on the degree of differentiation[4]. Gastric SRC (GSRC) is considered a special type of gastric adenocarcinoma and characterized by the accumulation of mucin in the cytoplasm and the displacement of the nucleus to the periphery of the cells[5]. GSRC is divided into

diffuse type in the light of Lauren classification[6], infiltrating type by Ming staging[7], and undifferentiated by Nakamura staging[8].

GSRC is considered to be the histological type with the worst prognosis because of the low survival rate and high recurrence rate. However, with further research on GSRC, we found that it has unique clinicopathological characteristics and prognosis. It has been demonstrated that GSRC was highly aggressive, and once found, most of it had been in progressive stage with lymph node metastasis[9]. Other studies also showed that the histological characteristics of GSRC were weak cohesion, and early tumor cells mainly spread in the mucosa or submucosa[10]. Moreover, the survival prognosis and treatment options for GSRC were controversial. Early Western studies have shown that GSRC or diffuse gastric cancer (DGC) had a poor prognosis[11]. However, scholars in Asian had expressed doubts and believed that research should be carried out according to the different processes of tumors. Studies have showed that the prognosis of early SRC is good, while the prognosis of advanced stage SRC is poor[9, 12]. Most recently, abundant evidence in the United States suggested that GSRC was not necessarily a risk factor affecting prognosis[13]. Combination of postoperative adjuvant radiotherapy, chemotherapy, and targeted therapy had an excellent effect in improving the survival rate of gastric cancer patients. However, a number of studies and retrospective analyses have shown that GSRC was resistant to chemotherapy, and patients could not benefit from postoperative adjuvant treatment[14]. For this conclusion, more research is needed to confirm.

In order to more accurately study the pathological features and survival prognosis of GSRC, the pathological features and prognosis of a large number of postoperative gastric cancer patients need to be analysed and compared with non-GSRC according to different tumour progression and different tumour differentiation types. Therefore, we explored the differences in pathological features and survival prognosis between GSRC and different differentiation types of gastric adenocarcinoma by analysing information related to pathological features and survival prognosis of surgically resected specimens from a large United States National Registry database [Surveillance, Epidemiology and End Results (SEER) database] and the Second Affiliated Hospital of Nanchang University.

MATERIALS AND METHODS

Study population

We applied the SEER database from the National Cancer Institute which recorded essential information of around 28% of United States cases. Since the database is available to the public, and we had achieved authorization from the database (account number: 12846-Nov2019), there is no need to acquire patients' informed consent. Furthermore, the hospital ethics committee has approved the study to conduct. Some concerned information was obtained from the database, mainly including general characteristics, pathological characteristics, and clinical tumor characteristics, as well as treatment methods, survival, and prognosis. Meanwhile, we collected the clinical data of patients with gastric cancer who underwent surgery from January 2014 to December 2017 at the Second Affiliated Hospital of Nanchang University. The inclusion criteria were: (1) Postoperative diagnosis of gastric adenocarcinoma; (2) Complete survival information; and (3) Gastric cancer as the first primary tumor. The exclusion criteria were: (1) Suffering from multiple tumors *in situ*; (2) Incomplete tumor staging; (3) Distant metastasis; (4) Not undergoing surgical resection; and (5) Incomplete information. Tumor histology, site, and grade were classified based on the International Classification of Diseases for Oncology, version 3. Stage of tumor was identified on the basis of the AJCC tumor-node-metastasis staging system, 7th edition[15]. The details of screening is displayed in Figure 1.

Data classification

Ages were divided into < 40 years old, 40-60 years old, 60-80 years old, and > 80 years old. Race included Blacks, Whites, Asian Pacific Islanders (API), and American Indians (AI). Size of tumor was classified into < 2 cm, 2-5 cm, > 5 cm, and not available (NA). The type of differentiation comprised SRC, differentiated type (highly differentiated and moderately differentiated), and undifferentiated (poorly differentiated and undifferentiated). T stage included T1a, T1b, T2, T3, T4a, and T4b. N stage included N0, N1, N2, and N3. AJCC stage included I, II, and III. Primary tumors could be located at different parts, divided into the fundus, gastric body, pylorus, antrum, greater curvature, lesser curvature, and overlapping/not otherwise specified (NOS). Tumor progression could be sorted into advanced gastric cancer (AGC) and early gastric cancer (EGC).

Statistical methods

We applied the Fisher exact probability method or chi-square test to analyse categorical variables for descriptive statistics. Univariate and binary logistic regression was employed to conduct analysis of the risk factors for gastric cancer survival and prognosis, after which the result was suggested as 95% confidence intervals (CIs) and odds ratios (ORs). The R software (version 4.0.5) was adopted to establish the survival prognostic model for advanced GSRC. The area under curve (AUC) value and C-index were used to assess the accuracy of the model. The Kaplan-Meier method was used to conduct




Figure 1 Flow chart of data screening in this study. A: Surveillance, Epidemiology and End Results data set; B: External verification group. AGC: Advanced gastric cancer; EGC: Early gastric cancer; SRC: Signet ring cell carcinoma.

comparative analysis about the difference in survival [overall survival (OS)] and the efficacy of adjuvant therapy between early and advanced GSRC and gastric adenocarcinoma with different levels of differentiation, and the results were validated using the external data set. $P < 0.05$ was considered statistically significant.

RESULTS

General characteristics

According to the inclusion and exclusion criteria, finally 5200 patients were chosen in the SEER database, and 603 patients were collected from The Second Affiliated Hospital of Nanchang University finally. Table 1 summarizes the pathological and clinical variables of the two data sets.

Analysis of the clinical characteristics of GSRC

Compared with the differentiated type, GSRC was more common in young patients (< 60 years) (47.36% vs 16.39%, $P < 0.05$), and the same results were obtained when compared with the undifferentiated type (47.36% vs 30.42%, $P < 0.05$). In addition, GSRC was more frequent in female patients (52.87% vs 38.13% vs 41.95%, $P < 0.05$). In the case of race, compared with the differentiated type, GSRC was more common in Whites (61.74% vs 51.87%, $P < 0.05$), but less in White Blacks and API (Black: 14.19% vs 19.03%; API: 23.09% vs 28.07%, $P < 0.05$). Compared with the undifferentiated type, GSRC was also more common in Whites (61.74% vs 56.97%, $P < 0.05$), and less in API (23.09% vs 27.11%, $P < 0.05$).

At the initial diagnosis, 28.05% of GSRC patients had stage I disease, while 47.10% of differentiated patients and 20.57% of undifferentiated patients were diagnosed as stage I ($P < 0.05$). In terms of T stage and N stage, compared with the differentiated type, the proportion of EGC (23.60% vs 39.10%, $P < 0.05$) and N0 (38.61% vs 61.03%, $P < 0.05$) in GSRC patients was less; compared with the undifferentiated type, the proportion of EGC (23.60% vs 16.62%, $P < 0.05$) and N0 (38.61% vs 33.35%, $P < 0.05$) in GSRC patients was higher. Regarding tumor size, compared with the differentiated type, GSRC patients had more tumors > 5 cm (31.15% vs 27.10%, $P < 0.05$), and showed the opposite result when compared to the undifferentiated type (31.15% vs 37.31%, $P < 0.05$) (Table 2).

Survival and prognostic analysis of GSRC


As shown in the survival curve of Figure 2, no significant difference existed between the 5-year OS of SRC and non-SRC patients (44.6% vs 46.7%, $P > 0.05$) (Figure 2A). Interestingly, when gastric cancer patients were divided into EGC and AGC, the 5-year OS of SRC patients was obviously higher than that of non-SRC patients in EGC (89.0% vs 71.4%, $P < 0.05$) (Figure 2C), while the result indicated the

Table 1 General characteristics of Surveillance, Epidemiology and End Results and validation data sets

Variable	SEER data set (%), <i>n</i> = 5200	Validation data set (%), <i>n</i> = 603
Age (yr)		
< 40	200 (3.85)	41 (6.80)
40-60	1389 (26.71)	313 (51.91)
60-80	2690 (51.73)	236 (39.14)
> 80	921 (17.71)	13 (2.15)
Sex		
Male	2933 (56.40)	423 (70.15)
Female	2267 (43.60)	180 (29.85)
Race		
White	2947 (56.67)	
Black	830 (15.96)	
Black-AI	52 (1.00)	
Black-API	1371 (26.37)	
T stage		
T1a	597 (11.48)	247 (40.96)
T1b	708 (13.62)	
T2	693 (13.33)	70 (11.61)
T3	1697 (32.63)	8 (1.33)
T4a	1155 (22.21)	278 (46.10)
T4b	350 (6.73)	
N stage		
N0	2233 (42.94)	266 (44.11)
N1	951 (18.29)	184 (30.52)
N2	855 (16.44)	89 (14.76)
N3	1161 (22.33)	64 (10.61)
AJCC stage		
I	1580 (30.38)	262 (43.45)
II	1449 (27.87)	71 (11.77)
III	2171 (41.75)	270 (44.78)
Radiotherapy		
Yes	1416 (27.23)	16 (2.66)
No	3784 (72.77)	587 (97.34)
Chemotherapy		
Yes	2599 (49.98)	445 (73.80)
No	2601 (50.02)	158 (26.20)
Tumor size (cm)		
< 2	820 (15.77)	
2-5	2198 (42.27)	94 (15.59)
> 5	1700 (32.69)	393 (65.17)
NA	482 (9.27)	116 (19.24)
Histology		

SRC	1326 (25.50)	84 (13.93)
Differentiated	1550 (29.81)	241 (39.97)
Undifferentiated	2324 (44.69)	278 (46.10)

SEER: Surveillance, Epidemiology and End Results; SRC: Signet ring cell carcinoma; API: Asian Pacific Islanders; AI: American Indians.



DOI: 10.12998/wjcc.v10.i29.10451 Copyright ©The Author(s) 2022.

Figure 2 Kaplan-Meier overall survival curves of each group of patients in the Surveillance, Epidemiology and End Results database. A: Signet ring cell carcinoma vs non-signet cell carcinoma (NSRC) in the overall population; B: SRC vs differentiated and undifferentiated carcinoma in the overall population; C: SRC vs NSRC in early gastric cancer (EGC) patients; D: SRC vs differentiated and undifferentiated carcinoma in EGC patients; E: SRC vs NSRC in advanced gastric cancer (AGC) patients; F: SRC vs differentiated and undifferentiated carcinoma in AGC patients. OS: Overall survival; SRC: Signet ring cell carcinoma; NSRC: Non-signet cell carcinoma.

opposite conclusion in AGC patients (30.6% vs 38.2%, $P < 0.05$) (Figure 2E).

Afterwards, the comparison was conducted again when non-SRC was divided into differentiated type and undifferentiated type. We found that the 5-year OS of SRC was lower than that of differentiated patients (44.6% vs 55.3%, $P < 0.05$), but higher than that of undifferentiated patients (44.6% vs 40.8%) (Figure 2B). Interestingly, different results were obtained when gastric cancer patients were divided into EGC and AGC patients. In EGC patients, the 5-year OS of SRC was better than that of differentiated and undifferentiated types (89.0% vs 71.2% and 71.9%, $P < 0.05$), while there was no significant difference between differentiated and undifferentiated types (Figure 2D). And in the external data set, no obvious difference existed in survival rates between SRC and undifferentiated or differentiated types (Figure 3A). In AGC, the 5-year OS of SRC and undifferentiated type was worse than that of differentiated type (30.6% and 34.7% vs 45.3%, $P < 0.05$). However, in pairwise comparison, no obvious difference existed in survival rate between SRC and undifferentiated type (Figure 2F). It was the same as the verification result of the external data set (Figure 3B).

At the same time, our survival analysis of patients with advanced SRC after surgery and chemotherapy demonstrated that postoperative adjuvant chemotherapy or chemotherapy could significantly increase the 5-year OS of patients (34.6% and 36.2% vs 18.6%, $P < 0.05$) (Figure 4A). The external data set analysis showed that postoperative adjuvant chemotherapy had no effect in improving the survival rate of patients ($P > 0.05$) (Figure 4B).

Table 2 Comparison of general characteristics of patients with different differentiated types of gastric cancer

Variable	Differentiated (%), <i>n</i> = 1550	<i>P</i> value	SRC (%), <i>n</i> = 1326	<i>P</i> value	Undifferentiated (%), <i>n</i> = 2324
Age (yr)					
< 40	13 (0.84)	< 0.001	110 (8.30)	< 0.001	77 (3.31)
40-60	241 (15.55)	< 0.001	518 (39.06)	< 0.001	630 (27.11)
60-80	920 (59.35)	< 0.001	583 (43.97)	< 0.001	1187 (51.08)
> 80	376 (24.26)	< 0.001	115 (8.67)	< 0.001	430 (18.50)
Sex					
		< 0.001		< 0.001	
Male	959 (61.87)		625 (47.13)		1349 (58.05)
Female	591 (38.13)		701 (52.87)		975 (41.95)
Race					
White	804 (51.87)	< 0.001	819 (61.74)	0.005	1324 (56.97)
Black	295 (19.03)	0.001	188 (14.19)	0.536	347 (14.93)
Black-AI	16 (1.03)	0.890	13 (0.98)	0.978	23 (0.99)
Black-API	435 (28.07)	0.002	306 (23.09)	0.007	630 (27.11)
T stage					
T1a	281 (18.13)	0.004	187 (14.10)	< 0.001	129 (5.56)
T1b	325 (20.97)	< 0.001	126 (9.51)	0.140	257 (11.06)
T2	242 (15.61)	0.002	155 (11.69)	0.355	296 (12.74)
T3	450 (29.03)	0.930	383 (28.89)	< 0.001	864 (37.17)
T4a	171 (11.03)	< 0.001	385 (29.03)	0.033	599 (25.77)
T4b	81 (5.23)	0.078	90 (6.78)	0.309	179 (7.70)
N stage					
N0	946 (61.03)	< 0.001	512 (38.61)	0.001	775 (33.35)
N1	278 (17.94)	0.097	207 (15.61)	0.001	466 (20.05)
N2	181 (11.68)	0.010	198 (14.93)	< 0.001	476 (20.48)
N3	145 (9.35)	< 0.001	409 (30.85)	0.002	607 (26.12)
AJCC stage					
I	730 (47.10)	< 0.001	372 (28.05)	< 0.001	478 (20.57)
II	452 (29.16)	< 0.001	305 (23.00)	< 0.001	692 (29.78)
III	368 (23.74)	< 0.001	649 (48.95)	0.679	1154 (49.65)
Primary site					
Fundus	67 (4.32)	0.084	41 (3.09)	0.665	78 (3.36)
Body	198 (12.77)	0.527	180 (13.58)	0.898	319 (13.73)
Antrum	640 (41.29)	< 0.001	434 (32.73)	0.061	832 (35.80)
Pylorus	85 (5.48)	0.879	71 (5.36)	0.687	132 (5.68)
Lesser curve	228 (14.71)	0.694	202 (15.23)	0.452	376 (16.18)
Lesser curve-Greater curve	85 (5.48)	0.294	85 (6.41)	0.218	126 (5.42)
Lesser curve-Overlapping/NOS	247 (15.95)	< 0.001	313 (23.60)	0.007	461 (19.83)
Radiotherapy					
		< 0.001		0.410	
Yes	304 (19.61)		415 (31.30)		697 (29.99)
No	1246 (80.39)		911 (68.70)		1627 (70.01)
Chemotherapy					
		< 0.001		0.015	

Yes	535 (34.52)		785 (59.20)		1279 (55.03)
No	1015 (65.48)		541 (40.80)		1045 (44.97)
Tumor size (cm)					
< 2	354 (22.84)	< 0.001	212 (15.99)	< 0.001	254 (10.93)
2-5	674 (43.48)	0.008	512 (38.61)	0.004	1012 (43.55)
> 5	420 (27.10)	0.017	413 (31.15)	< 0.001	867 (37.31)
NA	102 (6.58)	< 0.001	189 (14.25)	< 0.001	191 (8.21)

SRC: Signet ring cell carcinoma; API: Asian Pacific Islanders; AI: American Indians.




Figure 3 Kaplan-Meier overall survival curves of external validation group. A: Early gastric cancer; B: Advanced gastric cancer. OS: Overall survival; SRC: Signet ring cell carcinoma.




Figure 4 Kaplan-Meier overall survival curves for adjuvant therapy in signet ring cell carcinoma. A: Surveillance, Epidemiology and End Results data set; B: External validation group. OS: Overall survival.

Prognosis prediction model of GSRC

In EGC, SRC was a favorable factor affecting patients' prognosis, and suggested a better prognosis when compared to the differentiated type (HR: 0.636, 95%CI: 0.426-0.950, $P < 0.05$) and non-SRC (HR: 0.626, 95%CI: 0.427-0.919, $P < 0.05$). In AGC, SRC was an unfavorable factor, and the results showed the prognosis was worse when compared to the differentiated type (HR: 1.276, 95%CI: 1.117-1.458, $P < 0.05$) and non-SRC (HR: 1.139, 95%CI: 1.030-1.258, $P < 0.05$) (Tables 3 and 4).

Among EGC, there was no obvious difference in SRC patients' prognosis. In AGC, the prognosis of GSRC was related to tumor size, age, race, AJCC stage, T stage, and postoperative adjuvant therapy (Tables 3 and 4). Finally, we established a prognostic prediction model for advanced GSRC; the C-index of this model was 0.766, and the AUC values of the model for predicting 3-year and 5-year OS were 0.787 and 0.806, respectively (Figures 5 and 6). Table 5 shows the risk score of each factor.

Table 3 Analysis of survival and prognosis of patients with early gastric cancer

Variable	OS (EGC = 1305) or (95%CI)	P value	OS (SRC = 313) or (95%CI)	P value
Age (yr)		< 0.001		0.128
< 40	1 (Reference)		1 (Reference)	
40-60	1.189 (0.282-5.015)	0.813	1.071 (0.128-8.947)	0.950
60-80	2.246 (0.550-9.174)	0.260	2.072 (0.268-16.012)	0.485
> 80	5.516 (1.337-22.753)	0.018	4.204 (0.463-38.177)	0.202
Race		< 0.001		0.093
White	1 (Reference)		1 (Reference)	
Black	1.459 (1.053-2.022)	0.023	2.052 (0.878-4.793)	0.097
Black-AI	1.947 (0.476-7.960)	0.354	0.000	0.990
Black-API	0.497 (0.358-0.691)	< 0.001	0.471 (0.137-1.271)	0.124
Sex				
Male	1 (Reference)		1 (Reference)	
Female	0.610 (0.467-0.797)	< 0.001	0.791 (0.383-1.632)	0.526
Tumor size (cm)		0.057		0.700
< 2	1 (Reference)		1 (Reference)	
2-5	1.317 (0.982-1.764)	0.066	1.579 (0.707-3.525)	0.265
> 5	1.859 (1.161-2.978)	0.010	0.000	0.974
AJCC stage				
I	1 (Reference)		1 (Reference)	
II	1.836 (1.100-3.064)	0.020	0.965 (0.231-4.027)	0.961
Depth				
T1a	1 (Reference)		1 (Reference)	
T1b	1.280 (0.962-1.703)	0.090	1.786 (0.810-3.938)	0.151
LN¹				
N0	1 (Reference)		1 (Reference)	
N1-3	1.271 (0.859-1.880)	0.230	2.134 (0.883-5.154)	0.151
Histology		0.045		
Differentiated	1(Reference)			
SRC	0.636 (0.426-0.950)	0.027		
Undifferentiated	1.039 (0.780-1.384)	0.794		
SRC <i>vs</i> non-SRC	0.626 (0.427-0.919)	0.017		

¹LN: Lymph node metastasis.

OS: Overall survival; SRC: Signet ring cell carcinoma; EGC: Early gastric cancer; API: Asian Pacific Islanders; AI: American Indians.

DISCUSSION

The incidence of GSRC is 3.4%-39% in primary gastric cancer[16,17]. In this study, GSRC patients in the SEER data set accounted for 25.5% of all patients undergoing gastrectomy. GSRC patients in the external validation set accounted for 13.9%. The clinical characteristics and prognosis of GSRC are still controversial. Eastern researchers believed that GSRC was not necessarily an unfavorable prognostic factor, while Western researchers considered that the prognosis of GSRC was poor and the incidence rate continued to increase worldwide[9,12,18,19]. However, most previous studies only included a small number of heterogeneous patients, and did not distinguish distinct differentiation types. Comparing SRC and non-SRC together will inevitably cause a certain impact on the final results. Therefore, during our research, a large database was utilized to analyze, compare, and verify the GSRC patients

Table 4 Analysis of survival and prognosis of patients with advanced gastric cancer

Variable	OS (AGC = 3895), or (95%CI)	P value	OS (SRC = 1013), or (95%CI)	P value
Age (yr)		< 0.001		< 0.001
< 40	1 (Reference)		1 (Reference)	
40-60	1.130 (0.882-1.447)	0.333	1.267 (0.898-1.788)	0.178
60-80	1.543 (1.212-1.985)	< 0.001	1.578 (1.123-2.218)	0.009
> 80	2.936 (2.281-3.780)	< 0.001	2.290 (1.528-3.431)	< 0.001
Race		< 0.001		< 0.001
White	1 (Reference)		1 (Reference)	
Black	1.096 (0.973-1.234)	0.133	1.326 (1.057-1.663)	0.015
Black-AI	1.382 (0.949-2.015)	0.092	2.624 (1.423-4.838)	0.002
Black-API	0.726 (0.651-0.810)	< 0.001	0.800 (0.867-1.205)	0.041
Sex				
Male	1 (Reference)		1 (Reference)	
Female	0.964 (0.883-1.052)	0.409	1.022 (0.867-1.205)	0.791
Tumor size (cm)		< 0.001		0.001
< 2	1 (Reference)		1 (Reference)	
2-5	1.026 (0.823-1.280)	0.817	1.031 (0.680-1.565)	0.885
> 5	1.211 (0.971-1.511)	0.089	1.415 (0.930-2.152)	0.105
AJCC stage		0.019		0.028
I	1 (Reference)		1 (Reference)	
II	1.227 (0.903-1.667)	0.191	1.946 (0.923-4.102)	0.080
III	1.584 (1.076-2.330)	0.020	3.050 (1.247-7.458)	0.015
T stage		< 0.001		< 0.001
T2	1 (Reference)		1 (Reference)	
T3	1.384 (1.135-1.689)	0.001	1.857 (1.189-2.899)	0.006
T4	2.025 (1.633-2.512)	< 0.001	2.615 (1.644-4.161)	< 0.001
N stage		< 0.001		0.086
N0	1 (Reference)		1 (Reference)	
N1	1.241 (1.056-1.459)	0.009	1.061 (0.744-1.512)	0.745
N2	1.295 (1.050-1.597)	0.016	1.034 (0.665-1.608)	0.881
N3	1.904 (1.549-2.341)	< 0.001	1.341 (0.882-2.037)	0.170
Histology		0.002		
Differentiated	1 (Reference)			
SRC	1.276 (1.117-1.458)	< 0.001		
Undifferentiated	1.164 (1.037-1.306)	0.010		
SRC <i>vs</i> non-SRC	1.139 (1.030-1.258)	0.011		
Adjuvant therapy				< 0.001
Radiotherapy			0.635 (0.367-1.099)	0.105
Chemotherapy			0.481 (0.388-0.595)	< 0.001
Radiotherapy + Chemotherapy			0.414 (0.334-0.512)	< 0.001
None			1 (Reference)	

OS: Overall survival; SRC: Signet ring cell carcinoma; AGC: Advanced gastric cancer; API: Asian Pacific Islanders; AI: American Indians.

Table 5 Scores of risk factors in the survival prognosis model of advanced gastric signet ring cell carcinoma

Variable	Points	Total points	3/5-yr survival	Total points
Age (yr)		0-56	0.9-1	0-33
< 40	0	56-111	0.8-0.9	33-88
40-60	18	111-146	0.7-0.8	33-123
60-80	36	146-172	0.6-0.7	123-149
> 80	60	172-195	0.5-0.6	149-172
Race		195-215	0.4-0.5	172-192
White	15	215-235	0.3-0.4	192-212
Black	37	235-257	0.2-0.3	212-234
Black-AI	92	257-283	0.1-0.2	234-260
Black-API	0	> 283	< 0.1	> 260
Tumor size (cm)				
< 2	0			
2-5	5			
> 5	31			
NA	42			
AJCC stage				
I	0			
II	54			
III	100			
T stage				
T2	0			
T3	41			
T4	67			
Adjuvant therapy				
Radiotherapy	33			
Chemotherapy	10			
Radiotherapy + Chemotherapy	0			
None	67			

API: Asian Pacific Islanders; AI: American Indians.

undergoing surgical resection in terms of different progression and differentiation degrees, so as to obtain the pathological characteristics of GSRC and the survival differences between gastric adenocarcinoma with different differentiation degrees to provide more accurate guidance for clinical treatment.

Our research showed that GSRC, as a special pathological type of gastric adenocarcinoma, had different clinical characteristics from differentiated and undifferentiated adenocarcinoma. GSRC tended to occur in young and female patients, which was consistent with previous studies[9]. Although gastric cancer was considered to be a disease more frequently seen in men, a large number of studies had shown that the incidence of GSRC in women was higher[13]. At the same time, the age of onset of GSRC was significantly earlier than that of gastric adenocarcinoma. In terms of race, our research showed that GSRC was more frequent in Whites, while there were fewer patients in API. Part of the etiology of young patients may be attributed to genetic factors. Such patients should be diagnosed as hereditary diffuse gastric cancer (HDGC)[20]. The 2015 multidisciplinary symposium defined HDGC as "early-




Figure 5 Nomogram for predicting the 3- and 5-year overall survival of patients with advanced gastric signet ring cell carcinoma. The total score is calculated by adding the scores of each factor. The total score corresponds to the patient's 3- and 5-year survival probabilities.

onset diffuse gastric cancer", where multiple generations of people in the family have a history of DGC or lobular breast cancer[21]. The gender differences of manifestations of GSRC may be related to the level of estrogen. Studies have shown that more frequent appearance of progesterone and estrogen receptors in the tissues of female patients suggests that they might be associated with the manifestations and progression of tumors[22]. The high-level CLDN18-ARHGAP26/6 fusion in GSRC results in genetic differences from other diffuse gastric cancer subtypes. These genetic types develop at a young age and are associated with a high proportion of females, high tumor stages, poor survival outcomes, and chemoresistance[23].

The microscopic features of GSRC are scattered malignant cells containing intracytoplasmic mucin, accounting for more than 50% of tumors[16,24]. GSRC is inert in the early stage and does not show strong invasiveness. When the tumor breaks through the submucosa, it shows strong aggression, rapidly invading the muscle layer, serosal layer, and surrounding lymph nodes[25]. Large-scale data studies have found that GSRC showed a higher proportion of serosal layer invasion and distant metastasis in the advanced stage, and it was prone to lymph node metastasis[25]. Our results found that compared to gastric adenocarcinoma, GSRC had a higher proportion of advanced stage, and showed larger tumor size and more lymph node metastases. This is consistent with the results of most studies.

The prognosis of GSRC is unanimously controversial. Previous studies have suggested that GSRC had a poorer prognosis than non-GSRC. In this study, we divided non-GSRC into differentiated gastric adenocarcinoma and undifferentiated gastric adenocarcinoma, and compared EGC and AGC separately. The results showed that in EGC, the prognosis of differentiated and undifferentiated gastric adenocarcinoma was not significantly different, and was both worse than that of SRC. The external validation indicated that the difference in the prognosis between the early gastric adenocarcinoma and GSRC was not obvious. Interestingly, the prognosis of patients with early GSRC had nothing to do with age, gender, and tumor size. In AGC patients, the prognosis of SRC and undifferentiated gastric cancer patients did not have much difference, and was both worse than that of differentiated type. The external validation set also reached the same conclusion. Through data analysis, we found that the prognosis of patients with advanced GSRC was related to tumor size, age, race, AJCC stage, T stage, and postoperative adjuvant treatment. Undoubtedly, as the tumour progresses, the patient's prognosis deteriorates. Elderly patients had a poorer prognosis due to reduced immunity and poor tolerance. Interestingly, race was also an independent risk factor for the prognosis of GSRC patients, and AI patients had a poorer prognosis. Intrinsic molecular and biological differences between different ethnic groups and living environments may be responsible for the differences in survival among heterogeneous Western populations. Finally, we established a survival prognostic prediction model based on the prognostic risk factors of advanced GSRC. The AUC values of the model for predicting 3-year and 5-year OS were 0.787 and 0.806, respectively, which indicated that the model had accurate predictive ability. Unfortunately, the model cannot be externally validated due to the small amount of data in the external validation set.

Regarding the treatment of GSRC, surgical treatment methods are also controversial for different stages[26]. Endoscopic submucosal dissection is an optional treatment for EGC. On the basis of the




Figure 6 Prediction of 3- and 5-year overall survival. A and B: Nomogram calibration plots to predict 3- and 5-year overall survival (OS); C and D: Nomogram ROC curves for predicting 3- and 5-year OS.

guidelines of the Japanese Gastric Cancer Association, endoscopic therapy could be applied for the cases with well-differentiated and non-ulcerated carcinoma whose diameter is smaller than 2 cm, but the therapy might not be so feasible for ulcerated and undifferentiated submucosal carcinomas[27]. Research has demonstrated that tumor size and lymph node metastasis are important factors for not recommending endoscopic treatment in the early stage of GSRC[28]. According to the clinical characteristics of GSRC in our study, early GSRC showed a higher lymph node metastasis rate and larger tumor size, so endoscopic treatment of GSRC is also not recommended. For AGC, adjuvant radiotherapy and chemotherapy after surgical resection can significantly enhance the survival rate of patients. However, the specific treatment regimen for GSRC is still uncertain. Many studies have confirmed that GSRC was resistant to chemical agents. However, it is still controversial about the efficiency of chemotherapy for GSRC. Voron *et al*[29] showed that postoperative chemotherapy has no significant effect on the survival rate of GSRC patients. In multivariate analysis, GSRC is an independent poor prognostic factor[29]. Another study also proved that GSRC patients cannot benefit from postoperative chemotherapy[30]. Shi *et al*[31] indicated that postoperative chemotherapy could still be effective for the patients with stage IV GSRC[31]. A recent large-scale data study based on the SEER confirmed that surgical resection combined with adjuvant radiotherapy and chemotherapy provides a favorable prognosis for GSRC[32]. Our research showed that postoperative adjuvant radiotherapy and chemotherapy can improve the survival rate of advanced GSRC patients. However, the external validation set showed that postoperative adjuvant chemotherapy cannot benefit patients. This may be because our sample is small,

which makes it impossible to conduct accurate verification. Therefore, the effectiveness of adjuvant therapy for GSRC still needs clinical verification.

Certainly, our analysis was convincing, because we adopted a staged analysis method for GSRC and a large population-based study. Furthermore, we conducted reasonable and effective verification through an external validation set. However, certain limitations still exist in our research. First of all, the SEER database lacks information related to postoperative adjuvant treatment, and information about adjuvant chemotherapy, duration, and neoadjuvant treatment is not available. Second, the analysis of SRC patients with metastasis was not included in the research. Therefore, another study of the particular population may need to be conducted. Finally, although we have established a prognostic prediction model for the advanced GSRC, it cannot be effectively externally verified because the external validation set was insufficient.

CONCLUSION

In summary, our research analysis showed that GSRC is more common in young female patients, and the clinical characteristics of GSRC are significantly different from those of gastric adenocarcinoma. The prognosis of early GSRC is not worse, even better than that of differentiated gastric adenocarcinoma. GSRC should be diagnosed early, and radical surgical resection with adjuvant radiotherapy and chemotherapy can significantly improve the survival rate of patients, though it still needs more clinical data to verify.

ARTICLE HIGHLIGHTS

Research background

The clinicopathological features and prognosis of gastric signet ring cell carcinoma (GSRC) remain controversial, particularly with regard to sensitivity to postoperative adjuvant therapy.

Research motivation

To our knowledge, this study is the first to analyse and compare the clinicopathological features and prognosis of GSRC with gastric adenocarcinoma of different degrees of differentiation and includes both Eastern and Western populations.

Research objectives

The aim of this study was to compare the pathological features of GSRC with those of gastric adenocarcinoma of different degrees of differentiation and the differences in survival prognosis between the different disease processes.

Research methods

This study was first conducted by analysing the differences in clinicopathological features between GSRC and gastric adenocarcinoma in Western populations and comparing the survival prognosis of the different processes. Finally, validation was performed using an Eastern population.

Research results

GSRC was more commonly seen in younger female patients and was more aggressive in the progressive stage, showing more common lymph node metastasis and larger tumour size. However, the prognosis of early GSRC was relatively good, even better than that of differentiated adenocarcinoma. The prognosis of advanced GSRC was not significantly different from that of undifferentiated gastric adenocarcinoma and was worse than that of differentiated gastric adenocarcinoma. Postoperative adjuvant radiotherapy can improve the survival rate of GPC.

Research conclusions

The prognosis of GSRC is better than that of undifferentiated type, especially in EGC, and its prognosis is even better than that of differentiated type. GSRC patients can benefit from early detection, surgical resection, and aggressive adjuvant therapy.

Research perspectives

GSRC has unique clinicopathological features and prognosis, and early diagnosis and treatment can significantly improve survival rates. Patients may benefit from postoperative adjuvant radiotherapy, but further validation is needed.

FOOTNOTES

Author contributions: Tian HK, Zhang Z, and Ning ZK contributed equally to this work; Tian HK and Li H conceived and designed this study; Zhang Z and Ning ZK collected and assembled the data; Zhang Z and Liu J analysed and interpreted the data; Tian HK and Liu J drafted the manuscript; Zong Z, Liu ZT, and Huang HY prepared the figures and tables; all authors read and approved the final manuscript.

Supported by National Natural Science Foundation of China, No. 81860433; Natural Science Youth Foundation of Jiangxi Province, No. 20192BAB215036; Jiangxi Province Natural Science Key R&D Project-General Project, No. 20202BBG73024; Training Plan for Academic and Technical Young Leaders of Major Disciplines in Jiangxi Province, No. 20204BCJ23021; and Jiangxi Provincial Education Department, Science and Technology Research Project-Youth Project, No. GJJ210252.

Institutional review board statement: The Ethics Committee of the Second Affiliated Hospital of Nanchang University approved the study [No. Review[2018]No.(100)].

Informed consent statement: Since this is a retrospective clinical study, no informed consent form was signed.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Hua-Kai Tian 0000-0002-4414-2970; Zuo Zhang 0000-0003-2647-1241; Zhi-Kun Ning 0000-0001-5357-3943; Jiang Liu 0000-0002-1986-0778; Zi-Tao Liu 0000-0002-7039-8689; Hao-Yu Huang 0000-0003-4099-4751; Zhen Zong 0000-0001-8953-7792; Hui Li 0000-0001-9232-2215.

S-Editor: Fan JR

L-Editor: Wang TQ

P-Editor: Yu HG

REFERENCES

- 1 Smyth EC, Nilsson M, Grabsch HI, van Grieken NC, Lordick F. Gastric cancer. *Lancet* 2020; **396**: 635-648 [PMID: 32861308 DOI: 10.1016/S0140-6736(20)31288-5]
- 2 Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* 2021; **71**: 209-249 [PMID: 33538338 DOI: 10.3322/caac.21660]
- 3 Nagtegaal ID, Odze RD, Klimstra D, Paradis V, Rugge M, Schirmacher P, Washington KM, Carneiro F, Cree IA; WHO Classification of Tumours Editorial Board. The 2019 WHO classification of tumours of the digestive system. *Histopathology* 2020; **76**: 182-188 [PMID: 31433515 DOI: 10.1111/his.13975]
- 4 Lai JF, Xu WN, Noh SH, Lu WQ. Effect of World Health Organization (WHO) Histological Classification on Predicting Lymph Node Metastasis and Recurrence in Early Gastric Cancer. *Med Sci Monit* 2016; **22**: 3147-3153 [PMID: 27595490 DOI: 10.12659/msm.897311]
- 5 Bürger ME, Gosens MJ, Jeuken JW, van Kempen LC, van de Velde CJ, van Krieken JH, Nagtegaal ID. Signet ring cell differentiation in mucinous colorectal carcinoma. *J Pathol* 2007; **212**: 278-286 [PMID: 17471475 DOI: 10.1002/path.2181]
- 6 Lauren P. The two histological main types of gastric carcinoma: Diffuse and so-called intestinal-type carcinoma. An attempt at a histo-clinical classification. *Acta Pathol Microbiol Scand* 1965; **64**: 31-49 [PMID: 14320675 DOI: 10.1111/apm.1965.64.1.31]
- 7 Ming SC. Gastric carcinoma. A pathobiological classification. *Cancer* 1977; **39**: 2475-2485 [PMID: 872047 DOI: 10.1002/1097-0142(197706)39:6<2475::aid-cnrcr2820390626>3.0.co;2-I]
- 8 Sugano H, Nakamura K, Kato Y. Pathological studies of human gastric cancer. *Acta Pathol Jpn* 1982; **32** Suppl 2: 329-347 [PMID: 6765003 DOI: 10.1002/1097-0142(19811101)48:9<2084::aid-cnrcr2820480928>3.0.co;2-8]
- 9 Chon HJ, Hyung WJ, Kim C, Park S, Kim JH, Park CH, Ahn JB, Kim H, Chung HC, Rha SY, Noh SH, Jeung HC. Differential Prognostic Implications of Gastric Signet Ring Cell Carcinoma: Stage Adjusted Analysis From a Single High-volume Center in Asia. *Ann Surg* 2017; **265**: 946-953 [PMID: 27232252 DOI: 10.1097/SLA.0000000000001793]
- 10 Chu PG, Weiss LM. Immunohistochemical characterization of signet-ring cell carcinomas of the stomach, breast, and colon. *Am J Clin Pathol* 2004; **121**: 884-892 [PMID: 15198362 DOI: 10.1309/a09erymfr64nerdw]

- 11 **Piessen G**, Messenger M, Leteurtre E, Jean-Pierre T, Mariette C. Signet ring cell histology is an independent predictor of poor prognosis in gastric adenocarcinoma regardless of tumoral clinical presentation. *Ann Surg* 2009; **250**: 878-887 [PMID: 19855261 DOI: [10.1097/SLA.0b013e3181b21c7b](https://doi.org/10.1097/SLA.0b013e3181b21c7b)]
- 12 **Bamboat ZM**, Tang LH, Vinuela E, Kuk D, Gonen M, Shah MA, Brennan MF, Coit DG, Strong VE. Stage-stratified prognosis of signet ring cell histology in patients undergoing curative resection for gastric adenocarcinoma. *Ann Surg Oncol* 2014; **21**: 1678-1685 [PMID: 24394986 DOI: [10.1245/s10434-013-3466-8](https://doi.org/10.1245/s10434-013-3466-8)]
- 13 **Wei Q**, Gao Y, Qi C, Yuan X, Li J, Xu Q, Luo C, Chen L, Zhuo W, Xu Z, Ying J. Clinicopathological Characteristics and Prognosis of Signet Ring Gastric Cancer: A Population-Based Study. *Front Oncol* 2021; **11**: 580545 [PMID: 34490073 DOI: [10.3389/fonc.2021.580545](https://doi.org/10.3389/fonc.2021.580545)]
- 14 **Charalampakis N**, Nogueras González GM, Elimova E, Wadhwa R, Shiozaki H, Shimodaira Y, Blum MA, Rogers JE, Harada K, Matamoros A Jr, Sagebiel T, Das P, Minsky BD, Lee JH, Weston B, Bhutani MS, Estrella JS, Badgwell BD, Ajani JA. The Proportion of Signet Ring Cell Component in Patients with Localized Gastric Adenocarcinoma Correlates with the Degree of Response to Pre-Operative Chemoradiation. *Oncology* 2016; **90**: 239-247 [PMID: 27046280 DOI: [10.1159/000443506](https://doi.org/10.1159/000443506)]
- 15 **Edge SB**, Compton CC. The American Joint Committee on Cancer: the 7th edition of the AJCC cancer staging manual and the future of TNM. *Ann Surg Oncol* 2010; **17**: 1471-1474 [PMID: 20180029 DOI: [10.1245/s10434-010-0985-4](https://doi.org/10.1245/s10434-010-0985-4)]
- 16 **Kwon KJ**, Shim KN, Song EM, Choi JY, Kim SE, Jung HK, Jung SA. Clinicopathological characteristics and prognosis of signet ring cell carcinoma of the stomach. *Gastric Cancer* 2014; **17**: 43-53 [PMID: 23389081 DOI: [10.1007/s10120-013-0234-1](https://doi.org/10.1007/s10120-013-0234-1)]
- 17 **Pernot S**, Voron T, Perkins G, Lagorce-Pages C, Berger A, Taieb J. Signet-ring cell carcinoma of the stomach: Impact on prognosis and specific therapeutic challenge. *World J Gastroenterol* 2015; **21**: 11428-11438 [PMID: 26523107 DOI: [10.3748/wjg.v21.i40.11428](https://doi.org/10.3748/wjg.v21.i40.11428)]
- 18 **Peng J**, Xiao P, Liao B, Ye J, He Y. [Analysis of clinicopathological features of 1879 cases of gastric cancer in Southern China: a single center experience]. *Zhonghua Wai Ke Za Zhi* 2014; **52**: 168-170 [PMID: 24785452 DOI: [10.1016/j.ejca.2014.03.066](https://doi.org/10.1016/j.ejca.2014.03.066)]
- 19 **Mariette C**, Carneiro F, Grabsch HI, van der Post RS, Allum W, de Manzoni G; European Chapter of International Gastric Cancer Association. Consensus on the pathological definition and classification of poorly cohesive gastric carcinoma. *Gastric Cancer* 2019; **22**: 1-9 [PMID: 30167905 DOI: [10.1007/s10120-018-0868-0](https://doi.org/10.1007/s10120-018-0868-0)]
- 20 **JONES EG**. FAMILIAL GASTRIC CANCER. *N Z Med J* 1964; **63**: 287-296 [PMID: 14158754 DOI: [10.1071/mu963287](https://doi.org/10.1071/mu963287)]
- 21 **van der Post RS**, Vogelaar IP, Carneiro F, Guilford P, Huntsman D, Hoogerbrugge N, Caldas C, Schreiber KE, Hardwick RH, Ausems MG, Bardram L, Benusiglio PR, Bisseling TM, Blair V, Bleiker E, Boussioutas A, Cats A, Coit D, DeGregorio L, Figueiredo J, Ford JM, Heijkoop E, Hermens R, Humar B, Kaurah P, Keller G, Lai J, Ligtenberg MJ, O'Donovan M, Oliveira C, Pinheiro H, Ragunath K, Rasenberg E, Richardson S, Roviello F, Schackert H, Seruca R, Taylor A, Ter Huurne A, Tischkowitz M, Joe ST, van Dijk B, van Grieken NC, van Hillegersberg R, van Sandick JW, Vehof R, van Krieken JH, Fitzgerald RC. Hereditary diffuse gastric cancer: updated clinical guidelines with an emphasis on germline CDH1 mutation carriers. *J Med Genet* 2015; **52**: 361-374 [PMID: 25979631 DOI: [10.1136/jmedgenet-2015-103094](https://doi.org/10.1136/jmedgenet-2015-103094)]
- 22 **Liu S**, He L, Wang Z, Wen G. [Expression of sex hormone receptor in gastric cancer with synchronous ovarian metastasis and its significance]. *Zhonghua Yi Xue Za Zhi* 2014; **94**: 1861-1865 [PMID: 25154989 DOI: [10.3816/coc.2009.n.016](https://doi.org/10.3816/coc.2009.n.016)]
- 23 **Shu Y**, Zhang W, Hou Q, Zhao L, Zhang S, Zhou J, Song X, Zhang Y, Jiang D, Chen X, Wang P, Xia X, Liao F, Yin D, Zhou X, Zhang D, Yin S, Yang K, Liu J, Fu L, Zhang L, Wang Y, Zhang J, An Y, Cheng H, Zheng B, Sun H, Zhao Y, Xie D, Ouyang L, Qiu M, Fu X, Dai L, He G, Yang H, Cheng W, Yang L, Liu B, Li W, Dong B, Zhou Z, Wei Y, Peng Y, Xu H, Hu J. Prognostic significance of frequent CLDN18-ARHGAP26/6 fusion in gastric signet-ring cell cancer. *Nat Commun* 2018; **9**: 2447 [PMID: 29961079 DOI: [10.1038/s41467-018-04907-0](https://doi.org/10.1038/s41467-018-04907-0)]
- 24 **Fléjou JF**. [WHO Classification of digestive tumors: the fourth edition]. *Ann Pathol* 2011; **31**: S27-S31 [PMID: 22054452 DOI: [10.1016/j.annpat.2011.08.001](https://doi.org/10.1016/j.annpat.2011.08.001)]
- 25 **Su WP**, Wang WJ, Chang JY, Ho PC, Liu TY, Wen KY, Kuo HL, Chen YJ, Huang SS, Subhan D, Chen YA, Lu CY, Wu CY, Lin SR, Lee MH, Chiang MF, Sze CI, Chang NS. Therapeutic Zfra4-10 or WWOX7-21 Peptide Induces Complex Formation of WWOX with Selective Protein Targets in Organs that Leads to Cancer Suppression and Spleen Cytotoxic Memory Z Cell Activation In Vivo. *Cancers (Basel)* 2020; **12** [PMID: 32764489 DOI: [10.3390/cancers12082189](https://doi.org/10.3390/cancers12082189)]
- 26 **Li Y**, Zhu Z, Ma F, Xue L, Tian Y. Gastric Signet Ring Cell Carcinoma: Current Management and Future Challenges. *Cancer Manag Res* 2020; **12**: 7973-7981 [PMID: 32943931 DOI: [10.2147/CMAR.S268032](https://doi.org/10.2147/CMAR.S268032)]
- 27 **Japanese Gastric Cancer Association**. Japanese gastric cancer treatment guidelines 2018 (5th edition). *Gastric Cancer* 2021; **24**: 1-21 [PMID: 32060757 DOI: [10.1007/s10120-020-01042-y](https://doi.org/10.1007/s10120-020-01042-y)]
- 28 **Yin X**, Xia J, Sun Y, Zhang Z. CHCHD2 is a potential prognostic factor for NSCLC and is associated with HIF-1α expression. *BMC Pulm Med* 2020; **20**: 40 [PMID: 32054470 DOI: [10.1186/s12890-020-1079-0](https://doi.org/10.1186/s12890-020-1079-0)]
- 29 **Voron T**, Messenger M, Duhamel A, Lefevre J, Mabrut JY, Goere D, Meunier B, Brigand C, Hamy A, Glehen O, Mariette C, Paye F. Is signet-ring cell carcinoma a specific entity among gastric cancers? *Gastric Cancer* 2016; **19**: 1027-1040 [PMID: 26606931 DOI: [10.1007/s10120-015-0564-2](https://doi.org/10.1007/s10120-015-0564-2)]
- 30 **Li Y**, Ma FH, Xue LY, Tian YT. Neoadjuvant chemotherapy vs upfront surgery for gastric signet ring cell carcinoma: A retrospective, propensity score-matched study. *World J Gastroenterol* 2020; **26**: 818-827 [PMID: 32148379 DOI: [10.3748/wjg.v26.i8.818](https://doi.org/10.3748/wjg.v26.i8.818)]
- 31 **Shi T**, Song X, Liu Q, Yang Y, Yu L, Liu B, Wei J. Survival benefit of palliative gastrectomy followed by chemotherapy in stage IV gastric signet ring cell carcinoma patients: A large population-based study. *Cancer Med* 2019; **8**: 6010-6020 [PMID: 31448584 DOI: [10.1002/cam4.2521](https://doi.org/10.1002/cam4.2521)]
- 32 **Zhang S**, Liu Y, Jiao Z, Li Z, Wang J, Li C, Qu X, Xu L. Development and Validation of a Prognostic Nomogram for Gastric Signet Ring Cell Carcinoma: A Multicenter Population-Based Study. *Front Oncol* 2021; **11**: 603031 [PMID: 33763350 DOI: [10.3389/fonc.2021.603031](https://doi.org/10.3389/fonc.2021.603031)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

