

# World Journal of *Clinical Cases*

*World J Clin Cases* 2022 October 26; 10(30): 10823-11213



**REVIEW**

- 10823** New insights into the interplay between intestinal flora and bile acids in inflammatory bowel disease  
*Zheng L*
- 10840** Role of visfatin in obesity-induced insulin resistance  
*Abdalla MMI*

**MINIREVIEWS**

- 10852** Hyperthermic intraperitoneal chemotherapy and colorectal cancer: From physiology to surgery  
*Ammerata G, Filippo R, Laface C, Memeo R, Solaini L, Cavaliere D, Navarra G, Ranieri G, Currò G, Ammendola M*
- 10862** New-onset diabetes secondary to acute pancreatitis: An update  
*Yu XQ, Zhu Q*
- 10867** Ketosis-prone diabetes mellitus: A phenotype that hospitalists need to understand  
*Boike S, Mir M, Rauf I, Jama AB, Sunesara S, Mushtaq H, Khedr A, Nitesh J, Surani S, Khan SA*
- 10873** 2022 Monkeypox outbreak: Why is it a public health emergency of international concern? What can we do to control it?  
*Ren SY, Li J, Gao RD*

**ORIGINAL ARTICLE****Retrospective Cohort Study**

- 10882** Clinical characteristics and prognosis of non-small cell lung cancer patients with liver metastasis: A population-based study  
*Wang JF, Lu HD, Wang Y, Zhang R, Li X, Wang S*

**Retrospective Study**

- 10896** Prevalence and risk factors for *Candida* esophagitis among human immunodeficiency virus-negative individuals  
*Chen YH, Jao TM, Shiue YL, Feng IJ, Hsu PI*

- 10906** Prognostic impact of number of examined lymph nodes on survival of patients with appendiceal neuroendocrine tumors  
*Du R, Xiao JW*

**Observational Study**

- 10921** Clinical and epidemiological features of ulcerative colitis patients in Sardinia, Italy: Results from a multicenter study  
*Magri S, Demurtas M, Onidi MF, Picchio M, Elisei W, Marzo M, Miculan F, Manca R, Dore MP, Quarta Colosso BM, Cicu A, Cugia L, Carta M, Binaghi L, Usai P, Lai M, Chicco F, Fantini MC, Armuzzi A, Mocci G*

- 10931** Clinical observation of laparoscopic cholecystectomy combined with endoscopic retrograde cholangiopancreatography or common bile duct lithotripsy

*Niu H, Liu F, Tian YB*

### Prospective Study

- 10939** Patient reported outcome measures in anterior cruciate ligament rupture and reconstruction: The significance of outcome score prediction

*Al-Dadah O, Shepstone L, Donnell ST*

### SYSTEMATIC REVIEWS

- 10956** Body mass index and outcomes of patients with cardiogenic shock: A systematic review and meta-analysis

*Tao WX, Qian GY, Li HD, Su F, Wang Z*

### META-ANALYSIS

- 10967** Impact of being underweight on peri-operative and post-operative outcomes of total knee or hip arthroplasty: A meta-analysis

*Ma YP, Shen Q*

- 10984** Branched-chain amino acids supplementation has beneficial effects on the progression of liver cirrhosis: A meta-analysis

*Du JY, Shu L, Zhou YT, Zhang L*

### CASE REPORT

- 10997** Wells' syndrome possibly caused by hematologic malignancy, influenza vaccination or ibrutinib: A case report

*Šajin M, Luzar B, Zver S*

- 11004** Giant cutaneous squamous cell carcinoma of the popliteal fossa skin: A case report

*Wang K, Li Z, Chao SW, Wu XW*

- 11010** Right time to detect urine iodine during papillary thyroid carcinoma diagnosis and treatment: A case report

*Zhang SC, Yan CJ, Li YF, Cui T, Shen MP, Zhang JX*

- 11016** Two novel mutations in the *VPS33B* gene in a Chinese patient with arthrogryposis, renal dysfunction and cholestasis syndrome 1: A case report

*Yang H, Lin SZ, Guan SH, Wang WQ, Li JY, Yang GD, Zhang SL*

- 11023** Effect of electroacupuncture for Pisa syndrome in Parkinson's disease: A case report

*Lu WJ, Fan JQ, Yan MY, Mukaeda K, Zhuang LX, Wang LL*

- 11031** Neonatal Cri du chat syndrome with atypical facial appearance: A case report

*Bai MM, Li W, Meng L, Sang YF, Cui YJ, Feng HY, Zong ZT, Zhang HB*

- 11037** Complete colonic duplication presenting as hip fistula in an adult with pelvic malformation: A case report

*Cai X, Bi JT, Zheng ZX, Liu YQ*

- 11044** Autoimmune encephalitis with posterior reversible encephalopathy syndrome: A case report  
*Dai SJ, Yu QJ, Zhu XY, Shang QZ, Qu JB, Ai QL*
- 11049** Hypophysitis induced by anti-programmed cell death protein 1 immunotherapy in non-small cell lung cancer: Three case reports  
*Zheng Y, Zhu CY, Lin J, Chen WS, Wang YJ, Fu HY, Zhao Q*
- 11059** Different intraoperative decisions for undiagnosed paraganglioma: Two case reports  
*Kang D, Kim BE, Hong M, Kim J, Jeong S, Lee S*
- 11066** Hepatic steatosis with mass effect: A case report  
*Hu N, Su SJ, Li JY, Zhao H, Liu SF, Wang LS, Gong RZ, Li CT*
- 11074** Bone marrow metastatic neuroendocrine carcinoma with unknown primary site: A case report and review of the literature  
*Shi XB, Deng WX, Jin FX*
- 11082** Child with adenylosuccinate lyase deficiency caused by a novel complex heterozygous mutation in the *ADSL* gene: A case report  
*Wang XC, Wang T, Liu RH, Jiang Y, Chen DD, Wang XY, Kong QX*
- 11090** Recovery of brachial plexus injury after bronchopleural fistula closure surgery based on electrodiagnostic study: A case report and review of literature  
*Go YI, Kim DS, Kim GW, Won YH, Park SH, Ko MH, Seo JH*
- 11101** Severe *Klebsiella pneumoniae* pneumonia complicated by acute intra-abdominal multiple arterial thrombosis and bacterial embolism: A case report  
*Bao XL, Tang N, Wang YZ*
- 11111** Spontaneous bilateral femur neck fracture secondary to grand mal seizure: A case report  
*Senocak E*
- 11116** Favorable response after radiation therapy for intraductal papillary mucinous neoplasms manifesting as acute recurrent pancreatitis: A case report  
*Harigai A, Kume K, Takahashi N, Omata S, Umezawa R, Jingu K, Masamune A*
- 11122** Acute respiratory distress syndrome following multiple wasp stings treated with extracorporeal membrane oxygenation: A case report  
*Cai ZY, Xu BP, Zhang WH, Peng HW, Xu Q, Yu HB, Chu QG, Zhou SS*
- 11128** Morphological and electrophysiological changes of retina after different light damage in three patients: Three case reports  
*Zhang X, Luo T, Mou YR, Jiang W, Wu Y, Liu H, Ren YM, Long P, Han F*
- 11139** Perirectal epidermoid cyst in a patient with sacrococcygeal scoliosis and anal sinus: A case report  
*Ji ZX, Yan S, Gao XC, Lin LF, Li Q, Yao Q, Wang D*

- 11146** Synchronous gastric cancer complicated with chronic myeloid leukemia (multiple primary cancers): A case report  
*Zhao YX, Yang Z, Ma LB, Dang JY, Wang HY*
- 11155** Giant struma ovarii with pseudo-Meigs' syndrome and raised cancer antigen-125 levels: A case report  
*Liu Y, Tang GY, Liu L, Sun HM, Zhu HY*
- 11162** Longest survival with primary intracranial malignant melanoma: A case report and literature review  
*Wong TF, Chen YS, Zhang XH, Hu WM, Zhang XS, Lv YC, Huang DC, Deng ML, Chen ZP*
- 11172** Spontaneous remission of hepatic myelopathy in a patient with alcoholic cirrhosis: A case report  
*Chang CY, Liu C, Duan FF, Zhai H, Song SS, Yang S*
- 11178** Cauda equina syndrome caused by the application of DuraSeal™ in a microlaminectomy surgery: A case report  
*Yeh KL, Wu SH, Fuh CS, Huang YH, Chen CS, Wu SS*
- 11185** Bioceramics utilization for the repair of internal resorption of the root: A case report  
*Riyahi AM*
- 11190** Fibrous hamartoma of infancy with bone destruction of the tibia: A case report  
*Qiao YJ, Yang WB, Chang YF, Zhang HQ, Yu XY, Zhou SH, Yang YY, Zhang LD*
- 11198** Accidental esophageal intubation *via* a large type C congenital tracheoesophageal fistula: A case report  
*Hwang SM, Kim MJ, Kim S, Kim S*
- 11204** Ventral hernia after high-intensity focused ultrasound ablation for uterine fibroids treatment: A case report  
*Park JW, Choi HY*

**LETTER TO THE EDITOR**

- 11210** C-Reactive protein role in assessing COVID-19 deceased geriatrics and survivors of severe and critical illness  
*Nori W*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Rajeev Gurunath Redkar, FRCS, FRCS (Ed), FRCS (Gen Surg), MBBS, MCh, MS, Dean, Professor, Surgeon, Department of Pediatric Surgery, Lilavati Hospital and Research Centre, Mumbai 400050, Maharashtra, India. rajeev.redkar@gmail.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

*WJCC* mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for *WJCC* as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The *WJCC*'s CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

October 26, 2022

**COPYRIGHT**

© 2022 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

# Severe *Klebsiella pneumoniae* pneumonia complicated by acute intra-abdominal multiple arterial thrombosis and bacterial embolism: A case report

Xiao-Li Bao, Nan Tang, Yang-Zhong Wang

**Specialty type:** Critical care medicine

**Provenance and peer review:** Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B, B  
Grade C (Good): C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Saki M, Iran; Shelat VG, Singapore; Velikova TV, Bulgaria

**Received:** May 31, 2022

**Peer-review started:** May 31, 2022

**First decision:** July 12, 2022

**Revised:** July 22, 2022

**Accepted:** September 12, 2022

**Article in press:** September 12, 2022

**Published online:** October 26, 2022



**Xiao-Li Bao, Nan Tang, Yang-Zhong Wang**, Department of Respiratory and Critical Care Medicine, Chongqing University Fuling Hospital, Chongqing 408000, China

**Corresponding author:** Yang-Zhong Wang, MD, Doctor, Department of Respiratory and Critical Care Medicine, Chongqing University Fuling Hospital, No. 2 Gaosuntang Road, Fuling District, Chongqing 408000, China. [632551432@qq.com](mailto:632551432@qq.com)

## Abstract

### BACKGROUND

*Klebsiella pneumoniae* (*K. pneumoniae*) is a clinically common Gram-negative bacillus that can cause community- and hospital-acquired infections and lead to pneumonia, liver abscesses, bloodstream infections, and other infectious diseases; however, severe pneumonia caused by hypervirulent *K. pneumoniae* (hvKp) complicated by acute intra-abdominal multiple arterial thrombosis and bacterial embolism is rarely seen in the clinical setting and has not been reported in the literature.

### CASE SUMMARY

A 51-year-old man was hospitalized with fever and dyspnea. Persistent mild pain in the middle and upper abdomen began at dawn on the 3<sup>rd</sup> day following admission and developed into persistent severe pain in the left upper abdomen 8 h later. Based on chest computed tomography (CT), bronchoscopy, bronchoalveolar lavage fluid metagenomic next-generation sequencing, abdominal aortic CT angiography (CTA), and culture of the superior mesenteric artery embolus, adult community-acquired severe hvKp pneumonia complicated by acute intra-abdominal multiple arterial thrombosis and bacterial embolism was diagnosed. Notably, he recovered and was discharged from the hospital after receiving effective meropenem anti-infection, endovascular contact thrombolytic, and systemic anticoagulant therapies and undergoing percutaneous thrombus aspiration. Ten days later, the patient returned to the hospital for abdominal CTA examination, which indicated blocked initial common pathway of the celiac trunk and superior mesenteric artery, and local stenosis. Therefore, celiac trunk artery stenting was performed in Chongqing Hospital, and postoperative recovery was good.

### CONCLUSION

We report a case of hvKp severe pneumonia complicated by acute intra-abdominal multiple arterial thrombosis and bacterial embolism and suggest that clinicians should consider the possibility of a Gram-negative bacillus infection and conduct effective pathogen detection in a timely fashion when managing patients with severe community-acquired pneumonia before obtaining bacteriologic and drug sensitivity results. At the same time, when patients have severe pulmonary infection complicated by severe abdominal pain, an acute mesenteric artery embolism should be considered to avoid delays in treatment.

**Key Words:** *Klebsiella pneumoniae*; Severe pneumonia; Acute mesenteric ischemia; Superior mesenteric artery embolism; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** The community-acquired *Klebsiella pneumoniae* (*K. pneumoniae*) infection rate is relatively low, but a hypervirulent *K. pneumoniae* infection can develop into a severe illness or even death, which deserves the attention of clinicians. At the same time, although current thrombolytic therapy for the arterial system complicated by infectious bacterial embolism is controversial, we believe that local thrombolytic therapy is still effective under effective anti-infection therapy and warrants further study.

**Citation:** Bao XL, Tang N, Wang YZ. Severe *Klebsiella pneumoniae* pneumonia complicated by acute intra-abdominal multiple arterial thrombosis and bacterial embolism: A case report. *World J Clin Cases* 2022; 10(30): 11101-11110

**URL:** <https://www.wjgnet.com/2307-8960/full/v10/i30/11101.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v10.i30.11101>

## INTRODUCTION

Hypervirulent *Klebsiella pneumoniae* (hvKp) are highly virulent variants of *K. pneumoniae*, and these strains can cause purulent liver abscesses and bacteremia in the absence of biliary or intestinal pathology, as well as spread to the lungs, eyes, and central nervous system, and cause severe community-acquired infections in previously healthy and young people, such as liver abscesses, pneumonia, meningitis, endophthalmitis, and necrotizing fasciitis[1-3].

Acute mesenteric ischemia (AMI) is one of the most severe acute abdominal diseases encountered in vascular surgery. AMI is caused by obstruction of the mesenteric artery or vein, which leads to a sudden interruption of the blood supply or reflux, obstruction of the blood supply and malnutrition of the intestinal canal, and finally the loss of bowel function and necrosis. AMI has an insidious onset and rapid progression and causes severe consequences. If diagnosis and treatment cannot be made in a timely fashion, the fatality rate can reach 50%-70%[4].

## CASE PRESENTATION

### Chief complaints

A 51-year-old nonsmoking male chemical plant worker with normal immune function was hospitalized on March 28, 2021 with fever and dyspnea for 4 d.

### History of present illness

Four days before admission, the patient developed a fever without any apparent trigger. His peak temperature was 39.3 °C, and he experienced progressive aggravation of dyspnea and fatigue. In addition, the patient had a paroxysmal, nonproductive cough.

### History of past illness

Hypertension existed for 10 d. Amlodipine besylate tablets (5 mg/d) were taken orally for antihypertensive treatment daily.

### Personal and family history

His personal and family histories were unremarkable.

**Physical examination**

The initial physical examination showed a heart rate of 107 beats/min, a respiratory rate of 33 breaths/min, a body temperature of 39.1 °C, and a blood pressure of 20.62/10.0 kPa. The body weight was 65 kg. Scratches were noted on both the forearms and hands, and skin ulcerations were occasionally observed (Figure 1). Coarse breath sounds and extensive rales were appreciated in both lungs. The remainder of the organ systems were unremarkable on physical examination.

**Laboratory examinations**

Laboratory testing revealed the following: WBC count,  $20.30 \times 10^9/L$ ; neutrophil percentage, 84.4%; hemoglobin concentration, 116 g/L; procalcitonin, 2.03 ng/mL; C-reactive protein (CRP), 380.00 mg/L; interleukin-6 (IL-6), 408.4 pg/mL; serum albumin, 25.0 g/L; D-dimer, 1.84 mg/L; arterial blood gas (FiO<sub>2</sub>, 33%; pH, 7.46; PaCO<sub>2</sub>, 35 mmHg; PaO<sub>2</sub>, 76 mmHg; HCO<sub>3</sub>, 24.9 mmol/L; lactic acid, 1.1 mmol/L; and P/F, 230 mmHg. The following measures and indicators were normal: liver function; coagulation function; HIV; HBV; A/B virus antigen; cytomegalovirus IgM/IgG; Epstein-Barr virus IgM/IgG; blood lipids; blood glucose; lymphocyte subsets; IgM, IgG, IgA, IgE, IgD; renal function; autoantibody spectrum; anti-neutrophil cytoplasmic antibody spectrum; brain natriuretic peptide; myocardial injury markers; ECG; and routine defecation. In addition, there were no abnormalities detected in the heart, abdominal and lower limb deep veins by ultrasonography, or by bilateral carotid artery ultrasonography.

**Imaging examinations**

On the day of admission, chest computed tomography (CT) showed multiple patchy, nodular, flocculent high-density shadows in both lungs, with blurred edges and small voids in some lesions (Figure 2A).

**Initial diagnosis**

We made an initial diagnosis of adult community-acquired severe pneumonia and type I respiratory failure; however, the pathogenesis was not determined.

**Initial treatment**

After admission, the patient was administered high-flow nasal cannula supportive therapy, piperacillin/tazobactam (4.5 g intravenous infusion every 8 h), and linezolid (0.6 g intravenous infusion every 12 h).

**Deterioration of patient condition**

The patient had recurrent high fevers, and the P/F decreased to 152 mmHg. During this period, sputum and blood cultures were repeatedly performed, but no causative organisms were isolated. In addition, in the early morning of the 3rd day following admission, the patient began to develop persistent mild pain in the middle and upper abdomen, although no positive signs were elicited on the abdominal physical examination. The pain was not relieved by intravenous proton pump inhibitor therapy. Eight hours later, the patient developed persistent severe pain in the left upper abdomen; abdominal signs on physical examination were still negative.

**Further diagnostic work-up**

An emergency determination of myocardial injury markers, serum amylase, serum lipase, and re-examination of the cardiac and whole abdominal ultrasounds were normal, but the plasma D-dimer increased to 25.5 mg/L. The results of repeat chest and abdominal CT angiography (CTA) showed that the proximal segment of the celiac trunk and superior mesenteric artery were embolized, and the distal branch appeared reduced in size. The hepatic parenchyma around the gallbladder was enhanced in the arterial stage, with uneven local perfusion and a few calcified plaques in the abdominal aorta (Figure 3A). The bilateral lung lesions increased, and the solid parts around some nodules increased, with reverse halo signs and trophovascular signs (Figure 2B).

---

**MULTIDISCIPLINARY EXPERT CONSULTATION**

---

**Ke Li, MD, Professor and Chief, Department of General Surgery, Chongqing University Fuling Hospital**

The patient had no signs of intestinal necrosis and peritonitis and no indications for a laparotomy; however, to avoid bowel resection, multiple intra-abdominal artery emboli require urgent restoration of the blood supply.



DOI: 10.12998/wjcc.v10.i30.11101 Copyright ©The Author(s) 2022.

Figure 1 Physical examination: Skin scratches on both forearms and hands with occasional skin ulcerations.

***Wei Wei, MD, Assistant Professor of Interventional Vascular Radiology, Department of Radiology, Chongqing University Fuling Hospital***

The acute intra-abdominal multiple arterial emboli were clear, and endovascular catheter treatment was recommended; however, the patient had severe pneumonia, and it is not clear what type the embolus was. Indeed, if the embolus is bacterial, thrombolytic therapy may cause systemic spread of bacteria, thus aggravating the patient's overall status.

***Yong-sheng Shi, MD, Attending Physician, Department of Vascular Surgery, Chongqing University Fuling Hospital***

The patient was in an early stage of acute abdominal multiple artery emboli without obvious peritoneal irritation, and the lesions were located in the main trunk. Therefore, intracavity catheter therapy combined with systemic anticoagulation could be a first treatment choice; however, the patient had severe pneumonia, and if the intra-abdominal artery was bacterially embolized, thrombolytics increased the risk of systemic bacterial spread, resulting in further aggravation of the infection or life-threatening complications.

***Wu Guo, MD, Professor and Chief, Department of Infectious Diseases, Chongqing University Fuling Hospital***

The patient had severe pneumonia with acute intra-abdominal multiple arterial emboli; thus, his current condition was severe. Although the etiology was unknown, it is not possible to know whether there were bacterial emboli. If the blood supply was not restored in a timely fashion, irreversible loss of the intestinal canal would occur, further aggravating the patient's condition. Therefore, endovascular catheter treatment and local intracatheter arterial thrombolysis were recommended on the basis of the patient undergoing strong anti-infection therapy. During the operation, aspirated emboli could be sent for culture. In addition, bronchoalveolar lavage fluid metagenomic next-generation sequencing (BALF-mNGS) was performed to further identify the pathogen(s) after the patient's general condition improved.

---


## FINAL DIAGNOSIS

We made a final diagnosis of community-acquired severe pneumonia with acute intra-abdominal multiple arterial emboli.

---

## TREATMENT

We replaced the antibiotics with meropenem (1 g intravenously every 8 h) combined with vancomycin (1 g intravenously every 12 h) for anti-infection, initiated low-molecular weight heparin sodium (100 IU/kg subcutaneously for anticoagulation every 12 h), withheld food, and provided fluid rehydration




DOI: 10.12998/wjcc.v10.i30.11101 Copyright ©The Author(s) 2022.

**Figure 2** Chest computed tomography imaging examinations. A: Multiple patchy, nodular, and flocculent high-density shadows in both lungs, with blurred edges and small voids in some lesions; B: The bilateral lung lesions increased, and the solid parts around some nodules increased, with reverse halo and trophovascular signs; C: The bilateral lung lesions were significantly absorbed and reduced compared with previous imaging.

and parenteral nutrition. We immediately performed artery digital subtraction angiography (DSA) and found that the embolus in the superior mesenteric artery opening had almost completely blocked the blood vessels.

We implanted a percutaneous endovascular catheter to aspirate the superior mesenteric artery emboli. The aspirated emboli were sent for microbial culture, and endovascular contact thrombolysis [urokinase (300000 U)] was performed. After surgery, urokinase was pumped with 100000 U arterial microbeam for 4 h, followed by a conventional heparin (1250 U) arterial micropump for 4 h by alternate intra-arterial catheter administration, and systemic low-molecular weight heparin was injected subcutaneously for anticoagulation.

Postoperative abdominal pain was significantly relieved, but abdominal distention appeared on the first day after surgery; thus, the alternate intra-arterial catheter administration of intra-arterial thrombolysis and anticoagulant was continued as before. After 72 h of thrombolytic therapy, intra-arterial thrombolytic therapy and intracatheter anticoagulation were discontinued due to positive fecal occult blood, but subcutaneous injection of 100 IU/kg low-molecular-weight heparin every 12 h was continued.



DOI: 10.12998/wjcc.v10.i30.11101 Copyright ©The Author(s) 2022.

**Figure 3** Abdominal aortic computed tomography angiography imaging examinations. A: The proximal segment of the celiac trunk and superior mesenteric artery were embolized (as indicated by the red arrow), and the distal branch appeared small. The hepatic parenchyma around the gallbladder was enhanced in the arterial stage, with uneven local perfusion and a few calcified plaques in the abdominal aorta; B: Reperfusion of the proper hepatic artery, partial infarction of the spleen and cystic changes, blocked initial common pathway of the celiac trunk and superior mesenteric artery, but the embolization improved (as indicated by the red arrow), and obvious local stenosis.

On the 5<sup>th</sup> d following admission, the superior mesenteric artery embolus culture suggested *K. pneumoniae* growth (PHOENIX100 Fully Automatic Microbial Analysis System, from BioMérieux, France). On the 7<sup>th</sup> d following admission, the patient still had intermittent fevers. Therefore, BALF-mNGS (Kingmed Diagnostics, China) was performed to further clarify the etiology, and the results showed that the sequence number of *Klebsiella* was 13636, among which the sequence number of *K. pneumoniae* detected was 4812. Therefore, we discontinued vancomycin and continued meropenem to fight the infection, and low-molecular-weight heparin 100 IU/kg subcutaneous injection was continued every 12 h during hospitalization.

## OUTCOME AND FOLLOW-UP

The dyspnea was relieved, the body temperature returned to normal, the abdominal pain resolved, abdominal distension improved, and a semifluid diet was resumed. On the 23<sup>rd</sup> d following admission, a chest CT re-examination showed that the bilateral lung lesions were significantly absorbed and reduced compared with the former CT findings (Figure 2C). The patient's condition improved, and he was discharged home. He was treated with rivaroxaban (15 mg twice daily) and clopidogrel (75 mg once daily). Although the abdominal distention improved, he still had symptoms of abdominal distention after discharge.

Ten days later, the patient returned to the hospital for abdominal CTA examination, which indicated reperfusion of the proper hepatic artery, partial infarction of the spleen, cystic changes, blocked initial common pathway of the celiac trunk and superior mesenteric artery, and local stenosis (Figure 3B). Therefore, celiac trunk artery stenting was performed in Chongqing Hospital, and postoperative recovery was good.

## DISCUSSION

In recent years, the *K. pneumoniae* detection rate has been increasing steadily. *K. pneumoniae* belongs to *Klebsiella Enterobacteriaceae*. According to its virulence and pathogenic characteristics, *K. pneumoniae* can be divided into classic *K. pneumoniae* (cKp) and hvKp[5]. One of its main characteristics is that cKp causes nosocomial infection and usually infects immunocompromised or immunocompromised people since cKp carry fewer virulence genes and are less virulent but can show high levels of multiresistance to antibiotics. However, hvKp often manifests as a community-acquired infection, which is more likely to cause disease in healthy people and is often accompanied by multisite invasive infection, which can be life-threatening in severe cases[6]. Compared with cKp strains, hvKp strains showed stronger virulence in various infection modes. Recent studies suggested the existence of hvKp strains with a negative string test. The K1 and K2 capsular serotypes *rmpA*, *rmpA2*, aerobactin and yersiniabactin are the common factors responsible for hypervirulent phenotypes[7].

hvKp was initially found mainly in Southeast Asia, but an increasing number of cases have been reported around the world, including in Europe and the United States, and China is a high-incidence area of hvKp infection[8]. A bloodstream source is currently the most recognized route of infection for hvKp. Due to the ability of hvKp to resist neutrophil phagocytosis, it can flow to various tissues and organs in the body through the blood circulation, thereby leading to infection[9].

Chest CT in patients with community-acquired *K. pneumoniae* pneumonia may show large, honeycomb, patchy consolidation shadows or be accompanied by abscesses, cavities, and liquefaction necrosis, and > 50% of patients have pleural effusions[10]. Pneumonia caused by hematogenous disseminated *K. pneumoniae* infections is similar to *Staphylococcus aureus* (*S. aureus*) infection, such as multiple nodules near the pleura of both lungs, or accompanied by hollow nodules, trophovascular signs, anti-halo signs, and other CT imaging features[11]. Therefore, there is a risk of failure of antibiotic therapy based only on the characteristics of chest CT images.

The patient presented herein was a healthy Asian man who worked as a cleaner in a factory. Before the disease onset, his skin was exposed to sewage pollutants due to touching it with both hands during cleaning, and the skin of his forearms and hands was scratched, resulting in local suppurative infections, after which he developed fevers and dyspnea. His procalcitonin, CRP, and IL-6 Levels were elevated. Chest CT showed multiple patchy, nodular, and flocculent bilateral lung high-density shadows, with blurred edges, and small cavities in some lesions. During chest and abdominal CTA, the lung lesions showed reverse halo and trophovascular signs.

An *S. aureus* infection was suspected as the initial clinical diagnosis, and treatment with piperacillin/tazobactam combined with linezolid failed. In recent years, mNGS has been shown to be able to rapidly and objectively detect various pathogenic microorganisms in clinical samples and conduct drug resistance gene detection of bacteria; thus, mNGS is widely used in clinical practice[12]. In our case, when early anti-infection treatment was ineffective, BALF was obtained for mNGS detection, and *K. pneumoniae* was detected, which was consistent with the results of the DSA aspiration embolus culture, proving that meropenem treatment was effective and reducing the adverse effects caused by the blind use of other broad-spectrum antibiotics.

Although the traditional research view is that high virulence and multidrug resistance do not overlap in the hvKp genome, most hvKp remain highly sensitive to commonly used antibiotics except ampicillin, and their resistance rate is extremely low[7,13]. However, with the extensive use of antibiotics, drug-resistant hvKp has emerged, especially carbapenem-resistant hvKp (CR-hvKp), which has brought great challenges to clinical treatment. Although our case was not detected as CR-hvKp, the application of meropenem was effective, but in the future, while treating hvKp invasion syndrome in a timely and standardized manner, we should also strengthen CR-hvKp detection and nosocomial infection prevention and control measures to prevent the spread of CR-hvKp infection[14,15].

AMI has an insidious onset and rapid progression. Various causes lead to acute obstruction of the mesenteric artery or vein, which leads to sudden interruption of the blood supply or reflux, intestinal blood supply disorder and malnutrition, and eventually loss of function and necrosis[4]. Although the clinical incidence is only 1%-2%, the fatality rate is very high[16,17]. Sixty-five percent of AMI patients have acute superior mesenteric artery emboli[18]. Mesenteric artery emboli mostly originate from the left atrium and are often associated with arrhythmias, such as atrial fibrillation, and can also be caused by heart valve dysfunction or bacterial emboli shedding. It has been reported that high-risk factors include atherosclerosis, dyslipidemia, hypertension, dehydration, diabetes, estrogen, and antiphospholipid syndrome[19].

Although the chief complaint of severe abdominal pain inconsistent with the physical examination findings is a classic presentation of early AMI, it is not sufficient for its diagnosis, and there are no specific biomarkers to confirm AMI[20]. CTA imaging technology has officially replaced angiography as the imaging examination of choice, with a sensitivity of 93%, a specificity of 100%, a positive rate of 94%, and an exclusion rate of 100%[21].

Currently, the "4R" treatment strategy (resuscitation, rapid diagnosis, revascularization, and reassessment of bowel) is recommended internationally for the treatment of AMI[22]. In patients with early acute arterial embolism or thrombosis, the onset of intestinal ischemia is < 12 h, intestinal injury is usually in the reversible stage without obvious peritoneal irritation, and the lesions are located in the

main trunk or branches. Endovascular catheter therapy combined with systemic anticoagulation can be the first choice for treatment[17,23]. In this case, the patient had acute abdominal pain, high blood pressure, and high-risk factors for dehydration, and the plasma D-dimer levels increased sharply.

After acute intra-abdominal multiple artery emboli were confirmed, although it was impossible to know whether there was bacterial embolism, timely intracatheter contact thrombolysis, percutaneous thrombolysis aspiration, and systemic anticoagulation therapy were performed on the basis of strong anti-infection treatment, which avoided intestinal necrosis and diffuse peritonitis and improved the patient's prognosis. Although hvKp infection can cause sepsis, the risk of infective endocarditis is very low[1], and the patient had no history of atrial fibrillation, which does not support mesenteric artery embolism due to shedding of infective endocarditis bacterial emboli.

Therefore, we speculated that hvKp entered his bloodstream through the damaged and exposed skin and then circulated to the mesenteric artery. Due to sepsis, high fever, and dehydration, the patient developed a hypercoagulable state, which resulted in the formation of bacterial emboli and thrombi. At present, the efficacy and safety of arterial thrombolysis are mainly reported for acute ischemic stroke secondary to infective endocarditis. Previously, infective endocarditis represented a classical contraindication to thrombolysis for acute ischemic stroke due to a potential increased risk of intracranial hemorrhage. However, some case reports have suggested the safety and potential efficacy of intravenous or intra-arterial thrombolysis in stroke related to infective endocarditis[24-27]. Notably, McCollom and Zwirko[28] reported a case of infected iliofemoral deep venous thrombosis that was successfully treated with catheter-directed thrombolysis, angioplasty, and stent placement. Although there are no reports of bacterial thrombolysis in mesenteric arteries, this case demonstrates that thrombolysis can be accomplished safely with adequate antibiotic coverage.

---

## CONCLUSION

At present, community-acquired severe pneumonia complicated with acute intra-abdominal multiple arterial thrombosis and bacterial embolism caused by hvKp has not been reported in the literature, and there is no rapid and reliable clinical test to identify hvKp. In summary, we suggest that clinicians should consider the possibility of Gram-negative bacilli infection and conduct effective pathogen detection in a timely fashion, such as BALF or tissue mNGS, when managing patients with severe community-acquired pneumonia without waiting for bacteriologic and drug sensitivity results. At the same time, in cases of acute abdominal multiple artery embolism and organ dysfunction, thrombolysis and systemic anticoagulation need to be applied along with vascular radiologic intervention. Although thrombolytic therapy for the arterial system complicated with infectious bacteria thrombolysis is currently controversial, we believe that local thrombolytic therapy is still effective under effective anti-infection therapy; however, further research is warranted in the future.

---

## ACKNOWLEDGEMENTS

We acknowledge the contributions of Mr. Xiu-Qing Liao, who endorsed the data and conclusions.

---

## FOOTNOTES

**Author contributions:** Bao XL analyzed the data and wrote the manuscript; Tang N made suggestions; Wang YZ revised the manuscript; all authors have read and approve the final manuscript.

**Supported by** the Chongqing Regional Key Discipline Construction Project, No. zdxk201702.

**Informed consent statement:** Informed written consent was obtained from the patient for publication of this report and any accompanying images.

**Conflict-of-interest statement:** All the authors report no relevant conflicts of interest for this article.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and this manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Xiao-Li Bao 0000-0001-9702-9173; Nan Tang 0000-0003-0956-2782; Yang-Zhong Wang 0000-0001-5780-2651.

S-Editor: Gong ZM

L-Editor: A

P-Editor: Gong ZM

## REFERENCES

- Russo TA, Marr CM. Hypervirulent *Klebsiella pneumoniae*. *Clin Microbiol Rev* 2019; **32** [PMID: 31092506 DOI: 10.1128/CMR.00001-19]
- Gu D, Dong N, Zheng Z, Lin D, Huang M, Wang L, Chan EW, Shu L, Yu J, Zhang R, Chen S. A fatal outbreak of ST11 carbapenem-resistant hypervirulent *Klebsiella pneumoniae* in a Chinese hospital: a molecular epidemiological study. *Lancet Infect Dis* 2018; **18**: 37-46 [PMID: 28864030 DOI: 10.1016/S1473-3099(17)30489-9]
- Sánchez-López J, Garcia-Caballero A, Navarro-San Francisco C, Quereda C, Ruiz-Garbjosa P, Navas E, Drona F, Morosini MI, Cantón R, Díez-Aguilar M. Hypermucoviscous *Klebsiella pneumoniae*: A challenge in community acquired infection. *IDCases* 2019; **17**: e00547 [PMID: 31193033 DOI: 10.1016/j.idcr.2019.e00547]
- Altintas Ü, Lawaetz M, Riazi H, de la Motte L, Lindh M, Lönn L, Sillesen H, Eiberg J. Early- and Long-term Outcome After Endovascular Treatment of Chronic and Acute on Chronic Mesenteric Ischemia in a Large National Cohort. *Eur J Vasc Endovasc Surg* 2019; **58**: e631-e632 [DOI: 10.1016/j.ejvs.2019.09.122]
- Lai YC, Lu MC, Hsueh PR. Hypervirulence and carbapenem resistance: two distinct evolutionary directions that led high-risk *Klebsiella pneumoniae* clones to epidemic success. *Expert Rev Mol Diagn* 2019; **19**: 825-837 [PMID: 31343934 DOI: 10.1080/14737159.2019.1649145]
- Hao Z, Duan J, Liu L, Shen X, Yu J, Guo Y, Wang L, Yu F. Prevalence of Community-Acquired, Hypervirulent *Klebsiella pneumoniae* Isolates in Wenzhou, China. *Microb Drug Resist* 2020; **26**: 21-27 [PMID: 31408411 DOI: 10.1089/mdr.2019.0096]
- Rodrigues C, Passet V, Rakotondrasoa A, Diallo TA, Criscuolo A, Brisse S. Description of *Klebsiella africanensis* sp. nov., *Klebsiella variicola* subsp. *tropicalensis* subsp. nov. and *Klebsiella variicola* subsp. *variicola* subsp. nov. *Res Microbiol* 2019; **170**: 165-170 [PMID: 30817987 DOI: 10.1016/j.resmic.2019.02.003]
- Liu C, Guo J. Hypervirulent *Klebsiella pneumoniae* (hypermucoviscous and aerobactin positive) infection over 6 years in the elderly in China: antimicrobial resistance patterns, molecular epidemiology and risk factor. *Ann Clin Microbiol Antimicrob* 2019; **18**: 4 [PMID: 30665418 DOI: 10.1186/s12941-018-0302-9]
- Palacios M, Miner TA, Frederick DR, Sepulveda VE, Quinn JD, Walker KA, Miller VL. Identification of Two Regulators of Virulence That Are Conserved in *Klebsiella pneumoniae* Classical and Hypervirulent Strains. *mBio* 2018; **9** [PMID: 30087173 DOI: 10.1128/mBio.01443-18]
- Evangelista V, Gonçalves CV, Almeida R, Henriques C, Baptista AM, da Graça JP, Araújo JL. *Klebsiella pneumoniae* Invasive Syndrome. *Eur J Case Rep Intern Med* 2018; **5**: 000800 [PMID: 30756018 DOI: 10.12890/2018\_000800]
- Morikawa K, Okada F, Ando Y, Ishii R, Matsushita S, Ono A, Maeda T, Mori H, Yamashita S, Kawahara K. Meticillin-resistant *Staphylococcus aureus* and meticillin-susceptible *S. aureus* pneumonia: comparison of clinical and thin-section CT findings. *Br J Radiol* 2012; **85**: e168-e175 [PMID: 21750126 DOI: 10.1259/bjr/65538472]
- Han D, Li Z, Li R, Tan P, Zhang R, Li J. mNGS in clinical microbiology laboratories: on the road to maturity. *Crit Rev Microbiol* 2019; **45**: 668-685 [PMID: 31691607 DOI: 10.1080/1040841X.2019.1681933]
- Khaertynov KS, Anokhin VA, Rizvanov AA, Davidiyuk YN, Semyenova DR, Lubin SA, Skvortsova NN. Virulence Factors and Antibiotic Resistance of *Klebsiella pneumoniae* Strains Isolated From Neonates With Sepsis. *Front Med (Lausanne)* 2018; **5**: 225 [PMID: 30155466 DOI: 10.3389/fmed.2018.00225]
- Liu C, Du P, Xiao N, Ji F, Russo TA, Guo J. Hypervirulent *Klebsiella pneumoniae* is emerging as an increasingly prevalent *K. pneumoniae* pathotype responsible for nosocomial and healthcare-associated infections in Beijing, China. *Virulence* 2020; **11**: 1215-1224 [PMID: 32921250 DOI: 10.1080/21505594.2020.1809322]
- Harada S, Aoki K, Yamamoto S, Ishii Y, Sekiya N, Kurai H, Furukawa K, Doi A, Tochitani K, Kubo K, Yamaguchi Y, Narita M, Kamiyama S, Suzuki J, Fukuchi T, Gu Y, Okinaka K, Shiiki S, Hayakawa K, Tachikawa N, Kasahara K, Nakamura T, Yokota K, Komatsu M, Takamiya M, Tateda K, Doi Y. Clinical and Molecular Characteristics of *Klebsiella pneumoniae* Isolates Causing Bloodstream Infections in Japan: Occurrence of Hypervirulent Infections in Health Care. *J Clin Microbiol* 2019; **57** [PMID: 31434721 DOI: 10.1128/JCM.01206-19]
- Salsano A, Salsano G, Spinella G, Palombo D, Santini F. Acute Mesenteric Ischemia: Have the Guidelines of the World Society of Emergency Surgery Analyzed All the Available Evidence? *Cardiovasc Intervent Radiol* 2018; **41**: 358-359 [PMID: 29086055 DOI: 10.1007/s00270-017-1817-8]
- Kerzmann A, Haumann A, Boesmans E, Detry O, Defraigne JO. [Acute mesenteric ischemia]. *Rev Med Liege* 2018; **73**: 300-303 [PMID: 29926570]
- Liao G, Chen S, Cao H, Wang W, Gao Q. Review: Acute superior mesenteric artery embolism: A vascular emergency cannot be ignored by physicians. *Medicine (Baltimore)* 2019; **98**: e14446 [PMID: 30732209 DOI: 10.1097/MD.00000000000014446]
- Pay L, Kolak Z, Çakır B, Kamber T, Yazıcı S. Atrial fibrillation-related acute myocardial infarction and acute mesenteric ischemia. *Turk Kardiyol Dern Ars* 2021; **49**: 410-413 [PMID: 34308875 DOI: 10.5543/tkda.2021.96533]
- Salsano G, Salsano A, Sportelli E, Petrocelli F, Dahmane M, Spinella G, Pane B, Mambrini S, Palombo D, Santini F. What

- is the Best Revascularization Strategy for Acute Occlusive Arterial Mesenteric Ischemia: Systematic Review and Meta-analysis. *Cardiovasc Intervent Radiol* 2018; **41**: 27-36 [PMID: 28752257 DOI: 10.1007/s00270-017-1749-3]
- 21 **Obmann MM**, Punjabi G, Obmann VC, Boll DT, Heye T, Benz MR, Yeh BM. Dual-energy CT of acute bowel ischemia. *Abdom Radiol (NY)* 2022; **47**: 1660-1683 [PMID: 34191075 DOI: 10.1007/s00261-021-03188-4]
  - 22 **Ben Abdallah I**, Castier Y, Corcos O. [Mesenteric arterial ischemia: from diagnosis to decision]. *Rev Prat* 2021; **71**: 853-859 [PMID: 35147338]
  - 23 **Altintas Ü**, Lawaetz M, de la Motte L, Riazi H, Lönn L, Lindh M, Sillesen H, Eiberg J. Endovascular Treatment of Chronic and Acute on Chronic Mesenteric Ischaemia: Results From a National Cohort of 245 Cases. *Eur J Vasc Endovasc Surg* 2021; **61**: 603-611 [PMID: 33589326 DOI: 10.1016/j.ejvs.2021.01.003]
  - 24 **Sicoli M**, Benninger D, Schuknecht B, Jenni R, Valavanis A, Bassetti C. Successful intra-arterial thrombolysis in basilar thrombosis secondary to infective endocarditis. *Cerebrovasc Dis* 2003; **16**: 295-297 [PMID: 12865620 DOI: 10.1159/000071131]
  - 25 **Junna M**, Lin CC, Espinosa RE, Rabinstein AA. Successful intravenous thrombolysis in ischemic stroke caused by infective endocarditis. *Neurocrit Care* 2007; **6**: 117-120 [PMID: 17522795 DOI: 10.1007/s12028-007-0017-9]
  - 26 **Tan M**, Armstrong D, Birken C, Bitnun A, Caldarone CA, Cox P, Kahr W, Macgregor D, Askalan R. Bacterial endocarditis in a child presenting with acute arterial ischemic stroke: should thrombolytic therapy be absolutely contraindicated? *Dev Med Child Neurol* 2009; **51**: 151-154 [PMID: 19191846 DOI: 10.1111/j.1469-8749.2008.03188.x]
  - 27 **Sontineni SP**, Mooss AN, Andukuri VG, Schima SM, Esterbrooks D. Effectiveness of Thrombolytic Therapy in Acute Embolic Stroke due to Infective Endocarditis. *Stroke Res Treat* 2010; **2010** [PMID: 20798840 DOI: 10.4061/2010/841797]
  - 28 **McCullom VE**, Zwirko RM. Catheter-directed thrombolysis in acute, infected iliofemoral venous thrombosis. *J Vasc Interv Radiol* 1998; **9**: 941-944 [PMID: 9840038 DOI: 10.1016/s1051-0443(98)70427-0]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

