World Journal of *Gastrointestinal Oncology*

World J Gastrointest Oncol 2022 November 15; 14(11): 2088-2301





Published by Baishideng Publishing Group Inc

WU

GO World Journal of Gastrointestinal

Contents

Monthly Volume 14 Number 11 November 15, 2022

MINIREVIEWS

Portal vein embolization failure: Current strategies and future perspectives to improve liver hypertrophy 2088 before major oncological liver resection

Cassese G, Han HS, Lee B, Cho JY, Lee HW, Guiu B, Panaro F, Troisi RI

ORIGINAL ARTICLE

Basic Study

- 2097 Proteomic signatures of infiltrative gastric cancer by proteomic and bioinformatic analysis Zhang LH, Zhuo HQ, Hou JJ, Zhou Y, Cheng J, Cai JC
- 2108 Potential role of long noncoding RNA RP5-881L22.5 as a novel biomarker and therapeutic target of colorectal cancer

Zong H, Zou JQ, Huang JP, Huang ST

2122 Synaptophysin-like 2 expression correlates with lymph node metastasis and poor prognosis in colorectal cancer patients

Zhao ZX, Liu QL, Yuan Y, Wang FS

2138 Comprehensive analysis of the potential role and prognostic value of sine oculis homeobox homolog family in colorectal cancer

Fang ZX, Li CL, Wu Z, Hou YY, Wu HT, Liu J

2157 KLF16 promotes pancreatic adenocarcinoma cell proliferation and migration by positively regulating SMAD6

Mi W, Zheng Z, Lu JD, Duan SQ, Zhang J, Zhang HQ, Ding YX, Yin J, Cao F, Zhang J, Li F

2170 MiR-30e-3p inhibits gastric cancer development by negatively regulating THO complex 2 and PI3K/AKT/mTOR signaling

Gu XJ, Li YJ, Wang F, Ye T

E3 ubiquitin ligase TRIM55 promotes metastasis of gastric cancer cells by mediating epithelial-2183 mesenchymal transition

Li WW, Yuan H, Kong S, Tian SB

Clinical and Translational Research

2195 Missed colorectal cancers in a fecal immunochemical test-based screening program: Molecular profiling of interval carcinomas

van der Vlugt M, Carvalho B, Fliers J, Montazeri N, Rausch C, Grobbee EJ, Engeland MV, Spaander MCW, Meijer GA, Dekker E



Contents

Monthly Volume 14 Number 11 November 15, 2022

Case Control Study

2208 Oxidative imbalance increases the risk for colonic polyp and colorectal cancer development

Tsounis D, Villiotou V, Melpidou A, Pantsiou C, Argyrou A, Giannopoulou C, Grigoratou A, Rontogianni D, Mantzaris GJ, Papatheodoridis G

Retrospective Study

2224 Predictive value of indirect bilirubin before neoadjuvant chemoradiotherapy in evaluating prognosis of local advanced rectal cancer patients

Li SF, Wei R, Yu GH, Jiang Z

Observational Study

2238 Features of gastric cancer by anatomic subsite in northern China: A multi-center Health Science Report database study

Qu RZ, Ma YP, Bao XY, Tao LY, Zhou X, Lu SY, Zhang Y, Wang BY, Li F, Tuo L, Zhang ZP, Fu W

CASE REPORT

2253 A rare synchrony of adenocarcinoma of the ampulla with an ileal gastrointestinal stromal tumor: A case report

Matli VVK, Zibari GB, Wellman G, Ramadas P, Pandit S, Morris J

2266 Silent advanced large cell neuroendocrine carcinoma with synchronous adenocarcinoma of the colon: A case report

Baek HS, Kim SW, Lee ST, Park HS, Seo SY

2273 Surgical management of monomorphic epitheliotropic intestinal T-cell lymphoma followed by chemotherapy and stem-cell transplant: A case report and review of the literature

Bissessur AS, Zhou JC, Xu L, Li ZQ, Ju SW, Jia YL, Wang LB

- 2288 Surgical treatment of liver inflammatory pseudotumor-like follicular dendritic cell sarcoma: A case report Fu LY, Jiang JL, Liu M, Li JJ, Liu KP, Zhu HT
- Rare squamous cell carcinoma of the jejunum causing perforated peritonitis: A case report 2295 Xiao L, Sun L, Zhang JX, Pan YS



Contents

Monthly Volume 14 Number 11 November 15, 2022

ABOUT COVER

Associate Editor of World Journal of Gastroenterology Oncology, Keun-Yeong Jeong, PhD, Research Assistant Professor, Chief Executive Officer, PearlsinMires, Seoul 03690, South Korea. alvirus@naver.com

AIMS AND SCOPE

The primary aim of World Journal of Gastrointestinal Oncology (WJGO, World J Gastrointest Oncol) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, etc.

INDEXING/ABSTRACTING

The WJGO is now abstracted and indexed in PubMed, PubMed Central, Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJGO as 3.404; IF without journal self cites: 3.357; 5-year IF: 3.250; Journal Citation Indicator: 0.53; Ranking: 162 among 245 journals in oncology; Quartile category: Q3; Ranking: 59 among 93 journals in gastroenterology and hepatology; and Quartile category: Q3. The WJGO's CiteScore for 2021 is 3.6 and Scopus CiteScore rank 2021: Gastroenterology is 72/149; Oncology is 203/360.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Xiang-Di Zhang, Production Department Director: Xiang Li, Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Gastrointestinal Oncology	https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 1948-5204 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
February 15, 2009	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Monthly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Monjur Ahmed, Florin Burada	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/1948-5204/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
November 15, 2022	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2022 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2022 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: bpgoffice@wjgnet.com https://www.wjgnet.com



0 W U

World Journal of **Gastrointestinal** Oncology

Submit a Manuscript: https://www.f6publishing.com

World J Gastrointest Oncol 2022 November 15; 14(11): 2157-2169

DOI: 10.4251/wjgo.v14.i11.2157

ISSN 1948-5204 (online)

ORIGINAL ARTICLE

Basic Study KLF16 promotes pancreatic adenocarcinoma cell proliferation and migration by positively regulating SMAD6

Wei Mi, Zhi Zheng, Jiong-Di Lu, Shu-Quan Duan, Jie Zhang, Hai-Qiao Zhang, Yi-Xuan Ding, Jie Yin, Feng Cao, Jun Zhang, Fei Li

Specialty type: Oncology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): 0 Grade C (Good): C, C Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Miceli V, Italy; Tantau AI, Romania

Received: July 15, 2022 Peer-review started: July 15, 2022 First decision: July 28, 2022 Revised: September 10, 2022 Accepted: October 18, 2022 Article in press: October 18, 2022 Published online: November 15, 2022



Wei Mi, Zhi Zheng, Hai-Qiao Zhang, Jie Yin, Jun Zhang, Department of General Surgery, Beijing Friendship Hospital, Capital Medical University, Beijing 100050, China

Jiong-Di Lu, Yi-Xuan Ding, Feng Cao, Fei Li, Department of General Surgery, Xuanwu Hospital, Capital Medical University, Beijing 100053, China

Shu-Quan Duan, Department of Digestive Minimally Invasive Surgery, The Second Affiliated Hospital of Baotou Medical College, Baotou 014030, Inner Mongolia Autonomous Region, China

Jie Zhang, Department of Radiology, Beijing Friendship Hospital, Capital Medical University, Beijing 100050, China

Corresponding author: Jun Zhang, MD, PhD, Chief Doctor, General Surgery Department, Beijing Friendship Hospital, Capital Medical University, No. 95 Yongan Road, Xi-Cheng District, Beijing 100050, China. zhangjun5986@ccmu.edu.cn

Abstract

BACKGROUND

Pancreatic adenocarcinoma (PAAD) is a cancerous tumor with an extremely poor 5-year survival rate. The exploration of biomarkers for the diagnosis and treatment of PAAD is crucial in clinical practice. Krüppel-like factors (KLFs) are involved in a variety of biological functions in cells. According to multiple studies, KLF16 behave as an oncogene in prostate, breast and gastric cancers. However, no research has been done on the significance of KLF16 in PAAD.

AIM

To explore the molecular mechanisms of KLF16 in PAAD.

METHODS

KLF16 was identified in the tumor specimens and normal tissues by GEPIA database and verified by quantitative real-time PCR (qRT-PCR). Knockdown or exogenous expression of KLF16, combined with in vitro and in vivo assays, was performed to show the functional significance of KLF16. The molecular mechanism of KLF16 was demonstrated by qRT-PCR, western blotting, immunoprecipitation assay and flow cytometry.



RESULTS

We showed that KLF16 was highly expressed in PAAD patients based on the GEPIA database. KLF16 silencing suppressed while KLF16 overexpression promoted the malignant function of PAAD cells. Based on RNA sequencing, we discovered that KLF16 potentiated the expression of SMAD6 in PAAD cells. SMAD6 transcript abundance was increased and positively correlated with KLF16 expression in PAAD samples. In addition, inhibiting SMAD6 was able to mitigate the effects of KLF16 overexpression on PAAD cell processes, suggesting the importance of SMAD6 in the development of KLF16-triggered PAAD.

CONCLUSION

KLF16/SMAD6 axis might be explored as a therapeutic target for PAAD therapy.

Key Words: KLF16; Pancreatic adenocarcinoma; SMAD6; Tumorigenesis; Therapeutic target

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Our study provided the evidence that KLF16 acted as an oncogene in pancreatic adenocarcinoma (PAAD). We also identified that SMAD6 served as the downstream substrate of KLF16 and this signaling cascade has never been reported. This mechanism indicated a novel insight into the pathological events during the development of PAAD.

Citation: Mi W, Zheng Z, Lu JD, Duan SQ, Zhang J, Zhang HQ, Ding YX, Yin J, Cao F, Zhang J, Li F. KLF16 promotes pancreatic adenocarcinoma cell proliferation and migration by positively regulating SMAD6. World J Gastrointest Oncol 2022; 14(11): 2157-2169

URL: https://www.wjgnet.com/1948-5204/full/v14/i11/2157.htm DOI: https://dx.doi.org/10.4251/wjgo.v14.i11.2157

INTRODUCTION

Pancreatic adenocarcinoma (PAAD) ranks fourteenth in incidence with about 496000 new cases (2.6%) per year, but seventh in mortality with roughly 466000 (4.7%) deaths per year, according to GLOBOCAN 2020[1]. Due to its dismal prognosis, experts expect that pancreatic cancer will rank third in terms of fatality by 2025[2]. Currently, the only way to improve the outcomes of this condition is to diagnose it early[3]. However, since the early signs of PAAD are not well-defined, it is difficult to diagnose, even when it progresses fast to neighboring organs^[4]. As a result, new unique and useful biomarkers for the diagnosis and prognosis of PAAD are required.

Three zinc finger DNA binding domains distinguish the Krüppel-like factor (KLF) which belongs to the SP/KLF transcription factor family^[5]. KLF binds to GT or GC-rich regions and regulates transcription as an activator or repressor, depending on the promoter type[6]. The KLF16 gene is located on chromosome 19 that participates in various cellular activities such as proliferation, metabolism and death[7]. Various studies on the role of KLF16 in cancer have been published in recent years with mixed results. It has been reported that KLF16 promoted progression of breast cancer via activating MAGT1 [8]. Ma et al[9] found that KLF16 enhanced colorectal carcinoma progression by modulating nucleolar homeostasis and translational reprogramming. On the other hand, in some malignancies, KLF16 functions as an oncogenic suppressor. It has been reported that KLF16 suppressed human glioma cell proliferation and tumorigenesis by regulating TFAM[10]. Nevertheless, understanding the molecular mechanisms of KLF16 in PAAD is still limited.

MATERIALS AND METHODS

The Cancer Genome Atlas provided clinical samples

A total of 350 cases which encompassed 171 normal and 179 tumor specimens were analyzed from the database of Gene Expression Profiling Interactive Analysis (GEPIA) (http://gepia.cancerpku.cn/index.html) database, which combines the TCGA database and GEO database.

Cell culture

Shanghai Institutes for Biological Sciences (Shanghai, China) provided the AsPC-1 and MIA PaCa-2 cell lines. Cells were incubated in Eagle's minimal essential medium (MEM) with 10% fetal bovine serum



(FBS), in a 37°C incubator supplemented with 5% CO₂. L-glutamine and sodium pyruvate were both supplied by Gibco and added to the medium to ensure constant cell growth.

Quantitative real-time PCR (gRT-PCR)

Cultivated cell lines were lysed in TRIzol (Ambion, Austin, TX, United States). To reverse transcribe 500 ng of RNA template, the Takara PrimeScript RT Reagent Kit was employed. The following primers were used: KLF16-F: 5'-TGGGCAAACCCTGAAGACA-3' and KLF16-R, 5'-GTTGCACAGATGGGAAGAAA-3'; SMAD6-F:5'- CCTCCCTACTCTCGGCTGTC-3', and SMAD6-R, 5'- GGTAGCCTCCGTTTCAGTGTA-3'; and β-actin-F: 5'-CATGTACGTTGCTATCCAGGC-3', β-actin-R, 5'-CTCCTTAATGTCACGCACGAT-3'. RT-qPCR was conducted on real-time PCR system (7500 Fast; Life Technologies Holdings Pte Ltd, Singapore). The expression of targeted genes was adjusted to β-actin.

Knockdown and overexpression assay

Specific small interfering RNAs (siRNAs) targeting KLF16 (siKLF16#1, siKIF16#2) and SMAD6 (siSMAD6#1, siSMAD6#2) and negative controls siRNA (siCtrl) were provided by Genechem (Shanghai, China). For transient transfection, Invitrogen's Lipofectamine RNAiMAX transfection reagent was employed. The sequences used were as follows: siKLF16#1 (5'-GGGUUCUUC-CAAAGAACAU-3'), siKLF16#2 (5'-GGUAUCACGUGACAAUCAA-3'), siSMAD6#1 (5'-CCACAT-TGTCTTACACTGAAA-3'), siSMAD6#2 (5'-CTCCATCAAGGTGTTCGACTT-3') and siCtrl (5'-UUCUCCGAACGUGUCACGU-3').

KLF16-pCDNA3.1 was overexpressed using Lipofectamine 3000 (Thermo Fisher Scientific, Waltham, MA, United States). shRNA targeting SMAD6 (shSMAD6) purchased from Genechem (Shanghai, China) was used for lentiviral transduction.

Western blotting

To lyse the cells, the RIPA lysis buffer (Beyotime, Jiangsu, China) was used. This solution contains the protease inhibitor cocktail (Roche, Basel, Switzerland). To detect protein amount present in the lysates, a Bradford reagent manufactured by Sigma was used. After that, 30 µg of protein per row was run on SDS-PAGE gels and then immunoblotted onto a 0.22 m polyvinylidene fluoride membrane (Roche). The membranes were incubated for an hour with 5% nonfat milk solution that dissolved in TBST. After adding primary antibodies against KLF16 (1:1000; sc-377519; Santa Cruz Biotechnology, Dallas, TX, United States), SMAD6 (1:1000; ab273106; Abcam), and GAPDH (1:1000; ab8245; Abcam), the membrane was incubated at 4°C overnight. After that, the secondary antibody specific to primary antibody was added. An Odessey CLx system uncovered evidence of the presence of protein bands (LI-COR, Lincoln, NE, United States).

Cell Counting Kit-8 and cell colony

The CCK-8 (Beyotime) was applied to determine cell proliferation rate. In all, 1 × 10³ transfected cells were planted in 200 μ L complete medium. A total of 20 μ L of CCK-8 was used to treat the cells and cultured for a further two hours at 37°C, followed by OD450 measurement. For cell colony assay, 1.2 × 10³ transfected cells were planted in six-well plates with 2 mL of complete media and cultivated for 14 d. Every 3 d, the medium was replaced. The cells were fixed with methanol and stained with 0.1% crystal violet (Solarbio, Beijing, China).

Flow cytometry

In a six-well plate, 2 × 10⁵ AsPC-1 and MIApaca-2 cells were planted per well. Flow cytometry was performed on cells 48 h after transfection with siKLF16#1, siKLF16#2, siCtrl, or KLF16 overexpressed plasmid and Ctrl plasmid.

A cell cycle staining kit (Cat No. 4040301; Yeasen, Shanghai, China) was used to detect the cell cycle. The cells were fixed for two hours at 4°C in 75% ethanol. The cells were stained with a staining solution containing 10 µL propidium iodide and 10 µL RNaseA in a volume of 0.5 mL. The cells were stained for 30 mins at 25°C and then subjected to analysis with the Guava easyCyte HT system (Millipore) after passing through a screen with a mesh size of 400. The cell cycle was analyzed by a flow cytometry system.

The Annexin V Apoptosis Detection Kit (Cat No. 88-8007; Invitrogen, Waltham, MA, United States) was used to examine apoptosis. The cells (1 to 5×10^5) were collected in 1X binding buffer and were incubated with staining solution in the dark for 15 min. The samples were analyzed using the Guava easyCyte HT system (Millipore, Burlington, MA, United States) within 1 h.

Cell migration

5.0 × 10⁴ AsPC-1 or MIApaca-2 cells in 300 µL medium were planted in the upper chambers of 24-well Corning® FluoroBlokTM Cell Culture Inserts (Corning, NY, United States). The bottom chamber was added to 600 µL medium which contained 10% FBS. The migrating cells were stained by 0.1% crystal violet. Images of migrating cells were collected under a microscope.



WJGO | https://www.wjgnet.com



Figure 1 Krüppel-like factor 16 expression in pancreatic adenocarcinoma. A: Expression of Krüppel-like factor (KLF)16 in various The Cancer Genome Atlas (TCGA) cancers; B: Expression of KLF16 in pancreatic adenocarcinoma (PAAD) from the Gene Expression Profiling Interactive Analysis (GEPIA) database. ^aP < 0.05.

RNA sequencing

RNA sequencing (RNA-seq) was carried out on MIA PaCa-2 cells using the Illumina HiSeq TM 4000 (Illumina, San Diego, CA, United States. The sequencing reads were processed using the Fast-QC program. The investigation of the signaling pathway was carried out by KEGG after the KLF16 knockdown.

Tumor xenograft

Vital River Company (Beijing, China) provided the Balb/c nude mice (4-6 wk, female). MIA PaCa-2 cells were subcutaneously implanted into three different groups of nude mice (n = 5). On days 20, 25, 29, 33, 38, and 43, the tumor's volume was assessed. Tumor volume = $4/3 \times \pi \times [(\text{length (l)} + \text{width (w)}) / 4]^{3[11]}$. The mice were euthanized after 43 d, and the tumor tissues were surgically removed and studied. Every experiment that included animals was carried out in compliance with the Principles on the Protection of Experimental Animals that are outlined by the Beijing Friendship Hospital.

Analysis of the data

Data were presented as mean standard deviation and analyzed by using GraphPad Prism 8.0 software (La Jolla, CA, United States). When comparing the difference between one or more groups, Student's *t* test or one-way ANOVA followed by Tukey's multiple comparison test was applied. Statistical significance was considered when P < 0.05.

RESULTS

KLF16 is upregulated in pancreatic adenocarcinoma samples

To examine the functions of KLF16 in PAAD, we interrogated the GEPIA database (http://gepia.cancerpku.cn) and assessed the expression of KLF16 across different tumor histocytes. We found that KLF16 was overexpressed in almost all tumors (Figure 1A). More importantly, we found that KLF16 upregulation was observed in 179 PAAD patients compared with 171 healthy subjects (P < 0.05; Figure 1B).

KLF16 regulates pancreatic adenocarcinoma cell proliferation

To examine if KLF16 could influence the progression of PAAD cells, we performed knockdown and overexpression studies. Firstly, we showed that KLF16 expression was significantly reduced in PAAD cells transfected with siRNAs against KLF16, indicating that the knockdown was effective (Figure 2A and 2B). The number of cell colonies and viability of PAAD cells were significantly reduced when KLF16 was knocked down (Figure 2C-F). Then overexpression experiments were conducted. The results showed that KLF16 overexpression was successful (Figure 2G and 2H). When KLF16 was overexpressed in AsPC-1 and MIA PaCa-2 cells, both the cell viability and the number of cell colonies considerably increased (Figure 2I-L). These findings indicated that KLF16 accelerated the proliferation of PAAD cells.

KLF16 regulates apoptosis and the cell cycle

We then investigated the impact of KLF16 on the PAAD cell cycle as well as the apoptotic process. Following KLF16 knockdown, an enhanced number of the cells at G0/G1 phase and a reduced number of the cells at G2/M phase were observed (Figure 3A). When KLF16 was overexpressed in cells, the opposite effects were found (Figure 3B). Apoptosis in the cells was inhibited after KLF16 overex-





Figure 2 Krüppel-like factor 16 regulates pancreatic adenocarcinoma cell proliferation. A, B: The efficiency of Krüppel-like factor (KLF)16 knockdown in pancreatic adenocarcinoma (PAAD) cells by (A) qPCR and (B) western blotting. GAPDH used as the internal control; C, D: Cell proliferation of PAAD cells after KLF16 knockdown. (E, F) Colony numbers of PAAD cells after KLF16 knockdown; G, H: The efficiency of KLF16 overexpression in the cells by (G) qPCR and (H) western blotting. GAPDH used as the internal control; I, J: Cell proliferation of PAAD cells after KLF16 overexpression by the CCK-8 assay; K, L: Colony numbers of PAAD cells after KLF16 overexpression by the CCK-8 assay; K, L: Colony numbers of PAAD cells after KLF16 overexpression by colony formation assay. $^{a}P < 0.05$; $^{b}P < 0.01$; $^{c}P < 0.001$.

pression, whereas it was significantly promoted by KLF16 downregulation (Figure 3C and 3D).

KLF16 regulates pancreatic adenocarcinoma cell migration

Following that, we investigated whether KLF16 influenced the migration of cells. After KLF16 was knocked down, the amount of cell migration was dramatically reduced (Figure 4A). When KLF16 was overexpressed, the number of migrating cells increased (Figure 4B). According to these findings, KLF16 could regulate the migration of the PAAD cell.





Figure 3 Krüppel-like factor 16 regulates the cell cycle progression and apoptosis in pancreatic adenocarcinoma cells. A, B: Cell cycle

Jaishideng® WJGO | https://www.wjgnet.com

November 15, 2022 Volume 14 Issue 11



progression of pancreatic adenocarcinoma (PAAD) cells with (A) Krüppel-like factor (KLF)16 knockdown or (B) KLF16 overexpression; C, D: Cell apoptosis of PAAD cells with (C) KLF16 knockdown or (D) KLF16 overexpression. ^aP < 0.05; ^bP < 0.01; ^cP < 0.001.

Figure 4 Krüppel-like factor 16 regulates pancreatic adenocarcinoma cell migration. A, B: Cell migration of pancreatic adenocarcinoma (PAAD) cells with (A) knockdown of Krüppel-like factor (KLF16) or (B) overexpression of KLF16 by transwell assay. Bar, 100 μm. ^b*P* < 0.01, ^c*P* < 0.001.

SMAD6 participates in the tumor promoting function of KLF16 in pancreatic adenocarcinoma cells

RNA-seq studies were carried out so that we could determine the pathways that KLF16 is responsible for in the development of PAAD. As shown in Figure 5A, the suppression of KLF16 in MIA PaCa-2 cells resulted in downregulation and upregulation of several genes. Figure 5B displays the top ten pathways that were discovered by cluster analysis to be associated with these genes. Autophagy-related genes were found to be among the most highly upregulated genes, while ribosome-related genes were the most highly downregulated. A highly expressed SMAD6 gene that had not been characterized before was found in PAAD. As a result, we investigated the connection between KLF16 and SMAD6. After KLF16 was either knocked down or overexpressed, respectively, we found that there was either a considerable rise or reduction in the levels of SMAD6 (Figure 5C and 5D). In addition, SMAD6 expression was increased in PAAD tissues retrieved from the TCGA database (Figure 5E). According to the findings of the study of correlation, the level of expression of KLF16 was shown to have a positive association with the level of expression of SMAD6 (r = 0.37; Figure 5F). Collectively, KLF16 promoted the expression of SMAD6 in PAAD cells and tissues.

SMAD6 regulated PAAD cell proliferation and migration

Following that, we explored the impact of SMAD6 on PAAD cells. qPCR was used to test the efficiency of SMAD6 knockdown or overexpression, as shown in Figure 6A. The ability of PAAD cells was suppressed by SMAD6 downregulation, while it was increased when SMAD6 was overexpressed (Figure 6B). This oncogenic function of SMAD6 was further validated by the findings that the number of migrating cells was greatly inhibited by downregulation of SMAD6 (Figure 6C) but was stimulated after SMAD6 overexpression (Figure 6D). According to these findings, SMAD6 was responsible for PAAD cell proliferation and migration.

SMAD6 knockdown rescues the effects of KLF16 overexpression in pancreatic adenocarcinoma cells

To understand how SMAD6 interacted with KLF16 overexpression, we investigated the impact of SMAD6 knockdown on KLF16-mediated PAAD. The protein level of SMAD6 was moderately elevated in MIA PaCa-2 cells transfected with KLF16 and was drastically reduced in cells transfected with KLF16+shSMAD6 (Figure 7A). The viability of MIA PaCa-2 cells improved when KLF16 was overex-





Figure 5 SMAD6 is a downstream target of Krüppel-like factor 16 in pancreatic adenocarcinoma. A: Volcano analysis of differentially expressed genes with Krüppel-like factor (KLF)16 knockdown in MIA PaCa-2 cells by RNA-seq; B: The top 10 pathways or biological processes of upregulated or downregulated genes; C and D: SMAD6 mRNA and protein levels in pancreatic adenocarcinoma (PAAD) cells with KLF16 knockdown or overexpression. GAPDH used as the internal control; E: The expression of SMAD6 in PAAD from The Cancer Genome Atlas; F: Spearman relationship between KLF16 and SMAD6 in PAAD. ^bP < 0.01; ^cP < 0.001.

pressed, but SMAD6 knockdown had a significant adverse effect on that viability (Figure 7B). Moreover, we overexpressed SMAD6 in the KLF16 knockdown AsPC1 cells. SMAD6 was significantly downregulated in the KLF16 knockdown cells, while the protein level of SMAD6 was rescued after SMAD6 overexpression (Figure 7C). The viability of AsPC1 cells was impaired when KLF16 was knocked down, but SMAD6 overexpression had a significant adverse effect on that viability (Figure 7D). *In vivo* tumor transplantation investigations validated these results as well, showing that the tumor size and volume of xenografts increased after KLF16 overexpression in nude mice injected with MIA PaCa-2 cells, but decreased following SMAD6 knockdown (Figure 7E-G). Overall, SMAD6 knockdown rescued the effects of KLF16 overexpression in PAAD cells.

DISCUSSION

KLF16 was highly expressed in PAAD patients based on the GEPIA database. KLF16 silencing suppressed, while KLF16 overexpression promoted the malignant function of PAAD cells. Based on RNA sequencing, we discovered that KLF16 potentiated the expression of SMAD6 in PAAD cells. SMAD6 transcript abundance was increased and positively correlated with KLF16 expression in PAAD





Figure 6 SMAD6 regulates pancreatic adenocarcinoma cell malignancy. A: The efficiency of SMAD6 knockdown or SMAD6 overexpression in pancreatic adenocarcinoma (PAAD) cells by qPCR; B: Cell proliferation of PAAD cells with knockdown or overexpression of SMAD6; C, D: The migration capacity of PAAD cells with (C) SMAD6 knockdown or (D) SMAD6 overexpression by transwell assay. Bar, 100 µm. ^bP < 0.01; ^oP < 0.001.

samples, and inhibiting SMAD6 was able to mitigate the effects of KLF16 overexpression on PAAD cell processes, suggesting the importance of SMAD6 in the development of KLF16-triggered PAAD.

Pancreatic adenocarcinoma, a very aggressive tumor of the digestive system, is difficult to diagnose and treat. PAAD is also known as pancreatic ductal adenocarcinoma, a kind of pancreatic cancer that accounts for the vast majority of cases (95%)[12]. Accumulation of genetic alterations, such as KRAS, CDKN2A/P16, TP53 and SMAD4 contributes to the progression of PAAD[13,14]. These abnormalities may be used to develop a novel effective therapy targeting PAAD.

KLFs refer to zinc finger transcription factors involved in various developmental processes through enhancing and/or repressing the expression of several genes[15]. KLF3, KLF4, KLF12 and KLF15 are only a few of the KLFs that participate in the formation and advancement of PAAD. KLF3 is a pancreatic cancer cell growth inhibitor, according to the study conducted by Wan *et al*[16]. KLF12 was found to be a miR-137 target and inhibited the cancer stem cell phenotype in pancreatic cancer cells[17]. In contrast, Zhu *et al*[18] found that downregulated Caveolin-1 by KLF4 maintained pancreatic cancer epithelial-mesenchymal transition and metastasis. As a member of KLFs, KLF16's function in PAAD is less clear. During our research, we observed that pancreatic adenocarcinoma samples and cells had



WJGO https://www.wjgnet.com

Mi W et al. KLF16 tumor biomarker for pancreatic cancer



Figure 7 SMAD6 knockdown rescues the effects of Krüppel-like factor 16 overexpression on pancreatic adenocarcinoma. (A) Protein levels of Krüppel-like factor (KLF)16 and SMAD6 in control (Ctrl), KLF16, and KLF16+shSAMD6 cells by western blotting in MIA PaCa-2 cells. GAPDH used as the internal control; B: Cell viability of Ctrl, KLF16, and KLF16+shSAMD6 cells by CCK-8 assay in MIA PaCa-2 cells; C: Protein levels of KLF16 and SMAD6 in siCtrl, siKLF16, and siKLF16+SAMD6 cells by western blotting in AsPC-1 cells. GAPDH used as the internal control; D: Cell viability of siCtrl, siKLF16, and siKLF16+SAMD6 cells by the CCK-8 assay in AsPC-1 cells; E: Tumor size from three groups of nude mice that were injection with MIA PaCa-2 cells transfected with Ctrl, KLF16, and KLF16+shSAMD6, separately; F: Tumor growth curve in three groups; G: Tumor weight among the three groups. NS: Not significant. ^bP < 0.01; ^cP < 0.001.

elevated levels of KLF16. By completing knockdown and overexpression studies on KLF16, we were able to show that it controlled PAAD cell malignancy. These studies were carried out using PAAD cells.

SMAD6, a member of the SMAD family, was discovered in the 1990s and serves as a key mediator of the TGF signaling pathway[19,20]. In mammalian cells, eight SMAD have been discovered[21]. SMAD4 Loss predicts poor prognosis in PAAD[14], while SMAD7 has been discovered to be adversely regulated in PAAD[22]. SMAD6 was shown to be downregulated in patients with colorectal cancer[23], while its upregulation was associated with poor patient survival from lung cancer[24]. Experiments using RNA sequencing were carried out so that we could determine the specific mechanism by which KLF16 is involved in the development of PAAD. It has been reported that KLF16 acted as a transcriptional repressor[25]. In the present study, KLF16 expression was discovered to have a favorable positive correlation with SMAD6, which was considerably upregulated. In this context, it was suggested that SMAD6 is not the directed target gene of KLF16. Hence, the role of KLF16 in regulating SMAD6 expression required further experimentation. Finally, we showed that SMAD6 expression was elevated in PAAD samples as well. SMAD6 knockdown inhibited PAAD cell proliferation and migration, but SMAD6 overexpression increased PAAD cell advancement. In addition, inhibiting SMAD6 in PAAD cells reversed the effects of KLF16 overexpression. Overall, KLF16 is an oncogenic protein in PAAD that positively influence SMAD6 expression.

Although this study can fully explain the mechanism of KLF16 on the proliferation and migration of PAAD cells, the lack of prognosis information of patients makes this study limited to a certain extent. It will be further verified in human pancreatic cancer tissues in the future, so as to better support the experimental results and enhance the credibility of the study.

CONCLUSION

The role that KLF16 and SMAD6 play in PAAD was investigated in this work. Both genes carried out their oncogenic functions and contributed to the development of PAAD. Furthermore, we identified a strong correlation between KLF16 and SMAD6 expression levels. KLF16 and SMAD6 appear to have the potential to act as both a novel prognostic marker and a potential therapeutic target for PAAD, based on our findings.

WJGO https://www.wjgnet.com

ARTICLE HIGHLIGHTS

Research background

Pancreatic adenocarcinoma (PAAD) is a cancerous tumor with an extremely poor 5-year survival rate. The exploration of biomarkers for the diagnosis and treatment of PAAD is crucial in clinical practice.

Research motivation

KLF16 behaves as an oncogene in prostate, breast and gastric cancers. However, no research has been done on the significance of Krüppel-like factor 16 (KLF16) in PAAD.

Research objectives

This study aimed to explore the molecular mechanisms of KLF16 in PAAD.

Research methods

KLF16 was identified in the tumor specimens and normal tissues by GEPIA database and verified by quantitative real-time PCR (qRT-PCR). Knockdown or exogenous expression of KLF16, combined with in vitro and in vivo assays, was performed to show the functional significance of KLF16. The molecular mechanism of KLF16 was demonstrated by qRT-PCR, western blotting, immunoprecipitation assay and flow cytometry.

Research results

KLF16 was highly expressed in PAAD patients based on the GEPIA database. KLF16 silencing suppressed, while KLF16 overexpression promoted the malignant function of PAAD cells. Based on RNA sequencing, we discovered that KLF16 potentiated the expression of SMAD6 in PAAD cells. SMAD6 transcript abundance was increased and positively correlated with KLF16 expression in PAAD samples. In addition, inhibiting SMAD6 was able to mitigate the effects of KLF16 overexpression on PAAD cell processes, suggesting the importance of SMAD6 in the development of KLF16-triggered PAAD.

Research conclusions

KLF16/SMAD6 axis might be explored as a therapeutic target for PAAD therapy.

Research perspectives

KLF16 and SMAD6 appear to have the potential to act as both a novel prognostic marker and a potential therapeutic target for PAAD, based on our findings.

ACKNOWLEDGEMENTS

We would like to thank the Clinical Data and Biobank Resource of Beijing Friendship Hospital for their help with data processing.

FOOTNOTES

Author contributions: Mi W, Zheng Z, Lu JD, Duan SQ, Zhang J and Zhang HQ all contributed equally to this work; Mi W, Zheng Z, Lu JD and Duan SQ designed the study; Zhang HQ, Yin J and Zhang J performed the experiments; Zheng Z and Lu JD wrote the manuscript; Ding YX and Cao F performed statistical analysis; Li F and Zhang J revised the manuscript; All authors read and approved the final manuscript.

Institutional review board statement: The research was approved by the Ethics Committee of Beijing Friendship Hospital, Capital Medical University (No. 2018-P2-015-02). The principles outlined in the Declaration of Helsinki were adhered to throughout the course of this research project (as revised in 2013).

Institutional animal care and use committee statement: Every experiment that included animals was carried out in compliance with the Principles on the Protection of Experimental Animals that are outlined by the Beijing Friendship Hospital.

Conflict-of-interest statement: All authors report having no relevant conflict of interest for this article.

Data sharing statement: Data are available from the corresponding authors upon request.

ARRIVE guidelines statement: The authors have read the ARRIVE guidelines, and the manuscript was prepared and revised according to the ARRIVE guidelines.



Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is noncommercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: China

ORCID number: Wei Mi 0000-0002-1330-9926; Zhi Zheng 0000-0003-0390-9466; Hai-Qiao Zhang 0000-0002-4224-7146; Jie Yin 0000-0003-2708-0111; Jun Zhang 0000-0001-5411-1273.

S-Editor: Wu YXJ L-Editor: Filipodia P-Editor: Yuan YY

REFERENCES

- Sung H. Ferlay J. Siegel RL, Laversanne M. Soeriomataram I. Jemal A. Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. CA Cancer J Clin 2021; 71: 209-249 [PMID: 33538338 DOI: 10.3322/caac.21660]
- 2 Ferlay J, Partensky C, Bray F. More deaths from pancreatic cancer than breast cancer in the EU by 2017. Acta Oncol 2016; 55: 1158-1160 [PMID: 27551890 DOI: 10.1080/0284186X.2016.1197419]
- 3 Ottenhof NA, Milne AN, Morsink FH, Drillenburg P, Ten Kate FJ, Maitra A, Offerhaus GJ. Pancreatic intraepithelial neoplasia and pancreatic tumorigenesis: of mice and men. Arch Pathol Lab Med 2009; 133: 375-381 [PMID: 19260743 DOI: 10.5858/133.3.375]
- Goral V. Pancreatic Cancer: Pathogenesis and Diagnosis. Asian Pac J Cancer Prev 2015; 16: 5619-5624 [PMID: 26320426 DOI: 10.7314/apjcp.2015.16.14.5619]
- 5 Philipsen S, Suske G. A tale of three fingers: the family of mammalian Sp/XKLF transcription factors. Nucleic Acids Res 1999; 27: 2991-3000 [PMID: 10454592 DOI: 10.1093/nar/27.15.2991]
- 6 Pearson R, Fleetwood J, Eaton S, Crossley M, Bao S. Krüppel-like transcription factors: a functional family. Int J Biochem Cell Biol 2008; 40: 1996-2001 [PMID: 17904406 DOI: 10.1016/j.biocel.2007.07.018]
- 7 Sun N, Shen C, Zhang L, Wu X, Yu Y, Yang X, Yang C, Zhong C, Gao Z, Miao W, Yang Z, Gao W, Hu L, Williams K, Liu C, Chang Y, Gao Y. Hepatic Krüppel-like factor 16 (KLF16) targets PPARα to improve steatohepatitis and insulin resistance. Gut 2021; 70: 2183-2195 [PMID: 33257471 DOI: 10.1136/gutjnl-2020-321774]
- 8 Bang S, Li J, Zhang M, Cui R, Wu X, Xin Z, Ma D, Zhang J, Zhang H. The Clinical Relevance and Function of Krüppel-Like Factor 16 in Breast Cancer. Cancer Manag Res 2020; 12: 6373-6383 [PMID: 32821156 DOI: 10.2147/CMAR.S256490]
- Ma XD, Xu SD, Hao SH, Han K, Chen JW, Ling H, Chen RX, Jin XH, Cao JH, Lin JL, Ou QJ, Fang YJ, Pan ZZ, Xie D, Wang FW. KLF16 enhances stress tolerance of colorectal carcinomas by modulating nucleolar homeostasis and translational reprogramming. Mol Ther 2022; 30: 2828-2843 [PMID: 35524408 DOI: 10.1016/j.ymthe.2022.04.022]
- 10 Chen X, Li S, Ke Y, Wu S, Huang T, Hu W, Fu H, Guo X. KLF16 suppresses human glioma cell proliferation and tumourigenicity by targeting TFAM. Artif Cells Nanomed Biotechnol 2018; 46: 608-615 [PMID: 29374989 DOI: 10.1080/21691401.2018.1431654]
- 11 Lee MD, She Y, Soskis MJ, Borella CP, Gardner JR, Hayes PA, Dy BM, Heaney ML, Philips MR, Bornmann WG, Sirotnak FM, Scheinberg DA. Human mitochondrial peptide deformylase, a new anticancer target of actinonin-based antibiotics. J Clin Invest 2004; 114: 1107-1116 [PMID: 15489958 DOI: 10.1172/JCI22269]
- Tanaka S. Molecular Pathogenesis and Targeted Therapy of Pancreatic Cancer. Ann Surg Oncol 2016; 23 Suppl 2: S197-12 S205 [PMID: 25749932 DOI: 10.1245/s10434-015-4463-x]
- Vincent A, Herman J, Schulick R, Hruban RH, Goggins M. Pancreatic cancer. Lancet 2011; 378: 607-620 [PMID: 13 21620466 DOI: 10.1016/S0140-6736(10)62307-0]
- Jones S, Zhang X, Parsons DW, Lin JC, Leary RJ, Angenendt P, Mankoo P, Carter H, Kamiyama H, Jimeno A, Hong SM, Fu B, Lin MT, Calhoun ES, Kamiyama M, Walter K, Nikolskaya T, Nikolsky Y, Hartigan J, Smith DR, Hidalgo M, Leach SD, Klein AP, Jaffee EM, Goggins M, Maitra A, Iacobuzio-Donahue C, Eshleman JR, Kern SE, Hruban RH, Karchin R, Papadopoulos N, Parmigiani G, Vogelstein B, Velculescu VE, Kinzler KW. Core signaling pathways in human pancreatic cancers revealed by global genomic analyses. Science 2008; 321: 1801-1806 [PMID: 18772397 DOI: 10.1126/science.1164368]
- Swamynathan SK. Krüppel-like factors: three fingers in control. Hum Genomics 2010; 4: 263-270 [PMID: 20511139 15 DOI: 10.1186/1479-7364-4-4-263]
- 16 Wan Y, Luo H, Yang M, Tian X, Peng B, Zhan T, Chen X, Ding Y, He J, Cheng X, Huang X, Zhang Y. miR-324-5p Contributes to Cell Proliferation and Apoptosis in Pancreatic Cancer by Targeting KLF3. Mol Ther Oncolytics 2020; 18: 432-442 [PMID: 32913892 DOI: 10.1016/j.omto.2020.07.011]
- 17 He Z, Guo X, Tian S, Zhu C, Chen S, Yu C, Jiang J, Sun C. MicroRNA-137 reduces stemness features of pancreatic cancer cells by targeting KLF12. J Exp Clin Cancer Res 2019; 38: 126 [PMID: 30866999 DOI: 10.1186/s13046-019-1105-3]
- 18 Zhu Z, Yu Z, Wang J, Zhou L, Zhang J, Yao B, Dou J, Qiu Z, Huang C. Krüppel-Like Factor 4 Inhibits Pancreatic Cancer Epithelial-to-Mesenchymal Transition and Metastasis by Down-Regulating Caveolin-1 Expression. Cell Physiol Biochem



2018; 46: 238-252 [PMID: 29587259 DOI: 10.1159/000488426]

- 19 Sekelsky JJ, Newfeld SJ, Raftery LA, Chartoff EH, Gelbart WM. Genetic characterization and cloning of mothers against dpp, a gene required for decapentaplegic function in Drosophila melanogaster. Genetics 1995; 139: 1347-1358 [PMID: 7768443 DOI: 10.1093/genetics/139.3.1347]
- 20 Derynck R, Gelbart WM, Harland RM, Heldin CH, Kern SE, Massagué J, Melton DA, Mlodzik M, Padgett RW, Roberts AB, Smith J, Thomsen GH, Vogelstein B, Wang XF. Nomenclature: vertebrate mediators of TGFbeta family signals. Cell 1996; 87: 173 [PMID: 8861901 DOI: 10.1016/s0092-8674(00)81335-5]
- 21 Singh P, Wig JD, Srinivasan R. The Smad family and its role in pancreatic cancer. Indian J Cancer 2011; 48: 351-360 [PMID: 21921337 DOI: 10.4103/0019-509X.84939]
- Wang Y, Yang Y, Wang Y, Li X, Xiao Y, Wang W. High Cancer Susceptibility Candidate 8 Expression Is Associated 22 With Poor Prognosis of Pancreatic Adenocarcinoma: Validated Analysis Based on Four Cancer Databases. Front Cell Dev Biol 2020; 8: 392 [PMID: 32582694 DOI: 10.3389/fcell.2020.00392]
- 23 Zhang Z, Wang L, Wang Q, Zhang M, Wang B, Jiang K, Ye Y, Wang S, Shen Z. Molecular Characterization and Clinical Relevance of RNA Binding Proteins in Colorectal Cancer. Front Genet 2020; 11: 580149 [PMID: 3319370] DOI: 10.3389/fgene.2020.580149]
- Jeon HS, Dracheva T, Yang SH, Meerzaman D, Fukuoka J, Shakoori A, Shilo K, Travis WD, Jen J. SMAD6 contributes to 24 patient survival in non-small cell lung cancer and its knockdown reestablishes TGF-beta homeostasis in lung cancer cells. Cancer Res 2008; 68: 9686-9692 [PMID: 19047146 DOI: 10.1158/0008-5472.CAN-08-1083]
- Yao W, Jiao Y, Zhou Y, Luo X. KLF13 suppresses the proliferation and growth of colorectal cancer cells through 25 transcriptionally inhibiting HMGCS1-mediated cholesterol biosynthesis. Cell Biosci 2020; 10: 76 [PMID: 32523679 DOI: 10.1186/s13578-020-00440-0





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

