

World Journal of *Clinical Cases*

World J Clin Cases 2022 November 6; 10(31): 11214-11664



Contents

Thrice Monthly Volume 10 Number 31 November 6, 2022

REVIEW

- 11214** Diabetes and skin cancers: Risk factors, molecular mechanisms and impact on prognosis
Dobrică EC, Banciu ML, Kipkorir V, Khazeei Tabari MA, Cox MJ, Simhachalam Kutikuppala LV, Găman MA
- 11226** Endocrine disruptor chemicals as obesogen and diabetogen: Clinical and mechanistic evidence
Kurşunoğlu NE, Sarer Yurekli BP
- 11240** Intestinal microbiota in the treatment of metabolically associated fatty liver disease
Wang JS, Liu JC

MINIREVIEWS

- 11252** Lactation mastitis: Promising alternative indicators for early diagnosis
Huang Q, Zheng XM, Zhang ML, Ning P, Wu MJ
- 11260** Clinical challenges of glycemic control in the intensive care unit: A narrative review
Sreedharan R, Martini A, Das G, Aftab N, Khanna S, Ruetzler K
- 11273** Concise review on short bowel syndrome: Etiology, pathophysiology, and management
Lakkasani S, Seth D, Khokhar I, Touza M, Dacosta TJ
- 11283** Role of nickel-regulated small RNA in modulation of *Helicobacter pylori* virulence factors
Freire de Melo F, Marques HS, Fellipe Bueno Lemos F, Silva Luz M, Rocha Pinheiro SL, de Carvalho LS, Souza CL, Oliveira MV
- 11292** Surgical intervention for acute pancreatitis in the COVID-19 era
Su YJ, Chen TH

ORIGINAL ARTICLE

Clinical and Translational Research

- 11299** Screening of traditional Chinese medicine monomers as ribonucleotide reductase M2 inhibitors for tumor treatment
Qin YY, Feng S, Zhang XD, Peng B

Case Control Study

- 11313** Covered transjugular intrahepatic portosystemic stent-shunt *vs* large volume paracentesis in patients with cirrhosis: A real-world propensity score-matched study
Dhaliwal A, Merhzad H, Karkhanis S, Tripathi D

Retrospective Cohort Study

- 11325** Endoscopic submucosal tunnel dissection for early esophageal squamous cell carcinoma in patients with cirrhosis: A propensity score analysis
Zhu LL, Liu LX, Wu JC, Gan T, Yang JL

Retrospective Study

- 11338** Nomogram for predicting overall survival in Chinese triple-negative breast cancer patients after surgery
Lin WX, Xie YN, Chen YK, Cai JH, Zou J, Zheng JH, Liu YY, Li ZY, Chen YX
- 11349** Early patellar tendon rupture after total knee arthroplasty: A direct repair method
Li TJ, Sun JY, Du YQ, Shen JM, Zhang BH, Zhou YG
- 11358** Coxsackievirus A6 was the most common enterovirus serotype causing hand, foot, and mouth disease in Shiyang City, central China
Li JF, Zhang CJ, Li YW, Li C, Zhang SC, Wang SS, Jiang Y, Luo XB, Liao XJ, Wu SX, Lin L
- 11371** Dynamic changes of estimated glomerular filtration rate are conversely related to triglyceride in non-overweight patients
Liu SQ, Zhang XJ, Xue Y, Huang R, Wang J, Wu C, He YS, Pan YR, Liu LG
- 11381** C-reactive protein as a non-linear predictor of prolonged length of intensive care unit stay after gastrointestinal cancer surgery
Yan YM, Gao J, Jin PL, Lu JJ, Yu ZH, Hu Y

Clinical Trials Study

- 11391** Dan Bai Xiao Formula combined with glucocorticoids and cyclophosphamide for pediatric lupus nephritis: A pilot prospective study
Cao TT, Chen L, Zhen XF, Zhao GJ, Zhang HF, Hu Y

Observational Study

- 11403** Relationship between lipids and sleep apnea: Mendelian randomization analysis
Zhang LP, Zhang XX
- 11411** Efficacy and safety profile of two-dose SARS-CoV-2 vaccines in cancer patients: An observational study in China
Cai SW, Chen JY, Wan R, Pan DJ, Yang WL, Zhou RG

Prospective Study

- 11419** Pressure changes in tapered and cylindrical shaped cuff after extension of head and neck: A randomized controlled trial
Seol G, Jin J, Oh J, Byun SH, Jeon Y

Randomized Controlled Trial

- 11427** Effect of intradermal needle therapy at combined acupoints on patients' gastrointestinal function following surgery for gastrointestinal tumors
Guo M, Wang M, Chen LL, Wei FJ, Li JE, Lu QX, Zhang L, Yang HX

SYSTEMATIC REVIEWS

- 11442** Video-assisted bystander cardiopulmonary resuscitation improves the quality of chest compressions during simulated cardiac arrests: A systemic review and meta-analysis

Pan DF, Li ZJ, Ji XZ, Yang LT, Liang PF

META-ANALYSIS

- 11454** Efficacy of the femoral neck system in femoral neck fracture treatment in adults: A systematic review and meta-analysis

Wu ZF, Luo ZH, Hu LC, Luo YW

- 11466** Prevalence of polymyxin-induced nephrotoxicity and its predictors in critically ill adult patients: A meta-analysis

Wang JL, Xiang BX, Song XL, Que RM, Zuo XC, Xie YL

CASE REPORT

- 11486** Novel compound heterozygous variants in the LHX3 gene caused combined pituitary hormone deficiency: A case report

Lin SZ, Ma QJ, Pang QM, Chen QD, Wang WQ, Li JY, Zhang SL

- 11493** Fatal bleeding due to an aorto-esophageal fistula: A case report and literature review

Ćeranić D, Nikolić S, Lučev J, Slanić A, Bujas T, Ocepek A, Skok P

- 11500** Tolvaptan ameliorated kidney function for one elderly autosomal dominant polycystic kidney disease patient: A case report

Zhou L, Tian Y, Ma L, Li WG

- 11508** Extensive right coronary artery thrombosis in a patient with COVID-19: A case report

Dall'Orto CC, Lopes RPF, Cancela MT, de Sales Padilha C, Pinto Filho GV, da Silva MR

- 11517** Yokoyama procedure for a woman with heavy eye syndrome who underwent multiple recession-resection operations: A case report

Yao Z, Jiang WL, Yang X

- 11523** Rectal cancer combined with abdominal tuberculosis: A case report

Liu PG, Chen XF, Feng PF

- 11529** Malignant obstruction in the ileocecal region treated by self-expandable stent placement under the fluoroscopic guidance: A case report

Wu Y, Li X, Xiong F, Bao WD, Dai YZ, Yue LJ, Liu Y

- 11536** Granulocytic sarcoma with long spinal cord compression: A case report

Shao YD, Wang XH, Sun L, Cui XG

- 11542** Aortic dissection with epileptic seizure: A case report

Zheng B, Huang XQ, Chen Z, Wang J, Gu GF, Luo XJ

- 11549** Multiple bilateral and symmetric C1-2 ganglioneuromas: A case report
Wang S, Ma JX, Zheng L, Sun ST, Xiang LB, Chen Y
- 11555** Acute myocardial infarction due to Kounis syndrome: A case report
Xu GZ, Wang G
- 11561** Surgical excision of a large retroperitoneal lymphangioma: A case report
Park JH, Lee D, Maeng YH, Chang WB
- 11567** Mass-like extragonadal endometriosis associated malignant transformation in the pelvis: A rare case report
Chen P, Deng Y, Wang QQ, Xu HW
- 11574** Gastric ulcer treated using an elastic traction ring combined with clip: A case report
Pang F, Song YJ, Sikong YH, Zhang AJ, Zuo XL, Li RY
- 11579** Novel liver vein deprivation technique that promotes increased residual liver volume (with video): A case report
Wu G, Jiang JP, Cheng DH, Yang C, Liao DX, Liao YB, Lau WY, Zhang Y
- 11585** Linear porokeratosis of the foot with dermoscopic manifestations: A case report
Yang J, Du YQ, Fang XY, Li B, Xi ZQ, Feng WL
- 11590** Primary hepatic angiosarcoma: A case report
Wang J, Sun LT
- 11597** Hemorrhagic shock due to ruptured lower limb vascular malformation in a neurofibromatosis type 1 patient: A case report
Shen LP, Jin G, Zhu RT, Jiang HT
- 11607** Gastric linitis plastica with autoimmune pancreatitis diagnosed by an endoscopic ultrasonography-guided fine-needle biopsy: A case report
Sato R, Matsumoto K, Kanzaki H, Matsumi A, Miyamoto K, Morimoto K, Terasawa H, Fujii Y, Yamazaki T, Uchida D, Tsutsumi K, Horiguchi S, Kato H
- 11617** Favorable response of primary pulmonary lymphoepithelioma-like carcinoma to sintilimab combined with chemotherapy: A case report
Zeng SY, Yuan J, Lv M
- 11625** Benign paroxysmal positional vertigo with congenital nystagmus: A case report
Li GF, Wang YT, Lu XG, Liu M, Liu CB, Wang CH
- 11630** Secondary craniofacial necrotizing fasciitis from a distant septic emboli: A case report
Lee DW, Kwak SH, Choi HJ
- 11638** Pancreatic paraganglioma with multiple lymph node metastases found by spectral computed tomography: A case report and review of the literature
Li T, Yi RQ, Xie G, Wang DN, Ren YT, Li K

- 11646** Apnea caused by retrobulbar anesthesia: A case report
Wang YL, Lan GR, Zou X, Wang EQ, Dai RP, Chen YX
- 11652** Unexplained septic shock after colonoscopy with polyethylene glycol preparation in a young adult: A case report
Song JJ, Wu CJ, Dong YY, Ma C, Gu Q
- 11658** Metachronous isolated penile metastasis from sigmoid colon adenocarcinoma: A case report
Yin GL, Zhu JB, Fu CL, Ding RL, Zhang JM, Lin Q

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Muhammad Hamdan Gul, MD, Assistant Professor, Department of Internal Medicine, University of Kentucky, Chicago, IL 60657, United States.
hamdan3802@hotmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Xu Guo*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

November 6, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Benign paroxysmal positional vertigo with congenital nystagmus: A case report

Gui-Fang Li, Yue-Tang Wang, Xin-Ge Lu, Man Liu, Chao-Bing Liu, Chun-Hua Wang

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): C, C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Bhandari A, United States; Faraji N, Iran

Received: July 20, 2022

Peer-review started: July 20, 2022

First decision: August 4, 2022

Revised: August 15, 2022

Accepted: September 22, 2022

Article in press: September 22, 2022

Published online: November 6, 2022



Gui-Fang Li, Department of Otolaryngology, Hebei Provincial Eye Hospital, Hebei Provincial Key Laboratory of Ophthalmology, Hebei Provincial Eye Institute, Xingtai 054001, Hebei Province, China

Yue-Tang Wang, Xin-Ge Lu, Man Liu, Chao-Bing Liu, Chun-Hua Wang, Department of Ophthalmology, Hebei Provincial Eye Hospital, Hebei Provincial Key Laboratory of Ophthalmology, Hebei Provincial Eye Institute, Xingtai 054001, Hebei Province, China

Corresponding author: Gui-Fang Li, MPhil, Chief Physician, Department of Otolaryngology, Hebei Provincial Eye Hospital, Hebei Provincial Key Laboratory of Ophthalmology, Hebei Provincial Eye Institute, No. 399 Quanbei East Street, Xingtai 054001, Hebei Province, China. lifang9922@163.com

Abstract

BACKGROUND

Benign paroxysmal positional vertigo (BPPV) is a form of temporary vertigo induced by moving the head to a specific position. It is a self-limited, peripheral, vestibular disease and can be divided into primary and secondary forms. Congenital nystagmus (CN), an involuntary, rhythmic, binocular-symmetry, conjugated eye movement, is found at birth or within 3 mo of birth. According to the pathogenesis, CN can be divided into sensory-defect nystagmus and motor-defect nystagmus. The coexistence of BPPV and CN is rarely seen in the clinic.

CASE SUMMARY

A 62-year-old woman presented to our clinic complaining of a 15-d history of recurrent positional vertigo. The vertigo lasting less than 1 min occurred when she turned over, sometimes accompanied by nausea and vomiting. Both the patient and her father had CN. Her spontaneous nystagmus was horizontal to right; however, the gaze test revealed variable horizontal nystagmus with the same degree when the eyes moved. The patient's Dix-Hallpike test was normal, except for persistent nystagmus, and the roll test showed severe variable horizontal nystagmus, which lasted for about 20 s in the same direction as her head movement to the right and left, although the right-side nystagmus was stronger than the left-side. Since these symptoms were accompanied by nausea, she was diagnosed with BPPV with CN and treated by manual reduction.

CONCLUSION

Though rare, if BPPV with CN is correctly identified and diagnosed, reduction treatment is comparably effective to other vertigo types.

Key Words: Congenital; Nystagmus; Benign paroxysmal positional vertigo; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Benign paroxysmal positional vertigo (BPPV) is defined as a disorder of the inner ear characterized by repeated episodes of positional vertigo. Congenital nystagmus (CN), an involuntary, rhythmic, binocular-symmetry, conjugated eye movement, is found at birth or within 3 mo of birth. BPPV with CN is rarely seen in the clinic. CN should be distinguished from other pathologic nystagmus types. BPPV can be accurately determined through postural nystagmus. We report the characteristics of BPPV with CN, further explaining how to identify nystagmus.

Citation: Li GF, Wang YT, Lu XG, Liu M, Liu CB, Wang CH. Benign paroxysmal positional vertigo with congenital nystagmus: A case report. *World J Clin Cases* 2022; 10(31): 11625-11629

URL: <https://www.wjgnet.com/2307-8960/full/v10/i31/11625.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i31.11625>

INTRODUCTION

Benign paroxysmal positional vertigo (BPPV) is defined as a disorder of the inner ear characterized by repeated episodes of positional vertigo[1]. BPPV is among the common diseases that cause aural vertigo, and 24.1% of patients with dizziness or vertigo are diagnosed with BPPV[2]. Spontaneous nystagmus refers to a continuous, involuntary, and rhythmic movement of the eyeball in the absence of inducing factors and is divided into congenital nystagmus (CN) and acquired nystagmus. The latter is a type of nystagmus commonly seen in the clinic. CN is an ocular motor disorder in which patients are afflicted by periodic involuntary ocular oscillations affecting both eyes[3,4]. It develops during the first 3 to 6 mo of a patient's life and has a prevalence of 14 per 10000 people in the United Kingdom[5]. The etiology of CN is largely unknown, but we know that most patients have lifelong nystagmus, although it can gradually relieve with age in some patients. Some BPPV patients present with spontaneous nystagmus, but BPPV with CN is rare: To date, we have not seen such cases reported. In this report, we present the case of a BPPV patient with CN and her nystagmus findings.

CASE PRESENTATION

Chief complaints

A 62-year-old woman presented at our clinic, complaining of positional vertigo that had recurred for 15 d.

History of present illness

Fifteen days previously, the patient's symptoms had begun with severe dizziness when she rose, which recurred when she rolled over or lied down. She sometimes experienced nausea and vomiting at the onset of these symptoms.

History of past illness

The patient was physically healthy in the past.

Personal and family history

Both the patient and her father had a history of CN.

Physical examination

The patient's physical examination revealed no abnormal findings.

Laboratory examinations

The patient did not undergo laboratory examinations.

Imaging examinations

The patient did not undergo imaging examinations.

FINAL DIAGNOSIS

The patient was ultimately diagnosed with BPPV with CN.

TREATMENT

The patient was prescribed a barbecue roll maneuver to treat her right, lateral, semicircular canal BPPV of geotropic type. This treatment required her to lie down with the affected ear facing downward. Then, she rolled over to the opposite side for 90 degrees until she returned to the original position. Her vertigo symptoms disappeared after the therapy.

OUTCOME AND FOLLOW-UP

The patient's follow-up comprised three telephone appointments at 1 wk, 1 mo, and 6 mo after treatment. She was asymptomatic, without any recurrence of vertigo.

DISCUSSION

BPPV is generally categorized as posterior semicircular canal, anterior semicircular canal, and horizontal semicircular canal types. Of these categories, posterior semicircular canal BPPV is the most common (affecting 80%-90% of patients), followed by horizontal semicircular canal BPPV (10%-20%), while anterior semicircular canal BPPV is rare (3%)[6,7]. The Dix-Hallpike maneuver is considered the gold standard test to diagnose posterior canal BPPV, and the supine roll test is considered the gold standard for diagnosing horizontal semicircular canal BPPV[8]. Upbeat-torsional nystagmus is provoked by vertical semicircular canal BPPV.

The pathogenesis of BPPV remains unclear; however, risk factors include age, mental stress, osteoporosis, insomnia, and hypertension[9,10]. Currently, the following two theories are widely accepted. First, canalithiasis suggests that when the head is moved relative to gravity, otoliths residing on the macula utriculi migrate into the semicircular canal and are displaced relative to the semicircular canal wall because of gravity, causing endolymph flow and resulting in the deviation of the cupula terminalis and, in turn, corresponding signs and symptoms. When the otolith moves due to gravity to the lowest point in the semicircular canal lumen, the endolymph stops, the cupula terminalis returns to its original position, and signs and symptoms disappear.

Second, eupulolithiasis suggests that the detached otoliths on the macula utriculi adhere to the cupula terminalis, changing the density of the latter relative to the endolymph and making it sensitive to gravity, resulting in the corresponding symptoms and signs[11]. CN usually occurs at birth or within 3 mo of birth. Although this nystagmus persists throughout most patients' lives, some patients' symptoms gradually relieve with age. CN is divided into two categories. The first is congenital motor defect nystagmus, in which eye movement includes fast and slow phases. The second is congenital sensory defect nystagmus, also known as "pendular nystagmus", in which the eye moves at one speed. CN is clinically rare, with an incidence of about 0.005%-0.286%[3]. To our knowledge, BPPV with CN has not been previously reported.

In the United States, according to statistics, about 5.6 million patients clinically complain of dizziness per year, and 17%-42% of patients with vertigo are diagnosed with BPPV[12-14]. BPPV treatment can be categorized as canalith repositioning maneuvers or vestibular rehabilitation[1]. Diagnosis of horizontal semicircular canal BPPV relies on the supine roll test. During examinations, clinicians should observe whether the direction of the nystagmus is geotropic or apogeotropic and which side of the nystagmus is stronger to enable identification of the patient's affected side. In our case, the Dix-Hallpike test showed the signs of horizontal nystagmus without vertigo and geotropic nystagmus, which was stronger on her right side during the roll test (Figure 1). Unlike other patients with BPPV, she exhibited persistent horizontal nystagmus on the right side after intense nystagmus lasting more than 10 s, which was accompanied by vertigo. The lasting nystagmus is suggestive of CN, which is similar to that observed in patients with spontaneous nystagmus but absent in BPPV patients without spontaneous nystagmus. Spontaneous nystagmus is very common clinically among patients with vertigo.

The following aspects should be used to distinguish CN from other central and peripheral spontaneous forms of nystagmus. The first and most important is a patient's medical history. Nystagmus is always present during CN; someone in the patient's family will have CN because of the heritability of the disease, while other types of spontaneous nystagmus only appear at the onset of the disease. Second, the direction of peripheral nystagmus is constant, while that of CN is variable. The test can be conducted with Frenzel glasses to observe the nystagmus accurately. The direction of the nystagmus will be seen to remain the same in the peripheral nystagmus; however, the direction of the




Figure 1 Nystagmus in the roll test. A and B: When the head was to the right, intensive nystagmus occurred between 13 s and 32 s with dizziness and nausea; the highest slow phase velocity (SPV) was 162°/s and then weakened gradually to 8°/s, and vertigo disappeared; C and D: When the head was to the left, the intensive nystagmus occurred between 9 s and 30 s with dizziness; the highest SPV was 62°/s and then weakened gradually to 9°/s, and vertigo disappeared.




Figure 2 Spontaneous nystagmus and nystagmus in the gaze test. A: The intensity of spontaneous nystagmus was 20°/s, and it was constant; B: When the eyes moved to right, the direction of nystagmus changed to left, and the slow phase velocity (SPV) of nystagmus was 7°/s; C: When the eyes moved to left, the nystagmus was still to the left, and the SPV of nystagmus was 8°/s.

nystagmus is consistent with the eye movement in central nystagmus and variable nystagmus in CN (Figure 2). Third, a CN patient usually experiences horizontal nystagmus of variable intensity, while other pathologic central nystagmus types may entail vertical and horizontal nystagmus with generally persistent intensity.

CONCLUSION

CN is rare in the clinic. If individuals experience spontaneous nystagmus with constant intensity and variable direction, a careful medical history should be taken to eliminate CN, which may influence the diagnosis. The treatment for BPPV with CN is the same as that for BPPV. In the case reported here, the patient was diagnosed with BPPV with CN, and the result was good.

FOOTNOTES

Author contributions: Liu M and Liu CB were the otolaryngologists for the patient; Li GF and Wang CH analyzed the case, reviewed the literature, and contributed to drafting the manuscript; Wang YT and Lu XG were responsible for manual reduction; and all authors gave their approval for the submission of the final version of the manuscript.

Informed consent statement: Informed written consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Gui-Fang Li 0000-0001-7326-3491.

S-Editor: Wang JJ

L-Editor: Wang TQ

P-Editor: Wang JJ

REFERENCES

- 1 **Bhattacharyya N**, Gubbels SP, Schwartz SR, Edlow JA, El-Kashlan H, Fife T, Holmberg JM, Mahoney K, Hollingsworth DB, Roberts R, Seidman MD, Steiner RW, Do BT, Voelker CC, Waguespack RW, Corrigan MD. Clinical Practice Guideline: Benign Paroxysmal Positional Vertigo (Update). *Otolaryngol Head Neck Surg* 2017; **156**: S1-S47 [PMID: 28248609 DOI: 10.1177/0194599816689667]
- 2 **Kim HJ**, Lee JO, Choi JY, Kim JS. Etiologic distribution of dizziness and vertigo in a referral-based dizziness clinic in South Korea. *J Neurol* 2020; **267**: 2252-2259 [PMID: 32300888 DOI: 10.1007/s00415-020-09831-2]
- 3 **Casteels I**, Harris CM, Shawkat F, Taylor D. Nystagmus in infancy. *Br J Ophthalmol* 1992; **76**: 434-437 [PMID: 1627515 DOI: 10.1136/bjo.76.7.434]
- 4 **Abadi RV**, Bjerre A. Motor and sensory characteristics of infantile nystagmus. *Br J Ophthalmol* 2002; **86**: 1152-1160 [PMID: 12234898 DOI: 10.1136/bjo.86.10.1152]
- 5 **Proudlock FA**, Gottlob I. Nystagmus in childhood. In: Hoyt C, Taylor D. *Pediatric Ophthalmology and Strabismus*, 4th ed. Philadelphia: Saunders, 2013: 922-932
- 6 **Kim JS**, Zee DS. Clinical practice. Benign paroxysmal positional vertigo. *N Engl J Med* 2014; **370**: 1138-1147 [PMID: 24645946 DOI: 10.1056/NEJMcpl309481]
- 7 **Hornibrook J**. Benign Paroxysmal Positional Vertigo (BPPV): History, Pathophysiology, Office Treatment and Future Directions. *Int J Otolaryngol* 2011; **2011**: 835671 [PMID: 21808648 DOI: 10.1155/2011/835671]
- 8 **Fife TD**, Iverson DJ, Lempert T, Furman JM, Baloh RW, Tusa RJ, Hain TC, Herdman S, Morrow MJ, Gronseth GS; Quality Standards Subcommittee, American Academy of Neurology. Practice parameter: therapies for benign paroxysmal positional vertigo (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2008; **70**: 2067-2074 [PMID: 18505980 DOI: 10.1212/01.wnl.0000313378.77444.ac]
- 9 **Byun H**, Chung JH, Lee SH, Park CW, Kim EM, Kim I. Increased risk of benign paroxysmal positional vertigo in osteoporosis: a nationwide population-based cohort study. *Sci Rep* 2019; **9**: 3469 [PMID: 30837524 DOI: 10.1038/s41598-019-39830-x]
- 10 **Chen J**, Zhang S, Cui K, Liu C. Risk factors for benign paroxysmal positional vertigo recurrence: a systematic review and meta-analysis. *J Neurol* 2021; **268**: 4117-4127 [PMID: 32839838 DOI: 10.1007/s00415-020-10175-0]
- 11 **Editorial Board of Chinese Journal of Otorhinolaryngology Head and Neck Surgery**; Chinese Medical Association of Otorhinolaryngology Head and Neck Surgery. Clinical Guidelines: Benign Paroxysmal Positional Vertigo. *Chin J Otorhinolaryngol Head Neck Surg* 2017; **52**: 173-177
- 12 **Schappert SM**. National Ambulatory Medical Care Survey: 1989 summary. *Vital Health Stat* 13 1992; **1-80** [PMID: 1376543]
- 13 **Katsarkas A**. Benign paroxysmal positional vertigo (BPPV): idiopathic versus post-traumatic. *Acta Otolaryngol* 1999; **119**: 745-749 [PMID: 10687929 DOI: 10.1080/00016489950180360]
- 14 **Hanley K**, O'Dowd T, Considine N. A systematic review of vertigo in primary care. *Br J Gen Pract* 2001; **51**: 666-671 [PMID: 11510399]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

