

# World Journal of *Clinical Cases*

*World J Clin Cases* 2022 November 6; 10(31): 11214-11664



## Contents

Thrice Monthly Volume 10 Number 31 November 6, 2022

## REVIEW

- 11214 Diabetes and skin cancers: Risk factors, molecular mechanisms and impact on prognosis  
*Dobrică EC, Banciu ML, Kipkorir V, Khazeei Tabari MA, Cox MJ, Simhachalam Kutikuppala LV, Găman MA*
- 11226 Endocrine disruptor chemicals as obesogen and diabetogen: Clinical and mechanistic evidence  
*Kurşunoğlu NE, Sarer Yurekli BP*
- 11240 Intestinal microbiota in the treatment of metabolically associated fatty liver disease  
*Wang JS, Liu JC*

## MINIREVIEWS

- 11252 Lactation mastitis: Promising alternative indicators for early diagnosis  
*Huang Q, Zheng XM, Zhang ML, Ning P, Wu MJ*
- 11260 Clinical challenges of glycemic control in the intensive care unit: A narrative review  
*Sreedharan R, Martini A, Das G, Aftab N, Khanna S, Ruetzler K*
- 11273 Concise review on short bowel syndrome: Etiology, pathophysiology, and management  
*Lakkasani S, Seth D, Khokhar I, Touza M, Dacosta TJ*
- 11283 Role of nickel-regulated small RNA in modulation of *Helicobacter pylori* virulence factors  
*Freire de Melo F, Marques HS, Fellipe Bueno Lemos F, Silva Luz M, Rocha Pinheiro SL, de Carvalho LS, Souza CL, Oliveira MV*
- 11292 Surgical intervention for acute pancreatitis in the COVID-19 era  
*Su YJ, Chen TH*

## ORIGINAL ARTICLE

## Clinical and Translational Research

- 11299 Screening of traditional Chinese medicine monomers as ribonucleotide reductase M2 inhibitors for tumor treatment  
*Qin YY, Feng S, Zhang XD, Peng B*

## Case Control Study

- 11313 Covered transjugular intrahepatic portosystemic stent-shunt vs large volume paracentesis in patients with cirrhosis: A real-world propensity score-matched study  
*Dhaliwal A, Merhzad H, Karkhanis S, Tripathi D*

**Retrospective Cohort Study**

- 11325** Endoscopic submucosal tunnel dissection for early esophageal squamous cell carcinoma in patients with cirrhosis: A propensity score analysis  
*Zhu LL, Liu LX, Wu JC, Gan T, Yang JL*

**Retrospective Study**

- 11338** Nomogram for predicting overall survival in Chinese triple-negative breast cancer patients after surgery  
*Lin WX, Xie YN, Chen YK, Cai JH, Zou J, Zheng JH, Liu YY, Li ZY, Chen YX*
- 11349** Early patellar tendon rupture after total knee arthroplasty: A direct repair method  
*Li TJ, Sun JY, Du YQ, Shen JM, Zhang BH, Zhou YG*
- 11358** Coxsackievirus A6 was the most common enterovirus serotype causing hand, foot, and mouth disease in Shiyan City, central China  
*Li JF, Zhang CJ, Li YW, Li C, Zhang SC, Wang SS, Jiang Y, Luo XB, Liao XJ, Wu SX, Lin L*
- 11371** Dynamic changes of estimated glomerular filtration rate are conversely related to triglyceride in non-overweight patients  
*Liu SQ, Zhang XJ, Xue Y, Huang R, Wang J, Wu C, He YS, Pan YR, Liu LG*
- 11381** C-reactive protein as a non-linear predictor of prolonged length of intensive care unit stay after gastrointestinal cancer surgery  
*Yan YM, Gao J, Jin PL, Lu JJ, Yu ZH, Hu Y*

**Clinical Trials Study**

- 11391** Dan Bai Xiao Formula combined with glucocorticoids and cyclophosphamide for pediatric lupus nephritis: A pilot prospective study  
*Cao TT, Chen L, Zhen XF, Zhao GJ, Zhang HF, Hu Y*

**Observational Study**

- 11403** Relationship between lipids and sleep apnea: Mendelian randomization analysis  
*Zhang LP, Zhang XX*
- 11411** Efficacy and safety profile of two-dose SARS-CoV-2 vaccines in cancer patients: An observational study in China  
*Cai SW, Chen JY, Wan R, Pan DJ, Yang WL, Zhou RG*

**Prospective Study**

- 11419** Pressure changes in tapered and cylindrical shaped cuff after extension of head and neck: A randomized controlled trial  
*Seol G, Jin J, Oh J, Byun SH, Jeon Y*

**Randomized Controlled Trial**

- 11427** Effect of intradermal needle therapy at combined acupoints on patients' gastrointestinal function following surgery for gastrointestinal tumors  
*Guo M, Wang M, Chen LL, Wei FJ, Li JE, Lu QX, Zhang L, Yang HX*

**SYSTEMATIC REVIEWS**

- 11442** Video-assisted bystander cardiopulmonary resuscitation improves the quality of chest compressions during simulated cardiac arrests: A systemic review and meta-analysis  
*Pan DF, Li ZJ, Ji XZ, Yang LT, Liang PF*

**META-ANALYSIS**

- 11454** Efficacy of the femoral neck system in femoral neck fracture treatment in adults: A systematic review and meta-analysis  
*Wu ZF, Luo ZH, Hu LC, Luo YW*
- 11466** Prevalence of polymyxin-induced nephrotoxicity and its predictors in critically ill adult patients: A meta-analysis  
*Wang JL, Xiang BX, Song XL, Que RM, Zuo XC, Xie YL*

**CASE REPORT**

- 11486** Novel compound heterozygous variants in the LHX3 gene caused combined pituitary hormone deficiency: A case report  
*Lin SZ, Ma QJ, Pang QM, Chen QD, Wang WQ, Li JY, Zhang SL*
- 11493** Fatal bleeding due to an aorto-esophageal fistula: A case report and literature review  
*Ćeranić D, Nikolić S, Lučev J, Slanić A, Bujas T, Ocepek A, Skok P*
- 11500** Tolvaptan ameliorated kidney function for one elderly autosomal dominant polycystic kidney disease patient: A case report  
*Zhou L, Tian Y, Ma L, Li WG*
- 11508** Extensive right coronary artery thrombosis in a patient with COVID-19: A case report  
*Dall'Orto CC, Lopes RPF, Cancela MT, de Sales Padilha C, Pinto Filho GV, da Silva MR*
- 11517** Yokoyama procedure for a woman with heavy eye syndrome who underwent multiple recession-resection operations: A case report  
*Yao Z, Jiang WL, Yang X*
- 11523** Rectal cancer combined with abdominal tuberculosis: A case report  
*Liu PG, Chen XF, Feng PF*
- 11529** Malignant obstruction in the ileocecal region treated by self-expandable stent placement under the fluoroscopic guidance: A case report  
*Wu Y, Li X, Xiong F, Bao WD, Dai YZ, Yue LJ, Liu Y*
- 11536** Granulocytic sarcoma with long spinal cord compression: A case report  
*Shao YD, Wang XH, Sun L, Cui XG*
- 11542** Aortic dissection with epileptic seizure: A case report  
*Zheng B, Huang XQ, Chen Z, Wang J, Gu GF, Luo XJ*

- 11549** Multiple bilateral and symmetric C1-2 ganglioneuromas: A case report  
*Wang S, Ma JX, Zheng L, Sun ST, Xiang LB, Chen Y*
- 11555** Acute myocardial infarction due to Kounis syndrome: A case report  
*Xu GZ, Wang G*
- 11561** Surgical excision of a large retroperitoneal lymphangioma: A case report  
*Park JH, Lee D, Maeng YH, Chang WB*
- 11567** Mass-like extragonadal endometriosis associated malignant transformation in the pelvis: A rare case report  
*Chen P, Deng Y, Wang QQ, Xu HW*
- 11574** Gastric ulcer treated using an elastic traction ring combined with clip: A case report  
*Pang F, Song YJ, Sikong YH, Zhang AJ, Zuo XL, Li RY*
- 11579** Novel liver vein deprivation technique that promotes increased residual liver volume (with video): A case report  
*Wu G, Jiang JP, Cheng DH, Yang C, Liao DX, Liao YB, Lau WY, Zhang Y*
- 11585** Linear porokeratosis of the foot with dermoscopic manifestations: A case report  
*Yang J, Du YQ, Fang XY, Li B, Xi ZQ, Feng WL*
- 11590** Primary hepatic angiosarcoma: A case report  
*Wang J, Sun LT*
- 11597** Hemorrhagic shock due to ruptured lower limb vascular malformation in a neurofibromatosis type 1 patient: A case report  
*Shen LP, Jin G, Zhu RT, Jiang HT*
- 11607** Gastric linitis plastica with autoimmune pancreatitis diagnosed by an endoscopic ultrasonography-guided fine-needle biopsy: A case report  
*Sato R, Matsumoto K, Kanzaki H, Matsumi A, Miyamoto K, Morimoto K, Terasawa H, Fujii Y, Yamazaki T, Uchida D, Tsutsumi K, Horiguchi S, Kato H*
- 11617** Favorable response of primary pulmonary lymphoepithelioma-like carcinoma to sintilimab combined with chemotherapy: A case report  
*Zeng SY, Yuan J, Lv M*
- 11625** Benign paroxysmal positional vertigo with congenital nystagmus: A case report  
*Li GF, Wang YT, Lu XG, Liu M, Liu CB, Wang CH*
- 11630** Secondary craniofacial necrotizing fasciitis from a distant septic emboli: A case report  
*Lee DW, Kwak SH, Choi HJ*
- 11638** Pancreatic paraganglioma with multiple lymph node metastases found by spectral computed tomography: A case report and review of the literature  
*Li T, Yi RQ, Xie G, Wang DN, Ren YT, Li K*

- 11646** Apnea caused by retrobulbar anesthesia: A case report  
*Wang YL, Lan GR, Zou X, Wang EQ, Dai RP, Chen YX*
- 11652** Unexplained septic shock after colonoscopy with polyethylene glycol preparation in a young adult: A case report  
*Song JJ, Wu CJ, Dong YY, Ma C, Gu Q*
- 11658** Metachronous isolated penile metastasis from sigmoid colon adenocarcinoma: A case report  
*Yin GL, Zhu JB, Fu CL, Ding RL, Zhang JM, Lin Q*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Muhammad Hamdan Gul, MD, Assistant Professor, Department of Internal Medicine, University of Kentucky, Chicago, IL 60657, United States.  
hamdan3802@hotmail.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

*WJCC* mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for *WJCC* as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The *WJCC*'s CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Xu Guo*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

November 6, 2022

**COPYRIGHT**

© 2022 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Secondary craniofacial necrotizing fasciitis from a distant septic emboli: A case report

Da-Woon Lee, Si-Hyun Kwak, Hwan-Jun Choi

**Specialty type:** Medicine, research and experimental

**Provenance and peer review:** Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B, B  
Grade C (Good): 0  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Jian X, China; Shelat VG, Singapore

**Received:** July 22, 2022

**Peer-review started:** July 22, 2022

**First decision:** August 22, 2022

**Revised:** August 23, 2022

**Accepted:** September 22, 2022

**Article in press:** September 22, 2022

**Published online:** November 6, 2022



**Da-Woon Lee, Si-Hyun Kwak, Hwan-Jun Choi**, Department of Plastic and Reconstructive Surgery, College of Medicine, Soonchunhyang University, Cheonan-si 31151, South Korea

**Hwan-Jun Choi**, Institute of Tissue Regeneration, College of Medicine, Soonchunhyang University, Cheonan-si 31151, South Korea

**Corresponding author:** Hwan-Jun Choi, MD, PhD, Professor, Department of Plastic and Reconstructive Surgery, College of Medicine, Soonchunhyang University, Soonchunhyang University Cheonan Hospital, 31, Sooncheonyang 6-Gil, Dongnam-gu, Cheonan-si 31151, South Korea. [iprskorea@gmail.com](mailto:iprskorea@gmail.com)

### Abstract

#### BACKGROUND

Craniofacial necrotizing fasciitis (CNF) is an uncommon but fatal infection that can spread rapidly through the subfascial planes in the head and neck region. Symptoms usually progress rapidly, and early management is necessary to optimize outcomes.

#### CASE SUMMARY

A 43-year-old man visited our hospital with left hemifacial swelling involving the buccal and submandibular areas. The patient had fever for approximately 10 d before visiting the hospital, but did not report any other systemic symptoms. Computed tomography scan demonstrated an abscess with gas formation. After surgical drainage of the facial abscess, the patient's systemic condition worsened and progressed to septic shock. Further examination revealed pulmonary and renal abscesses. Renal percutaneous catheter drainage was performed at the renal abscess site, which caused improvement of symptoms. The patient showed no evidence of systemic complications during the 4-mo post-operative follow-up period.

#### CONCLUSION

As the patient did not improve with conventional CNF treatment and symptoms only resolved after controlling the infection, the final diagnosis was secondary CNF with septic emboli. Aggressive surgical decompression is important for CNF management. However, if symptoms worsen despite early diagnosis and management, such as pus drainage and surgical intervention, clinicians should consider the possibility of a secondary abscess from internal organs.

**Key Words:** Craniofacial necrotizing fasciitis; Septic emboli; *Klebsiella pneumoniae*; Secondary; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Craniofacial necrotizing fasciitis (CNF) is a severe infection that can rapidly spread and progress to life-threatening conditions; therefore, early diagnosis and appropriate management are essential. It can occur mainly due to odontogenic infection, but many other causes lead to development of symptoms. This report describes a rare case of secondary necrotizing fasciitis originating from internal organ abscesses. It should be considered that septic emboli from other internal organs can cause a secondary CNF if there is no improvement of symptoms after radical surgical management.

**Citation:** Lee DW, Kwak SH, Choi HJ. Secondary craniofacial necrotizing fasciitis from a distant septic emboli: A case report. *World J Clin Cases* 2022; 10(31): 11630-11637

**URL:** <https://www.wjgnet.com/2307-8960/full/v10/i31/11630.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v10.i31.11630>

## INTRODUCTION

Craniofacial necrotizing fasciitis (CNF) is a rare and life-threatening condition characterized by the rapid progressive necrosis of the fascia, fat, and muscle. Fascial necrosis is a key feature of the disease progression. CNF is a lethal disease with a mortality rate of 20%-40%[1]. Although CNFs are rare, they are often associated with systemic conditions owing to the abundant vascularity of the head and neck region. Based on a literature review, CNF is mainly caused by odontogenic infections after tooth extraction, especially in patients with chronic diseases such as diabetes, immunosuppression, and elderly patients[2]. It can also be caused by peritonsillar abscesses, skin trauma due to insect bites, and fractures[3]. In most wound cultures of necrotizing fasciitis, group A *Streptococcus* bacteria are the most common pathogens[4,5]. Management of necrotizing fasciitis includes early detection, broad-spectrum antibiotics, and most importantly, radical surgical management. Early detection of CNF can depend on findings such as air bubbles in soft tissues on computed tomography (CT) and rapid progression of infection. The use of appropriate antibiotics based on bacterial cultures is vital in improving patient conditions[6]. However, CNFs are frequently difficult to manage despite aggressive treatment; thus, leading to significant morbidity and potential mortality[7].

In this report, we describe a rare case of CNF as a secondary presentation in a 43-year-old male patient who developed necrotizing fasciitis due to hematogenic spread from lung and renal abscesses. This patient met the diagnostic criteria of necrotizing fasciitis, unlike Ludwig's angina which is bilateral inflammation that invades the submandibular area, including evidence of necrotizing fascia, characteristic pathological features (extensive tissue necrosis, pattern of infection spreading along the fascia), and/or the clinical presentation of gas-forming air bubbles in the fascia or muscle necrosis invasion on imaging tests such as contrast-enhanced CT[8].

## CASE PRESENTATION

### Chief complaints


A 43-year-old man visited our hospital with left facial swelling involving the buccal and submandibular areas.

### History of present illness

A 43-year-old male patient was admitted to our outpatient department complaining of swelling and pain in the left hemifacial area which started two weeks prior to consult. The patient had fever for approximately 10 d, but no other systemic symptoms occurred. At the time of the visit, the patient had fever with a temperature of 38.0 °C.

### History of past illness

The patient was diagnosed with diabetes mellitus treated with insulin. In addition, the patient had no history of illness associated with CNF, such as dental procedures, including tooth extraction, skin trauma, head and neck infections, and other facial fractures.



DOI: 10.12998/wjcc.v10.i31.11630 Copyright ©The Author(s) 2022.

**Figure 1 Preoperative computed tomographic imaging.** A: Axial computed tomography (CT) showing multiple soft tissue abscesses with air bubbles (white arrow); B: In the enhanced phase, axial CT showing a polymorphic wall with enhancing lesions in the muscle (white arrow).

### Personal and family history

The patient had no medical history other than diabetes mellitus, and no family history of CNF.

### Physical examination

**Preoperative:** On physical examination, trismus and thinning of the skin with crepitation were observed on the left cheek. There were also no pathological findings on dental examination.

**Postoperative:** On postoperative day (POD) 1, after surgical decompression by incision and drainage, the patient's systemic condition worsened, accompanied by tachycardia, hypotension, and overall deterioration of the patient's condition; hence, septic shock was considered.

### Laboratory examinations

At the time of hospitalization, the glycated hemoglobin (HbA1C) level was elevated at 17.1, indicating uncontrolled diabetes. The Laboratory Risk Indicator for Necrotizing Fasciitis (LRINEC) scoring system was used for proper diagnosis of necrotizing fasciitis[9]. Laboratory findings showed a high white blood cell (WBC) count (15170/ $\mu$ L; LRINEC score 1 point), a low hemoglobin level (10.5 g/dL; LRINEC score 2 points), an elevated high-sensitivity C-reactive protein (hs-CRP) level (212.94 mg/L; LRINEC score 4 points), a high glucose level (242 mg/dL; LRINEC score 1 point), and a slightly low sodium level (134 mmol/L; LRINEC score 2 points). The following results were normal: Creatinine, bilirubin, aspartate aminotransferase, and alanine aminotransferase. The patient had a score of 10 points in the LRINEC, indicating a high risk of CNF[9]. In addition, the albumin level was low at 2.3 g/dL. Procalcitonin, measured on the 3<sup>rd</sup> d after admission, was elevated at 1.140 ng/mL (normal range: < 0.1 ng/mL). Blood cultures after admission, wound swabs, and tissue cultures of the surgical field showed gram-negative *Klebsiella pneumoniae* (*K. pneumoniae*).

### Imaging examinations

**Preoperative:** On facial CT, multiple abscesses with air bubble in the left temporal, masticator, and buccal spaces along the fascial layer were observed (Figure 1A). Pleomorphic wall-enhancing lesions were also observed in the left masseter and temporalis muscles (Figure 1B).

**Postoperative:** On POD 1, follow-up CT revealed that the abscess pocket had become smaller. However, after decompression of the CNF, the patient's general condition deteriorated and progressed to septic shock. Contrast-enhanced chest CT was performed because multiple nodules were observed on chest X-ray upon admission (Figure 2). On chest CT on POD 2, septic emboli and pneumonia were observed in both lungs (Figure 3A). In addition, a lesion suspected as renal abscess was observed on chest CT. On POD 3, contrast-enhanced abdominopelvic CT (APCT) was also done, which showed multiple variably-sized renal abscesses in both kidneys (Figure 3B). This raised the suspicion that septic pneumonia and renal abscesses were the sources of sepsis.

## FURTHER DIAGNOSTIC WORK-UP

Prior to blood culture results for identification of sepsis, intravenous antibiotics were shifted to meropenem 1 g every 8 h and levofloxacin 750 mg per day, after consultation with the Department of Infectious Diseases. On POD 2, after *K. pneumoniae* was identified from the blood culture and detection



DOI: 10.12998/wjcc.v10.i31.11630 Copyright ©The Author(s) 2022.

**Figure 2** Chest X ray taken at the time of admission of the patient. Multiple masses and nodules are observed as findings of septic pneumonia caused by hematogenous metastasis.

of septic pneumonia and renal abscess, the antibiotic was changed to cefotaxime (POD 3). Cytopenia due to sepsis persisted, and despite the addition of metronidazole (POD 6), the clinical symptoms did not improve. On POD 3, the patient developed seizures due to metabolic encephalopathy, but magnetic resonance imaging or electroencephalography showed no acute brain lesions. Renal percutaneous catheter drainage (PCD) was performed on POD 7 to treat the renal abscess. After PCD insertion, the clinical signs and laboratory findings improved rapidly (Figure 4). After two weeks of PCD, the signs of infection and laboratory findings gradually stabilized, and the abscesses of the left masseter and temporalis muscle further decreased in size on facial CT taken on POD 25 (Figure 5A). Figure 6 shows a timeline of the course of illness.

## FINAL DIAGNOSIS

The patient's CNF symptoms and septic condition did not improve after surgical decompression of CNF but improved rapidly after PCD. Therefore, the final diagnosis of this patient was secondary CNF due to septic pulmonary and renal emboli.


## TREATMENT

### ***Surgical decompression of CNF***

Under urgent general anesthesia, an incision was made transversely above the left temporal crest. The superficial and deep temporal fasciae were dissected and the temporalis muscle was identified. After confirming the position of the abscess pocket on CT, dissection of the temporalis muscle was performed to drain the abscess, followed by massive saline irrigation. A drainage line was inserted through the temporal incision into the abscess pocket. A left upper gingivobuccal sulcus incision was then made to drain the abscess pocket of the masseter muscle and the pre-zygomatic area.


### ***Treatment of internal organ septic emboli***

Intravenous empiric antibiotics were initially given, and after detection of *K. pneumoniae*, the appropriate antibiotic was administered based on the sensitivity results of the culture strain. The patient did not show improvement in clinical symptoms even after the administration of meropenem, levofloxacin, cefataxime, or metrodidazole, but eventually showed rapid improvement after PCD insertion. After PCD insertion, the antibiotic was again changed to ceftriaxone (2 g/d). In addition, since the patient had uncontrolled diabetes mellitus with a significantly elevated HbA1C level at the time of admission, blood glucose tests were monitored daily, and blood glucose was controlled with insulin. No further surgical procedures were performed after PCD insertion, and the patient was discharged on the POD 28 after blood glucose control, and the infection symptoms completely resolved.




DOI: 10.12998/wjcc.v10.i31.11630 Copyright ©The Author(s) 2022.

**Figure 3 Computed tomographic imaging of internal organ abscesses.** A: Septic emboli on chest computed tomography (white arrow); B: Large renal abscess lesions are observed on abdominopelvic computed tomography (black arrow).



DOI: 10.12998/wjcc.v10.i31.11630 Copyright ©The Author(s) 2022.

**Figure 4 The graph of quantitative change of high-sensitivity C-reactive protein, white blood cell and albumin in laboratory findings.** Before and after percutaneous catheter drainage insertion, high-sensitivity C-reactive protein and white blood cell decreased, and albumin recovered to 3 or higher as the patient's clinical symptoms improved. hs-CRP: High-sensitivity C-reactive protein; WBC: White blood cell; POD: Postoperative day.




DOI: 10.12998/wjcc.v10.i31.11630 Copyright ©The Author(s) 2022.

**Figure 5 Postoperative computed tomographic imaging.** A: At postoperative day 25 showing a greater decrease in the size of abscess formation in the left masseter and temporalis muscles (white arrow); B: Abdominopelvic computed tomographic images. On postoperative day 45, The abscess is in remission and is not observed (black arrow).

## OUTCOME AND FOLLOW-UP

No further renal abscesses were observed on APCT on POD 45 (Figure 5B). At outpatient follow-up after discharge, the patient's HbA1C decreased to 4.9. Trismus was no longer observed six months after surgical decompression. No other symptoms suggestive of complications were observed in the craniofacial area.



DOI: 10.12998/wjcc.v10.i31.11630 Copyright ©The Author(s) 2022.

**Figure 6 Timeline of reported events.** The chronological sequence of changes in clinical manifestation, examination and management was recorded. LRINEC: Laboratory risk indicator for necrotizing fasciitis; POD: Postoperative day; CT: Computed tomography; hs-CRP: High-sensitivity C-reactive protein; APCT: Abdominal and pelvic computed tomography; ICU: Intensive care unit; MRI: Magnetic resonance imaging; EEG: Electroencephalography; PCD: Percutaneous catheter drainage.

## DISCUSSION

It is easy to overlook CNF especially when only the initial clinical symptoms are present, and it is difficult to differentiate CNF at its initial stage from a simple abscess[10]. However, CNF progresses rapidly and have serious implications. Unlike simple abscesses, CNFs have several specific clinical features, including rapid progression of clinical infectious manifestations, an abscess with disproportionate pain, presence of subcutaneous air bubbles or crepitations, and accompanying hyperesthesia or anesthesia to pinpricks. Air bubble formation in tissues, although not pathognomonic, is a critical sign that may appear on soft tissue plain radiographs or CT scans[11].

CNF may occur as a monomicrobial or polymicrobial infection. Park *et al*[4] reported that the prognosis of CNF differs according to the gram-staining pattern of the bacterium. Among gram-negative organisms, *K. pneumoniae* is the second most commonly detected pathogen, and most monomicrobial *K. pneumoniae* infections have a fatal course[7]. It is a highly virulent pathogen with a high rate of mortality. Most necrotizing fasciitis rapidly progresses due to direct infiltration of bacteria into the lesion; however, in the case of *K. pneumoniae*, as in this case, necrotizing fasciitis is caused by hematogenous dissemination to a septic lesion with an unknown etiology without a traumatic skin lesion[12].

In the present case, severe infection occurred to the extent that abscesses developed in both the lungs and kidneys. It is difficult to determine the antecedent relationship between the two organs. As the patient's CNF symptoms did not improve after surgical decompression and the laboratory values worsened and progressed into severe sepsis, craniofacial infection was not the primary cause of sepsis. In addition, considering that the patient's systemic condition improved dramatically after PCD insertion into the kidney, it was suspected that the infection was renal in origin. The hypervirulent nature of *K. pneumoniae* induces the formation of multiple abscesses, which may have resulted in secondary CNF, and underlying diseases such as diabetes may have contributed to the spread of septic emboli. Thus, this patient appeared to have developed secondary CNF in the craniofacial area, probably due to an internal organ abscess.

The early diagnosis of CNF in this case was made using the LRINEC score, which can be used to distinguish CNF from severe cellulitis or abscess based on relevant hematologic values. The LRINEC score has been proven useful for detecting severe infections of the upper and lower extremities, which has improved the efficiency of the early diagnosis[13]. Some reports have shown that lower albumin levels in patients are associated with a more fatal prognosis[9,14]. Procalcitonin is synthesized in various

tissues and organs in response to bacterial, fungal, and parasitic infections, and it has become a useful marker for determining whether antibiotics should be used for the duration of treatment[15].

This case study has three implications. First, CNF is a possible rare secondary cause of internal organ infections. The most common etiology is odontogenic infection, and skin or blunt trauma have also been reported. To the best of our knowledge, this is the first reported case of secondary CNF caused by septic emboli from an internal organ. Second, the progression of gram-negative *K. pneumoniae* infection caused a pulmonary or renal abscess, consistent with a previous report showing a higher risk of mortality in cases of sepsis caused by gram-negative pathogens[4]. Third, the patient had no physical examination findings suggestive of an internal organ infection other than fever. However, internal organ abscesses were observed, and despite successful drainage of the CNF, the infection still progressed to septic shock. This suggests that despite absence of other symptoms, the possibility of progression to systemic infection should always be considered if fever is accompanied with low albumin and high WBC count. In cases where a strain of *K. pneumoniae* is identified and if symptoms do not improve even after surgical decompression, a primary or secondary problem in the internal organs should be investigated further.

---

## CONCLUSION

CNFs mainly occur after dental procedures; however, it may also be caused by other infections. In the present case, CNF was caused by a systemic infection, which showed a more severe course than a simple abscess or CNF due to other causes. The findings of this study suggest that if CNF is not caused by adjacent skin or teeth, further investigation is indicated to determine the primary source of infection, considering the possibility of an internal organ origin. If the clinical manifestations suggest CNF, active surgical intervention should be performed without delay to reduce the risk of mortality.

---

## FOOTNOTES

**Author contributions:** Lee DW was involved in the concept and design of this manuscript; Lee DW and Kwak SH drafted the manuscript; Kwak SH and Choi HJ reviewed and revised the literature; and all authors have read and issued final approval of the version for submission.

**Supported by** the National Research Foundation of Korea Grant funded by the Korea Government (MSIT), No. 2021R1G1A1008337; and the Soonchunhyang University Research Fund.

**Informed consent statement:** Informed written consent was obtained from the patient for publication of this report and any accompanying images.

**Conflict-of-interest statement:** All the authors report no relevant conflicts of interest for this article.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** South Korea

**ORCID number:** Da-Woon Lee 0000-0002-0266-8480; Si-Hyun Kwak 0000-0001-9594-700X; Hwan-Jun Choi 0000-0002-0752-0389.

**S-Editor:** Wang JJ

**L-Editor:** A

**P-Editor:** Wang JJ

---

## REFERENCES

- 1 **Golger A**, Ching S, Goldsmith CH, Pennie RA, Bain JR. Mortality in patients with necrotizing fasciitis. *Plast Reconstr Surg* 2007; **119**: 1803-1807 [PMID: 17440360 DOI: 10.1097/01.prs.0000259040.71478.27]
- 2 **Juncar M**, Bran S, Juncar RI, Baciut MF, Baciut G, Onisor-Gligor F. Odontogenic cervical necrotizing fasciitis, etiological aspects. *Niger J Clin Pract* 2016; **19**: 391-396 [PMID: 27022806 DOI: 10.4103/1119-3077.179278]

- 3 **Oguz H**, Yilmaz MS. Diagnosis and management of necrotizing fasciitis of the head and neck. *Curr Infect Dis Rep* 2012; **14**: 161-165 [PMID: [22281689](#) DOI: [10.1007/s11908-012-0240-1](#)]
- 4 **Park SY**, Yu SN, Lee EJ, Kim T, Jeon MH, Choo EJ, Park S, Chae JW, In Bang H, Kim TH. Monomicrobial gram-negative necrotizing fasciitis: An uncommon but fatal syndrome. *Diagn Microbiol Infect Dis* 2019; **94**: 183-187 [PMID: [30713116](#) DOI: [10.1016/j.diagmicrobio.2018.12.013](#)]
- 5 **Shin J**, Park SI, Cho JT, Jung SN, Byeon J, Seo BF. Necrotizing fasciitis of the masticator space with osteomyelitis of the mandible in an edentulous patient. *Arch Craniofac Surg* 2019; **20**: 270-273 [PMID: [31462021](#) DOI: [10.7181/acfs.2019.00311](#)]
- 6 **Ku I**, Park JU. Necrotizing fasciitis arisen from nose. *Arch Craniofac Surg* 2019; **20**: 279-280 [PMID: [31462023](#) DOI: [10.7181/acfs.2019.00262](#)]
- 7 **Wong CH**, Chang HC, Pasupathy S, Khin LW, Tan JL, Low CO. Necrotizing fasciitis: clinical presentation, microbiology, and determinants of mortality. *J Bone Joint Surg Am* 2003; **85**: 1454-1460 [PMID: [12925624](#)]
- 8 **Yahav D**, Duskin-Bitan H, Eliakim-Raz N, Ben-Zvi H, Shaked H, Goldberg E, Bishara J. Monomicrobial necrotizing fasciitis in a single center: the emergence of Gram-negative bacteria as a common pathogen. *Int J Infect Dis* 2014; **28**: 13-16 [PMID: [25220388](#) DOI: [10.1016/j.ijid.2014.05.024](#)]
- 9 **Lee DW**, Ryu H, Choi HJ, Heo NH. Early diagnosis of craniofacial necrotising fasciitis: Analysis of clinical risk factors. *Int Wound J* 2022; **19**: 1071-1084 [PMID: [34755456](#) DOI: [10.1111/iwj.13703](#)]
- 10 **Holt GR**, Young WC, Aufdemorte T, Mattox DE, Gates GA. Head and neck manifestations of uncommon infectious diseases. *Laryngoscope* 1982; **92**: 634-639 [PMID: [7087624](#) DOI: [10.1002/Lary.1982.92.6.634](#)]
- 11 **Yamaoka M**, Furusawa K, Uematsu T, Yasuda K. Early evaluation of necrotizing fasciitis with use of CT. *J Craniomaxillofac Surg* 1994; **22**: 268-271 [PMID: [7798356](#) DOI: [10.1016/s1010-5182\(05\)80074-5](#)]
- 12 **Wang JH**, Liu YC, Lee SS, Yen MY, Chen YS, Wang JH, Wann SR, Lin HH. Primary liver abscess due to Klebsiella pneumoniae in Taiwan. *Clin Infect Dis* 1998; **26**: 1434-1438 [PMID: [9636876](#) DOI: [10.1086/516369](#)]
- 13 **Su YC**, Chen HW, Hong YC, Chen CT, Hsiao CT, Chen IC. Laboratory risk indicator for necrotizing fasciitis score and the outcomes. *ANZ J Surg* 2008; **78**: 968-972 [PMID: [18959694](#) DOI: [10.1111/j.1445-2197.2008.04713.x](#)]
- 14 **Huang KC**, Weng HH, Yang TY, Chang TS, Huang TW, Lee MS. Distribution of Fatal *Vibrio Vulnificus* Necrotizing Skin and Soft-Tissue Infections: A Systematic Review and Meta-Analysis. *Medicine (Baltimore)* 2016; **95**: e2627 [PMID: [26844475](#) DOI: [10.1097/MD.0000000000002627](#)]
- 15 **Wacker C**, Prkno A, Brunkhorst FM, Schlattmann P. Procalcitonin as a diagnostic marker for sepsis: a systematic review and meta-analysis. *Lancet Infect Dis* 2013; **13**: 426-435 [PMID: [23375419](#) DOI: [10.1016/S1473-3099\(12\)70323-7](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

