

World Journal of *Clinical Cases*

World J Clin Cases 2022 December 16; 10(35): 12804-13147



EVIDENCE REVIEW

- 12804 Principle and progress of radical treatment for locally advanced esophageal squamous cell carcinoma
Zhang XF, Liu PY, Zhang SJ, Zhao KL, Zhao WX

REVIEW

- 12812 Minimally invasive techniques in benign and malignant adrenal tumors
Dogrul AB, Cennet O, Dincer AH
- 12822 Planning issues on linac-based stereotactic radiotherapy
Huang YY, Yang J, Liu YB

MINIREVIEWS

- 12837 Hepatitis of unknown etiology in children: Current evidence and association
Zhong R, Yi F, Xiang F, Qiu YF, Zhu L, Zou YH, Wang W, Zhang Q
- 12844 Anatomical basis for pancreas transplantation *via* isolated splenic artery perfusion: A literature review
Dmitriev I, Oganesyan M, Popova A, Orlov E, Sinelnikov M, Zharikov Y
- 12854 Antenatal imaging: A pictorial review
Ece B, Aydın S, Kantarci M
- 12875 Real role of growth factor receptor-binding protein 10: Linking lipid metabolism to diabetes cardiovascular complications
Yang Y, Yao HJ, Lin WJ, Huang SC, Li XD, He FZ

ORIGINAL ARTICLE**Retrospective Study**

- 12880 Radiological and clinical outcomes of midline lumbar fusion on sagittal lumbar-pelvic parameters for degenerative lumbar diseases
Wang YT, Li BX, Wang SJ, Li CD, Sun HL
- 12890 Clinical features of elderly patients with COVID-19 in Wuhan, China
Wei S, Chen G, Ouyang XC, Hong YC, Pan YH

Observational Study

- 12899 Do inflammatory bowel disease patient preferences from treatment outcomes differ by ethnicity and gender? A cross-sectional observational study
Naftali T, Richter V, Mari A, Khoury T, Shirin H, Broide E

- 12909** Lipoprotein (a) variability is associated with mean follow-up C-reactive protein in patients with coronary artery disease following percutaneous coronary intervention
Zhang SS, Hu WY, Li YJ, Yu J, Sang S, Alsalman ZM, Xie DQ
- 12920** Efficacy evaluation of neuroendoscopy *vs* burr hole drainage in the treatment of chronic subdural hematoma: An observational study
Wang XJ, Yin YH, Wang ZF, Zhang Y, Sun C, Cui ZM
- 12928** Optimal approach for total endoscopic discectomy and its effect on lumbar and leg function in patients with disc herniation
Zhang ZH, Du Q, Wu FJ, Liao WB
- 12936** Value of inflammatory mediator profiles and procalcitonin in predicting postoperative infection in patients with hypertensive cerebral hemorrhage
Yin RH, Zhang B, Zhou XH, Cao LP, Li M

SYSTEMATIC REVIEWS

- 12946** *De novo* non-alcoholic fatty liver disease after pancreatectomy: A systematic review
Shah P, Patel V, Ashkar M

META-ANALYSIS

- 12959** Comparative effectiveness of first-line therapies for eradication of antibiotic-resistant *Helicobacter pylori* strains: A network meta-analysis
Zou SP, Cheng Q, Feng CY, Xu C, Sun MH

CASE REPORT

- 12971** Malignant atrophic papulosis: Two case reports
Li ZG, Zhou JM, Li L, Wang XD
- 12980** Endoscopic treatment of urothelial encrusted pyelo-ureteritis disease: A case series
Liu YB, Xiao B, Hu WG, Zhang G, Fu M, Li JX
- 12990** Nearly-complete labial adhesions diagnosed with repetitive cystitis in postmenopausal women: A case report
Kwon H
- 12996** Congenital dysfibrinogenemia misdiagnosed and inappropriately treated as acute fatty liver in pregnancy: A case report and review of literature
Jia Y, Zhang XW, Wu YS, Wang QY, Yang SL
- 13006** Lung squamous cell carcinoma presenting as rare clustered cystic lesions: A case report and review of literature
Shen YY, Jiang J, Zhao J, Song J
- 13015** Management of ductal spasm in a neonate with pulmonary atresia and an intact ventricular septum during cardiac catheterization: A case report
Zhang X, Zhang N, Song HC, Ren YY

- 13022** Symptomatic accessory soleus muscle: A cause for exertional compartment syndrome in a young soldier: A case report
Woo I, Park CH, Yan H, Park JJ
- 13028** Multiple myeloma presenting with amyloid arthropathy as the first manifestation: Two case reports
He C, Ge XP, Zhang XH, Chen P, Li BZ
- 13038** Kawasaki disease without changes in inflammatory biomarkers: A case report
Yamashita K, Kanazawa T, Abe Y, Naruto T, Mori M
- 13044** Atypical Whipple's disease with special endoscopic manifestations: A case report
Chen S, Zhou YC, Si S, Liu HY, Zhang QR, Yin TF, Xie CX, Yao SK, Du SY
- 13052** Acute limb ischemia after minimally invasive cardiac surgery using the ProGlide: A case series
Lee J, Huh U, Song S, Lee CW
- 13058** Genetic changes in refractory relapsed acute myeloid leukemia with *NPM1* mutation: A case report
Wang SL
- 13064** Successful surgical treatment of polybacterial gas gangrene confirmed by metagenomic next-generation sequencing detection: A case report
Lu HY, Gao YB, Qiu XW, Wang Q, Liu CM, Huang XW, Chen HY, Zeng K, Li CX
- 13074** Pulmonary sarcoidosis: A novel sequelae of drug reaction with eosinophilia and systemic symptoms: A case report
Hu YQ, Lv CY, Cui A
- 13081** Hammered silver appearance of the corneal endothelium in Fuchs uveitis syndrome: A case report
Cheng YY, Wang CY, Zheng YF, Ren MY
- 13088** Tracheostomy and venovenous extracorporeal membrane oxygenation for difficult airway patient with carinal melanoma: A case report and literature review
Liu IL, Chou AH, Chiu CH, Cheng YT, Lin HT
- 13099** Surgery combined with antibiotics for thoracic vertebral *Escherichia coli* infection after acupuncture: A case report
Mo YF, Mu ZS, Zhou K, Pan D, Zhan HT, Tang YH
- 13108** Multidisciplinary treatment of a patient with severe immune checkpoint inhibitor-induced colitis: A case report
Lu L, Sha L, Feng Y, Yan L
- 13115** Systemic combined with intravitreal methotrexate for relentless placoid chorioretinitis: A case report
Luo L, Chen WB, Zhao MW, Miao H
- 13122** Response to roxadustat in a patient undergoing long-term dialysis and allergic to erythropoiesis-stimulating agents: A case report
Xu C, Luo DG, Liu ZY, Yang D, Wang DD, Xu YZ, Yang J, Fu B, Qi AR

13129 Liver collision tumor of primary hepatocellular carcinoma and neuroendocrine carcinoma: A rare case report

Jeng KS, Huang CC, Chung CS, Chang CF

13138 Unexpected delayed reversal of rocuronium-induced neuromuscular blockade by sugammadex: A case report and review of literature

Wang HC, Lu CW, Lin TY, Chang YY

LETTER TO THE EDITOR

13146 Immunoglobulin G4 associated autoimmune cholangitis and pancreatitis and nivolumab

Joob B, Wiwanitkit V

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Lovenish Bains, FACS, FICS, FRCS, MBBS, MS, Associate Professor, Surgeon, Teacher, Department of Surgery, Maulana Azad Medical College, New Delhi 110002, India. lovenishbains@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for *WJCC* as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The *WJCC*'s CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Hua-Ge Yu*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

December 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Management of ductal spasm in a neonate with pulmonary atresia and an intact ventricular septum during cardiac catheterization: A case report

Xu Zhang, Ning Zhang, Hai-Cheng Song, Yue-Yi Ren

Specialty type: Anesthesiology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: El-Gendy HA, Egypt;
Teragawa H, Japan

Received: August 1, 2022

Peer-review started: August 1, 2022

First decision: October 17, 2022

Revised: October 29, 2022

Accepted: November 22, 2022

Article in press: November 22, 2022

Published online: December 16, 2022



Xu Zhang, Hai-Cheng Song, Yue-Yi Ren, Department of Heart Center, Qingdao Women's and Children's Hospital Affiliated to Qingdao University, Qingdao 266034, Shandong Province, China

Ning Zhang, Department of Medical Technology, Qingdao Women's and Children's Hospital Affiliated to Qingdao University, Qingdao 266034, Shandong Province, China

Corresponding author: Yue-Yi Ren, MD, Chief Physician, Department of Heart Center, Qingdao Women's and Children's Hospital Affiliated to Qingdao University, No. 217 Liaoyang West Road, Shibei District, Qingdao 266034, Shandong Province, China.

xxgmz1173@126.com

Abstract

BACKGROUND

Ductal spasm is a rare but life-threatening complication of cardiac catheterization in neonates with pulmonary atresia and an intact ventricular septum. In patients with ductal-dependent pulmonary blood flow, ductal spasm may lead to refractory hypoxemia and severe hemodynamic instability, which need to be treated in perfect order.

CASE SUMMARY

We present a male infant with a gestational age of 39 wk, and his fetal echocardiography showed pulmonary atresia. At 28 d of age, transcatheter pulmonary valvuloplasty with balloon dilatation was performed. Two hours after the operation, the patient's pulse oxygen saturation continued to decrease. The patient was then transferred to receive cardiac catheterization. During catheterization, the invasive blood pressure and pulse oxygen saturation suddenly decreased, and repeated aortography revealed partial occlusion of the ductus arteriosus. It no longer changed when pulse oxygen saturation rose to 51% after approximately 20 min of maintenance therapy. Therefore, a ductal stent was used for implantation. Hemodynamics and hypoxemia were improved.

CONCLUSION

We should know that ductal spasm may occur during pulmonary atresia and intact ventricular septum cardiac catheterization. Understand the pathophysiology of ductal-dependent pulmonary blood flow and make comprehensive perioperative preparations essential to deal with hemodynamic disorders caused

by ductal spasm.

Key Words: Cardiac catheterization; Ductus arteriosus; Ductal stent; Ductal spasm; Intact ventricular septum; Pulmonary atresia; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: In patients with ductal-dependent pulmonary blood flow, ductal spasm may lead to refractory hypoxemia and severe hemodynamic instability. Ductal stenting has been widely developed as an initial palliative strategy to secure pulmonary blood flow in such patients. We report a case of a neonate with pulmonary atresia and an intact ventricular septum who developed ductal spasm during ductal stenting. We discussed the appropriate treatment for ductal spasm in a neonate during cardiac catheterization.

Citation: Zhang X, Zhang N, Song HC, Ren YY. Management of ductal spasm in a neonate with pulmonary atresia and an intact ventricular septum during cardiac catheterization: A case report. *World J Clin Cases* 2022; 10(35): 13015-13021

URL: <https://www.wjgnet.com/2307-8960/full/v10/i35/13015.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i35.13015>

INTRODUCTION

Percutaneous ductal stenting has been widely developed as an initial palliative strategy to secure patients with pulmonary atresia and intact ventricular septum (PA/IVS)[1]. Ductal stenting can be performed simultaneously or in stages after pulmonary valvuloplasty. The management of interventional surgery in such patients is challenging. Ductal spasm (DS) is a rare but life-threatening complication of cardiac catheterization in neonates with PA/IVS. In patients with ductal-dependent pulmonary blood flow (PBF), DS may lead to refractory hypoxemia and severe hemodynamic instability [2]. We report a case of a neonate with PA/IVS who developed DS during ductal stenting. Changes in hemodynamics and respiratory parameters were recorded, and the treatment process was summarized and analyzed.

CASE PRESENTATION

Chief complaints

The patient was a male infant with a gestational age of 39 wk, and his fetal echocardiography showed pulmonary atresia.

History of present illness

Fetal echocardiography of the neonatal patient was performed before delivery (24 wk pregnant) to show pulmonary atresia. Due to abnormal umbilical blood flow, the patient was delivered by cesarean section, without a history of intrauterine distress and premature rupture of membranes. Then, the patient was transferred to the cardiac intensive care unit (CICU) for further treatment. The patient received prostaglandin E1 (PGE1) at 3 ng/kg/min to maintain patency of the ductus arteriosus (DA) after delivery, and the pulse oxygen saturation (SPO₂) was 84%–88%.

History of past illness

The patient had no history of other illnesses.

Personal and family history

The patient had no family history of illness.

Physical examination

The physical examination revealed normal development (height, 50 cm; weight, 3.26 kg), a blood pressure of 85/47 mmHg, a pulse rate of 145 beats per min, and a respiratory rate (RR) of 30 breaths/min. His pulse oxygen saturation (SPO₂) was 84%–88% in room air.

Laboratory examinations

Blood tests showed an Hb level of 12.5 g/dL, hematocrit (Hct) of 37%, platelet count of 266×10^9 cells/L, leukocyte count of 11.22×10^9 /L, C-reactive protein of 4.16 mg/L, and N-terminal pro-brain natriuretic peptide (NT-proBNP) of 982.50 pg/mL. Other blood tests showed no significant abnormalities.

Imaging examinations

Postnatal echocardiography revealed a foramen ovale of approximately 5.5 mm, a right-to-left shunt, and an intact ventricular septum. The pulmonary valve ring was 6.2 mm, with membranous atresia. The descending aorta was connected to the bifurcation of the main pulmonary artery through the patent ductus arteriosus (PDA) (the inner diameter of the end of the pulmonary artery was 3 mm) and a continuous shunt.

FINAL DIAGNOSIS

Pulmonary atresia and intact ventricular septum; Patent foramen ovale; Patent ductus arteriosus.

TREATMENT

At 28 d of age, the patient planned to undergo pulmonary valve perforation and percutaneous balloon pulmonary valvuloplasty (PBPV). After induction of anesthesia, the patient inhaled 30% oxygen; left internal jugular vein and left radial artery catheterization were performed, and PGE1 was continuously infused at 3 ng/kg/min. Descending aortography showed that the DA was tubular with an internal diameter of 3.0 mm. After pulmonary valve perforation and balloon dilatation, right ventriculography showed that the pulmonary valve was open, and SPO₂ was 89% when the patient was returned to the CICU with an endotracheal tube. Two hours later, the patient's SPO₂ dropped to 75%. Echocardiography showed a significantly enlarged right atrium, with a right-to-left shunt at the atrial level, patent foramen ovale (diameter, 7 mm), PDA (diameter, 3 mm at the pulmonary end), severe tricuspid regurgitation, and a little forward flow through the pulmonary valve. Intensive care doctors considered the patient to have developed right-sided heart failure. The treatment included infusion of vasoactive drugs and diuretics, inhalation of nitric oxide, intravenous infusion of treprostinil to reduce pulmonary artery pressure, and adjustment of the PGE1 dose to 10 ng/kg/min. However, the SPO₂ continued to decrease to as low as 60%. The patient was then transferred from the CICU to the catheterization room. General anesthesia was maintained by inhaling 1.5% sevoflurane in 40% oxygen and cisatracurium 0.5 mg administered intravenously. Descending aortography revealed that the PDA was tortuous and flat (Figure 1A). When the catheter was adjusted, the invasive blood pressure (IBP) suddenly decreased from 68/40 mmHg to 44/28 mmHg (Figure 2A), SPO₂ decreased from 68% to 40%, and end-tidal carbon dioxide (PETCO₂) decreased from 36 mmHg to 18 mmHg. Repeated aortography revealed partial occlusion of the DA (Figure 1B). The catheter was withdrawn because of a suspected DS. PGE1 was increased to 20 ng/kg/min, and norepinephrine was increased to 0.2 µg/kg/min. Simultaneously, the concentration of inhaled oxygen was reduced to inhaled air. It no longer changed when SPO₂ rose to 51% after approximately 20 min of maintenance therapy. Therefore, a ductal stent was used for implantation. After successfully implanting a 4.5 mm × 21 mm stent, the change in invasive arterial pressure was 65/42 mmHg (Figure 2B), SPO₂ was 85%, and ETCO₂ was 33 mmHg. PGE1 treatment was then discontinued. Aortography showed that the stent was unobstructed without displacement (Figure 1C). The guidewire and balloon were withdrawn, and the patient was returned to the CICU.

OUTCOME AND FOLLOW-UP

The patient was hemodynamically in the immediate postoperative period, and his SPO₂ while breathing room air was 85%. Then, the endotracheal tube was removed on day 3 after ductal stenting intervention, and the patient was discharged from the CICU 2 days later. One month later, the patient underwent balloon dilatation of the pulmonary valve again, and catheter angiography showed ductal stent patency without stenosis or occlusion.


DISCUSSION

A 28-day-old neonate with PA/IVS who underwent PBPV developed persistent hypoxemia and could not be improved after a series of treatments. The patient had to be transferred to receive cardiac catheterization. During catheterization, the invasive blood pressure and pulse oxygen saturation suddenly



DOI: 10.12998/wjcc.v10.i35.13015 Copyright ©The Author(s) 2022.

Figure 1 Angiographic images of ductus arteriosus before and after spasm. A: Descending aortography revealed that the patent ductus arteriosus (PDA) was tortuous and flat. The orange arrow showed the position of PDA; B: Repeated aortography revealed partial occlusion of the ductus arteriosus. The orange arrow showed PDA markedly curved, S-shaped; C: Aortography showed that the stent was unobstructed without displacement.



DOI: 10.12998/wjcc.v10.i35.13015 Copyright ©The Author(s) 2022.

Figure 2 Hemodynamic changes of ductal spasm and stent implantation. A: The IBP suddenly decreased; B: After successful stent implantation, the invasive blood pressure changed to an instant increase.

decreased, and repeated aortography revealed DS. We adopted routine treatment methods, including inhaling air, increasing the infusion dose of PGE1 and vasoactive drugs. It no longer changed when SPO₂ rose to 51% after approximately 20 min of maintenance therapy. Therefore, a ductal stent was used for implantation. Hemodynamics and hypoxemia were improved. Pulmonary circulation in patients with PA/IVS depends on DA patency. Postpartum contraction of DA will lead to severe hypoxemia, cyanosis and death[3]. Ductal stenting for DA is usually used for patients who have successfully undergone pulmonary valve perforation but still need pulmonary circulation support because PGE1 infusion cannot be relieved[4]. Compared with a modified Blalock-Taussig (BT) shunt, a meta-analysis demonstrated that ductal stenting had fewer procedural complications and shorter CICU stays[5]. According to the Catheterization for Congenital Heart Disease Adjustment for Risk Method, pulmonary valve perforation and any catheterization within 4 days after surgery are classified as the most high-risk procedures. This classification is because most of these patients are newborns or premature infants with long operation times and high skill requirements. The procedure can easily cause drastic changes in hemodynamics and SPO₂. Most ductal stenting procedures are performed later after balloon pulmonary valvuloplasty. As a staged procedure, it was helpful to evaluate the ductal contraction time after PGE1 discontinuation and can avoid stent displacement and dislocation. In addition, some patients had to undergo cardiac catheterization and stent implantation a few days after balloon pulmonary valvuloplasty because they could not tolerate PGE1 infusion or SPO₂ continued to decrease[1].

In previous reports, the most common complication of PA/IVS cardiac catheterization was vascular-related injury. Other complications include DS, dissection, acute stent thrombosis, stent migration or dislodgment. However, the incidence of DS has not been reported in detail in previous ductal stenting studies. Chen and colleagues reported a case of ductal dissection and spasm during stenting of a PDA [6]. The catheter was considered sandwiched because the guidewire could not pass through the DA smoothly, and the end was slightly deformed. After pulling out the guidewire, the SPO₂ decreased to

60%. Repeated angiography revealed a severe DS. PGE1 was administered again (20 ng/kg/min). The hemodynamics were stable, and SPO₂ recovered to 85% after 20 min of PGE1 infusion. Finally, a ductal stent was implanted. According to the literatures[7-9], there are many factors leading to DS. One possible factor is mechanical stimulation. A cardiac interventionalist may unintentionally induce spasm when performing catheter manipulation to enter the aortic arch, main or branch pulmonary artery, or when crossing the PDA[8]. Another possible factor is biochemical stimulation. DS may be stimulated by stress-induced catecholamine (nitric oxide and prostaglandins) release[9] and inhalation of high concentrations of oxygen[10]. In this case, the concentration of inhaled oxygen did not change, and PGE1 was continuously infused before and during the catheterization procedure. Therefore, we considered that the catheter stimulated the DA during aortic angiography, causing contraction and spasm.

Evidence of DS in patients with ductal-dependent PBF includes confirmation of PDA presence by preoperative echocardiography and intraoperative angiography. Second, intraoperative echocardiography and angiography showed that the catheter's ampulla gradually became thinner and completely or nearly occluded[7]. Finally, significant reductions in IBP, SPO₂, and ETCO₂ (manifestations of decreased PBF) are important clinical indications. Therefore, the prevention of DS is important. In patients with ductal contractions, it is necessary to consider the possibility of further ductal contractions during the procedure. Determining continuous infusion of PGE1 and performing appropriate operations are important. To reduce the DS or expand the stenosis area, Bahaidarah and colleagues recommended that PGE1 should be infused intravenously until the guidewire passes through the DA [11]. After DS, if the catheter or wire has crossed the PDA, the stent can be implanted quickly to re-establish stable PBF[2]. If the stent cannot be implanted as soon as possible, the catheter should be removed immediately to alleviate DS[5]. In this case, the reason for this patient's ductal stenting was that SPO₂ continued to decrease, and the pulmonary artery had little forward blood flow, which was not enough to meet the needs of the body. DS during the procedure occurred suddenly, not slowly. The most appropriate response was to withdraw the catheter and provide conservative treatment. It was wise to wait for a few minutes and take an angiogram again to observe the size of the DA to decide the next treatment[12,13]. If the stent is directly implanted, it may be disastrous to further stimulate the ductus arteriosus and cause the ductus arteriosus to close directly. To avoid further aggravation of the DS, we did not immediately cross the wire and catheter to the PDA to implant the stent. Of course, DS is a functional abnormality that can be reversed through treatment, but it takes time to reverse before patient's condition continue to deteriorate. However, there was no significant relief of DS after conservative treatment. The patient could not continue to tolerate long-term hypoxia and hypotension, so we performed ductal stenting.

Specific anesthetic management at this time is not decisive, but measures should be taken to relieve DS and maintain hemodynamic stability as much as possible. Preoperative safe central venous access, inotropes, vasopressors, and sodium bicarbonate are necessary to increase blood pressure, improve acidosis, and help blood flow through the PDA during DS, thereby increasing PBF[14]. IBP monitoring is important to quickly observe the direct hemodynamic changes following DS. Batlivala and colleagues reported 2 cases of DS after angiography. Continuous arterial pressure monitoring helps patients quickly diagnose DS[15]. When DS occurs, the infusion dose of PGE1 should be increased immediately. Some studies suggest that when aggravating cyanosis compatible with impending closure occurs in patients with ductal-dependent PBF, an initial dose of 20 ng/kg/minute PGE1 should be started[16,17]. If there is no improvement in oxygenation, stepwise dose increases of 5 ng/kg/minute are administered until no further increase in oxygenation occurs or SPO₂ has reached the preclosure level of the ductus arteriosus, even to 100 ng/kg/min. If the hemodynamics are stable, the catheter is generally reopened within 30 mins to 2 h[18]. Of course, an initial dose of PGE1 of 20 ng/kg/min was given after DS in this case. It was undeniable that our approach was conservative, and we did not continuously increase the dose of PGE1. When hypoxaemia occurs in DS, some researchers believe that increasing the inhaled oxygen concentration cannot improve the condition. Instead, air should be inhaled for ventilation to avoid hyperoxia, stimulating the further development of DA occlusion. Historical animal experiments have shown that when the same isolated DA rings were transferred from low oxygen partial pressure (PO₂) to high PO₂ and restimulated by administering PGE1, the relaxation effect of PGE1 on DA rings at high PO₂ was significantly increased. However, no further study has confirmed whether the DA response to PGE1 is the same from low PO₂ to high PO₂. Decker and colleagues suggested that inhalation anesthesia should be changed to total intravenous anesthesia with propofol because 8 children with DS were well relieved in their cases[19]. However, there is no clear explanation for this phenomenon. Heparin administration to achieve an activated clotting time of > 250 s should be confirmed. If a stent cannot be implanted and hemodynamics deteriorate rapidly, cardiopulmonary bypass or extracorporeal membrane oxygenation can be performed as soon as possible to save time and life.

CONCLUSION

Ductal spasm may occur during PA/IVS cardiac catheterization. Understand the pathophysiology of

ductal-dependent PBF and make comprehensive perioperative preparations are essential to deal with hemodynamic disorders caused by ductal spasm.

FOOTNOTES

Author contributions: Zhang X and Ren YY drafted and revised the manuscript; Zhang N and Song HC collected all the data related to the case report; all authors read and approved the final manuscript.

Informed consent statement: Written informed consent was obtained from the patient's parent or guardian for publication of this case report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflicts of interest.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Xu Zhang 0000-0002-6481-2505; Ning Zhang 0000-0001-8358-2197; Hai-Cheng Song 0000-0002-4444-3562; Yue-Yi Ren 0000-0002-0410-7978.

S-Editor: Xing YX

L-Editor: A

P-Editor: Xing YX

REFERENCES

- Haddad RN, Hanna N, Charbel R, Daou L, Chehab G, Saliba Z. Ductal stenting to improve pulmonary blood flow in pulmonary atresia with intact ventricular septum and critical pulmonary stenosis after balloon valvuloplasty. *Cardiol Young* 2019; **29**: 492-498 [PMID: 31030705 DOI: 10.1017/S1047951119000118]
- Aggarwal V, Petit CJ, Glatz AC, Goldstein BH, Qureshi AM. Stenting of the ductus arteriosus for ductal-dependent pulmonary blood flow-current techniques and procedural considerations. *Congenit Heart Dis* 2019; **14**: 110-115 [PMID: 30811792 DOI: 10.1111/chd.12709]
- Eilers L, Qureshi AM. Advances in Pediatric Ductal Intervention: an Open or Shut Case? *Curr Cardiol Rep* 2020; **22**: 14 [PMID: 31997085 DOI: 10.1007/s11886-020-1266-x]
- Ratnayaka K, Nageotte SJ, Moore JW, Guyon PW, Bhandari K, Weber RL, Lee JW, You H, Griffin DA, Rao RP, Nigro JJ, El-Said HG. Patent Ductus Arteriosus Stenting for All Ductal-Dependent Cyanotic Infants: Waning Use of Blalock-Taussig Shunts. *Circ Cardiovasc Interv* 2021; **14**: e009520 [PMID: 33685211 DOI: 10.1161/CIRCINTERVENTIONS.120.009520]
- Alsagheir A, Koziaz A, Makhdoum A, Contreras J, Alraddadi H, Abdalla T, Benson L, Chaturvedi RR, Honjo O. Duct stenting vs modified Blalock-Taussig shunt in neonates and infants with duct-dependent pulmonary blood flow: A systematic review and meta-analysis. *J Thorac Cardiovasc Surg* 2021; **161**: 379-390.e8 [PMID: 32727685 DOI: 10.1016/j.jtcvs.2020.06.008]
- Chen TY, Chen PW, Wang JN. Patent ductus arteriosus stenting: ductal dissection and spasm in pulmonary atresia with intact ventricular septum. *Cardiol Young* 2022; **32**: 679-680 [PMID: 34493357 DOI: 10.1017/S1047951121003711]
- Batlivala SP, Glatz AC, Gillespie MJ, Dori Y, Rome JJ. Ductal spasm during performance of transcatheter ductal occlusion. *Catheter Cardiovasc Interv* 2014; **83**: 762-767 [PMID: 23832584 DOI: 10.1002/ccd.25120]
- Méot M, Haddad RN, Patkai J, Abu Zahira I, Di Marzio A, Szezepanski I, Bajolle F, Kermorvant E, Lapillonne A, Bonnet D, Malekzadeh-Milani S. Spontaneous Closure of the Arterial Duct after Transcatheter Closure Attempt in Preterm Infants. *Children (Basel)* 2021; **8** [PMID: 34943334 DOI: 10.3390/children8121138]
- Ovali F. Molecular and Mechanical Mechanisms Regulating Ductus Arteriosus Closure in Preterm Infants. *Front Pediatr* 2020; **8**: 516 [PMID: 32984222 DOI: 10.3389/fped.2020.00516]
- Coceani F. Oxygen sensing in the ductus arteriosus-A unifying vision for two concepts. *Br J Pharmacol* 2022; **179**: 3325-3329 [PMID: 35302234 DOI: 10.1111/bph.15828]
- Bahaidarah S, Al-Ata J, Alkhusi N, Azhar A, Zaher Z, Alnahdi B, Abdelsalam M, Elakaby A, Dohain A, Abdelmohsen G. Outcome of ductus arteriosus stenting including vertical tubular and convoluted tortuous ducts with emphasis on technical considerations. *Egypt Heart J* 2021; **73**: 83 [PMID: 34542737 DOI: 10.1186/s43044-021-00210-4]
- MacDonald ST, Bhindi R, Ormerod O, Wilson N. Ductus arteriosus spasm. *JACC Cardiovasc Interv* 2009; **2**: 73 [PMID: 19463401 DOI: 10.1016/j.jcin.2008.04.013]

- 13 **Alkamali A**, Alasrawi S. Dealing with patent ductus arteriosus (PDA) spasm in the cath lab. *J Cardiol Curr Res* 2018; **11**: 135-138 [DOI: [10.15406/jccr.2018.11.00386](https://doi.org/10.15406/jccr.2018.11.00386)]
- 14 **Udink Ten Cate FE**, Sreeram N, Hamza H, Agha H, Rosenthal E, Qureshi SA. Stenting the arterial duct in neonates and infants with congenital heart disease and duct-dependent pulmonary blood flow: a multicenter experience of an evolving therapy over 18 years. *Catheter Cardiovasc Interv* 2013; **82**: E233-E243 [PMID: [23420699](https://pubmed.ncbi.nlm.nih.gov/23420699/) DOI: [10.1002/ccd.24878](https://doi.org/10.1002/ccd.24878)]
- 15 **Sarquella-Brugada G**, Mivelaz Y, Dahdah N. Hemodynamic changes alert to spontaneous ductus arteriosus spasm. *Rev Esp Cardiol (Engl Ed)* 2013; **66**: 743 [PMID: [24773683](https://pubmed.ncbi.nlm.nih.gov/24773683/) DOI: [10.1016/j.rec.2011.11.018](https://doi.org/10.1016/j.rec.2011.11.018)]
- 16 **Huang FK**, Lin CC, Huang TC, Weng KP, Liu PY, Chen YY, Wang HP, Ger LP, Hsieh KS. Reappraisal of the prostaglandin E1 dose for early newborns with patent ductus arteriosus-dependent pulmonary circulation. *Pediatr Neonatol* 2013; **54**: 102-106 [PMID: [23590954](https://pubmed.ncbi.nlm.nih.gov/23590954/) DOI: [10.1016/j.pedneo.2012.10.007](https://doi.org/10.1016/j.pedneo.2012.10.007)]
- 17 **Khalil M**, Jux C, Rueblinger L, Behrje J, Esmacili A, Schranz D. Acute therapy of newborns with critical congenital heart disease. *Transl Pediatr* 2019; **8**: 114-126 [PMID: [31161078](https://pubmed.ncbi.nlm.nih.gov/31161078/) DOI: [10.21037/tp.2019.04.06](https://doi.org/10.21037/tp.2019.04.06)]
- 18 **Cucerea M**, Simon M, Moldovan E, Ungureanu M, Marian R, Suci L. Congenital Heart Disease Requiring Maintenance of Ductus Arteriosus in Critically Ill Newborns Admitted at a Tertiary Neonatal Intensive Care Unit. *J Crit Care Med (Targu Mures)* 2016; **2**: 185-191 [PMID: [29967858](https://pubmed.ncbi.nlm.nih.gov/29967858/) DOI: [10.1515/jccm-2016-0031](https://doi.org/10.1515/jccm-2016-0031)]
- 19 **De Decker R**, Comitis G, Thomas J, van der Merwe E, Lawrenson J. A novel approach to ductal spasm during percutaneous device occlusion of patent ductus arteriosus. *Cardiol Young* 2016; **26**: 1352-1358 [PMID: [26692358](https://pubmed.ncbi.nlm.nih.gov/26692358/) DOI: [10.1017/S1047951115002619](https://doi.org/10.1017/S1047951115002619)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

