World Journal of *Clinical Cases*

World J Clin Cases 2022 December 6; 10(34): 12462-12803





Published by Baishideng Publishing Group Inc

W J C C World Journal of Clinical Cases

Contents

Thrice Monthly Volume 10 Number 34 December 6, 2022

FIELD OF VISION

12462 Problematics of neurosurgical service during the COVID-19 pandemic in Slovenia Munda M, Bosnjak R, Velnar T

MINIREVIEWS

- 12470 Circulating angiotensin converting enzyme 2 and COVID-19 Leowattana W, Leowattana T, Leowattana P
- 12484 Evaluation of gut dysbiosis using serum and fecal bile acid profiles Monma T, Iwamoto J, Ueda H, Tamamushi M, Kakizaki F, Konishi N, Yara S, Miyazaki T, Hirayama T, Ikegami T, Honda A
- 12494 Pediatric kidney transplantation during the COVID-19 pandemic Tamura H

ORIGINAL ARTICLE

Clinical and Translational Research

12500 Coptis, Pinellia, and Scutellaria as a promising new drug combination for treatment of Helicobacter pylori infection

Yu Z, Sheng WD, Yin X, Bin Y

Case Control Study

12515 Effects of illness perception on negative emotions and fatigue in chronic rheumatic diseases: Rumination as a possible mediator

Lu Y, Jin X, Feng LW, Tang C, Neo M, Ho RC

Retrospective Study

12532 Significance of incidental focal fluorine-18 fluorodeoxyglucose uptake in colon/rectum, thyroid, and prostate: With a brief literature review

Lee H, Hwang KH

12543 Follow-up study on ThinPrep cytology test-positive patients in tropical regions

Chen YC, Liang CN, Wang XF, Wang MF, Huang XN, Hu JD

- 12551 Effect of teach-back health education combined with structured psychological nursing on adverse emotion and patient cooperation during 99mTc-3PRGD2.SPECT/CT Gong WN, Zhang YH, Niu J, Li XB
- Nosocomial infection and spread of SARS-CoV-2 infection among hospital staff, patients and caregivers 12559 Cheng CC, Fann LY, Chou YC, Liu CC, Hu HY, Chu D



World Journal of Clinical Cases

Contents

Thrice Monthly Volume 10 Number 34 December 6, 2022

Observational Study

- 12566 Effectiveness and safety of generic and brand direct acting antivirals for treatment of chronic hepatitis C Abdulla M, Al Ghareeb AM, Husain HAHY, Mohammed N, Al Qamish J
- 12578 Influence of group B streptococcus and vaginal cleanliness on the vaginal microbiome of pregnant women Liao Q, Zhang XF, Mi X, Jin F, Sun HM, Wang QX

Randomized Controlled Trial

12587 Clinical study on tri-tongue acupuncture combined with low-frequency electrical stimulation for treating post-stroke dysarthria

Man B, Li WW, Xu JF, Wang Q

META-ANALYSIS

12594 Three-dimensional time-of-flight magnetic resonance angiography combined with high resolution T2weighted imaging in preoperative evaluation of microvascular decompression

Liang C, Yang L, Zhang BB, Guo SW, Li RC

CASE REPORT

- 12605 Acute cytomegalovirus hepatitis in an immunocompetent patient: A case report Wang JP, Lin BZ, Lin CL, Chen KY, Lin TJ
- 12610 Long-term results of extended Boari flap technique for management of complete ureteral avulsion: A case report

Zhong MZ, Huang WN, Huang GX, Zhang EP, Gan L

12617 Amyloid β -related angiitis of the central nervous system occurring after COVID-19 vaccination: A case report

Kizawa M, Iwasaki Y

12623 Pseudoileus caused by primary visceral myopathy in a Han Chinese patient with a rare MYH11 mutation: A case report

Li N, Song YM, Zhang XD, Zhao XS, He XY, Yu LF, Zou DW

12631 Emergent use of tube tip in pharynx technique in "cannot intubate cannot oxygenate" situation: A case report Lin TC, Lai YW, Wu SH

12637 Inflammatory myofibroblastic tumor of the central nervous system: A case report Su ZJ, Guo ZS, Wan HT, Hong XY

- 12648 Atypical aggressive vertebral hemangioma of the sacrum with postoperative recurrence: A case report Wang GX, Chen YQ, Wang Y, Gao CP
- 12654 Closed reduction of hip dislocation associated with ipsilateral lower extremity fractures: A case report and review of the literature Xu Y, Lv M, Yu SO, Liu GP

• •	World Journal of Clinical Cases
Conten	ts Thrice Monthly Volume 10 Number 34 December 6, 2022
12665	Repair of a large patellar cartilage defect using human umbilical cord blood-derived mesenchymal stem cells: A case report
	Song JS, Hong KT, Song KJ, Kim SJ
12671	Abdominal bronchogenic cyst: A rare case report
	Li C, Zhang XW, Zhao CA, Liu M
12678	Malignant fibrous histiocytoma of the axilla with breast cancer: A case report
	Gao N, Yang AQ, Xu HR, Li L
12684	Rapid hemostasis of the residual inguinal access sites during endovascular procedures: A case report
	Kim H, Lee K, Cho S, Joh JH
12690	Formation of granulation tissue on bilateral vocal cords after double-lumen endotracheal intubation: A case report
	Xiong XJ, Wang L, Li T
12696	Giant cellular leiomyoma in the broad ligament of the uterus: A case report
	Yan J, Li Y, Long XY, Li DC, Li SJ
12703	Pomolidomide for relapsed/refractory light chain amyloidosis after resistance to both bortezomib and daratumumab: A case report
	Li X, Pan XH, Fang Q, Liang Y
12711	Ureteral- artificial iliac artery fistula: A case report
	Feng T, Zhao X, Zhu L, Chen W, Gao YL, Wei JL
12717	How to manage isolated tension non-surgical pneumoperitonium during bronchoscopy? A case report
	Baima YJ, Shi DD, Shi XY, Yang L, Zhang YT, Xiao BS, Wang HY, He HY
12726	Amiodarone-induced muscle tremor in an elderly patient: A case report
	Zhu XY, Tang XH, Yu H
12734	Surgical treatment of Pitt-Hopkins syndrome associated with strabismus and early-onset myopia: Two case reports
	Huang Y, Di Y, Zhang XX, Li XY, Fang WY, Qiao T
12742	Massive low-grade myxoid liposarcoma of the floor of the mouth: A case report and review of literature
	Kugimoto T, Yamagata Y, Ohsako T, Hirai H, Nishii N, Kayamori K, Ikeda T, Harada H
12750	Gingival enlargement induced by cyclosporine in Medullary aplasia: A case report
	Victory Rodríguez G, Ruiz Gutiérrez ADC, Gómez Sandoval JR, Lomelí Martínez SM
12761	Compound heterozygous mutations in PMFBP1 cause acephalic spermatozoa syndrome: A case report
	Deng TQ, Xie YL, Pu JB, Xuan J, Li XM
12768	Colonic tubular duplication combined with congenital megacolon: A case report
	Zhang ZM, Kong S, Gao XX, Jia XH, Zheng CN



Combon	World Journal of Clinical Cases
Conten	Thrice Monthly Volume 10 Number 34 December 6, 2022
12775	Perforated duodenal ulcer secondary to deferasirox use in a child successfully managed with laparoscopic drainage: A case report
	Alshehri A, Alsinan TA
12781	Complication after nipple-areolar complex tattooing performed by a non-medical person: A case report
	Byeon JY, Kim TH, Choi HJ
12787	Interventional urethral balloon dilatation before endoscopic visual internal urethrotomy for post-traumatic bulbous urethral stricture: A case report
	Ha JY, Lee MS
12793	Regression of gastric endoscopic submucosal dissection induced polypoid nodular scar after <i>Helicobacter pylori</i> eradication: A case report
	Jin BC, Ahn AR, Kim SH, Seo SY
12799	Congenital absence of the right coronary artery: A case report
	Zhu XY, Tang XH



Contents

Thrice Monthly Volume 10 Number 34 December 6, 2022

ABOUT COVER

Editorial Board Member of World Journal of Clinical Cases, Giuseppe Lanza, MD, MSc, PhD, Associate Professor, Department of Surgery and Medical-Surgical Specialties, University of Catania, Catania 95123, Italy. glanza@oasi.en.it

AIMS AND SCOPE

The primary aim of World Journal of Clinical Cases (WJCC, World J Clin Cases) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xu Guo; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Clinical Cases	https://www.wignet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 2307-8960 (online)	https://www.wignet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
April 16, 2013	https://www.wignet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Thrice Monthly	https://www.wignet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku	PUBLICATION MISCONDUCT https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wignet.com/2307-8960/editorialboard.htm	https://www.wignet.com/bpg/gerinfo/242
PUBLICATION DATE December 6, 2022	STEPS FOR SUBMITTING MANUSCRIPTS https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2022 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2022 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: bpgoffice@wjgnet.com https://www.wjgnet.com



W J C C World Journal of Clinical Cases

Submit a Manuscript: https://www.f6publishing.com

World J Clin Cases 2022 December 6; 10(34): 12750-12760

DOI: 10.12998/wjcc.v10.i34.12750

ISSN 2307-8960 (online)

CASE REPORT

Gingival enlargement induced by cyclosporine in Medullary aplasia: A case report

Giovanna Victory Rodríguez, Alondra del Carmen Ruiz Gutiérrez, Juan Ramón Gómez Sandoval, Sarah Monserrat Lomelí Martínez

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): 0 Grade C (Good): C, C Grade D (Fair): 0 Grade E (Poor): E

P-Reviewer: Yang J, China; Kakarmath SV, India; Tanaka H, Japan

Received: September 11, 2022 Peer-review started: September 11, 2022 First decision: September 26, 2022 Revised: October 8, 2022 Accepted: November 8, 2022 Article in press: November 8, 2022 Published online: December 6, 2022



Giovanna Victory Rodríguez, Alondra del Carmen Ruiz Gutiérrez, Periodontics Program, Department of Integrated Dentistry Clinics, Centro Universitario de Ciencias de la Salud, Universidad de Guadalajara, Guadalajara 44340, Jalisco, Mexico

Juan Ramón Gómez Sandoval, Research Institute of Dentistry, Department of Integrated Dentistry Clinics, Centro Universitario de Ciencias de la Salud, Universidad de Guadalajara, Guadalajara 44340, Jalisco, Mexico

Sarah Monserrat Lomelí Martínez, Department of Medical and Life Sciences, Centro Universitario de la Ciénega, Universidad de Guadalajara, Ocotlán 47810, Jalisco, Mexico

Corresponding author: Sarah Monserrat Lomelí Martínez, Doctor, Academic Research, Associate Research Scientist, Department of Medical and Life Sciences, Centro Universitario de la Ciénega, Universidad de Guadalajara, Universidad No. 1115, Lindavista, Ocotlán 47810, Jalisco, Mexico. sarah.lomeli@academicos.udg.mx

Abstract

BACKGROUND

Cyclosporine is an immunosuppressive agent used effectively for treatment of a rare haematological disorder known as medullary aplasia. This drug prevents several side effects, including gingival enlargement (GE) which compromises aesthetics, phonetics and chewing, and also predisposes patients to periodontitis.

CASE SUMMARY

This clinical case reports a 41-year-old woman who presented with cyclosporineinduced GE with underlying periodontitis and medullary aplasia. The management of the disease was approached through multidisciplinary strategy which allowed for accurate diagnosis and a strategic treatment based on the systemic condition and severity of oral pathology. The diagnosis was confirmed through histopathological analysis. The treatment was carried out in phases: Initial (oral hygiene motivation, mechanical supragingival plaque control, and non-surgical therapy); systemic treatment, corrective treatment, and maintenance.

CONCLUSION

Multidisciplinary management of cyclosporine-induced GE and medullary aplasia allows for correct diagnosis and effective treatment of this pathological expression through a phased therapeutic approach.



Key Words: Cyclosporine; Gingival enlargement; Medullary aplasia; Periodontal surgery; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Medullary aplasia is a marrow failure characterized by a total or partial disappearance of hematopoietic precursors in bone marrow, resulting in pancytopenia which is usually accompanied by hemorrhagic syndrome. The treatment involves immunosuppressive therapy with cyclosporine. This drug suppresses the activity of immune cells that damage the bone marrow, although it may produce an inflammatory reaction, with increased fibrotic response in the extracellular matrix, and gingival enlargement (GE). We report a case of GE induced by cyclosporine in a patient with rare medullary aplasia, which was managed with a phased periodontal treatment plan.

Citation: Victory Rodríguez G, Ruiz Gutiérrez ADC, Gómez Sandoval JR, Lomelí Martínez SM. Gingival enlargement induced by cyclosporine in Medullary aplasia: A case report. World J Clin Cases 2022; 10(34): 12750-12760

URL: https://www.wjgnet.com/2307-8960/full/v10/i34/12750.htm DOI: https://dx.doi.org/10.12998/wjcc.v10.i34.12750

INTRODUCTION

Medullary aplasia is a disease characterized by failure of hematopoietic function and consequent peripheral cytopenias. Indeed, medullary aplasia is generally considered a medical emergency. It is characterized by bone marrow failure and total or partial disappearance of hematopoietic precursors which cause pancytopenia in the peripheral blood, as well as hemorrhagic syndrome[1]. Cyclosporine is one of the most widely used immunosuppressive drugs for this disease[2]. However, widespread clinical use of cyclosporine has been questioned due to its various inherent side effects such as nephrotoxicity, hepatotoxicity, hypertension, neurotoxicity, hirsutism, gastrointestinal toxicity, and gingival enlargement (GE)[3].

GE is a pathological condition characterized by excessive growth in mass and volume of gingival tissue, dense collagen stroma, and epithelial hyperplasia[4-6]. Cyclosporine interferes with the modulation of fibrotic response in gingival fibroblasts, and inhibits the secretion of proteases from the matrix which contributes to the accumulation of components of the extracellular matrix in the gingival connective tissue, resulting in gingival growth[7-10]. In addition, this immunosuppressant generates oral microbial dysbiosis in dental biofilm formed by bacteria, fungi and viruses[10].

The therapeutic procedure used in the management of GE with underlying periodontitis in a patient with medullary aplasia is not fixed because it can be modified according to the conditions, behavior, and severity of the oral pathological changes, as well as the patient's systemic condition. This article presents the therapeutic management and maintenance of a clinical case of severe cyclosporine-induced GE with underlying periodontitis in a patient with medullary aplasia.

CASE PRESENTATION

Chief complaints

A female patient aged 41 years was referred to the Periodontics Clinic of the University of Guadalajara by the Hematology Department of the Fray Antonio Alcalde Civil Hospital of Guadalajara, Jalisco. The reason for consultation was evaluation of enlarged gums, the patient perceived an increase in the sizes of the gums in both arches.

History of present illness

The patient had enlargement of gums without associated symptoms 1 year before coming to the Periodontics Clinic.

History of past illness

Her medical history showed that, in 2019, she was hospitalized for gingival and vaginal bleeding, and she received blood transfusion. She was diagnosed with hypothyroidism and medullary aplasia in 2019. The pharmacological treatment consisted of 100 mg Cyclosporine given in 3 daily doses, Tribedoce (1 daily, comprising 100 mg of thiamine, 5 mg of pyridoxine and 50 mg of cyanocobalamin); Bedoyecta Tri



(1 daily, comprising 100 mg of hydroxycobalamin, 50 mg of thiamine, and 2 mL of pyridoxine); Cilocid (5 mg folic acid, 1 daily), and 600 Autrin (1 daily, consisting of iron and vitamins), as well as 100 mg of levothyroxine (1 daily) for hypothyroidism.

Personal and family history

Her father was diagnosed with type 2 diabetes mellitus in 2020.

Physical examination

During the intraoral clinical examination, increased nodal gingival volume was observed in the upper and lower arcades located mainly in the anterior area through both vestibular and palatine/lingual access. There was erythema in all areas, with increased volume, biofilm, and generalized calculus; the root remained intact, and carious lesions were evident (Figure 1). The periodontal evaluation presented depths to the probe of up to 12 mm, mainly in anterior teeth; the generalized probing caused bleeding, and there was mobility in some of the dentition (Figure 2).

Laboratory examinations

There were normal levels of platelets ($275 \times 10^3/\mu$ L) and white cells ($5.68 \times 10^3/\mu$ L), while hemoglobin level was slightly low (11.6 g/dL), and blood glucose level was 93 mg/dL. No abnormality was found in urine analysis.

Imaging examinations

At the radiographic level, generalized bone loss was observed in a horizontal pattern, being more accentuated in teeth 11, 21, 31, and 41 where the loss extended to the middle third (Figure 3).

FINAL DIAGNOSIS

Based on all the information obtained from the clinical and radiographic examinations, the diagnosis was localized periodontitis stage III, grade C in teeth 11, 21, 31 and 41, and generalized periodontitis stage I, grade C. These were associated with a presumptive diagnosis of drug-induced GE. The periodontitis was classified according to the new classification system of periodontal diseases and conditions of the American Academy of Periodontology and the European Federation of Periodontology (2018).

The histopathology study showed that the specimen presented fibroconnective tissue growth, epithelial nails exhibited elongation and pseudo-epithelionomatous hyperplasia. The connective tissue presented large focal areas of chronic inflammatory infiltrations and lymphoplasmacytic dispersion in a stroma of collagen fibers and active fibroblasts (Figure 4). This analysis corroborated the presumptive diagnosis of drug-associated GE.

TREATMENT

Initial phase

The first periodontal phase began with generalized supragingival debridement and hygiene instructions which consisted of advising the patient on dental brushing, use of interproximal brushes, and rinsing with 0.12% chlorhexidine. In the following months, subgingival debridement was performed in the four quadrants, and due to the coronavirus disease 2019 pandemic, the patient was summoned after 3 mo. Upon the patient's return, control photographs were taken in which reductions in the color, size, and shape of the GE were clearly observed (Figure 5).

Systemic phase

Consultation with the hematologist was carried out. The hematologist approved the surgical procedure and adjusted daily dose of cyclosporine from 3 capsules daily to 1 capsule every other day. Clindamycin (600 mg) was indicated 1 h before the intervention.

Corrective phase

The surgical approach in the upper arch comprised premolar to premolar. Measurements were made to calculate the amount of gingival tissue above the clinical crown, and 3 mm coronal to the mucogingival line was considered as internal level for submarginal incisions to generate new papillae and thin the tissues, with a 15 C scalpel blade. Then, intrasulcular incisions were made to reflect full-thickness flaps in the vestibular and palatine (Figure 6).

Granulomatous tissue was removed with McCall 13-14 and 17-18 curettes, and the roots were scraped and smoothed. Osteoplasties were performed with carbide ball burs. Gingival nodules were thinned with LaGrange scissors, the flaps were adapted, and a tissue sample was taken for histopathological





DOI: 10.12998/wjcc.v10.i34.12750 Copyright ©The Author(s) 2022.

Figure 1 Gingival enlargement in the adult patient treated with cyclosporine. In general, severe inflammation, redness of tissue, nodules, and calculus deposits were observed. A: Inter-occlusal view; B: Left side view; C: Upper occlusal view; D: Lower occlusal view;

> analysis. The incision site was sutured using polyglycolic acid 4-0 with modified horizontal mattress stitches, and periodontal dressing was applied. Paracetamol (500 mg) was administered for 3 d, and the prescription indicated that it was to be continued along with 300 mg clindamycin for 7 d. Moreover, 0.12% chlorhexidine gel was placed in the treated area 3 times daily for 15 d.

> The patient's healing was reviewed at 2 postoperative weeks. When the sutures were removed, the tissue was in good condition, with slight inflammation corresponding to the healing process (Figure 7). At one month, surgery was performed on the lower arch from canine to canine, with the technique previously described, and in the same way, the sutures were removed at 2 wk (Figures 8 and 9). Histopathological analysis was performed at the Center for Research and Diagnosis of Pathology of the University of Guadalajara. The specimen presented fibroconnective tissue growth and pseudo-epithelionomatous hyperplasia. The connective tissue showed large focal areas of chronic inflammatory infiltration and lymphoplasmacytic dispersion in a stroma of collagen fibers and active fibroblasts (Figure 4). These lesions corroborated the presumptive diagnosis of drug-associated GE.

OUTCOME AND FOLLOW-UP

Maintenance phase

Three months after conclusion of the corrective phase, the periodontal status was evaluated. Inflammation was observed in marginal and papillary areas in teeth 12, 13, 22, 23, 41, 31, 42 and 33, as well as redness and moderate calculus. The maintenance stage was focused on maintaining control of dental biofilm. Therefore, supragingival debridement with ultrasound was performed, and indications of oral hygiene were reinforced. The next maintenance visit was at 6 mo post-surgery, at which period it was seen that the patient's hygiene was in constant control (Figure 10). A comparison of her initial photographs with the most recent photographs at 6 mo of the maintenance phase, revealed the changes obtained through periodontal therapy (Figure 11).

DISCUSSION

GE is currently defined as "gingival overgrowth" manifested as increase in size (hypertrophy) and increase in number (hyperplasia) of tissue cells. More than 20 prescription drugs are known to be used for GE. These drugs comprise dilanthine group medications, calcium channel blockers, and cyclosporine [11]. Enlargement of gingiva triggers inflammatory reactions in the periodontium, resulting in generation of periodontal disease, or aggravation of existing periodontal disease[6].

Recent studies revealed that the accumulation of gingival fibroblasts observed in drug-associated gingival hyperplasia resulted from inhibition of apoptosis[12]. It has been reported that the pathogenesis of cyclosporine-induced gingival hyperplasia is associated with up-regulated expression





Figure 2 Periodontograms. A: Upper; B: Lower.

levels of proinflammatory cytokines in saliva *i.e.*, interleukin (IL)-1α, IL-8, and IL-6[13].

Various risk factors for GE have been identified. These are plaque control, level of gingival inflammation, extent of periodontal destruction, dose and duration of cyclosporine therapy, plasma and tissue concentrations of cyclosporine and its metabolites, age of the patient, and perhaps the underlying comorbidities[14,15]. Although suspending the use of medication in most cyclosporine-induced GE cases is not a viable option, such was the case with the patient in this study. In previous studies, it was reported that maintenance of adequate oral hygiene in humans markedly reduced the severity of cyclosporine-induced gingival hyperplasia [16,17]. In this study, although a significant reduction in GE was observed after the initial phase, it did not prevent the development of gingival hyperplasia.

Periodontitis is a bacterial infection that, if not treated, may cause systemic complications such as bacteremia, systemic lesions by toxins from oral pathogens, and systemic inflammation due to soluble antigens of oral pathogens[18]. In addition, dental biofilm plays a role as a reservoir of multi-resistant pathogens such as S. aureus, P. aeruginosa, A. baumannii and C. albicans which are associated with severe infections such as meningitis, endocarditis, sepsis and necrotizing fasciitis, especially in immunocompromised patients [19]. This is relevant because patients with medullary aplasia have altered immune responses due to immunosuppressive therapy. Therefore, the presence of periodontitis may trigger serious opportunistic infections. Therefore, the main objective of the periodontal treatment carried out in the present case was to reduce the pathogenic bacterial load in the oral cavity, in addition to use of initial therapy with rigorous hygiene which made it possible to reduce inflammatory response





DOI: 10.12998/wjcc.v10.i34.12750 Copyright ©The Author(s) 2022.

Figure 3 Images from complete periapical study. Generalized bone loss is evident, with greater loss in the incisors. A: Upper; B: Lower.



DOI: 10.12998/wjcc.v10.i34.12750 Copyright ©The Author(s) 2022.

Figure 4 Gingival histopathology. A: Fibroconnective tissue growth, elongated epithelial nails, and pseudo-epithelionomatous hyperplasia; B: Connective tissue showing large focal areas of chronic inflammatory infiltration and lymphoplasmacytic dispersion in a stroma of collagen fibers and active fibroblasts.



DOI: 10.12998/wjcc.v10.i34.12750 Copyright ©The Author(s) 2022.

Figure 5 Tissue conditions 3 mo after initial phase. The features depict decreased inflammation, pink coloration in interproximal areas, and smaller nodules. A: Inter-occlusal view; B: Upper sextant.

in gingival tissues[20].

Pharmacological treatments cannot be modified in most cases of severe GE, as this would lead to noncontrol of the systemic disease. In the case of the patient in this study, treatment modification aggravated the medullary aplasia, resulting in reappearance of the lesions after surgical removal, thereby requiring repeated interventions. Evidence suggests that 34% of cases show recurrence during 18 mo following periodontal surgery, regardless of medication[21]. In our patient's case, consultation with a hematologist was carried out in a bid to modify the dose of cyclosporine administered, from the initial schedule of one 100-mg capsule three times a day, to one 100-mg capsule every other day. When compared with the initial phase, an important reduction in enlargement was observed. This reveals the importance of adequate multidisciplinary management for GE.

Gaishideng® WJCC | https://www.wjgnet.com



DOI: 10.12998/wjcc.v10.i34.12750 Copyright ©The Author(s) 2022.

Figure 6 Corrective phase. A: Vestibular reflected flap; B: Palatine reflected flap; C: Modified horizontal mattress sutures (vestibular view); D: Modified horizontal mattress sutures (palatal view).



DOI: 10.12998/wjcc.v10.i34.12750 Copyright ©The Author(s) 2022.

Figure 7 Healing at postoperative 2 wk. A slight inflammation was observed in line with the healing process, and the sutures were removed. A: Upper; B: Lower.

> Periodontal therapeutic strategies for GE are generally categorized into initial, systemic, corrective and maintenance phases, depending on the severity and conditions of the disease[21]. The initial phase consists of oral hygiene instructions, dental biofilm control, and supra- and sub-gingival debridement. The aim of the systemic phase was to reduce the influence of cyclosporine on the results of periodontal therapy. This was the reason for which the hematologist reduced the drug dose. The corrective phase allows the return of periodontal tissues to states compatible with health. This phase is necessary when GE is severe or persistent, notwithstanding strict control of the dental biofilm. In this case, surgical correction is suggested. In the present case, the patient was subjected to the two periodontal phases, and



Baisbideng® WJCC | https://www.wjgnet.com



DOI: 10.12998/wjcc.v10.i34.12750 Copyright ©The Author(s) 2022.

Figure 8 Postoperative view of the lower anterior sextant. Depicting modified horizontal mattress sutures.



DOI: 10.12998/wjcc.v10.i34.12750 Copyright ©The Author(s) 2022.

Figure 9 Lower anterior sextant at postoperative 2 wk. Inflammation was observed in line with the healing process, and the sutures were removed.



DOI: 10.12998/wjcc.v10.i34.12750 Copyright ©The Author(s) 2022.

Figure 10 Photographs 3 mo after corrective phase. The figure depicts a general improvement in tissue conditions. A: Inter-occlusal view; B: Right side view; C: Left side view.

at the end of the initial phase, a considerable decrease in GE was observed, the patient's hygiene control was improved, but the enlargement persisted. The periodontal corrective phase was performed with the flap surgical technique which allowed for reduction of these defects while eliminating excess tissue, thereby resulting in gingival harmony that facilitated oral hygiene to the patient[5,6].

Baishideng® WJCC | https://www.wjgnet.com



DOI: 10.12998/wjcc.v10.i34.12750 Copyright ©The Author(s) 2022.

Figure 11 Comparative photographs. A and C: Tissue conditions when the patient attended the periodontics clinic; B and D: Tissue status after three phases of periodontal treatment.

> On the other hand, the maintenance phase began as soon as the periodontitis and GE were treated via the initial and corrective phase where inflammation was controlled, destruction of supporting tissues was stopped, and the enlargements in gingival tissue was removed[22]. The goal of this phase was to keep the teeth functional, without pain, excessive mobility, or long-term persistent infection. For this purpose, periodic visits were scheduled for maintenance of excellent oral hygiene, in addition to frequent monitoring of healing process, or recurrence of the disease. In this particular case, the added risk due to the use of immunosuppressive drugs was taken into consideration, since it makes GE patients highly susceptible to recurrence if optimal bacterial oral control is not maintained[23].

> Microbial dysbiosis plays a role in GE. However, the accumulation of dental biofilm is another key factor in the severity of this pathology [5,7,8]. This case has revealed that at the end of the periodontal initial phase involving strict control of dental biofilm, administering the same initial dose of cyclosporine allowed the patient to control levels of dental biofilm and inflammation, resulting in a large decrease in GE. Moreover, at the level of soft tissue management, the reductions in inflammation and biofilm positively impacted the next phase of treatment.

CONCLUSION

Adequate multidisciplinary management of cyclosporine-induced GE patients with medullary aplasia allows for accurate diagnosis and treatment of the disease. The choice of therapeutic strategies depends on the tissue conditions and severity of the systemic disease. In the case studied, the patient's response to medullary aplasia treatment was favorable. The patient was stable, and she kept appointments regularly. The patient had difficulty in cleaning the interproximal areas of the lower teeth, but in each maintenance appointment, physiotherapy was reinforced.

FOOTNOTES

Author contributions: Victory Rodríguez G, Ruiz Gutiérrez ADC and Lomelí Martínez SM contributed to diagnosis, surgical treatment and manage of patient; Lomelí Martínez SM and Victory Rodríguez G contributed to acquisition and analysis of data; Ruiz Gutiérrez ADC and Gómez Sandoval JR drafted manuscript and critically revised manuscript; all authors have read and approved the final manuscript.

Informed consent statement: Informed written consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was



prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is noncommercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: Mexico

ORCID number: Giovanna Victory Rodríguez 0000-0001-5543-0604; Alondra del Carmen Ruiz Gutiérrez 0000-0001-9672-2494; Juan Ramón Gómez Sandoval 0000-0002-3336-5159; Sarah Monserrat Lomelí Martínez 0000-0002-0569-1387.

S-Editor: Wei ZH L-Editor: A P-Editor: Wei ZH

REFERENCES

- Vallejo Llamas JC. Management of aplastic anemia during the phase of defervescence of the COVID-19 pandemic. Med 1 Clin (Engl Ed) 2021; 157: 89-90 [PMID: 34179500 DOI: 10.1016/j.medcle.2021.02.003]
- 2 Lorand IG, Souza CA, Caliani JA, Fabron Júnior A. [Medullary aplasia: diagnostic criteria, clinical course and prognostic factors]. AMB Rev Assoc Med Bras 1983; 29: 215-218 [PMID: 6608757]
- Khoori AH, Einollahi B, Ansari G, Moozeh MB. The effect of cyclosporine with and without nifedipine on gingival overgrowth in renal transplant patients. J Can Dent Assoc 2003; 69: 236-241 [PMID: 12662462]
- 4 Hallmon WW, Rossmann JA. The role of drugs in the pathogenesis of gingival overgrowth. A collective review of current concepts. Periodontol 2000 1999; 21: 176-196 [PMID: 10551182 DOI: 10.1111/j.1600-0757.1999.tb00175.x]
- Morales D, González M. E., Rangel L. Agrandamiento Gingival Generalizado en un Paciente con Trasplante Renal. 5 Revista Cubana de Medicina General Integral 2019; 35:658
- 6 Ayanoglou CM, Lesty C. Cyclosporin A-induced gingival overgrowth in the rat: a histological, ultrastructural and histomorphometric evaluation. J Periodontal Res 1999; 34: 7-15 [PMID: 10086881 DOI: 10.1111/j.1600-0765.1999.tb02216.x
- Moffitt ML, Bencivenni D, Cohen RE. Drug-induced gingival enlargement: an overview. Compend Contin Educ Dent 7 2013; 34: 330-336 [PMID: 23991852]
- 8 Nakib N, Ashrafi SS. Drug-induced gingival overgrowth. Dis Mon 2011; 57: 225-230 [PMID: 21569887 DOI: 10.1016/j.disamonth.2011.03.010]
- Lauritano D, Palmieri A, Lucchese A, Di Stasio D, Moreo G, Carinci F. Role of Cyclosporine in Gingival Hyperplasia: An 9 In Vitro Study on Gingival Fibroblasts. Int J Mol Sci 2020; 21 [PMID: 31963361 DOI: 10.3390/ijms21020595]
- Triveni MG, Rudrakshi C, Mehta DS. Amlodipine-induced gingival overgrowth. J Indian Soc Periodontol 2009; 13: 160-10 163 [PMID: 20379416 DOI: 10.4103/0972-124X.60231]
- Agnihotri R, Bhat KM, Bhat GS, Pandurang P. Periodontal management of a patient with severe aplastic anemia: a case 11 report. Spec Care Dentist 2009; 29: 141-144 [PMID: 19938255 DOI: 10.1111/j.1754-4505.2009.00079.x]
- 12 Jung JY, Jeong YJ, Jeong TS, Chung HJ, Kim WJ. Inhibition of apoptotic signals in overgrowth of human gingival fibroblasts by cyclosporin A treatment. Arch Oral Biol 2008; 53: 1042-1049 [PMID: 18471799 DOI: 10.1016/j.archoralbio.2008.03.008]
- 13 Ruhl S, Hamberger S, Betz R, Sukkar T, Schmalz G, Seymour RA, Hiller KA, Thomason JM. Salivary proteins and cytokines in drug-induced gingival overgrowth. J Dent Res 2004; 83: 322-326 [PMID: 15044507 DOI: 10.1177/154405910408300410]
- 14 Parkar MH, Hussain F, Wickramaratna A, Olsen I. The immunosuppressant and hyperplasia-inducing drug cyclosporin A regulates the cell cycle and cyclin B1 gene expression in gingival fibroblasts in vitro. Cell Tissue Res 2004; 317: 221-225 [PMID: 15221438 DOI: 10.1007/s00441-004-0909-3]
- 15 Drozdzik M, Kurzawski M, Drozdzik A, Kotrych K, Banach J, Pawlik A. Interleukin-6 gene polymorphism in renal transplant patients with and without gingival overgrowth. J Clin Periodontol 2005; 32: 955-958 [PMID: 16104959 DOI: 10.1111/j.1600-051X.2005.00766.x
- 16 Seymour RA, Jacobs DJ. Cyclosporin and the gingival tissues. J Clin Periodontol 1992; 19: 1-11 [PMID: 1732303 DOI: 10.1111/j.1600-051x.1992.tb01140.x]
- 17 Seymour RA, Smith DG. The effect of a plaque control programme on the incidence and severity of cyclosporin-induced gingival changes. J Clin Periodontol 1991; 18: 107-110 [PMID: 2005223 DOI: 10.1111/j.1600-051x.1991.tb01698.x]
- 18 Teshome A, Girma B, Aniley Z. The efficacy of azithromycin on cyclosporine-induced gingival enlargement: Systematic review and meta-analysis. J Oral Biol Craniofac Res 2020; 10: 214-219 [PMID: 32489824 DOI: 10.1016/j.jobcr.2019.12.005
- Morton RS, Dongari-Bagtzoglou AI. Regulation of gingival fibroblast interleukin-6 secretion by cyclosporine A. J Periodontol 1999; 70: 1464-1471 [PMID: 10632522 DOI: 10.1902/jop.1999.70.12.1464]
- 20 Han YW, Wang X. Mobile microbiome: oral bacteria in extra-oral infections and inflammation. J Dent Res 2013; 92: 485-491 [PMID: 23625375 DOI: 10.1177/0022034513487559]
- Lindhe J, Lang N, Berglundh T, Giannobile W, Sanz M. Lindhe's Clinical Periodontology and Implant Dentistry. 7th ed. 21



2 Volume Set. Wiley-Blackwell, 2021: 663-761

- 22 Ilgenli T, Atilla G, Baylas H. Effectiveness of periodontal therapy in patients with drug-induced gingival overgrowth. Long-term results. J Periodontol 1999; 70: 967-972 [PMID: 10505798 DOI: 10.1902/jop.1999.70.9.967]
- 23 Manresa C, Sanz-Miralles EC, Twigg J, Bravo M. Supportive periodontal therapy (SPT) for maintaining the dentition in adults treated for periodontitis. Cochrane Database Syst Rev 2018; 1: CD009376 [PMID: 29291254 DOI: 10.1002/14651858.CD009376.pub2]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

