

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 January 26; 11(3): 487-718



## Contents

Thrice Monthly Volume 11 Number 3 January 26, 2023

## MINIREVIEWS

- 487** Protective effects of combined treatment with ciprofol and mild therapeutic hypothermia during cerebral ischemia-reperfusion injury  
*Wang YC, Wu MJ, Zhou SL, Li ZH*
- 493** Non-pulmonary involvement in COVID-19: A systemic disease rather than a pure respiratory infection  
*El-Kassas M, Alboraie M, Elbadry M, El Sheemy R, Abdellah M, Afify S, Madkour A, Zaghloul M, Awad A, Wifi MN, Al Balakosy A, Eltabbakh M*
- 506** Progress and expectation of stem cell therapy for diabetic wound healing  
*Xu ZH, Ma MH, Li YQ, Li LL, Liu GH*
- 514** Prevention, diagnostic evaluation, management and prognostic implications of liver disease in critically ill patients with COVID-19  
*Valsamaki A, Xanthoudaki M, Oikonomou KG, Vlachostergios PJ, Papadogoulas A, Katsiafylloudis P, Voulgaridi I, Skoura AL, Komnos A, Papamichalis P*
- 528** Exosomal miRNA in early-stage hepatocellular carcinoma  
*Wu ZQ, Zhu YX, Jin Y, Zhan YC*
- 534** Impact of multidrug resistance on the management of bacterial infections in cirrhosis  
*Terra C, de Mattos ÁZ, Chagas MS, Torres A, Wiltgen D, Souza BM, Perez RM*
- 545** Could there be an interplay between periodontal changes and pancreatic malignancies?  
*Ungureanu BS, Gheorghe DN, Nicolae FM, Râmboiu S, Radu PA, Șurlin VM, Strâmbu VDE, Gheonea DI, Roman A, Șurlin P*

## ORIGINAL ARTICLE

## Retrospective Study

- 556** Qixue Shuangbu decoction and acupuncture combined with Western medicine in acute severe stroke patients  
*Gou LK, Li C*
- 566** Successful treatment of patients with refractory idiopathic membranous nephropathy with low-dose Rituximab: A single-center experience  
*Wang YW, Wang XH, Wang HX, Yu RH*
- 576** Bowel inflammatory presentations on computed tomography in adult patients with severe aplastic anemia during flared inflammatory episodes  
*Zhao XC, Xue CJ, Song H, Gao BH, Han FS, Xiao SX*

- 598 Clinical outcomes of AngioJet pharmacomechanical thrombectomy *versus* catheter-directed thrombolysis for the treatment of filter-related caval thrombosis

*Li JY, Liu JL, Tian X, Jia W, Jiang P, Cheng ZY, Zhang YX, Liu X, Zhou M*

#### Clinical Trials Study

- 610 Efficacy and safety of propofol target-controlled infusion combined with butorphanol for sedated colonoscopy

*Guo F, Sun DF, Feng Y, Yang L, Li JL, Sun ZL*

#### Observational Study

- 621 Application of a hospital-community-family trinity rehabilitation nursing model combined with motor imagery therapy in patients with cerebral infarction

*Li WW, Li M, Guo XJ, Liu FD*

#### CASE REPORT

- 629 Congenital biliary atresia caused by *GPC1* gene mutation in Chinese siblings: A case report

*Kong YM, Yuan K, Wang CL*

- 635 Rescuing "hopeless" avulsed teeth using autologous platelet-rich fibrin following delayed reimplantation: Two case reports

*Yang Y, Liu YL, Jia LN, Wang JJ, Zhang M*

- 645 Acute diffuse peritonitis secondary to a seminal vesicle abscess: A case report

*Li K, Liu NB, Liu JX, Chen QN, Shi BM*

- 655 Young thoracic vertebra diffuse idiopathic skeletal hyperostosis with Scheuermann disease: A case report

*Liu WZ, Chang ZQ, Bao ZM*

- 662 Relapsed primary extraskeletal osteosarcoma of liver: A case report and review of literature

*Di QY, Long XD, Ning J, Chen ZH, Mao ZQ*

- 669 Heterotopic pregnancy after assisted reproductive techniques with favorable outcome of the intrauterine pregnancy: A case report

*Wang YN, Zheng LW, Fu LL, Xu Y, Zhang XY*

- 677 Periprosthetic knee joint infection caused by *Brucella melitensis* which was first -osteoarticular brucellosis or osteoarthritis: A case report

*Stumpner T, Kuhn R, Hochreiter J, Ortmaier R*

- 684 Recurrent intramuscular lipoma at extensor pollicis brevis: A case report

*Byeon JY, Hwang YS, Lee JH, Choi HJ*

- 692 Imaging features of retinal hemangioblastoma: A case report

*Tang X, Ji HM, Li WW, Ding ZX, Ye SL*

- 700** Clinical and genetic diagnosis of autosomal dominant osteopetrosis type II in a Chinese family: A case report  
*Gong HP, Ren Y, Zha PP, Zhang WY, Zhang J, Zhang ZW, Wang C*
- 709** Soft tissue tuberculosis detected by next-generation sequencing: A case report and review of literature  
*He YG, Huang YH, Yi XL, Qian KL, Wang Y, Cheng H, Hu J, Liu Y*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Baharudin Abdullah, MMed, Professor, Surgeon, Department of Otorhinolaryngology-Head and Neck Surgery, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan, Malaysia. profbaha@gmail.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

January 26, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Impact of multidrug resistance on the management of bacterial infections in cirrhosis

Carlos Terra, Ângelo Zambam de Mattos, Marcelo Souza Chagas, Andre Torres, Denusa Wiltgen, Barbara Muniz Souza, Renata Mello Perez

**Specialty type:** Gastroenterology and hepatology

**Provenance and peer review:** Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): A  
Grade B (Very good): B, B  
Grade C (Good): 0  
Grade D (Fair): D  
Grade E (Poor): 0

**P-Reviewer:** Bukhari SM, Pakistan; Kreisel W, Germany; Sahle Z, Ethiopia

**Received:** November 10, 2022

**Peer-review started:** November 10, 2022

**First decision:** November 26, 2022

**Revised:** December 22, 2022

**Accepted:** January 5, 2023

**Article in press:** January 5, 2023

**Published online:** January 26, 2023



**Carlos Terra, Marcelo Souza Chagas, Andre Torres, Barbara Muniz Souza,** Gastroenterology-Liver Unit, State University of Rio de Janeiro, Rio de Janeiro 20551-030, Rio de Janeiro, Brazil

**Carlos Terra,** Liver Unit, Casa de Saúde São José-Rede Santa Catarina, Rio de Janeiro 22271-080, Rio de Janeiro, Brazil

**Carlos Terra, Ângelo Zambam de Mattos, Marcelo Souza Chagas, Andre Torres, Denusa Wiltgen, Barbara Muniz Souza, Renata Mello Perez,** Alliance of Brazilian Centers for Cirrhosis Car, The ABC Group, Rio de Janeiro 20551-030, Rio de Janeiro, Brazil

**Carlos Terra,** Liver Unit, Federal Hospital of Lagoa, Rio de Janeiro 22470-050, Rio de Janeiro, Brazil

**Ângelo Zambam de Mattos,** Graduate Program in Medicine: Hepatology, Federal University of Health Sciences of Porto Alegre, Porto Alegre 90020-090, Rio Grande do Sul, Brazil

**Ângelo Zambam de Mattos,** Gastroenterology and Hepatology Unit, Irmandade Santa Casa de Misericórdia de Porto Alegre, Porto Alegre 90020-090, Rio Grande do Sul, Brazil

**Marcelo Souza Chagas,** Internal Medicine, Federal Hospital of Lagoa, Rio de Janeiro 22470-050, Rio de Janeiro, Brazil

**Denusa Wiltgen,** Department of Internal Medicine, Federal University of Health Sciences of Porto Alegre, Porto Alegre 90020-090, Brazil

**Renata Mello Perez,** Hepatology Division, Federal University of Rio de Janeiro, Rio de Janeiro 21941-617, Rio de Janeiro, Brazil

**Renata Mello Perez,** IDOR, D'Or Institute for Research and Education, Rio de Janeiro 22281-100, Rio de Janeiro, Brazil

**Corresponding author:** Carlos Terra, MD, PhD, Associate Professor, Doctor, Gastroenterology-Liver Unit, State University of Rio de Janeiro, Boulevard 28 Setembro, 77-3rd floor-Vila Isabel, Rio de Janeiro 20551-030, Rio de Janeiro, Brazil. [carlosterrior@hotmail.com](mailto:carlosterrior@hotmail.com)

### Abstract

Patients with cirrhosis have an increased risk of infection and differently from other complications, that over the years are improving in their outcomes, infections in cirrhotic patients are still a major cause of hospitalization and death

(up to 50% in-hospital mortality). Infections by multidrug-resistant organisms (MDRO) have become a major challenge in the management of cirrhotic patients with significant prognostic and cost-related impact. About one third of cirrhotic patients with bacterial infections is infected with MDR bacteria and their prevalence has increased in recent years. MDR infections have a worse prognosis compared to infections by non-resistant bacteria because they are associated with lower rate of infection resolution. An adequate management of cirrhotic patients with infections caused by MDR bacteria depends on the knowledge of some epidemiological aspects, such as the type of infection (spontaneous bacterial peritonitis, pneumonia, urinary tract infection and spontaneous bacteremia), bacteriological profile of antibiotic resistance at each health care unit and site of infection acquisition (community acquired, healthcare associated or nosocomial). Furthermore, regional variations in the prevalence of MDR infections determine that the choice of empirical antibiotic therapy must be adapted to the local microbiological epidemiology. Antibiotic treatment is the most effective measure to treat infections caused by MDRO. Therefore, optimizing antibiotic prescribing is critical to effectively treat these infections. Identification of risk factors for multidrug resistance is essential to define the best antibiotic treatment strategy in each case and the choice of an effective empirical antibiotic therapy and its early administration is cardinal to reduce mortality. On the other hand, the supply of new agents to treat these infections is very limited. Thus, specific protocols that include preventive measures must be implemented in order to limit the negative impact of this severe complication in cirrhotic patients.

**Key Words:** Cirrhosis; Infection; Multidrug-resistance; Bacterial; Antibiotics; Microbiota

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Infections by multidrug-resistant organisms (MDRO) have become a major challenge in the management of cirrhotic patients with significant prognostic and cost-related impact. This review presents the main epidemiological data, clinical impact, risk factors, and the best management of cirrhotic patients infected with MDR bacteria.

**Citation:** Terra C, de Mattos ÂZ, Chagas MS, Torres A, Wiltgen D, Souza BM, Perez RM. Impact of multidrug resistance on the management of bacterial infections in cirrhosis. *World J Clin Cases* 2023; 11(3): 534-544

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i3/534.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i3.534>

## INTRODUCTION

The rising prevalence of multidrug-resistant organisms (MDRO) (resistant at least to one agent in three or more antimicrobial categories), extensively drug-resistant organisms (a resistance profile that compromised at least one agent in all but two or fewer antimicrobial categories), and pan drug-resistant organisms (PDRO) (resistant to all known antimicrobial agents) represents a global threat to human health[1,2]. At same time, active agents against MDRO are limited despite an increase in the availability of novel antibiotics in recent years.

Reports from the United States and Europe estimate a death toll of 29-33 patients each year associated with antimicrobial resistant microorganisms, with a huge attributable healthcare cost[2-5]. The so called “ESKAPE” pathogens [*Enterococcus faecium*, *Staphylococcus aureus* (*S. aureus*), *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacteriaceae* species] are especially worrisome: The acquisition of antimicrobial resistance genes, in association to the classical mechanisms of antimicrobial resistance (inactivation or alteration of the antimicrobial molecule, bacterial target site modifications, reduced antibiotic penetration/accumulation, and the formation of bacterial biofilms), make ESKAPE pathogens challenging to surveillance and subsequent infections difficult to treat. The SKAPE pathogens have developed resistance against almost all antibiotics used in the clinical setting (oxazolidinones, lipopeptides, macrolides, fluoroquinolones, B-lactams, and also to antibiotics that are considered “the last line of defense”, like carbapenems, glycopeptides and polymyxins). Face to the increased burden of disease and death rates due to treatment failure, the World Health Organization has designated the SKAPE group as “priority organisms” to focus and guide research and development related to new antibiotics[1].

Over the last years cirrhosis complications are improving their outcomes with new strategies and technologies. However, infections in cirrhotic patients are still a major cause of hospitalization and death (up to 50% in-hospital mortality)[6]. Although the reasons for these imbalanced morbidity and



mortality are not totally elucidated, MDR infections may play an important role. In a recent multicentric study that assessed the epidemiology of bacterial infections in hospitalized cirrhotic patients, the overall prevalence of MDRO was 34%[7]. Another report concerning critically ill patients with decompensated cirrhosis points to 46% of isolates being MDROs at admissions. Meanwhile, MDRO isolates responsible for infection during intensive care unit stay were at 60%[8]. Different studies confirm the ominous prognosis of cirrhotic patients with infections by MDRO[9,10] (Table 1). Antimicrobial standard prescription for infection in liver cirrhosis doesn't routinely comprehend the MDRO spectrum. Cirrhosis itself increases the risk for sepsis and septic shock and when associated with an inappropriate choice of empirical antimicrobial treatment may be determinant of a worst outcome.

In this review, we describe the epidemiology, clinical settings and the current evidence-based strategies for early recognition and treatment alternatives for MDR infection in cirrhotic patients.

## EPIDEMIOLOGY

Bacterial infections affect 25%-35% of hospitalized patients with cirrhosis and lead to a four-fold increase in their mortality when compared to noninfected counterparts[11]. The prognostic impact of infections is such that they are considered defining events of state 6 (end-state, late decompensation) in the clinical course of cirrhosis[12]. Moreover, infections (together with alcoholic hepatitis) are the most important drivers of acute decompensation of cirrhosis and acute-on-chronic liver failure (ACLF)[9,13,14].

Despite the evolution in medical care, there is evidence suggesting that mortality associated with infections in individuals with cirrhosis might be increasing. Our group, for instance, has demonstrated that mortality associated with spontaneous bacterial peritonitis increased from 22% to 40% in a decade [15,16]. An explanation for this finding might be the growing importance of infections caused by MDRO. In our setting, after evaluating 5800 isolates from hospitalized patients, we have shown that 38% and 44% of individuals with and without cirrhosis respectively were infected with MDRO. Furthermore, in that study, 20% of *Escherichia coli* and *Klebsiella sp* strains infecting patients with cirrhosis were extended-spectrum beta-lactamase (ESBL)-producing bacteria, and 44% of *S. aureus* strains were methicillin-resistant[17].

Additionally, a recent European study has clearly demonstrated that infections associated with MDRO are increasing in individuals with cirrhosis. The study evaluated two prospective multicenter cohorts of patients hospitalized for acute decompensation of cirrhosis or ACLF. The first cohort consisted of 1146 individuals evaluated in 2011, of which 39.7% were infected. The second cohort consisted of 883 individuals evaluated in 2018, of which 32.2% were infected. In that study, infections associated with MDRO were diagnosed in 29.2% of subjects with positive cultures in the 2011 cohort and in 37.9% of those pertaining to the 2018 cohort[10].

On a global level, another prospective cohort study has demonstrated the relevance of infections associated with MDRO in individuals with cirrhosis worldwide. The authors included 1302 infected patients with cirrhosis from 46 different centers in Europe, America and Asia. The most common infections were spontaneous bacterial peritonitis [spontaneous bacterial peritonitis (SBP), 27%], urinary tract infection (22%) and pneumonia (19%), and 57% of isolates consisted of Gram-negative bacteria. Among individuals with positive cultures, 34% were infected with MDRO, most commonly ESBL-producing *Enterobacteriaceae*, methicillin-resistant *S. aureus*, vancomycin-resistant *Enterococci*, *Pseudomonas aeruginosa*, and *Acinetobacter baumannii*. The prevalence of infections caused by MDRO was higher in Asia (51%), than in Europe (29%) or America (27%). Independent risk factors for infections with MDRO were being from Asia (and mostly from India, where MDRO were present in 73% of isolates), using antibiotics in the three months previous to hospital admission, being exposed to healthcare facilities, and the site of infection (pneumonia, skin and soft tissue infection and urinary tract infection had higher odds of being caused by MDRO)[7].

Africa and Oceania are poorly represented in studies evaluating the prevalence of MDRO in patients with cirrhosis. However, a recent retrospective cohort study has demonstrated a low prevalence of MDRO in blood cultures of patients with cirrhosis hospitalized in Australia (5.6% of admissions). Despite the low prevalence, the study has shown a significant increase in infections caused by MDRO over a decade[18], similarly to what had been previously verified in Europe[10].

## PROGNOSTIC IMPACT OF MULTIDRUG RESISTANT BACTERIAL INFECTIONS

Different abnormalities related to the immune system and the occurrence of bacterial translocation from the intestinal lumen increase the susceptibility to infections in cirrhotic patients[19]. Bacterial infections are very common in cirrhosis affecting approximately 1/3 of patients with decompensated cirrhosis and are responsible for significant mortality. In a review study Arvaniti *et al*[11] evaluated 178 studies with more than 11000 patients with cirrhosis and found that infections increase mortality four-fold. In these patients delayed antibiotic treatment and inadequate empirical therapy are independently associated



**Table 1** Prevalence and prognosis of multidrug-resistant organisms infections in cirrhosis

Ref.	Study design	n	MDR infection (%)	Mortality (%)	Comments
Piano <i>et al</i> [7]	Single center study	75	35.0	86.0	ACLF grade 2 and 3 were more frequent in MDRO infected patients
Cassini <i>et al</i> [3]	Meta-analysis	671 689	NA	4.9	Estimate the incidence of infections caused by selected antibiotic-resistant bacteria in countries of the EU and EEA in 2015
Trebicka <i>et al</i> [9]	European multicenter	376	18.9	40.8 at 28.0 d; 48.7 at 90.0 d	In infection-induced ACLF, the prevalence of MDR strains was significantly higher; severe sepsis (40.7% <i>vs</i> 21.6%), ACLF (72.3% <i>vs</i> 42.0%) and 90-d mortality (48.7% <i>vs</i> 30.7%) were more frequent in infections caused by MDR strains compared to non-MDR strains
Costabeber <i>et al</i> [17]	Retrospective	474	37.5	-	To evaluate the resistance profile of bacteria isolated from cirrhotic patients admitted to a referral hospital in Brazil
Trebicka <i>et al</i> [9]	European multicenter	520	14.8	35.1 at 28.0 d	MDROs were not significantly different between specific infections in the different European regions; MDROs were more frequently isolated in the ICU (23.8% <i>vs</i> 12.2%) and nosocomial infections (21.3% <i>vs</i> 8.3% and 6.6% in CA and HCA infections, respectively); MDROs were more prevalent in infections causing severe sepsis/shock (30.3% <i>vs</i> 12.2%) or ACLF (20.5% <i>vs</i> 9.4%)
Johnson <i>et al</i> [18]	Retrospective	3951	5.6	27.7	Presence of MDR bacteria in the blood was not associated with in-hospital mortality

ACLF: Acute on chronic liver failure; EU: European Union; EEA: European Economic Area; MDR: Multi-drug resistant; MDRO: Multi drug resistant organism; ICU: Intensive care unit; CA: Community acquired; HCA: Health-care associated; NA: Not available.

with mortality[20,21].

As stated before, recent studies suggest that about 34% of cirrhotic patients with bacterial infections are infected with MDRO[7]. Infections by MDRO have a worse prognosis because they are associated with lower rate of infection resolution with traditional empirical antibiotic treatment. In the first series (2005-2007) of a Spanish study, failure to antibiotic treatment was higher (30% *vs* 8%) in MDR infections than in susceptible bacterial infections. In addition, this study found a higher frequency of septic shock (26% *vs* 10%) and higher hospital mortality rate (25% *vs* 12%) in MDR infections compared to infections by non-resistant bacteria. An important finding of the second series (2010-2011) of this study was the higher prevalence of MDRO in nosocomial infections (39%) compared to HCA (a type of infection that occurs in patients with a previous contact with a healthcare environment, *e.g.*, hospitalization or short-term admission for at least 2 d in the previous 90 d, residence in a nursing home or a long-term care facility, or chronic hemodialysis) and community-acquired infections (20% and 0%, respectively)[22]. Similarly, an intercontinental study evaluated 1302 infected cirrhotic patients and confirmed that infections caused by MDRO were associated with a lower efficacy (40% *vs* 68%) and a longer duration (12 d[7-18] *vs* 10 d[7-15]) of empirical antibiotic treatment. Furthermore, patients with bacterial infections by multi-resistant strains had a higher incidence of septic shock (27% *vs* 13%) and higher in-hospital and 28-d mortality rate (31% *vs* 21% and 34% *vs* 22%, respectively)[7]. Finally, a meta-analysis on the impact of infections by MDRO on mortality in cirrhosis found a four times increased risk of mortality associated with bacterial resistance compared to non-resistant bacterial infections[11].

## RISK FACTORS FOR MULTIDRUG RESISTANT BACTERIAL INFECTIONS

MDR infection results from an interaction of different risk factors that act synergistically. Although some risk factors have been identified, we still far from completely understand all the mechanisms involved, and there is still much to research in this field. Identifying risk factors for multidrug resistance is essential to define the best antibiotic treatment strategy in each case[21].

Previous use of antibiotics is a well-known driver for multidrug resistant infection in different clinical settings. A strong association of MDRO with previous antibiotic therapy was also observed in cirrhotic patients. Extended use of broad-spectrum antibiotics[23] and exposure to systemic antibiotics treatment for at least five days, especially in the previous three months, were highlighted as risk factors[7,24]. Prior use of beta-lactam antibiotics is especially important and has been identified as an independent predictor in the multivariate analysis[22,25]. These findings reinforce the importance of the judicious use of antibiotics in preventing MDRO emergence, with avoidance of overuse and early de-escalation strategies.

Another important risk factor for MDR infections is the previous occurrence of infections by resistant bacteria. In the multivariate analysis, infection caused by MDR bacteria in the last 6 mo increases the risk by 2.45 times[22].

Current or recent contact with the healthcare system is another important risk factor for MDR infections. A strong association between MDR infections and hospital admission has been shown in several studies[10,26,27]. Patients with nosocomial infection, hospitalization for more than 48 h, and those discharged in the last 30 d are at increased risk for MDR infections. In addition, an increased risk has been reported in patients admitted to the intensive care unit. The risk of MDR infections is also related to the duration of hospitalization and the invasiveness of the procedures performed.

Non-hospitalized patients with healthcare-associated (HCA) infections have an intermediate rate (14%-41%) of infections caused by MDRO, which is lower than in nosocomial infections (23%-39%) but higher than that observed in community-acquired infections (0%-16%). Therefore, the risk of MDR infection is directly related to where the infection was acquired (nosocomial or community-acquired)[22,26,28].

Although the association of quinolone prophylaxis with MDR infection was not identified with the use of norfloxacin for six months in a placebo-controlled trial[29] and in an epidemiological study[7], the prophylactic use of antibiotics has been pointed out as a risk factor for MDR infections in other studies [22,25,30].

The bacteriological profile of antibiotic resistance in each geographic region also influences the risk of multidrug resistance. Patients from India, other Asian centers, and South America had an increased risk of MDR bacteria[7]. The bacteriological profile of antibiotic resistance at each health care unit is also an important point, and this local antibiotic resistance profile should be considered in the estimation of the MDR risk of each patient[7,31].

MDR risk is also related to the site of the infection. For example, MDR infections were more commonly observed in patients with pneumonia, skin and soft tissue infections than in those with SBP or spontaneous bacteremia[7,27].

A higher prevalence of proton pump inhibitors (PPI) use among patients with MDR infection has been reported[24], suggesting that PPI could be a risk factor for infections caused by MDRO. However, this association still needs to be better explored in further studies.

MDR infections are more commonly observed in patients with worse liver function, however it is difficult to establish whether liver function is an independent risk factor for MDR because patients with more severe liver disease have more frequent hospitalizations and are more exposed to the use of antibiotics. In the study of Piano *et al*[7], liver function was not independently associated with MDR infections in the multivariate analysis although patients with MDR infections presented higher Child and MELD-Na scores. Therefore, even if it is not an independent risk factor, MDR infection is often associated with more severe liver disease.

Table 2 shows the main risk factors for MDR infections.

## MANAGEMENT

### Empirical antibiotic therapy

In the early phases of bacterial infections in patients with cirrhosis, typical signs of infection (like fever) may not be present. Bacterial infections can precipitate and/or constitute part of the process of acute decompensation of cirrhosis, and an appropriate work-up for infections (*e.g.*, diagnostic paracentesis; chest X-ray; urinalysis; blood, ascites and urine cultures) should be made in all patients hospitalized for decompensated liver disease[32].

Optimizing the prescription of antibiotics is cardinal to effectively treat infections, protect patients from harms caused by unnecessary antibiotic use, and combat antibiotic resistance. As the prevalence of MDRO differs throughout the world, the choice of an empirical antibiotic therapy should be tailored to the local microbiological epidemiology, and it should also be influenced by the type of infection (*e.g.*, SBP, urinary tract infection, pneumonia, soft tissue infection), the severity of infection, and the potential risk factors for infections caused by MDRO[32,33] (Table 1).

Effective antibiotics need to be administered as early as possible. In a retrospective cohort study of 126 cirrhotic patients with SBP-associated septic shock, each hour of delay in the appropriate antimicrobial therapy was associated with 1.86 times increase in hospital mortality[34]. In a worldwide study of hospitalized patients with cirrhosis, the administration of adequate empirical antibiotic treatment was found to be an independent and the only potentially modifiable predictor of in-hospital and 28-d mortality[7].

In the whole Canonic series, MDROs were more prevalent in infections causing severe sepsis/septic shock and/or ACLF and associated to lower resolution rate and higher mortality at 28 d, especially if treated with inadequate empirical antibiotic strategies[10]. A multicenter retrospective study of 865 consecutive patients with a first presentation of SBP in Korea pointed that empirical carbapenem treatment was significantly associated to lower in-hospital mortality than third-generation cephalosporins of among 314 critically ill patients (CLIF-SOFA scores  $\geq 7$ ; 23.1% *vs* 38.8%; aOR, 0.84;

**Table 2 Risk factors for multidrug-resistant infection in cirrhosis**

No.	Risk factors for MDR infection in cirrhosis
1	Prior (3 mo) use of broad-spectrum antibiotics
2	Prior infection by MDROs (6 mo)
3	Nosocomial infection
4	Recent contact with the healthcare system
5	Site of infection (pneumonia, skin, and soft tissue infections)
6	Geographic region
7	Prophylactic use of antibiotics (?)/proton pump inhibitors use?

MDR: Multi-drug resistant; MDRO: Multi drug resistant organism.

95%CI, 0.75-0.94;  $P = 0.002$ )[35].

In the treatment of HCA infections in an Italian population of cirrhotic patients, with a prevalence of MDRO of 40%-46%, empirical broad-spectrum therapy (imipenem/cilastatin  $\pm$  vancomycin) significantly reduced in-hospital mortality when compared to third generation cephalosporins, particularly in patients with sepsis. It also reduced the rate of treatment failure and length of stay[36]. In a prospective randomized controlled trial (RCT) that enrolled 32 patients with nosocomial SBP, the broad-spectrum antibiotic therapy (meropenem plus daptomycin) was more effective than ceftazidime (86.7% vs 25%,  $P < 0.001$ ). Furthermore, meropenem plus daptomycin was effective in 90% of nonresponders to ceftazidime. The response to first-line treatment was an independent predictor of survival[37].

The isolation of MDRO in rectal and nasal swabs could also guide empirical antibiotic strategies in cirrhotic patients. In a study of two European cohorts comprising a total of 907 critically ill patients, including 550 patients with cirrhosis, rectal colonization by MDRO was highly prevalent in cirrhotic patients, ranging from 28.7% to 31.1% in intensive care unit admissions, and MDRO carriage increased the short-term risk of subsequent infection by the colonizing organism[38].

Patients with cirrhosis and severe infections may benefit from therapeutic strategies aimed at optimizing the antibiotics' pharmacokinetic/pharmacodynamic target. The use of high antibiotic doses within the first 48-72 h after the diagnosis of infection and the continuous or extended infusions of beta-lactams are more likely to achieve and to maintain serum drug levels above the minimum inhibitory concentration compared to standard bolus administration. In a secondary analysis of a European prospective multicenter study of patients with cirrhosis and bloodstream infection, the empirical continuous/extended infusion of piperacillin-tazobactam or carbapenems was associated with lower mortality compared to traditional dosing schedules (adjusted hazard ratio, 0.41; 95%CI, 0.110-0.936;  $P = 0.04$ ), and it was particularly useful in those patients who were critically ill[39].

After completing 48-72 h of antibiotic therapy, early de-escalation can be considered based on clinical evolution and available antibiotic susceptibility tests. Short-term treatment is another key measure to prevent antibiotic resistance[7,40,41]. In the 1990s, a randomized controlled trial of 100 patients with SBP showed that a short-course (5-d) treatment is as effective as long-course (10-d) therapy and significantly less expensive[42]. In non-SBP infections, the optimal duration of antibiotic therapy has not been established, but data from the general population suggests that a 7-d course is adequate for most infections[32].

Figure 1 and Table 3 summarizes the management of patients with cirrhosis and bacterial infection.

### Non-antibiotic approach

The emergence and spread of MDRO in cirrhosis require the implementation of measures aimed to prevent its complications. Pharmacological and non-pharmacological strategies are needed, including hand hygiene and barrier precaution, restriction of antibiotic use to high-risk populations, de-escalating antibiotic therapy based on rapid microbiological tests, study of non-antibiotic prophylaxis measures, broad and strict infection control policies, and programs of epidemiological surveillance[25].

**Non-antibiotic drugs with potential benefit on infections in cirrhosis:** Although antibiotic treatment is the most effective measure for controlling established MDR infection, other drugs have shown potential benefits in preventing infections in cirrhosis. A lower occurrence of SBP has been demonstrated in patients using non-selective beta-blockers (NSBB)[43]. This benefit of NSBB has been related to its potential effect in improving intestinal motility, improving intestinal permeability, and reducing bacterial translocation[44,45]. Statins also seem to have a beneficial effect against bacterial infection[46] that is attributed to its anti-inflammatory and immunomodulatory properties.


**Table 3 Recommended empirical antibiotic treatment for bacterial infection in cirrhosis**

Type of infection	Community-acquired infection	Nosocomial and HCA infection or sepsis
SBP, spontaneous bacterial empyema and spontaneous bacteremia	Cefotaxime or Amoxicillin/clavulanic acid	Piperacillin/tazobactam or Meropenem ± Vancomycin or Daptomycin or Linezolid <sup>1</sup>
UTI	Fosfomycin or cotrimoxazole	Uncomplicated: Nitrofurantoin or Fosfomycin; if sepsis: Piperacillin/tazobactam or Meropenem ± Glycopeptide
Pneumonia	Amoxicillin/clavulanic acid; Ceftriaxone + Macrolide; Levofloxacin; Moxifloxacin	Piperacillin/tazobactam or Meropenem or Ceftazidime + Ciprofloxacin; Glycopeptides or Linezolid <sup>1</sup> should be added in patients with risk factors for MRSA <sup>2</sup>
Skin and soft tissue infections	Amoxicillin/clavulanic acid or ± Clindamycin	Meropenem or Piperacillin/tazobactam + Glycopeptide or Daptomycin or Linezolid <sup>1</sup> ± Clindamycin; if necrotizing fascitis: Meropenem + Daptomycin + Clindamycin

<sup>1</sup>Vancomycin, teicoplanin or daptomycin in areas with a high prevalence Methicillin-resistant *Staphylococcus aureus* and vancomycin-susceptible enterococci. Vancomycin must be replaced by linezolid in areas with a high prevalence of vancomycin-resistant enterococci.

<sup>2</sup>Ventilator-associated pneumonia, previous antibiotic therapy, nasal Methicillin-resistant *Staphylococcus aureus* carriage.

HCA: Health care associated; MRSA: Methicillin-resistant *Staphylococcus aureus*; SBP: Spontaneous bacterial peritonitis; UTI: Urinary tract infection.



DOI: 10.12998/wjcc.v11.i3.534 Copyright ©The Author(s) 2023.

**Figure 1 Suggested algorithm for the management of patients with cirrhosis and bacterial infections.** MDR: Multidrug-resistant.

**Non-pharmacological measures:** Non-pharmacological measures are based on preventive strategies and procedures focusing on intestinal colonization with MDRO. The most important preventive measure is the restrictive and judicious antibiotic use since the main driver for the emergence of MDR infections is the widespread use of antibiotics.

The main non-pharmacological measures focus on gut microbiota[27]. A healthy microbiome is essential to prevent colonization and infection by MDRO[25]. Different approaches focusing on the modulation of the intestinal microbiome were studied, such as probiotics, prebiotics/synbiotics dietary regimens, and fecal microbiota transplant (FMT)[45].

Although some studies have shown favorable results with probiotics, there are also negative studies. This controversy is probably related to different probiotics used and the different number and concentration of the species. Further studies are necessary to define the ideal combination, dose, and duration of administration.

FMT involves the safe transfer of exogenous bacterial flora from a healthy donor to another patient, in capsule or liquid formulations. The rationale for using this technique is the central role of the gastrointestinal colonization in the development of MDR infections. It has been demonstrated that colonization by MDRO is associated with increased risk of infection by the colonizing bacteria in the

short-term[38]. FMT has the potential effect of promoting MDRO decolonization.

A systematic review with meta-analysis[47] of five studies, with a total number of 52 patients, evaluated whether FMT decolonizes antibiotic-resistant bacteria from the gut of colonized adults. Evidence from this meta-analysis indicates a potential benefit of FMT as a decolonization intervention, with few adverse effects. Despite the low quality of evidence appointed by this meta-analysis, these preliminary results suggest that FMT is a promising approach that deserves further analysis in RCTs.

In cirrhotic patients, a preliminary study of FMT in patients with advanced cirrhosis on lactulose and rifaximin demonstrated that FMT restored antibiotic-associated disruption in microbial diversity and function[48]. The impact of FMT in the reduction of gut microbial antibiotic resistance genes was later reported in two trials: a capsule FMT trial and an enema FMT trial with 20 patients each. This study demonstrated that, despite differences in routes of administration, antibiotic resistance gene abundance was reduced after FMT compared to pre-FMT baseline and non-FMT groups in decompensated cirrhosis[49].

Another possible approach, although still requiring further studies, is phage therapy. This technique is based on the use of bacteriophages which are viruses that infect bacteria[50]. The bacteriophages replicate inside the bacteria leading to their destruction. It is an old technique that was left aside with the advent of antibiotics, but nowadays it has been considered again as a therapeutic option to face the serious problem of antibiotic multidrug resistance[51,52]. Although specific studies in cirrhotic patients are not available, phage therapy represents a possible future alternative therapy for controlling MDRO [53].

## CONCLUSION

In this review, we describe the epidemiology, clinical settings and the current evidence-based strategies for early recognition and treatment alternatives for MDR infection in cirrhotic patients.

## FOOTNOTES

**Author contributions:** Terra C, de Mattos ÂZ, Chagas MS, Torres A, Wiltgen D, Souza BM, and Perez RM performed the research, analyzed the data, and wrote the manuscript; all authors have read and approve the final manuscript.

**Conflict-of-interest statement:** All authors report no relevant conflicts of interest for this article.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** Brazil

**ORCID number:** Carlos Terra 0000-0003-3069-128X; Ângelo Zambam de Mattos 0000-0002-3063-0199; Marcelo Souza Chagas 0000-0002-1418-1696; Andre Torres 0000-0002-4994-7137; Denusa Wiltgen 0000-0002-7334-2779; Barbara Muniz Souza 0000-0001-7061-1788; Renata Mello Perez 0000-0002-3367-9991.

**S-Editor:** Chen YL

**L-Editor:** A

**P-Editor:** Chen YL

## REFERENCES

- 1 **Magiorakos AP**, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth S, Hindler JF, Kahlmeter G, Olsson-Liljequist B, Paterson DL, Rice LB, Stelling J, Struelens MJ, Vatopoulos A, Weber JT, Monnet DL. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012; **18**: 268-281 [PMID: 21793988 DOI: 10.1111/j.1469-0691.2011.03570.x]
- 2 **Kadri SS**. Key Takeaways From the U.S. CDC's 2019 Antibiotic Resistance Threats Report for Frontline Providers. *Crit Care Med* 2020; **48**: 939-945 [PMID: 32282351 DOI: 10.1097/CCM.0000000000004371]
- 3 **Cassini A**, Högberg LD, Plachouras D, Quattrocchi A, Hoxha A, Simonsen GS, Colomb-Cotinat M, Kretzschmar ME, Devleeschauwer B, Cecchini M, Ouakrim DA, Oliveira TC, Struelens MJ, Suetens C, Monnet DL; Burden of AMR Collaborative Group. Attributable deaths and disability-adjusted life-years caused by infections with antibiotic-resistant bacteria in the EU and the European Economic Area in 2015: a population-level modelling analysis. *Lancet Infect Dis*



- 2019; **19**: 56-66 [PMID: [30409683](#) DOI: [10.1016/S1473-3099\(18\)30605-4](#)]
- 4 **Weist K**, Högberg LD. ECDC publishes 2015 surveillance data on antimicrobial resistance and antimicrobial consumption in Europe. *Euro Surveill* 2016; **21** [PMID: [27918266](#) DOI: [10.2807/1560-7917.ES.2016.21.46.30399](#)]
- 5 **Serra-Burriel M**, Keys M, Campillo-Artero C, Agodi A, Barchitta M, Gikas A, Palos C, López-Casasnovas G. Impact of multi-drug resistant bacteria on economic and clinical outcomes of healthcare-associated infections in adults: Systematic review and meta-analysis. *PLoS One* 2020; **15**: e0227139 [PMID: [31923281](#) DOI: [10.1371/journal.pone.0227139](#)]
- 6 **Fernández J**, Piano S, Bartoletti M, Wey EQ. Management of bacterial and fungal infections in cirrhosis: The MDRO challenge. *J Hepatol* 2021; **75** Suppl 1: S101-S117 [PMID: [34039482](#) DOI: [10.1016/j.jhep.2020.11.010](#)]
- 7 **Piano S**, Singh V, Caraceni P, Maiwall R, Alessandria C, Fernandez J, Soares EC, Kim DJ, Kim SE, Marino M, Vorobioff J, Barea RCR, Merli M, Elkrief L, Vargas V, Krag A, Singh SP, Lesmana LA, Toledo C, Marciano S, Verhelst X, Wong F, Intagliata N, Rabinowich L, Colombato L, Kim SG, Gerbes A, Durand F, Roblero JP, Bhamidimarri KR, Boyer TD, Maevskaya M, Fassio E, Kim HS, Hwang JS, Gines P, Gadano A, Sarin SK, Angeli P; International Club of Ascites Global Study Group. Epidemiology and Effects of Bacterial Infections in Patients With Cirrhosis Worldwide. *Gastroenterology* 2019; **156**: 1368-1380.e10 [PMID: [30552895](#) DOI: [10.1053/j.gastro.2018.12.005](#)]
- 8 **Fischer P**, Pandrea S, Dan Grigorescu M, Stefanescu H, Tefas C, Hadade A, Procopet B, Ionescu D. The threat of carbapenem resistance in Eastern Europe in patients with decompensated cirrhosis admitted to intensive care unit. *Dig Liver Dis* 2022; **54**: 1385-1391 [PMID: [35732546](#) DOI: [10.1016/j.dld.2022.05.011](#)]
- 9 **Trebicka J**, Fernandez J, Arroyo V; PREDICT STUDY group of the EASL-CLIF CONSORTIUM. Reply to: Correspondence on 'The PREDICT study uncovers three clinical courses of acutely decompensated cirrhosis that have distinct pathophysiology'. *J Hepatol* 2021; **74**: 480-481 [PMID: [33279257](#) DOI: [10.1016/j.jhep.2020.11.019](#)]
- 10 **Fernández J**, Prado V, Trebicka J, Amoros A, Gustot T, Wiest R, Deulofeu C, Garcia E, Acevedo J, Fuhrmann V, Durand F, Sánchez C, Papp M, Caraceni P, Vargas V, Bañares R, Piano S, Janicko M, Albillos A, Alessandria C, Soriano G, Welzel TM, Laleman W, Gerbes A, De Gottardi A, Merli M, Coenraad M, Saliba F, Pavesi M, Jalan R, Ginès P, Angeli P, Arroyo V; European Foundation for the Study of Chronic Liver Failure (EF-Clif). Multidrug-resistant bacterial infections in patients with decompensated cirrhosis and with acute-on-chronic liver failure in Europe. *J Hepatol* 2019; **70**: 398-411 [PMID: [30391380](#) DOI: [10.1016/j.jhep.2018.10.027](#)]
- 11 **Arvaniti V**, D'Amico G, Fede G, Manousou P, Tsochatzis E, Pleguezuelo M, Burroughs AK. Infections in patients with cirrhosis increase mortality four-fold and should be used in determining prognosis. *Gastroenterology* 2010; **139**: 1246-1256.e1 [PMID: [20558165](#) DOI: [10.1053/j.gastro.2010.06.019](#)]
- 12 **D'Amico G**, Morabito A, D'Amico M, Pasta L, Malizia G, Rebora P, Valsecchi MG. Clinical states of cirrhosis and competing risks. *J Hepatol* 2018; **68**: 563-576 [PMID: [29111320](#) DOI: [10.1016/j.jhep.2017.10.020](#)]
- 13 **Moreau R**, Jalan R, Gines P, Pavesi M, Angeli P, Cordoba J, Durand F, Gustot T, Saliba F, Domenicali M, Gerbes A, Wendon J, Alessandria C, Laleman W, Zeuzem S, Trebicka J, Bernardi M, Arroyo V; CANONIC Study Investigators of the EASL-CLIF Consortium. Acute-on-chronic liver failure is a distinct syndrome that develops in patients with acute decompensation of cirrhosis. *Gastroenterology* 2013; **144**: 1426-1437, 1437.e1 [PMID: [23474284](#) DOI: [10.1053/j.gastro.2013.02.042](#)]
- 14 **O'Leary JG**, Reddy KR, Garcia-Tsao G, Biggins SW, Wong F, Fallon MB, Subramanian RM, Kamath PS, Thuluvath P, Vargas HE, Maliakkal B, Tandon P, Lai J, Thacker LR, Bajaj JS. NACSELD acute-on-chronic liver failure (NACSELD-ACLF) score predicts 30-day survival in hospitalized patients with cirrhosis. *Hepatology* 2018; **67**: 2367-2374 [PMID: [29315693](#) DOI: [10.1002/hep.29773](#)]
- 15 **Coral G**, de Mattos AA, Damo DF, Viégas AC. [Prevalence and prognosis of spontaneous bacterial peritonitis. Experience in patients from a general hospital in Porto Alegre, RS, Brazil (1991-2000)]. *Arq Gastroenterol* 2002; **39**: 158-162 [PMID: [12778307](#) DOI: [10.1590/s0004-28032002000300005](#)]
- 16 **Musskopf MI**, Fonseca FP, Gass J, de Mattos AZ, John JA, de Mello Brandão AB. Prognostic factors associated with in-hospital mortality in patients with spontaneous bacterial peritonitis. *Ann Hepatol* 2012; **11**: 915-920 [PMID: [23109456](#)]
- 17 **Costabeber AM**, Mattos AA, Sukiennik TC. Prevalence of bacterial resistance in hospitalized Cirrhotic patients in southern brazil: A new challenge. *Rev Inst Med Trop Sao Paulo* 2016; **58**: 36 [PMID: [27253738](#) DOI: [10.1590/S1678-9946201658036](#)]
- 18 **Johnson AL**, Ratnasekera IU, Irvine KM, Henderson A, Powell EE, Valery PC. Bacteraemia, sepsis and antibiotic resistance in Australian patients with cirrhosis: a population-based study. *BMJ Open Gastroenterol* 2021; **8** [PMID: [34876410](#) DOI: [10.1136/bmjgast-2021-000695](#)]
- 19 **Thulstrup AM**, Sørensen HT, Schönheyder HC, Møller JK, Tage-Jensen U. Population-based study of the risk and short-term prognosis for bacteremia in patients with liver cirrhosis. *Clin Infect Dis* 2000; **31**: 1357-1361 [PMID: [11096002](#) DOI: [10.1086/317494](#)]
- 20 **Bartoletti M**, Giannella M, Lewis R, Caraceni P, Tedeschi S, Paul M, Schramm C, Bruns T, Merli M, Cobos-Trigueros N, Seminari E, Retamar P, Muñoz P, Tumbarello M, Burra P, Torrani Cerenzia M, Barsic B, Calbo E, Maraolo AE, Petrosillo N, Galan-Ladero MA, D'Offizi G, Bar Sinai N, Rodríguez-Baño J, Verucchi G, Bernardi M, Viale P; ESGIBIS/BICHROME Study Group. A prospective multicentre study of the epidemiology and outcomes of bloodstream infection in cirrhotic patients. *Clin Microbiol Infect* 2018; **24**: 546.e1-546.e8 [PMID: [28818628](#) DOI: [10.1016/j.cmi.2017.08.001](#)]
- 21 **Arabi YM**, Dara SI, Memish Z, Al Abdulkareem A, Tamim HM, Al-Shirawi N, Parrillo JE, Dodek P, Lapinsky S, Feinstein D, Wood G, Dial S, Zanotti S, Kumar A; Cooperative Antimicrobial Therapy of Septic Shock (CATSS) Database Research Group. Antimicrobial therapeutic determinants of outcomes from septic shock among patients with cirrhosis. *Hepatology* 2012; **56**: 2305-2315 [PMID: [22753144](#) DOI: [10.1002/hep.25931](#)]
- 22 **Fernández J**, Acevedo J, Castro M, Garcia O, de Lope CR, Roca D, Pavesi M, Sola E, Moreira L, Silva A, Seva-Pereira T, Corradi F, Mensa J, Ginès P, Arroyo V. Prevalence and risk factors of infections by multiresistant bacteria in cirrhosis: a prospective study. *Hepatology* 2012; **55**: 1551-1561 [PMID: [22183941](#) DOI: [10.1002/hep.25532](#)]
- 23 **Zhong L**, Men TY, Li H, Peng ZH, Gu Y, Ding X, Xing TH, Fan JW. Multidrug-resistant gram-negative bacterial infections after liver transplantation - spectrum and risk factors. *J Infect* 2012; **64**: 299-310 [PMID: [22198738](#) DOI: [10.1016/j.jinf.2011.12.005](#)]

- 24 **Figueiredo LM**, Rafael MA, Alexandrino G, Branco JC, Carvalho R, Costa MN, Martins A. Risk factors for the emergence of multidrug-resistant organisms in liver cirrhosis. *Gastroenterol Hepatol* 2022; **45**: 186-191 [PMID: [34052400](#) DOI: [10.1016/j.gastrohep.2021.04.006](#)]
- 25 **Fernández J**, Bert F, Nicolas-Chanoine MH. The challenges of multi-drug-resistance in hepatology. *J Hepatol* 2016; **65**: 1043-1054 [PMID: [27544545](#) DOI: [10.1016/j.jhep.2016.08.006](#)]
- 26 **Jalan R**, Fernandez J, Wiest R, Schnabl B, Moreau R, Angeli P, Stadlbauer V, Gustot T, Bernardi M, Canton R, Albillos A, Lammert F, Wilmer A, Mookerjee R, Vila J, Garcia-Martinez R, Wendon J, Such J, Cordoba J, Sanyal A, Garcia-Tsao G, Arroyo V, Burroughs A, Ginès P. Bacterial infections in cirrhosis: a position statement based on the EASL Special Conference 2013. *J Hepatol* 2014; **60**: 1310-1324 [PMID: [24530646](#) DOI: [10.1016/j.jhep.2014.01.024](#)]
- 27 **Gallagher CE**, Shawcross DL. Management of Multidrug-Resistant Infections in Cirrhosis. *Semin Liver Dis* 2022; **42**: 173-187 [PMID: [35130574](#) DOI: [10.1055/a-1765-0056](#)]
- 28 **Merli M**, Lucidi C, Giannelli V, Giusto M, Riggio O, Falcone M, Ridola L, Attili AF, Venditti M. Cirrhotic patients are at risk for health care-associated bacterial infections. *Clin Gastroenterol Hepatol* 2010; **8**: 979-985 [PMID: [20621200](#) DOI: [10.1016/j.cgh.2010.06.024](#)]
- 29 **Moreau R**, Elkrief L, Bureau C, Perarnau JM, Thévenot T, Saliba F, Louvet A, Nahon P, Lannes A, Anty R, Hillaire S, Pasquet B, Ozenne V, Rudler M, Ollivier-Hourmand I, Robic MA, d'Alteroche L, Di Martino V, Ripault MP, Pauwels A, Grangé JD, Carbonell N, Bronowicki JP, Payancé A, Rautou PE, Valla D, Gault N, Lebrech D; NORFLOCIR Trial Investigators. Effects of Long-term Norfloxacin Therapy in Patients With Advanced Cirrhosis. *Gastroenterology* 2018; **155**: 1816-1827.e9 [PMID: [30144431](#) DOI: [10.1053/j.gastro.2018.08.026](#)]
- 30 **Salerno F**, Borzio M, Pedicino C, Simonetti R, Rossini A, Boccia S, Cacciola I, Burroughs AK, Manini MA, La Mura V, Angeli P, Bernardi M, Dalla Gasperina D, Dionigi E, Dibenedetto C, Arghittu M; AISF Investigators. The impact of infection by multidrug-resistant agents in patients with cirrhosis. A multicenter prospective study. *Liver Int* 2017; **37**: 71-79 [PMID: [27364035](#) DOI: [10.1111/Liv.13195](#)]
- 31 **Aguirre-García J**. Comment on: Ruelas-Villavicencio A L, et. al. "In whom, how and how often is surveillance for hepatocellular carcinoma cost-effective? *Ann Hepatol* 2005; **4**: 289 [PMID: [16432498](#) DOI: [10.1016/j.aohep.2022.100719](#)]
- 32 **Piano S**, Tonon M, Angeli P. Changes in the epidemiology and management of bacterial infections in cirrhosis. *Clin Mol Hepatol* 2021; **27**: 437-445 [PMID: [33504138](#) DOI: [10.3350/cmh.2020.0329](#)]
- 33 **Miranda-Zazueta G**, León-Garduño LAP, Aguirre-Valadez J, Torre-Delgadillo A. Bacterial infections in cirrhosis: Current treatment. *Ann Hepatol* 2020; **19**: 238-244 [PMID: [32317149](#) DOI: [10.1016/j.aohep.2019.09.011](#)]
- 34 **Karvellas CJ**, Abalde JG, Arabi YM, Kumar A; Cooperative Antimicrobial Therapy of Septic Shock (CATSS) Database Research Group. Appropriate and timely antimicrobial therapy in cirrhotic patients with spontaneous bacterial peritonitis-associated septic shock: a retrospective cohort study. *Aliment Pharmacol Ther* 2015; **41**: 747-757 [PMID: [25703246](#) DOI: [10.1111/apt.13135](#)]
- 35 **Kim SW**, Yoon JS, Park J, Jung YJ, Lee JS, Song J, Lee HA, Seo YS, Lee M, Park JM, Choi DH, Kim MY, Kang SH, Yang JM, Song DS, Chung SW, Kim MA, Jang HJ, Oh H, Lee CH, Lee YB, Cho EJ, Yu SJ, Kim YJ, Yoon JH, Lee JH. Empirical Treatment With Carbapenem vs Third-generation Cephalosporin for Treatment of Spontaneous Bacterial Peritonitis. *Clin Gastroenterol Hepatol* 2021; **19**: 976-986.e5 [PMID: [32623007](#) DOI: [10.1016/j.cgh.2020.06.046](#)]
- 36 **Merli M**, Lucidi C, Di Gregorio V, Lattanzi B, Giannelli V, Giusto M, Farcomeni A, Ceccarelli G, Falcone M, Riggio O, Venditti M. An empirical broad spectrum antibiotic therapy in health-care-associated infections improves survival in patients with cirrhosis: A randomized trial. *Hepatology* 2016; **63**: 1632-1639 [PMID: [26529126](#) DOI: [10.1002/hep.28332](#)]
- 37 **Piano S**, Fasolato S, Salinas F, Romano A, Tonon M, Morando F, Cavallin M, Gola E, Sticca A, Loregian A, Palù G, Zanusi G, Senzolo M, Burra P, Cillo U, Angeli P. The empirical antibiotic treatment of nosocomial spontaneous bacterial peritonitis: Results of a randomized, controlled clinical trial. *Hepatology* 2016; **63**: 1299-1309 [PMID: [26084406](#) DOI: [10.1002/hep.27941](#)]
- 38 **Prado V**, Hernández-Tejero M, Mücke MM, Marco F, Gu W, Amoros A, Toapanta D, Reverter E, Peña-Ramírez C, Altenpeter L, Bassegoda O, Mezzano G, Aziz F, Juanola A, Rodríguez-Tajes S, Chamorro V, López D, Reyes M, Hogardt M, Kempf VAJ, Ferstl PG, Zeuzem S, Martínez JA, Vila J, Arroyo V, Trebicka J, Fernandez J. Rectal colonization by resistant bacteria increases the risk of infection by the colonizing strain in critically ill patients with cirrhosis. *J Hepatol* 2022; **76**: 1079-1089 [PMID: [35074475](#) DOI: [10.1016/j.jhep.2021.12.042](#)]
- 39 **Bartoletti M**, Giannella M, Lewis RE, Caraceni P, Tedeschi S, Paul M, Schramm C, Bruns T, Merli M, Cobos-Trigueros N, Seminari E, Retamar P, Muñoz P, Tumbarello M, Burra P, Torrani Cerenza M, Barsic B, Calbo E, Maraolo AE, Petrosillo N, Galan-Ladero MA, D'Offizi G, Zak-Doron Y, Rodríguez-Baño J, Baldassarre M, Verucchi G, Domenicali M, Bernardi M, Viale P; ESGIB/BICHROME study group. Extended Infusion of  $\beta$ -Lactams for Bloodstream Infection in Patients With Liver Cirrhosis: An Observational Multicenter Study. *Clin Infect Dis* 2019; **69**: 1731-1739 [PMID: [30649218](#) DOI: [10.1093/cid/ciz032](#)]
- 40 **Fernández J**, Tandon P, Mensa J, Garcia-Tsao G. Antibiotic prophylaxis in cirrhosis: Good and bad. *Hepatology* 2016; **63**: 2019-2031 [PMID: [26528864](#) DOI: [10.1002/hep.28330](#)]
- 41 **Allaire M**, Cadranet JF, Nguyen TTN, Garioud A, Zougmore H, Heng R, Perignon C, Ollivier-Hourmand I, Dao T. Management of infections in patients with cirrhosis in the context of increasing therapeutic resistance: A systematic review. *Clin Res Hepatol Gastroenterol* 2020; **44**: 264-274 [PMID: [31706985](#) DOI: [10.1016/j.clinre.2019.10.003](#)]
- 42 **Runyon BA**, McHutchison JG, Antillon MR, Akriviadis EA, Montano AA. Short-course versus long-course antibiotic treatment of spontaneous bacterial peritonitis. A randomized controlled study of 100 patients. *Gastroenterology* 1991; **100**: 1737-1742 [PMID: [2019378](#) DOI: [10.1016/0016-5085\(91\)90677-d](#)]
- 43 **Senzolo M**, Cholongitas E, Burra P, Leandro G, Thalheimer U, Patch D, Burroughs AK. beta-Blockers protect against spontaneous bacterial peritonitis in cirrhotic patients: a meta-analysis. *Liver Int* 2009; **29**: 1189-1193 [PMID: [19508620](#) DOI: [10.1111/j.1478-3231.2009.02038.x](#)]
- 44 **Reiberger T**, Ferlitsch A, Payer BA, Mandorfer M, Heinisch BB, Hayden H, Lammert F, Trauner M, Peck-Radosavljevic M, Vogelsang H; Vienna Hepatic Hemodynamic Lab. Non-selective betablocker therapy decreases intestinal permeability and serum levels of LBP and IL-6 in patients with cirrhosis. *J Hepatol* 2013; **58**: 911-921 [PMID: [23262249](#) DOI: [10.1016/j.jhep.2013.03.011](#)]



- 10.1016/j.jhep.2012.12.011]
- 45 **Yan K**, Garcia-Tsao G. Novel prevention strategies for bacterial infections in cirrhosis. *Expert Opin Pharmacother* 2016; **17**: 689-701 [PMID: [26799197](#) DOI: [10.1517/14656566.2016.1145663](#)]
- 46 **Motzkus-Feagans C**, Pakyz AL, Ratliff SM, Bajaj JS, Lapane KL. Statin use and infections in Veterans with cirrhosis. *Aliment Pharmacol Ther* 2013; **38**: 611-618 [PMID: [23889738](#) DOI: [10.1111/apt.12430](#)]
- 47 **Tavoukjian V**. Faecal microbiota transplantation for the decolonization of antibiotic-resistant bacteria in the gut: a systematic review and meta-analysis. *J Hosp Infect* 2019; **102**: 174-188 [PMID: [30926290](#) DOI: [10.1016/j.jhin.2019.03.010](#)]
- 48 **Bajaj JS**, Kakiyama G, Savidge T, Takei H, Kassam ZA, Fagan A, Gavis EA, Pandak WM, Nittono H, Hylemon PB, Boonma P, Haag A, Heuman DM, Fuchs M, John B, Sikaroodi M, Gillevet PM. Antibiotic-Associated Disruption of Microbiota Composition and Function in Cirrhosis Is Restored by Fecal Transplant. *Hepatology* 2018; **68**: 1549-1558 [PMID: [29665102](#) DOI: [10.1002/hep.30037](#)]
- 49 **Bajaj JS**, Shamsaddini A, Fagan A, Sterling RK, Gavis E, Khoruts A, Fuchs M, Lee H, Sikaroodi M, Gillevet PM. Fecal Microbiota Transplant in Cirrhosis Reduces Gut Microbial Antibiotic Resistance Genes: Analysis of Two Trials. *Hepatol Commun* 2021; **5**: 258-271 [PMID: [33553973](#) DOI: [10.1002/hep4.1639](#)]
- 50 **Principi N**, Silvestri E, Esposito S. Advantages and Limitations of Bacteriophages for the Treatment of Bacterial Infections. *Front Pharmacol* 2019; **10**: 513 [PMID: [31139086](#) DOI: [10.3389/fphar.2019.00513](#)]
- 51 **Golkar Z**, Bagasra O, Pace DG. Bacteriophage therapy: a potential solution for the antibiotic resistance crisis. *J Infect Dev Ctries* 2014; **8**: 129-136 [PMID: [24518621](#) DOI: [10.3855/jidc.3573](#)]
- 52 **Kortright KE**, Chan BK, Koff JL, Turner PE. Phage Therapy: A Renewed Approach to Combat Antibiotic-Resistant Bacteria. *Cell Host Microbe* 2019; **25**: 219-232 [PMID: [30763536](#) DOI: [10.1016/j.chom.2019.01.014](#)]
- 53 **Fauconner A**, Nagel TE, Fauconner C, Verbeken G, De Vos D, Merabishvili M, Pirnay JP. The Unique Role That WHO Could Play in Implementing Phage Therapy to Combat the Global Antibiotic Resistance Crisis. *Front Microbiol* 2020; **11**: 1982 [PMID: [33013742](#) DOI: [10.3389/fmicb.2020.01982](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

