

World Journal of *Gastroenterology*

World J Gastroenterol 2023 April 14; 29(14): 2064-2221



REVIEW

- 2064 Insight into the liver dysfunction in COVID-19 patients: Molecular mechanisms and possible therapeutic strategies
Khullar N, Bhatti JS, Singh S, Thukral B, Reddy PH, Bhatti GK
- 2078 Role of prebiotics, probiotics, and synbiotics in management of inflammatory bowel disease: Current perspectives
Roy S, Dhaneshwar S
- 2101 Changing trends in the minimally invasive surgery for chronic pancreatitis
Kalayarasan R, Shukla A
- 2114 Challenges in pediatric inherited/metabolic liver disease: Focus on the disease spectrum, diagnosis and management of relatively common disorders
Zou YG, Wang H, Li WW, Dai DL

MINIREVIEWS

- 2127 COVID-19 related biliary injury: A review of recent literature
Yadlapati S, Jarrett SA, Baik D, Chaaya A

ORIGINAL ARTICLE**Basic Study**

- 2134 Anti-inflammatory effect and antihepatoma mechanism of carrimycin
Li XY, Luo YT, Wang YH, Yang ZX, Shang YZ, Guan QX
- 2153 Effect and mechanism of reactive oxygen species-mediated NOD-like receptor family pyrin domain-containing 3 inflammasome activation in hepatic alveolar echinococcosis
Chen CS, Zhang YG, Wang HJ, Fan HN
- 2172 Changes in the gut mycobiome in pediatric patients in relation to the clinical activity of Crohn's disease
Krawczyk A, Salamon D, Kowalska-Duplaga K, Zapala B, Książek T, Drażniuk-Warchol M, Gosiewski T

Retrospective Cohort Study

- 2188 Acoustic radiation force impulse predicts long-term outcomes in a large-scale cohort: High liver cancer, low comorbidity in hepatitis B virus
Tai J, Harrison AP, Chen HM, Hsu CY, Hsu TH, Chen CJ, Jeng WJ, Chang ML, Lu L, Tai DI

SYSTEMATIC REVIEWS

- 2202 Effectiveness of *Helicobacter pylori* eradication in the treatment of early-stage gastric mucosa-associated lymphoid tissue lymphoma: An up-to-date meta-analysis

Lemos FFB, Castro CT, Calmon MS, Silva Luz M, Pinheiro SLR, Faria Souza Mendes dos Santos C, Correa Santos GL, Marques HS, Delgado HA, Teixeira KN, Souza CL, Oliveira MV, Freire de Melo F

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Masataka Kikuyama, MD, PhD, Associate Professor, Department of Gastroenterology, Tokyo Women's Medical University, 8-1 Kawadacho, Shinjuku-ku, Tokyo 162-8666, Japan. kikuyama.masataka@twmu.ac.jp

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports, Index Medicus, MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJG as 5.374; IF without journal self cites: 5.187; 5-year IF: 5.715; Journal Citation Indicator: 0.84; Ranking: 31 among 93 journals in gastroenterology and hepatology; and Quartile category: Q2. The WJG's CiteScore for 2021 is 8.1 and Scopus CiteScore rank 2021: Gastroenterology is 18/149.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Hua-Ge Yu*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jia-Ru Fan*.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

April 14, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Changing trends in the minimally invasive surgery for chronic pancreatitis

Raja Kalayarasan, Ankit Shukla

Specialty type: Surgery

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): A
Grade B (Very good): B, B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Hassan MA, Pakistan;
Liu H, China; Wang ZX, China

Received: December 17, 2022

Peer-review started: December 17, 2022

First decision: February 8, 2023

Revised: February 21, 2023

Accepted: March 23, 2023

Article in press: March 23, 2023

Published online: April 14, 2023



Raja Kalayarasan, Ankit Shukla, Surgical Gastroenterology, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Puducherry 605006, India

Corresponding author: Raja Kalayarasan, DNB, FRCS (Ed), MCh, MS, Professor, Surgical Gastroenterology, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Dhanvantari Nagar, Gorimedu, Puducherry 605006, India.

kalayarasanraja@yahoo.com

Abstract

Chronic pancreatitis is a debilitating pancreatic inflammatory disease characterized by intractable pain resulting in poor quality of life. Conventional management of pancreatic pain consists of a step-up approach with medications and lifestyle modifications followed by endoscopic intervention. Traditionally surgery is reserved for patients who do not improve with other interventions. However, recent studies suggest that early surgical intervention is more beneficial as it can mitigate the progression of the pathological process and prevent loss of pancreatic function. Despite the widespread adoption of minimally invasive approaches in various gastrointestinal surgical disorders, minimally invasive surgery for chronic pancreatitis is slow to evolve. Technical difficulty due to severe inflammatory changes has been the major impediment to the widespread usage of minimally invasive surgery in chronic pancreatitis. With this background, the present review aimed to critically analyze the available evidence on the minimally invasive treatment of chronic pancreatitis. A Pub Med search of all relevant articles was performed using the appropriate keywords, parentheses, and Boolean operators. Most initial laparoscopic series have reported the feasibility of lateral pancreaticojejunostomy, considered an adequate procedure only in a small proportion of patients. The pancreatic head is the pacemaker of pain, so adequate decompression is critical for long-term pain relief. Recent studies have documented the feasibility of minimally invasive duodenum-preserving pancreatic head resection. With improvements in laparoscopic instrumentation and technological advances, minimally invasive surgery for chronic pancreatitis is gaining momentum. However, more high-quality evidence is required to document the superiority of minimally invasive surgery for chronic pancreatitis.

Key Words: Robotics; Laparoscopy; Surgery; Chronic pancreatitis; Pancreas; Pancreatitis

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Pain in chronic pancreatitis is a significant symptom that demands utmost attention as it compromises the quality of life and inherently risks narcotic addiction. Surgical management for chronic pancreatitis began with various open surgical drainage and resection procedures. Since pain is the primary indication for intervention, a minimally invasive approach is an attractive proposition in chronic pancreatitis. Despite the slow adoption of laparoscopic and robotic surgery in chronic pancreatitis, safety, and feasibility have been documented in recent studies. The challenges and limitations highlighted in the present review could guide future research on minimally invasive surgery in chronic pancreatitis.

Citation: Kalayarasan R, Shukla A. Changing trends in the minimally invasive surgery for chronic pancreatitis. *World J Gastroenterol* 2023; 29(14): 2101-2113

URL: <https://www.wjgnet.com/1007-9327/full/v29/i14/2101.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v29.i14.2101>

INTRODUCTION

Chronic pancreatitis is a progressive pancreatic inflammatory disease that leads to fibrosis and parenchymal tissue loss resulting in impaired endocrine and exocrine function. While consumption of alcohol is the leading cause of chronic pancreatitis worldwide, idiopathic pancreatitis remains common in India and China[1,2]. The most common and dominant symptom of chronic pancreatitis is pain, which can be persistent, severe, or recurrent episodes with pain-free intervals significantly impacting the quality of life[2,3].

Traditionally, pain in chronic pancreatitis is managed initially with analgesics, pancreatic enzyme replacement therapy, and lifestyle modifications followed by endoscopic or surgical intervention. Almost half of the patients who do not respond to medical or endoscopic management are referred for surgical intervention[4]. The most common indication for surgery is intractable pain. Studies have reported that early surgical management has better outcomes than intervention in the advanced disease stage[3,5]. While the minimally invasive approach is widely used for various gastrointestinal and pancreatobiliary disorders, its application in chronic pancreatitis is disproportionately low. As chronic pancreatitis is a benign disorder with pain as the primary indication for intervention, a minimally invasive approach is an attractive proposition. The technical difficulty, combined with the potential for vascular injury and bleeding associated with pancreatic inflammation and fibrosis, is the primary reason for the slow adoption of minimally invasive techniques in chronic pancreatitis. However, recent studies have shown the feasibility of laparoscopic and robotic surgery for chronic pancreatitis. Also, with advancements in endoscopic treatment, there is a trend towards the less invasive treatment of chronic pancreatitis. The present review focuses on the challenges, evolution, and changing trends in the minimally invasive management of chronic pancreatitis.

SEARCH STRATEGY

All the authors did a PubMed search of relevant articles. Further, the articles' reference lists were also searched for additional appropriate studies. The keywords and combinations included in the search were: "Pancreatitis"; "chronic pancreatitis"; "idiopathic pancreatitis"; endoscopic management" and "chronic pancreatitis"; "Frey's procedure" and "Laparoscopic"; "Frey's procedure" and "robotic"; "Puestow procedure" and "Laparoscopic"; "Puestow procedure" and "robotic"; "Beger procedure" and "Laparoscopic"; "Beger procedure" and "robotic"; "chronic pancreatitis" and "total pancreatectomy" and "laparoscopic"; "chronic pancreatitis" and "total pancreatectomy", and "robotic". The search was limited to publications in English literature. All the authors agreed that the articles selected for review were relevant.

ENDOSCOPIC MANAGEMENT OF CHRONIC PANCREATITIS

Endoscopic intervention is recommended as a minimally invasive alternative to surgery in patients who do not improve with medical management. The journey of stone removal from the pancreatic duct dates back to 1891 when Alfred Pearce Gould retrieved calculi from the Wirsung duct in a London hospital [6]. Berkeley G Moynihan performed transduodenal removal of pancreatic stones in 1902, followed by transpancreatic stone removal in 1908 by Mayo Robson. The development of the fiberoptic endoscope for diagnosis in 1958 by Basil Isaac Hirschowitz changed the trends of endoscopic management[7]. Watson *et al*[8] succeeded in developing the technique for papillotomy with an energy source leading to

the endoscopic extraction of calculi and subsequent stent placement. Pathogenesis of pain in chronic pancreatitis is multifactorial and includes anatomical and neuropathic factors. Anatomical alterations include ductal hypertension, raised pancreatic parenchymal pressure, acute inflammation, and pancreatic ischemia[9-11]. The lack of correlation between pain severity and anatomical changes suggests neurological factors' role. Neuropathophysiology of pain in chronic pancreatitis includes peripheral sensitization-induced pain, neuropathic remodeling, and central sensitization of pancreatic pain[10,11]. Endoscopic therapy aims to relieve pain by clearance of intraductal stones, thereby decompressing the pancreatic duct. Proponents of endoscopic therapy suggest that patients with complete ductal clearance by endoscopic approach have shown similar pain relief compared to the surgical group[12]. At the same time, the ability to modify the disease progression and prevent loss of pancreatic function with early surgical intervention was proposed in favor of surgical management[13, 14].

Traditionally chronic pancreatitis patients were managed initially with analgesics and pancreatic enzyme replacement therapy, followed by endoscopic intervention with or without extracorporeal shockwave therapy[15,16]. Patients with an inadequate clinical response following endoscopic treatment should be discussed in the multidisciplinary meeting and considered for surgical management[10,16]. The potential benefit of this approach is that surgery could be avoided in some patients with successful endoscopic management. However, various studies comparing the efficacy of endoscopic management with surgery have shown better pain control with the early surgical intervention[17-19]. Randomized trials comparing the endoscopic and surgical approaches have shown a better quality of life and pain relief in the surgical group, especially with early surgery[12,20,21] (Table 1). Also, patients with inflammatory pancreatic head mass, distal pancreatic duct stricture, and extensive parenchyma calcifications of the pancreatic head might be difficult to treat by endoscopy. Despite the available evidence favoring surgical treatment, advancements in endoscopic lithotripsy techniques and extracorporeal shockwave lithotripsy (ESWL) are continuously improving the ductal clearance rate, and the endoscopic approach is helpful in a subset of chronic pancreatitis patients[16]. Patients with dominant stricture in the pancreatic head with upstream dilatation and those with intraductal calculi in the pancreatic head or proximal body are ideal candidates for endoscopic intervention.

Pancreatic lithotripsy

Worldwide, ESWL is commonly used for pancreatic lithotripsy, especially in Asia and Europe. While recent advancements like intraductal endoscopic laser or electrohydraulic lithotripsy might improve ductal clearance, they are not widely available. Hence, most of the available data are for ESWL[22]. A meta-analysis of various studies has reported a 70% stone clearance rate with ESWL[23]. With ESWL, stone clearance is more favorable for solitary calculus in the head of the pancreas. However, recurrence of stones after ESWL was seen in 14% to 23% of patients mandating further intervention[24]. ESWL is combined with pancreatic duct stenting in patients with associated pancreatic duct stricture[25]. European Society of Gastrointestinal Endoscopy (ESGE) recommends ESWL for the removal of radiopaque obstructive main pancreatic duct calculi greater than 5 mm found in the head or body of the pancreas and endoscopic retrograde cholangiopancreatography (ERCP) for main pancreatic duct calculi that are radiolucent or smaller than 5 mm[26].

Endoscopic stenting

Main pancreatic duct strictures are often seen in half of the chronic pancreatitis patients and are usually located in the pancreatic head region. The standard management of these strictures is balloon dilatation and placement of a temporary stent for at least a year[27]. Pain relief after a long-term follow-up of 5 years has been seen in almost half of patients after stent withdrawal[28,29]. However, stricture recurrence has been reported in up to 38% of patients after two years[28,30]. Complications related to stenting include stent migration and occlusion. Distal stent migration toward the pancreatic tail and proximal stent migration to the duodenum was reported in 7.5% and 5.2% of patients, respectively[31]. Stent migration can be prevented with large-winged or pigtail catheters[31,32]. The ESGE recommends managing painful pancreatic duct strictures with the help of a single 10 Fr stent for one uninterrupted year if symptoms improve after placement[26]. The stent should be exchanged based on symptoms or signs of stent dysfunction on imaging at least six months intervals.

Endoscopic ultrasound-guided intervention

Endoscopic ultrasound (EUS) guided drainage of the main pancreatic duct is used as a second-line procedure after the failure of ERCP. Despite high success rates of 68% to 75%, the complications like perforation, bleeding, and pancreatitis reported in 5% to 43% of patients were key drawbacks for EUS-guided drainage[33,34]. Another EUS-guided procedure is the celiac plexus block, whereby a steroid with a local anesthetic agent is injected at the celiac plexus to block the pancreatic nerve fibers. EUS-guided celiac plexus block is preferred over the traditional percutaneous method. However, despite the high technical success and short-term pain relief in 55%-70% of patients, long-term outcomes are discouraging[35-37].

Table 1 Studies comparing endoscopic and surgical management of chronic pancreatitis

Ref.	Country	No. of patients (n), surgery/endoscopy	Complete and partial pain relief (%), surgery/endoscopy	Complication (%), surgery/endoscopy	Mortality (n), surgery/endoscopy	Follow up (mo), surgery/endoscopy
Dite <i>et al</i> [12], 2003	Czech Republic	36/36	86/61	Not reported	0/0	60
Cahen <i>et al</i> [20], 2011	Netherlands	16/15	80/38	0/25	0/1	79
Hong <i>et al</i> [19], 2011	China	27/35	77/47	14/22	1/0	60
Kawashima <i>et al</i> [17], 2018	Japan	41/10	100/100	20/27	0/0	-
Jiang <i>et al</i> [18], 2018	China	40/46	83/80	26/8	0/0	63.5/57.3
Issa <i>et al</i> [21], 2020	Netherlands	44/44	58/39	27/25	0/0	18

SURGICAL MANAGEMENT OF CHRONIC PANCREATITIS

Evolution of open surgical procedures

The systemic reviews and meta-analyses of the available studies comparing surgical management with endoscopic intervention continue to support the superiority of surgical treatment in chronic pancreatitis [38-40]. The intriguing voyage of surgical management of chronic pancreatitis has witnessed a huge transformation from surgical decompression of the obstructed duct and denervation of the pancreas to pancreatic head resection, total pancreatectomy, and various other hybrid procedures. In 1911, Link[41], a gynecologist from Indiana, was the first to describe external tube drainage of the pancreatic duct for ductal calculi or stricture with good long-term pain relief. Desjardins and Coffey, around similar times, proposed drainage of the pancreas using the intestine after conducting animal studies[42,43]. This empiricism was materialized by Duval[44] and separately by Zollinger *et al*[45] in 1954 by describing the first caudal end-to-end pancreaticojejunostomy using a Roux-en-Y jejunal loop for chronic pancreatitis. Further, modification in the decompressive procedure was done by Puestow and Gillesby[46], in 1958, by invaginating caudal pancreaticojejunostomy after longitudinally opening the pancreatic duct from the body to the tail region of the pancreas. Their procedure was further modified by Partington and Rochelle[47], in 1960, by creating side-to-side Roux-en-Y pancreaticojejunostomy, commonly called the Puestow procedure. Though the Puestow procedure was considered standard drainage procedure for almost 40 years, the long-term benefits were not befitting. Despite short-term pain relief in 80% of patients, the pain recurred in 30% on long-term follow-up[48-51]. The foremost reason for recurrent pain was undealt nidus of inflammation and persistent ductal disease in the head of the pancreas. To tackle the head disease, German surgeon Hans Beger performed the first duodenum-preserving pancreatic head resection in 1971 and reported postoperative outcomes of 52 patients in 1980[52]. In 1984, Warren described splenopancreatic flap, a denervation procedure for chronic pancreatitis[53]. However, the long-term results of this complex procedure were never published or replicated by other surgeons[9,53].

In 1987, Frey and Smith[54] described a hybrid operation consisting of pancreatic head resection and longitudinal pancreaticojejunostomy, also known as Frey's procedure. Izbicki *et al*[55] modified Frey's procedure by doing a more extensive excavation of the pancreatic head and duct and named it Hamburg modification. Similarly, Gloor *et al*[56] from Berne modified the duodenum-preserving pancreatic head resection by omitting the challenging step of pancreatic neck transection. To minimize the risk of penetrating the posterior pancreatic capsule, Ho and Frey[57], and Frey and Amikura[58] recommended limiting the posterior extent of head coring to the back wall of opened Wirsung and the uncinat duct, also known as modified Frey's procedure. Sakata *et al*[59] described the minimum Frey procedure in which a small spindle-shaped anterior resection of the pancreatic head was performed and reported an equivalent outcome. However, a retrospective study by Tan *et al*[60] reported superior long-term pain relief and quality of life with the original Frey's procedure compared to the modified Frey's. A systematic review and meta-analysis of trials comparing various resectional and hybrid procedures reported similar postoperative pain relief. However, quality of life and other perioperative outcomes favor duodenum-preserving pancreatic head resection procedures[61].

Era of minimally invasive surgery

There is a trend towards minimally invasive procedures for various surgical disorders, and the change is inevitable for chronic pancreatitis[62-64]. Though the surgical procedures described for chronic

pancreatitis are complex and challenging due to inflamed gland they can be accomplished in selected patients[63-65]. In high-volume centers with expertise in advanced laparoscopic procedures, complex pancreatic procedures can be safely performed with comparable postoperative outcomes[63-69]. Also, with its ergonomic advantages, robotic surgery could overcome some of the technical limitations of laparoscopic surgery and potentially widen the use of a minimally invasive approach in chronic pancreatitis[70,71].

Minimally invasive Puestow procedure

Like open surgical procedures, minimally invasive surgery for chronic pancreatitis began with a modified Puestow procedure. Kurian and Gagner[72], in 1999, reported the first series of five patients who underwent a laparoscopic Puestow procedure. Subsequently, two series with 17 and 12 patients were published from India[73,74]. The first small case series of 5 patients from the United Kingdom was published by Khaled *et al*[75] in 2014. The feasibility and favorable short-term outcomes of the laparoscopic Puestow procedure were documented in multiple case series[76-81] (Table 2). In most series, the procedure was performed with five laparoscopic trocars. The initial entry to the lesser sac and exposure of the anterior surface of the pancreas can be technically challenging in patients with recent or recurrent acute episodes of pancreatitis. Hence, those patients should be avoided during the early phase of the minimally invasive Puestow procedure. In most laparoscopic series, two to three gastric retraction sutures are used to lift the stomach away from the pancreas and improve exposure. Needle aspiration is commonly used to identify the pancreatic duct, and intraoperative ultrasound is helpful in patients with undilated duct[74]. Extraction of all intraductal calculi, especially those in the head and tail region, is critical for long-term pain relief. Sahoo *et al*[76] reported the usefulness of cystoscope and endoscopic basket in clearing residual intraductal stones. Proficient intracorporeal suturing skill is critical to accomplish safe pancreaticojejunostomy. Kim *et al*[77] reported the benefits of using barbed sutures for laparoscopic pancreaticojejunostomy. Bhandarwar *et al*[78] used endostaplers for laparoscopic pancreaticojejunostomy anastomosis with the anvil part placed within the pancreatic duct. However, as highlighted by the authors, the technique was feasible only in seven out of 17 patients with pancreatic duct diameter of more than 10 mm[78]. Alternatively, a robotic platform can minimize the challenges associated with intracorporeal suturing. After the initial case reports of the robotic Puestow procedure documented its usefulness in a series of seven patients[82-87]. However, with the emerging evidence supporting some form of head resection to achieve long-term pain relief, the minimally invasive Puestow procedure is recommended only in a subset of chronic pancreatitis patients with an atrophic pancreas and dilated pancreatic duct with predominant intraductal calculi.

Minimally invasive Frey's procedure

Frey's procedure is one of the most commonly performed surgeries for chronic pancreatitis. Pancreatic head coring is the technically challenging step of this hybrid procedure, especially during minimally invasive surgery. The first series of laparoscopic Frey's procedure was published by Tan *et al*[88] in 2015. Subsequently, a small series of four patients reported the feasibility and short-term outcomes of laparoscopic Frey's procedure[89]. The largest published series to date had 15 patients in the laparoscopic Frey's group[90]. The relatively small number of studies with fewer patients highlight the technical challenges of laparoscopic Frey's procedure (Table 3). In open Frey's procedure, the surgeon's left hand, kept under the posterior surface of the pancreas head, guides the extent of posterior head coring. In the absence of a definite landmark, pancreatic head coring until the level of the posterior pancreatic capsule is challenging during laparoscopic surgery. Hence, in all the laparoscopic series, only modified Frey's procedure was performed using the main pancreatic duct as the landmark and coring to the posterior wall of the duct[88-90]. In the laparoscopic approach, ultrasonic shears and bipolar vessel sealing devices are commonly used for head coring. Lack of articulation and difficulty securing precise hemostatic sutures further increase the difficulty of laparoscopic pancreatic head coring, especially along uncinate ducts[90]. With its articulating instruments, the robotic platform could potentially overcome technical difficulties during head coring and pancreaticojejunostomy. Hamad *et al*[62] highlighted the usefulness of the robotic approach in bleeding control during head coring in their series of four patients. However, similar to the laparoscopic approach, due to the lack of a definite landmark, coring was limited to the posterior wall of the duct in the robotic modified Frey's procedure (Figure 1). The median operative time and blood loss were 372 min and 163 mL, respectively[62]. The parenchyma posterior to the main pancreatic duct in the pancreatic head was preserved to prevent injury to the superior mesenteric vein[62,91].

Shukla *et al*[92] reported the feasibility of robot assisted Frey's procedure in nine patients with chronic pancreatitis. Robotic approach is associated with less blood loss and shorter hospital stay compared to open Frey's procedure. Tile Pro technology in the robotic platform allows the surgeon to view ultrasound images in the console, thereby avoiding damage to the common bile duct and portal vein during dissection[62]. Bleeding is one of the common causes of conversion in minimally invasive Frey's procedure. Inflammatory pancreatic head mass and preoperative acute exacerbation of pancreatitis were identified as significant risk factors for intraoperative blood loss. Hence, minimally invasive Frey's procedure is recommended in patients with dilated pancreatic duct and enlarged pancreatic head on imaging without inflammatory mass, recent acute exacerbation, and pancreatitis-related complica-

Table 2 Studies on laparoscopic Puestow procedure

Ref.	Country	No. of patients (n)	Mean operative time (min)	Mean hospital stay (d)	Conversion (n)	Mortality	Follow up, (mo)	Complete pain relief (%)
Kurian <i>et al</i> [72], 1999	United States	5	240.0	3-7	Nil	Nil	30.0	80.0
Tantia <i>et al</i> [73], 2004	India	17	277.0	5.2	4	Nil	12.0	82.3
Palanivelu <i>et al</i> [74], 2006	India	12	172.0	5.0	Nil	Nil	52.8	83.3
Khaled <i>et al</i> [75], 2014	United Kingdom	6	278.0	7.0	Not reported	Nil	14.2	66.7
Sahoo <i>et al</i> [76], 2014	India	12	265.5	5.8	Nil	Nil	16.5	100 (follow up reported for 8 patients)
Kim and Hong [77], 2016	Korea	11	200.0	7.0	Nil	Nil	21.0	100
Bhandarwar, <i>et al</i> [78], 2019	India	28	189.7	5.8	4	Nil	12.0	87.5
Rege <i>et al</i> [79], 2019	India	32	131.2	5.2	1	Nil	14.2	75.0
Javed <i>et al</i> [80], 2020	India	41	180.0	5.0	Excluded	Nil	43.6	91.0
Nag <i>et al</i> [81], 2022	India	33	300.0	7.0	Nil	Nil	25.0	71.0


Table 3 Studies on laparoscopic Frey's procedure

Ref.	Country	Patients (n)	Mean operative time (min)	Mean hospital stay (d)	Conversion (n)	Mortality	Follow up (mo)	Complete and partial pain relief
Tan <i>et al</i> [88], 2015	China	9	323	7	2	Nil	3	Not reported
Kilburn <i>et al</i> [89], 2017	Australia	4	130	7	Nil	Nil	26	100%
Senthilnathan <i>et al</i> [90], 2019	India	15	271	6.4	10 out of 57 patients in different arms	Nil	60	88%

ations.


Minimally invasive duodenum preserving pancreatic head resection

In patients with inflammatory head mass, duodenum preserving pancreatic head resection (DPHR) is preferred over pancreatoduodenectomy because of favorable long-term outcomes. As previously highlighted, lack of definite landmarks precludes pancreatic head coring until the posterior pancreatic capsule in minimally invasive Frey's procedure. Minimally invasive duodenum-preserving pancreatic head resection could potentially overcome that limitation. After the initial case reports, it is reported that the feasibility of laparoscopic DPHR procedure in 5 patients with chronic pancreatitis[93-95]. The mean operative time and hospital stay were 275 min and 11 d, respectively. One patient had grade B postoperative pancreatic fistula, and pancreaticojejunostomy anastomotic site bleed in one patient[95]. In laparoscopic DPHR after the pancreatic neck transection, the pancreatic head is retracted to identify the plane between the pancreatic parenchyma and posterior pancreatic capsule. As the dissection proceeds along the superior border of the pancreas, the bile duct should be identified and preserved. Identifying the intrapancreatic duct is a significant technical challenge with minimally invasive DPHR. Indocyanine green (ICG) fluorescence facilitates bile duct identification in the triangle formed by the gastroduodenal artery, portal vein, and superior border of the pancreas (Figure 2). Energy sources should be judiciously used around the bile duct to prevent thermal damage[96-98]. Also, ischemia of the bile duct can be prevented by preserving the posterosuperior pancreaticoduodenal artery, a proximal branch of the gastroduodenal artery, and preserving pancreatic tissue medial to the bile duct. To prevent duodenal ischemia and delayed gastric emptying, the pancreaticoduodenal arcade along the medial border of the duodenum should be preserved. Hong *et al*[95] reported the usefulness of ICG in assessing vascular arcade and identifying common bile duct in a series of 22 patients with different pancreatic pathology. The mean operative time and blood loss of five patients with chronic pancreatitis



DOI: 10.3748/wjg.v29.i14.2101 Copyright ©The Author(s) 2023.

Figure 1 Robotic modified Frey's procedure. A: Pancreatic head coring is done till the level of the posterior wall of the pancreatic duct (marked with star). The bile duct can be seen on the medial wall of the cored-out tissue (arrow); B: Indocyanine green fluorescence demonstrates the bile duct on the medial wall of the cored-out tissue.



DOI: 10.3748/wjg.v29.i14.2101 Copyright ©The Author(s) 2023.

Figure 2 Robotic duodenum preserving pancreatic head resection. A: Dissection of the pancreatic parenchyma from the posterior pancreatic capsule; B: Identification of the common bile duct in the triangle formed by the gastroduodenal artery, superior border of the pancreas, and portal vein; C: Pancreatic duct (arrow) divided at its junction with the bile duct; D: Post pancreatic head resection, indocyanine green fluorescence demonstrates bile duct.

included in their series were 264 min and 215 mL, respectively. The mean postoperative hospital stay was 7.5 d, and there was no conversion to open surgery or postoperative mortality[96].

Inflammation and tissue adhesion in chronic pancreatitis can distort the anatomy of pancreaticoduodenal vessels resulting in vascular injury and significant bleeding during DPHR. A 3D reconstruction of preoperative cross-sectional imaging could help to better understand the anatomy of pancreaticoduodenal vessels and the relationship of the intrapancreatic common bile duct. Also, 3D printing technology can be helpful for surgical training and preoperative planning in patients undergoing minimally invasive DPHR[96-98]. As with other minimally invasive procedures for chronic pancreatitis, a robotic platform could minimize the technical challenges associated with DPHR. Peng *et al*[98] first reported the feasibility of robotic DPHR. However, in a recent series of 68 patients undergoing robotic DPHR for various pancreatic diseases, only three patients had chronic pancreatitis

[99]. Different published series on minimally invasive DPHR highlight technical development and challenges of the procedure in chronic pancreatitis. Minimally invasive DPHR is the preferred procedure in patients with inflammatory head mass and those with enlarged pancreatic head and extensive parenchymal calcifications. However, if the inflammatory changes preclude the safe creation of a retropancreatic tunnel over the portal vein alternative surgical procedure should be considered.

Minimal invasive total pancreatectomy with or without islet cell autotransplantation

Total pancreatectomy is primarily indicated in chronic pancreatitis patients with debilitating pain in whom all other measures are unsuccessful and those with recurrent acute pancreatitis[100]. However, total pancreatectomy should be combined with islet cell autotransplantation to minimize the risk of brittle diabetes. Some centers recommend total pancreatectomy early in the disease course before activation of neuropathic pain circuits, especially in patients with small duct disease or genetic etiology [101]. However, selecting suitable patients is critical as, despite islet cell autotransplantation, more than 50% of patients might require lifelong exogenous insulin. Literature on minimally invasive total pancreatectomy is sparse, with variations in technique[102-105]. Blair *et al*[103], in 2016, reported the feasibility and safety of laparoscopic total pancreatectomy with islet cell autotransplantation in 20 patients with chronic pancreatitis. The mean operative time and hospital stay were 430 min and 11 d, respectively, with no postoperative mortality. Similarly, Fan *et al*[101] reported the feasibility of laparoscopic total pancreatectomy with islet cell autotransplantation in 22 patients with two conversions. In both the laparoscopic series, the pancreatic neck was transected, and two-stage retrieval was used, with the pancreatic head and body retrieved separately[102,104]. However, studies have shown the importance of preserving pancreatic arterial and venous flow until retrieval to reduce warm ischemia time during the pancreatic dissection phase and improve islet yield[105]. In the robotic series reported by Galvani *et al*[102] and Zureikat *et al*[104] the feasibility of total pancreatectomy without pancreatic neck transection and preserving vascular flow till the final step to reduce warm ischemia was documented. Another technical challenge is dense retroperitoneal adhesions due to recurrent pancreatic inflammation. Although laparoscopic and robotic total pancreatectomy with islet autotransplantation is safe and feasible, appropriate patient selection is critical for deriving the benefit of a minimally invasive approach.

CHALLENGES AND FUTURE PERSPECTIVES

The available evidence suggests that minimally invasive surgery for chronic pancreatitis is feasible in selected patients. However, the poor quality of available evidence precludes definite conclusions. Also, the surgeon should adhere to surgery principles for chronic pancreatitis, irrespective of the approach. As the pancreatic head is the pacemaker of pain in most patients, adequate resection and decompression of the pancreatic head are critical. However, most of the reported series on minimally invasive approaches for chronic pancreatitis have focused on the feasibility of lateral pancreaticojejunostomy or modified Frey's procedure which may be appropriate only in a minority of chronic pancreatitis patients. Recent series have shown the feasibility of minimally invasive duodenal preserving pancreas head resection, which may be the ideal procedure for most chronic pancreatitis patients. However, a minimally invasive approach is feasible only in patients without extensive inflammatory adhesions or recent acute exacerbation. Preoperative cross-sectional imaging and biochemical parameters like serum amylase and lipase are not sensitive to predict inflammatory changes. Also, studies evaluating the predictive value of markers of systemic inflammation like white blood cell count, IL-6, and C reactive protein yielded disappointing results[88]. Future studies should focus on identifying reliable markers that can accurately predict ongoing pancreatic inflammation, thereby aiding patient selection for a minimally invasive approach. With recent evidence supporting early surgical intervention before the development of extensive fibrosis or local complications, more patients may be suitable for minimally invasive surgery. Also, the main problem with the existing procedures is they are primarily focused on pancreatic ductal and parenchymal decompression. However, it is well-documented that anatomical factors alone do not contribute to pancreatic pain in all patients. In a subgroup of chronic pancreatitis patients, neurological pathways of pain play a dominant role, which is not addressed by the commonly performed surgical procedures. Also, future studies should compare laparoscopic and robotic procedures for chronic pancreatitis to document the advantages of the robotic platform.

CONCLUSION

As pain is the primary indication for intervention in chronic pancreatitis use of a minimally invasive approach is an attractive proposition. However, due to technical challenges, both endoscopic intervention and minimally invasive surgery for chronic pancreatitis have lagged compared to other benign gastrointestinal orders. With improvements in laparoscopic instrumentation and technological

advances like ICG fluorescence, minimally invasive surgery for chronic pancreatitis is gaining momentum. Also, with its distinct advantages, the robotic platform can widen the adoption of minimally invasive surgery in chronic pancreatitis. However, well-designed trials with long-term follow-ups are required to document the superiority of minimally invasive surgery for chronic pancreatitis.

FOOTNOTES

Author contributions: Kalayarasan R and Shukla A did the literature search; Shukla A wrote the first draft of the review; Kalayarasan R conceptualized the work, supervised the writing, gave intellectual inputs, and critically revised the manuscript.

Conflict-of-interest statement: All authors have no conflicts of interest to report.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: India

ORCID number: Raja Kalayarasan 0000-0003-4056-8672; Ankit Shukla 0000-0002-5037-8525.

S-Editor: Chen YL

L-Editor: A

P-Editor: Chen YL

REFERENCES

- Garg PK, Tandon RK. Survey on chronic pancreatitis in the Asia-Pacific region. *J Gastroenterol Hepatol* 2004; **19**: 998-1004 [PMID: 15304116 DOI: 10.1111/j.1440-1746.2004.03426.x]
- Witt H, Apte MV, Keim V, Wilson JS. Chronic pancreatitis: challenges and advances in pathogenesis, genetics, diagnosis, and therapy. *Gastroenterology* 2007; **132**: 1557-1573 [PMID: 17466744 DOI: 10.1053/j.gastro.2007.03.001]
- Yang CJ, Bliss LA, Freedman SD, Sheth S, Vollmer CM, Ng SC, Callery MP, Tseng JF. Surgery for chronic pancreatitis: the role of early surgery in pain management. *Pancreas* 2015; **44**: 819-823 [PMID: 25882695 DOI: 10.1097/MPA.0000000000000333]
- Ray S, Basu C, Dhali A, Dhali GK. Frey procedure for chronic pancreatitis: A narrative review. *Ann Med Surg (Lond)* 2022; **80**: 104229 [PMID: 36045829 DOI: 10.1016/j.amsu.2022.104229]
- Nitesh PNB, Pottakkat B. Bile duct preserving pancreatic head resection (BDPPHR): Can we conclusively define the extent of head resection in surgery for chronic pancreatitis? *Ann Hepatobiliary Pancreat Surg* 2020; **24**: 309-313 [PMID: 32843597 DOI: 10.14701/ahbps.2020.24.3.309]
- Howard JM, Hess W. History of the pancreas: mysteries of a hidden organ. Nueva York: Kluwer Academic/Plenum Publishers, 2002 [DOI: 10.1016/s1091-255x(03)00078-7]
- Navarro S. The art of pancreatic surgery. Past, present and future. The history of pancreatic surgery. *Gastroenterol Hepatol* 2017; **40**: 648.e1-648.e11 [PMID: 28533021 DOI: 10.1016/j.gastrohep.2017.03.010]
- Watson WC. Direct vision of the ampulla of Vater through the gastroduodenal fiberscope. *Lancet* 1966; **1**: 902-903 [PMID: 4159621 DOI: 10.1016/s0140-6736(66)91578-9]
- Andersen DK, Frey CF. The evolution of the surgical treatment of chronic pancreatitis. *Ann Surg* 2010; **251**: 18-32 [PMID: 20009754 DOI: 10.1097/SLA.0b013e3181ae3471]
- Dumonceau JM. Endoscopic therapy for chronic pancreatitis. *Gastrointest Endosc Clin N Am* 2013; **23**: 821-832 [PMID: 24079792 DOI: 10.1016/j.giec.2013.06.004]
- Drewes AM, Krarup AL, Detlefsen S, Malmström ML, Dimcevski G, Funch-Jensen P. Pain in chronic pancreatitis: the role of neuropathic pain mechanisms. *Gut* 2008; **57**: 1616-1627 [PMID: 18566105 DOI: 10.1136/gut.2007.146621]
- Díte P, Ruzicka M, Zboril V, Novotný I. A prospective, randomized trial comparing endoscopic and surgical therapy for chronic pancreatitis. *Endoscopy* 2003; **35**: 553-558 [PMID: 12822088 DOI: 10.1055/s-2003-40237]
- Maartense S, Ledebøer M, Bemelman WA, Ringers J, Frolich M, Masclee AA. Effect of surgery for chronic pancreatitis on pancreatic function: pancreatico-jejunostomy and duodenum-preserving resection of the head of the pancreas. *Surgery* 2004; **135**: 125-130 [PMID: 14739846 DOI: 10.1016/j.surg.2003.09.004]
- Lamme B, Boermeester MA, Straatsburg IH, van Buijtenen JM, Boerma D, Offerhaus GJ, Gouma DJ, van Gulik TM. Early versus late surgical drainage for obstructive pancreatitis in an experimental model. *Br J Surg* 2007; **94**: 849-854 [PMID: 17335122 DOI: 10.1002/bjs.5722]
- Dumonceau JM, Kapral C, Aabakken L, Papanikolaou IS, Tringali A, Vanbiervliet G, Beyna T, Dinis-Ribeiro M, Hritz I, Mariani A, Paspatis G, Radaelli F, Lakhtakia S, Veitch AM, van Hooft JE. ERCP-related adverse events: European

- Society of Gastrointestinal Endoscopy (ESGE) Guideline. *Endoscopy* 2020; **52**: 127-149 [PMID: 31863440 DOI: 10.1055/a-1075-4080]
- 16 **Drewes AM**, Bouwense SAW, Campbell CM, Ceyhan GO, Delhaye M, Demir IE, Garg PK, van Goor H, Halloran C, Isaji S, Neoptolemos JP, Olesen SS, Palermo T, Pasricha PJ, Sheel A, Shimosegawa T, Szigethy E, Whitcomb DC, Yadav D; Working group for the International (IAP – APA – JPS – EPC) Consensus Guidelines for Chronic Pancreatitis. Guidelines for the understanding and management of pain in chronic pancreatitis. *Pancreatol* 2017; **17**: 720-731 [PMID: 28734722 DOI: 10.1016/j.pan.2017.07.006]
 - 17 **Kawashima Y**, Kawaguchi Y, Kawanishi A, Ogawa M, Hirabayashi K, Nakagohri T, Mine T. Comparison between Endoscopic Treatment and Surgical Drainage of the Pancreatic Duct in Chronic Pancreatitis. *Tokai J Exp Clin Med* 2018; **43**: 117-121 [PMID: 30191547]
 - 18 **Jiang L**, Ning D, Cheng Q, Chen XP. Endoscopic versus surgical drainage treatment of calcific chronic pancreatitis. *Int J Surg* 2018; **54**: 242-247 [PMID: 29684668 DOI: 10.1016/j.ijso.2018.04.027]
 - 19 **Hong J**, Wang J, Keleman AM, Imagawa DK, Xu K, Wang W, Liu E, Niu W, Sun Q, Peng C, Zhao W, Niu J. Endoscopic versus surgical treatment of downstream pancreatic duct stones in chronic pancreatitis. *Am Surg* 2011; **77**: 1531-1538 [PMID: 22196670 DOI: 10.1177/000313481107701146]
 - 20 **Cahen DL**, Gouma DJ, Laramée P, Nio Y, Rauws EA, Boermeester MA, Busch OR, Fockens P, Kuipers EJ, Pereira SP, Wonderling D, Dijkgraaf MG, Bruno MJ. Long-term outcomes of endoscopic vs surgical drainage of the pancreatic duct in patients with chronic pancreatitis. *Gastroenterology* 2011; **141**: 1690-1695 [PMID: 21843494 DOI: 10.1053/j.gastro.2011.07.049]
 - 21 **Issa Y**, Kempeneers MA, Bruno MJ, Fockens P, Poley JW, Ahmed Ali U, Bollen TL, Busch OR, Dejong CH, van Duijvendijk P, van Dullemen HM, van Eijck CH, van Goor H, Hadithi M, Haveman JW, Keulemans Y, Nieuwenhuijs VB, Poen AC, Rauws EA, Tan AC, Thijs W, Timmer R, Witteman BJ, Besselink MG, van Hooft JE, van Santvoort HC, Dijkgraaf MG, Boermeester MA; Dutch Pancreatitis Study Group. Effect of Early Surgery vs Endoscopy-First Approach on Pain in Patients With Chronic Pancreatitis: The ESCAPE Randomized Clinical Trial. *JAMA* 2020; **323**: 237-247 [PMID: 31961419 DOI: 10.1001/jama.2019.20967]
 - 22 **Fishman DS**, Tarnasky PR, Patel SN, Rajman I. Management of pancreaticobiliary disease using a new intra-ductal endoscope: the Texas experience. *World J Gastroenterol* 2009; **15**: 1353-1358 [PMID: 19294765 DOI: 10.3748/wjg.15.1353]
 - 23 **Guda NM**, Partington S, Freeman ML. Extracorporeal shock wave lithotripsy in the management of chronic calcific pancreatitis: a meta-analysis. *JOP* 2005; **6**: 6-12 [PMID: 15650279]
 - 24 **Tandan M**, Reddy DN, Santosh D, Vinod K, Ramchandani M, Rajesh G, Rama K, Lakhtakia S, Banerjee R, Pratap N, Venkat Rao G. Extracorporeal shock wave lithotripsy and endotherapy for pancreatic calculi-a large single center experience. *Indian J Gastroenterol* 2010; **29**: 143-148 [PMID: 20717860 DOI: 10.1007/s12664-010-0035-y]
 - 25 **Tandan M**, Nageshwar Reddy D. Endotherapy in chronic pancreatitis. *World J Gastroenterol* 2013; **19**: 6156-6164 [PMID: 24115811 DOI: 10.3748/wjg.v19.i37.6156]
 - 26 **Dumonceau JM**, Delhaye M, Tringali A, Arvanitakis M, Sanchez-Yague A, Vaysse T, Aithal GP, Anderloni A, Bruno M, Cantú P, Devière J, Domínguez-Muñoz JE, Lekkerkerker S, Poley JW, Ramchandani M, Reddy N, van Hooft JE. Endoscopic treatment of chronic pancreatitis: European Society of Gastrointestinal Endoscopy (ESGE) Guideline - Updated August 2018. *Endoscopy* 2019; **51**: 179-193 [PMID: 30654394 DOI: 10.1055/a-0822-0832]
 - 27 **Ponchon T**, Bory RM, Hedelius F, Roubein LD, Paliard P, Napoleon B, Chavaillon A. Endoscopic stenting for pain relief in chronic pancreatitis: results of a standardized protocol. *Gastrointest Endosc* 1995; **42**: 452-456 [PMID: 8566637 DOI: 10.1016/s0016-5107(95)70049-8]
 - 28 **Binmoeller KF**, Jue P, Seifert H, Nam WC, Izbicki J, Soehendra N. Endoscopic pancreatic stent drainage in chronic pancreatitis and a dominant stricture: long-term results. *Endoscopy* 1995; **27**: 638-644 [PMID: 8903975 DOI: 10.1055/s-2007-1005780]
 - 29 **Costamagna G**, Bulajic M, Tringali A, Pandolfi M, Gabbrielli A, Spada C, Petruzzello L, Familiari P, Mutignani M. Multiple stenting of refractory pancreatic duct strictures in severe chronic pancreatitis: long-term results. *Endoscopy* 2006; **38**: 254-259 [PMID: 16528652 DOI: 10.1055/s-2005-921069]
 - 30 **Moon SH**, Kim MH, Park DH, Song TJ, Eum J, Lee SS, Seo DW, Lee SK. Modified fully covered self-expandable metal stents with antimigration features for benign pancreatic-duct strictures in advanced chronic pancreatitis, with a focus on the safety profile and reducing migration. *Gastrointest Endosc* 2010; **72**: 86-91 [PMID: 20493483 DOI: 10.1016/j.gie.2010.01.063]
 - 31 **Johanson JF**, Schmalz MJ, Geenen JE. Incidence and risk factors for biliary and pancreatic stent migration. *Gastrointest Endosc* 1992; **38**: 341-346 [PMID: 1607087 DOI: 10.1016/s0016-5107(92)70429-5]
 - 32 **Sherman S**, Hawes RH, Savides TJ, Gress FG, Ikenberry SO, Smith MT, Zaidi S, Lehman GA. Stent-induced pancreatic ductal and parenchymal changes: correlation of endoscopic ultrasound with ERCP. *Gastrointest Endosc* 1996; **44**: 276-282 [PMID: 8885346 DOI: 10.1016/s0016-5107(96)70164-5]
 - 33 **Shah JN**, Marson F, Weilert F, Bhat YM, Nguyen-Tang T, Shaw RE, Binmoeller KF. Single-operator, single-session EUS-guided anterograde cholangiopancreatography in failed ERCP or inaccessible papilla. *Gastrointest Endosc* 2012; **75**: 56-64 [PMID: 22018554 DOI: 10.1016/j.gie.2011.08.032]
 - 34 **Kahaleh M**, Hernandez AJ, Tokar J, Adams RB, Shami VM, Yeaton P. EUS-guided pancreaticogastrostomy: analysis of its efficacy to drain inaccessible pancreatic ducts. *Gastrointest Endosc* 2007; **65**: 224-230 [PMID: 17141775 DOI: 10.1016/j.gie.2006.05.008]
 - 35 **Stevens T**, Costanzo A, Lopez R, Kapural L, Parsi MA, Vargo JJ. Adding triamcinolone to endoscopic ultrasound-guided celiac plexus blockade does not reduce pain in patients with chronic pancreatitis. *Clin Gastroenterol Hepatol* 2012; **10**: 186-191, 191.e1 [PMID: 21946121 DOI: 10.1016/j.cgh.2011.09.006]
 - 36 **Santosh D**, Lakhtakia S, Gupta R, Reddy DN, Rao GV, Tandan M, Ramchandani M, Guda NM. Clinical trial: a randomized trial comparing fluoroscopy guided percutaneous technique vs. endoscopic ultrasound guided technique of coeliac plexus block for treatment of pain in chronic pancreatitis. *Aliment Pharmacol Ther* 2009; **29**: 979-984 [PMID:

- 19222416 DOI: [10.1111/j.1365-2036.2009.03963.x](https://doi.org/10.1111/j.1365-2036.2009.03963.x)]
- 37 **Sahai AV**, Lemelin V, Lam E, Paquin SC. Central vs. bilateral endoscopic ultrasound-guided celiac plexus block or neurolysis: a comparative study of short-term effectiveness. *Am J Gastroenterol* 2009; **104**: 326-329 [PMID: [19174816](https://pubmed.ncbi.nlm.nih.gov/19174816/) DOI: [10.1038/ajg.2008.64](https://doi.org/10.1038/ajg.2008.64)]
 - 38 **Ahmed Ali U**, Pahlplatz JM, Nealon WH, van Goor H, Gooszen HG, Boermeester MA. Endoscopic or surgical intervention for painful obstructive chronic pancreatitis. *Cochrane Database Syst Rev* 2015; CD007884 [PMID: [25790326](https://pubmed.ncbi.nlm.nih.gov/25790326/) DOI: [10.1002/14651858.CD007884.pub3](https://doi.org/10.1002/14651858.CD007884.pub3)]
 - 39 **Jawad ZAR**, Kyriakides C, Pai M, Wadsworth C, Westaby D, Vlavianos P, Jiao LR. Surgery remains the best option for the management of pain in patients with chronic pancreatitis: A systematic review and meta-analysis. *Asian J Surg* 2017; **40**: 179-185 [PMID: [26778832](https://pubmed.ncbi.nlm.nih.gov/26778832/) DOI: [10.1016/j.asjsur.2015.09.005](https://doi.org/10.1016/j.asjsur.2015.09.005)]
 - 40 **Ma KW**, So H, Shin E, Mok JHM, Yuen KHK, Cheung TT, Park DH. Endoscopic versus Surgical Intervention for Painful Obstructive Chronic Pancreatitis: A Systematic Review and Meta-Analysis. *J Clin Med* 2021; **10** [PMID: [34203858](https://pubmed.ncbi.nlm.nih.gov/34203858/) DOI: [10.3390/jcm10122636](https://doi.org/10.3390/jcm10122636)]
 - 41 **Link G. V.** The Treatment of Chronic Pancreatitis by Pancreatostomy: A New Operation. *Ann Surg* 1911; **53**: 768-782 [PMID: [17862691](https://pubmed.ncbi.nlm.nih.gov/17862691/) DOI: [10.1097/00000658-191106000-00005](https://doi.org/10.1097/00000658-191106000-00005)]
 - 42 **Desjardins A.** Technique de la pancreatectomy. *Rev Chir Paris* 1907; **35**: 945-957
 - 43 **Coffey RC.** XVII. Pancreato-enterostomy and Pancreatectomy: A Preliminary Report. *Ann Surg* 1909; **50**: 1238-1264 [PMID: [17862458](https://pubmed.ncbi.nlm.nih.gov/17862458/) DOI: [10.1097/00000658-190912000-00017](https://doi.org/10.1097/00000658-190912000-00017)]
 - 44 **Duval MK Jr.** Caudal pancreatico-jejunosomy for chronic relapsing pancreatitis. *Ann Surg* 1954; **140**: 775-785 [PMID: [13208131](https://pubmed.ncbi.nlm.nih.gov/13208131/) DOI: [10.1097/00000658-195412000-00001](https://doi.org/10.1097/00000658-195412000-00001)]
 - 45 **Zollinger RM**, KEITH LM Jr, ELLISON EH. Pancreatitis. *N Engl J Med* 1954; **251**: 497-502 [PMID: [13203731](https://pubmed.ncbi.nlm.nih.gov/13203731/) DOI: [10.1056/NEJM195409232511301](https://doi.org/10.1056/NEJM195409232511301)]
 - 46 **Puestow CB**, GILLESBY WJ. Retrograde surgical drainage of pancreas for chronic relapsing pancreatitis. *AMA Arch Surg* 1958; **76**: 898-907 [PMID: [13532132](https://pubmed.ncbi.nlm.nih.gov/13532132/) DOI: [10.1001/archsurg.1958.01280240056009](https://doi.org/10.1001/archsurg.1958.01280240056009)]
 - 47 **Partington PF**, ROCHELLE RE. Modified Puestow procedure for retrograde drainage of the pancreatic duct. *Ann Surg* 1960; **152**: 1037-1043 [PMID: [13733040](https://pubmed.ncbi.nlm.nih.gov/13733040/) DOI: [10.1097/00000658-196012000-00015](https://doi.org/10.1097/00000658-196012000-00015)]
 - 48 **Bradley EL 3rd.** Long-term results of pancreatojejunosomy in patients with chronic pancreatitis. *Am J Surg* 1987; **153**: 207-213 [PMID: [3812895](https://pubmed.ncbi.nlm.nih.gov/3812895/) DOI: [10.1016/0002-9610\(87\)90816-6](https://doi.org/10.1016/0002-9610(87)90816-6)]
 - 49 **Mannell A**, Adson MA, McIlrath DC, Ilstrup DM. Surgical management of chronic pancreatitis: long-term results in 141 patients. *Br J Surg* 1988; **75**: 467-472 [PMID: [3390681](https://pubmed.ncbi.nlm.nih.gov/3390681/) DOI: [10.1002/bjs.1800750522](https://doi.org/10.1002/bjs.1800750522)]
 - 50 **Taylor RH**, Bagley FH, Braasch JW, Warren KW. Ductal drainage or resection for chronic pancreatitis. *Am J Surg* 1981; **141**: 28-33 [PMID: [7457724](https://pubmed.ncbi.nlm.nih.gov/7457724/) DOI: [10.1016/0002-9610\(81\)90007-6](https://doi.org/10.1016/0002-9610(81)90007-6)]
 - 51 **Fry WJ**, Child CG 3rd. Ninety-five per cent distal pancreatectomy for chronic pancreatitis. *Ann Surg* 1965; **162**: 543-549 [PMID: [5833584](https://pubmed.ncbi.nlm.nih.gov/5833584/) DOI: [10.1097/00000658-196510000-00001](https://doi.org/10.1097/00000658-196510000-00001)]
 - 52 **Beger HG**, Witte C, Krautzberger W, Bittner R. [Experiences with duodenum-sparing pancreas head resection in chronic pancreatitis]. *Chirurg* 1980; **51**: 303-307 [PMID: [7408575](https://pubmed.ncbi.nlm.nih.gov/7408575/)]
 - 53 **Warren WD**, Millikan WJ Jr, Henderson JM, Hersh T. A denervated pancreatic flap for control of chronic pain in pancreatitis. *Surg Gynecol Obstet* 1984; **159**: 581-583 [PMID: [6505945](https://pubmed.ncbi.nlm.nih.gov/6505945/)]
 - 54 **Frey CF**, Smith GJ. Description and rationale of a new operation for chronic pancreatitis. *Pancreas* 1987; **2**: 701-707 [PMID: [3438308](https://pubmed.ncbi.nlm.nih.gov/3438308/) DOI: [10.1097/00006676-198711000-00014](https://doi.org/10.1097/00006676-198711000-00014)]
 - 55 **Izbicki JR**, Yekebas EF. Chronic pancreatitis--lessons learned. *Br J Surg* 2005; **92**: 1185-1186 [PMID: [16175520](https://pubmed.ncbi.nlm.nih.gov/16175520/) DOI: [10.1002/bjs.5176](https://doi.org/10.1002/bjs.5176)]
 - 56 **Gloor B**, Friess H, Uhl W, Büchler MW. A modified technique of the Beger and Frey procedure in patients with chronic pancreatitis. *Dig Surg* 2001; **18**: 21-25 [PMID: [11244255](https://pubmed.ncbi.nlm.nih.gov/11244255/) DOI: [10.1159/000050092](https://doi.org/10.1159/000050092)]
 - 57 **Ho HS**, Frey CF. The Frey procedure: local resection of pancreatic head combined with lateral pancreaticojejunostomy. *Arch Surg* 2001; **136**: 1353-1358 [PMID: [11735858](https://pubmed.ncbi.nlm.nih.gov/11735858/) DOI: [10.1001/archsurg.136.12.1353](https://doi.org/10.1001/archsurg.136.12.1353)]
 - 58 **Frey CF**, Amikura K. Local resection of the head of the pancreas combined with longitudinal pancreaticojejunostomy in the management of patients with chronic pancreatitis. *Ann Surg* 1994; **220**: 492-504; discussion 504 [PMID: [7524454](https://pubmed.ncbi.nlm.nih.gov/7524454/) DOI: [10.1007/BF02348284](https://doi.org/10.1007/BF02348284)]
 - 59 **Sakata N**, Egawa S, Motoi F, Goto M, Matsuno S, Katayose Y, Unno M. How much of the pancreatic head should we resect in Frey's procedure? *Surg Today* 2009; **39**: 120-127 [PMID: [19198989](https://pubmed.ncbi.nlm.nih.gov/19198989/) DOI: [10.1007/s00595-008-3816-5](https://doi.org/10.1007/s00595-008-3816-5)]
 - 60 **Tan CL**, Zhang H, Yang M, Li SJ, Liu XB, Li KZ. Role of original and modified Frey's procedures in chronic pancreatitis. *World J Gastroenterol* 2016; **22**: 10415-10423 [PMID: [28058022](https://pubmed.ncbi.nlm.nih.gov/28058022/) DOI: [10.3748/wjg.v22.i47.10415](https://doi.org/10.3748/wjg.v22.i47.10415)]
 - 61 **Zhao X**, Cui N, Wang X, Cui Y. Surgical strategies in the treatment of chronic pancreatitis: An updated systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore)* 2017; **96**: e6220 [PMID: [28248878](https://pubmed.ncbi.nlm.nih.gov/28248878/) DOI: [10.1097/MD.00000000000006220](https://doi.org/10.1097/MD.00000000000006220)]
 - 62 **Hamad A**, Zenati MS, Nguyen TK, Hogg ME, Zeh HJ 3rd, Zureikat AH. Safety and feasibility of the robotic platform in the management of surgical sequelae of chronic pancreatitis. *Surg Endosc* 2018; **32**: 1056-1065 [PMID: [29273874](https://pubmed.ncbi.nlm.nih.gov/29273874/) DOI: [10.1007/s00464-017-6010-2](https://doi.org/10.1007/s00464-017-6010-2)]
 - 63 **Galvez D**, Sorber R, Javed AA, He J. Technical considerations for the fully robotic pancreaticoduodenectomy. *J Vis Surg* 2017; **3**: 81 [PMID: [29078644](https://pubmed.ncbi.nlm.nih.gov/29078644/) DOI: [10.21037/jovs.2017.05.08](https://doi.org/10.21037/jovs.2017.05.08)]
 - 64 **Giannini A**, D'Oria O, Chiantera V, Margioulas-Siakou C, Di Donna MC, Terzic S, Sleiman Z, Laganà AS. Minimally Invasive Surgery for Cervical Cancer: Should We Look beyond Squamous Cell Carcinoma? *J Invest Surg* 2022; **35**: 1602-1603 [PMID: [35549629](https://pubmed.ncbi.nlm.nih.gov/35549629/) DOI: [10.1080/08941939.2022.2075495](https://doi.org/10.1080/08941939.2022.2075495)]
 - 65 **Cuschieri A.** Laparoscopy for pancreatic cancer: does it benefit the patient? *Eur J Surg Oncol* 1988; **14**: 41-44 [PMID: [2964382](https://pubmed.ncbi.nlm.nih.gov/2964382/)]
 - 66 **Pietrabissa A**, Caramella D, Di Candio G, Carobbi A, Boggi U, Rossi G, Mosca F. Laparoscopy and laparoscopic ultrasonography for staging pancreatic cancer: critical appraisal. *World J Surg* 1999; **23**: 998-1002; discussion 1003

- [PMID: [10512938](#) DOI: [10.1007/s002689900614](#)]
- 67 **Gagner M**, Pomp A. Laparoscopic pylorus-preserving pancreatoduodenectomy. *Surg Endosc* 1994; **8**: 408-410 [PMID: [7915434](#) DOI: [10.1007/BF00642443](#)]
 - 68 **Cuschieri A**. Laparoscopic surgery of the pancreas. *J R Coll Surg Edinb* 1994; **39**: 178-184 [PMID: [7932341](#)]
 - 69 **Cuschieri A**, Jakimowicz JJ, van Spreuwel J. Laparoscopic distal 70% pancreatectomy and splenectomy for chronic pancreatitis. *Ann Surg* 1996; **223**: 280-285 [PMID: [8604908](#) DOI: [10.1097/00000658-199603000-00008](#)]
 - 70 **Giulianotti PC**, Coratti A, Angelini M, Sbrana F, Cecconi S, Balestracci T, Caravaglios G. Robotics in general surgery: personal experience in a large community hospital. *Arch Surg* 2003; **138**: 777-784 [PMID: [12860761](#) DOI: [10.1001/archsurg.138.7.777](#)]
 - 71 **Zureikat AH**, Moser AJ, Boone BA, Bartlett DL, Zenati M, Zeh HJ 3rd. 250 robotic pancreatic resections: safety and feasibility. *Ann Surg* 2013; **258**: 554-9; discussion 559 [PMID: [24002300](#) DOI: [10.1097/SLA.0b013e3182a4e87c](#)]
 - 72 **Kurian MS**, Gagner M. Laparoscopic side-to-side pancreaticojejunostomy (Partington-Rochelle) for chronic pancreatitis. *J Hepatobiliary Pancreat Surg* 1999; **6**: 382-386 [PMID: [10664286](#) DOI: [10.1007/s005340050135](#)]
 - 73 **Tantia O**, Jindal MK, Khanna S, Sen B. Laparoscopic lateral pancreaticojejunostomy: our experience of 17 cases. *Surg Endosc* 2004; **18**: 1054-1057 [PMID: [15156382](#) DOI: [10.1007/s00464-003-9210-x](#)]
 - 74 **Palanivelu C**, Shetty R, Jani K, Rajan PS, Sendhilkumar K, Parthasarathi R, Malladi V. Laparoscopic lateral pancreaticojejunostomy: a new remedy for an old ailment. *Surg Endosc* 2006; **20**: 458-461 [PMID: [16424983](#) DOI: [10.1007/s00464-005-0680-x](#)]
 - 75 **Khaled YS**, Ammori BJ. Laparoscopic lateral pancreaticojejunostomy and laparoscopic Berne modification of Beger procedure for the treatment of chronic pancreatitis: the first UK experience. *Surg Laparosc Endosc Percutan Tech* 2014; **24**: e178-e182 [PMID: [24710258](#) DOI: [10.1097/SLE.0b013e31829ce803](#)]
 - 76 **Sahoo MR**, Kumar A. Laparoscopic longitudinal pancreaticojejunostomy using cystoscope and endoscopic basket for clearance of head and tail stones. *Surg Endosc* 2014; **28**: 2499-2503 [PMID: [24962852](#) DOI: [10.1007/s00464-013-3330-8](#)]
 - 77 **Kim EY**, Hong TH. Laparoscopic Longitudinal Pancreaticojejunostomy Using Barbed Sutures: an Efficient and Secure Solution for Pancreatic Duct Obstructions in Patients with Chronic Pancreatitis. *J Gastrointest Surg* 2016; **20**: 861-866 [PMID: [26691145](#) DOI: [10.1007/s11605-015-3053-3](#)]
 - 78 **Bhandarwar A**, Arora E, Gajbhiye R, Gandhi S, Patel C, Wagh A, Kothari P, Jadhav S. Laparoscopic lateral pancreaticojejunostomy: an evolution to endostapled technique. *Surg Endosc* 2019; **33**: 1749-1756 [PMID: [30194645](#) DOI: [10.1007/s00464-018-6434-3](#)]
 - 79 **Rege S**, Singh A, Rewatkar A, Murugan J, Menezes R, Surpam S, Chiranjeev R. Laparoscopic parastomal hernia repair: A modified technique of mesh placement in Sugarbaker procedure. *J Minim Access Surg* 2019; **15**: 224-228 [PMID: [29794357](#) DOI: [10.4103/jmas.JMAS_17_18](#)]
 - 80 **Javed A**, Kiran SBD, Aravinda PS, Saravanan MN, Agarwal AK. Laparoscopic lateral pancreatico-jejunostomy: an experience from a tertiary care center. *Trop Gastroenterol* 2020; **41**: 66-72 [DOI: [10.7869/TG.581](#)]
 - 81 **Nag HH**, Nekarakanti PK, Arvinda PS, Sharma A. Laparoscopic versus open surgical management of patients with chronic pancreatitis: Amatched case-control study. *J Minim Access Surg* 2022; **18**: 191-196 [PMID: [33885009](#) DOI: [10.4103/jmas.JMAS_183_20](#)]
 - 82 **Criss CN**, Gadepalli SK. Sponsoring surgeons: An investigation on the influence of the da Vinci robot. *Am J Surg* 2018; **216**: 84-87 [PMID: [28859920](#) DOI: [10.1016/j.amjsurg.2017.08.017](#)]
 - 83 **Meehan JJ**, Sawin R. Robotic lateral pancreaticojejunostomy (Puestow). *J Pediatr Surg* 2011; **46**: e5-e8 [PMID: [21683190](#) DOI: [10.1016/j.jpedsurg.2011.03.003](#)]
 - 84 **Eid GM**, Entabi F, Watson AR, Zuckerbraun BS, Wilson MA. Robotic-assisted laparoscopic side-to-side lateral pancreaticojejunostomy. *J Gastrointest Surg* 2011; **15**: 1243 [PMID: [21584825](#) DOI: [10.1007/s11605-011-1495-9](#)]
 - 85 **Kirks RC**, Lorimer PD, Fruscione M, Cochran A, Baker EH, Iannitti DA, Vrochides D, Martinie JB. Robotic longitudinal pancreaticojejunostomy for chronic pancreatitis: Comparison of clinical outcomes and cost to the open approach. *Int J Med Robot* 2017; **13** [PMID: [28548233](#) DOI: [10.1002/rcs.1832](#)]
 - 86 **Khan AS**, Siddiqui I, Vrochides D, Martinie JB. Robotic pancreas drainage procedure for chronic pancreatitis: robotic lateral pancreaticojejunostomy (Puestow procedure). *J Vis Surg* 2018; **4**: 72 [PMID: [29780718](#) DOI: [10.21037/jovs.2017.10.10](#)]
 - 87 **Balduzzi A**, Zwart MJW, Kempeneers RMA, Boermeester MA, Busch OR, Besselink MG. Robotic Lateral Pancreaticojejunostomy for Chronic Pancreatitis. *J Vis Exp* 2019 [PMID: [31885388](#) DOI: [10.3791/60301](#)]
 - 88 **Tan CL**, Zhang H, Li KZ. Single center experience in selecting the laparoscopic Frey procedure for chronic pancreatitis. *World J Gastroenterol* 2015; **21**: 12644-12652 [PMID: [26640341](#) DOI: [10.3748/wjg.v21.i44.12644](#)]
 - 89 **Kilburn DJ**, Chiow AKH, Leung U, Siriwardhane M, Cavallucci DJ, Bryant R, O'Rourke NA. Early Experience with Laparoscopic Frey Procedure for Chronic Pancreatitis: a Case Series and Review of Literature. *J Gastrointest Surg* 2017; **21**: 904-909 [PMID: [28025771](#) DOI: [10.1007/s11605-016-3343-4](#)]
 - 90 **Senthilnathan P**, Subrahmaneswara Babu N, Vikram A, Sabnis SC, Srivatsan Gurumurthy S, Anand Vijai N, Nalankilli VP, Palanivelu C. Laparoscopic longitudinal pancreaticojejunostomy and modified Frey's operation for chronic calcific pancreatitis. *BJS Open* 2019; **3**: 666-671 [PMID: [31592076](#) DOI: [10.1002/bjs.5.50185](#)]
 - 91 **Zureikat AH**, Nguyen KT, Bartlett DL, Zeh HJ, Moser AJ. Robotic-assisted major pancreatic resection and reconstruction. *Arch Surg* 2011; **146**: 256-261 [PMID: [21079111](#) DOI: [10.1001/archsurg.2010.246](#)]
 - 92 **Shukla A**, Gnanasekaran S, Kalayarasan R, Pottakkat B. Early experience with robot-assisted Frey's procedure surgical outcome and technique: Indian perspective. *J Minim Invasive Surg* 2022; **25**: 145-151 [PMID: [36601487](#) DOI: [10.7602/jmis.2022.25.4.145](#)]
 - 93 **Izrailov RE**, Andrianov AV. Laparoscopic Beger procedure. *HPB* 2018; **20**: S118 [DOI: [10.1016/j.hpb.2018.02.096](#)]
 - 94 **Cai H**, Cai Y, Wang X, Peng B. Laparoscopic Beger procedure for the treatment of chronic pancreatitis: a single-centre first experience. *BMC Surg* 2020; **20**: 84 [PMID: [32349723](#) DOI: [10.1186/s12893-020-00750-7](#)]
 - 95 **Hong D**, Cheng J, Wu W, Liu X, Zheng X. How to Perform Total Laparoscopic Duodenum-Preserving Pancreatic Head

- Resection Safely and Efficiently with Innovative Techniques. *Ann Surg Oncol* 2021; **28**: 3209-3216 [PMID: 33123857 DOI: 10.1245/s10434-020-09233-8]
- 96 **Strobel O**, Büchler MW, Werner J. [Duodenum-preserving pancreatic head resection: technique according to Beger, technique according to Frey and Berne modifications]. *Chirurg* 2009; **80**: 22-27 [PMID: 18820883 DOI: 10.1007/s00104-008-1577-8]
- 97 **Cai Y**, Zheng Z, Gao P, Li Y, Peng B. Laparoscopic duodenum-preserving total pancreatic head resection using real-time indocyanine green fluorescence imaging. *Surg Endosc* 2021; **35**: 1355-1361 [PMID: 32221750 DOI: 10.1007/s00464-020-07515-6]
- 98 **Peng CH**, Shen BY, Deng XX, Zhan Q, Han B, Li HW. Early experience for the robotic duodenum-preserving pancreatic head resection. *World J Surg* 2012; **36**: 1136-1141 [PMID: 22415757 DOI: 10.1007/s00268-012-1503-6]
- 99 **Jiang Y**, Jin JB, Zhan Q, Deng XX, Peng CH, Shen BY. Robot-assisted duodenum-preserving pancreatic head resection with pancreaticogastrostomy for benign or premalignant pancreatic head lesions: a single-centre experience. *Int J Med Robot* 2018; **14**: e1903 [PMID: 29498195 DOI: 10.1002/ics.1903]
- 100 **Abu-El-Haija M**, Anazawa T, Beilman GJ, Besselink MG, Del Chiaro M, Demir IE, Dennison AR, Dudeja V, Freeman ML, Friess H, Hackert T, Kleeff J, Laukkarinen J, Levy MF, Nathan JD, Werner J, Windsor JA, Neoptolemos JP, Sheel ARG, Shimosegawa T, Whitcomb DC, Bellin MD. The role of total pancreatectomy with islet autotransplantation in the treatment of chronic pancreatitis: A report from the International Consensus Guidelines in chronic pancreatitis. *Pancreatolgy* 2020; **20**: 762-771 [PMID: 32327370 DOI: 10.1016/j.pan.2020.04.005]
- 101 **Fan CJ**, Hirose K, Walsh CM, Quartuccio M, Desai NM, Singh VK, Kalyani RR, Warren DS, Sun Z, Hanna MN, Makary MA. Laparoscopic Total Pancreatectomy With Islet Autotransplantation and Intraoperative Islet Separation as a Treatment for Patients With Chronic Pancreatitis. *JAMA Surg* 2017; **152**: 550-556 [PMID: 28241234 DOI: 10.1001/jamasurg.2016.5707]
- 102 **Galvani CA**, Rodriguez Rilo H, Samamé J, Porubsky M, Rana A, Gruessner RW. Fully robotic-assisted technique for total pancreatectomy with an autologous islet transplant in chronic pancreatitis patients: results of a first series. *J Am Coll Surg* 2014; **218**: e73-e78 [PMID: 24559970 DOI: 10.1016/j.jamcollsurg.2013.12.006]
- 103 **Blair AB**, Burkhart RA, Hirose K, Makary MA. Laparoscopic total pancreatectomy with islet autotransplantation for chronic pancreatitis. *J Vis Surg* 2016; **2**: 121 [PMID: 29321981 DOI: 10.21037/jovs.2016.07.05]
- 104 **Zureikat AH**, Nguyen T, Boone BA, Wijkstrom M, Hogg ME, Humar A, Zeh H 3rd. Robotic total pancreatectomy with or without autologous islet cell transplantation: replication of an open technique through a minimal access approach. *Surg Endosc* 2015; **29**: 176-183 [PMID: 25005012 DOI: 10.1007/s00464-014-3656-x]
- 105 **Nathan JD**, Yang Y, Eaton A, Witkowski P, Wijkstrom M, Walsh M, Trikudanathan G, Singh VK, Schwarzenberg SJ, Pruett TL, Posselt A, Naziruddin B, Mokshagundam SP, Morgan K, Lara LF, Kirchner V, He J, Gardner TB, Freeman ML, Ellery K, Conwell DL, Chinnakotla S, Beilman GJ, Ahmad S, Abu-El-Haija M, Hodges JS, Bellin MD. Surgical approach and short-term outcomes in adults and children undergoing total pancreatectomy with islet autotransplantation: A report from the Prospective Observational Study of TPIAT. *Pancreatolgy* 2022; **22**: 1-8 [PMID: 34620552 DOI: 10.1016/j.pan.2021.09.011]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

