World Journal of *Clinical Cases*

World J Clin Cases 2023 July 6; 11(19): 4458-4733





Published by Baishideng Publishing Group Inc

W J C C World Journal of Clinical Cases

Contents

Thrice Monthly Volume 11 Number 19 July 6, 2023

REVIEW

4458 Molecular signalling during cross talk between gut brain axis regulation and progression of irritable bowel syndrome: A comprehensive review

Singh SV, Ganguly R, Jaiswal K, Yadav AK, Kumar R, Pandey AK

4477 Diffusion tensor imaging in the courtroom: Distinction between scientific specificity and legally admissible evidence

van Velkinburgh JC, Herbst MD, Casper SM

MINIREVIEWS

4498 Inequity in the global distribution of monkeypox vaccines

Tovani-Palone MR, Doshi N, Pedersini P

4504 Long-term effectiveness, outcomes and complications of bariatric surgery

> Gulinac M, Miteva DG, Peshevska-Sekulovska M, Novakov IP, Antovic S, Peruhova M, Snegarova V, Kabakchieva P, Assyov Y, Vasilev G, Sekulovski M, Lazova S, Tomov L, Velikova T

ORIGINAL ARTICLE

Retrospective Cohort Study

4513 Age, blood tests and comorbidities and AIMS65 risk scores outperform Glasgow-Blatchford and preendoscopic Rockall score in patients with upper gastrointestinal bleeding

Morarasu BC, Sorodoc V, Haisan A, Morarasu S, Bologa C, Haliga RE, Lionte C, Marciuc EA, Elsiddig M, Cimpoesu D, Dimofte GM, Sorodoc L

Retrospective Study

4531 Application of cross-migration theory in limb rehabilitation of stroke patients with hemiplegia

Lu YH, Fu Y, Shu J, Yan LY, Shen HJ

4544 Analysis of characteristic features in ultrasound diagnosis of fetal limb body wall complex during 11-13⁺⁶ weeks

Ye CH, Li S, Ling L

Network pharmacology and molecular docking-based analyses to predict the potential mechanism of 4553 Huangqin decoction in treating colorectal cancer

Li YJ, Tang DX, Yan HT, Yang B, Yang Z, Long FX

4567 Assessment of functional prognosis of anterior cruciate ligament reconstruction in athletes based on a body shape index

Wang YJ, Zhang JC, Zhang YZ, Liu YH



World Journal of Clinical Cases

Contents

Thrice Monthly Volume 11 Number 19 July 6, 2023

EVIDENCE-BASED MEDICINE

Network pharmacology and molecular docking to explore Polygoni Cuspidati Rhizoma et Radix treatment 4579 for acute lung injury

Zheng JL, Wang X, Song Z, Zhou P, Zhang GJ, Diao JJ, Han CE, Jia GY, Zhou X, Zhang BQ

ORIGINAL ARTICLE

Randomized Controlled Trial

4601 Ulinastatin in the treatment of severe acute pancreatitis: A single-center randomized controlled trial

Wang SQ, Jiao W, Zhang J, Zhang JF, Tao YN, Jiang Q, Yu F

Fecal microbiota transplantation in patients with metabolic syndrome and obesity: A randomized 4612 controlled trial

da Ponte Neto AM, Clemente ACO, Rosa PW, Ribeiro IB, Funari MP, Nunes GC, Moreira L, Sparvoli LG, Cortez R, Taddei CR, Mancini MC, de Moura EGH

SYSTEMATIC REVIEWS

4625 Combined medial patellofemoral ligament and medial patellotibial ligament reconstruction in recurrent patellar instability: A systematic review and meta-analysis

Abbaszadeh A, Saeedi M, Hoveidaei AH, Dadgostar H, Razi S, Razi M

CASE REPORT

4635 Unique Roberts syndrome with bilateral congenital glaucoma: A case report

Almulhim A, Almoallem B, Alsirrhy E, Osman EA

4640 CK5/6-positive, P63-positive lymphoepithelioma-like hepatocellular carcinoma: A case report and literature review

Tang HT, Lin W, Zhang WQ, Qian JL, Li K, He K

4648 Edaravone administration and its potential association with a new clinical syndrome in cerebral infarction patients: Three case reports

Yang L, Xu X, Wang L, Zeng KB, Wang XF

- 4655 CDKN1C gene mutation causing familial Silver-Russell syndrome: A case report and review of literature Li J, Chen LN, He HL
- Hypothetical hypoxia-driven rapid disease progression in hepatocellular carcinoma post transarterial 4664 chemoembolization: A case report

Yeo KF, Ker A, Kao PE, Wang CC

4670 Metastatic colon cancer treated using traditional Chinese medicine combined with chemotherapy: A case report

Deng CG, Tang MY, Pan X, Liu ZH

4677 Rare cause of cerebral venous sinus thrombosis: Spontaneous intracranial hypotension syndrome: A case report

Huang P



0	World Journal of Clinical Cases
Conten	Thrice Monthly Volume 11 Number 19 July 6, 2023
4684	Integrated Chinese and Western medicine in the treatment of a patient with podocyte infolding glomerulopathy: A case report
	Chang MY, Zhang Y, Li MX, Xuan F
4692	Morbihan disease misdiagnosed as senile blepharoptosis and successfully treated with short-term minocycline and ketotifen: A case report
	Na J, Wu Y
4698	With two episodes of right retromandibular angle subcutaneous emphysema during right upper molar crown preparation: A case report
	Bai YP, Sha JJ, Chai CC, Sun HP
4707	Poststroke rehabilitation using repetitive transcranial magnetic stimulation during pregnancy: A case report
	Jo J, Kim H
4713	Tuberculosis-induced aplastic crisis and atypical lymphocyte expansion in advanced myelodysplastic syndrome: A case report and review of literature
	Sun XY, Yang XD, Xu J, Xiu NN, Ju B, Zhao XC
4723	Posterior reversible encephalopathy syndrome following uneventful clipping of an unruptured intracranial aneurysm: A case report
	Hwang J, Cho WH, Cha SH, Ko JK

ACADEMIC WRITING

4729 Revitalizing case reports: Standardized guidelines and mentorship

Jeyaraman M, Ramasubramanian S, Jeyaraman N, Nallakumarasamy A, Sharma S



Contents

Thrice Monthly Volume 11 Number 19 July 6, 2023

ABOUT COVER

Editorial Board Member of World Journal of Clinical Cases, Miroslav Vujasinovic, MD, PhD, Associate Professor, Department of Upper Abdominal Diseases, Karolinska University Hospital, Stockholm 14186, Sweden. mvujas@gmail.com

AIMS AND SCOPE

The primary aim of World Journal of Clinical Cases (WJCC, World J Clin Cases) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Clinical Cases	https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 2307-8960 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
April 16, 2013	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Thrice Monthly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku	PUBLICATION MISCONDUCT https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/2307-8960/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
July 6, 2023	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2023 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2023 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: bpgoffice@wjgnet.com https://www.wjgnet.com



W J C C World Journal of Clinical Cases

Submit a Manuscript: https://www.f6publishing.com

World J Clin Cases 2023 July 6; 11(19): 4458-4476

DOI: 10.12998/wjcc.v11.i19.4458

ISSN 2307-8960 (online)

REVIEW

Molecular signalling during cross talk between gut brain axis regulation and progression of irritable bowel syndrome: A comprehensive review

Shiv Vardan Singh, Risha Ganguly, Kritika Jaiswal, Aditya Kumar Yadav, Ramesh Kumar, Abhay K Pandey

Specialty type: Pharmacology and pharmacy

Provenance and peer review: Invited article; Externally peer reviewed

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): A Grade B (Very good): 0 Grade C (Good): C Grade D (Fair): D Grade E (Poor): 0

P-Reviewer: Losurdo G, Italy; Yu YB, China

Received: December 24, 2022 Peer-review started: December 24, 2022 First decision: March 23, 2023 Revised: May 9, 2023 Accepted: June 6, 2023 Article in press: June 6, 2023 Published online: July 6, 2023



Shiv Vardan Singh, Risha Ganguly, Kritika Jaiswal, Aditya Kumar Yadav, Ramesh Kumar, Abhay K Pandey, Department of Biochemistry, University of Allahabad, Allahabad (Prayagraj) 211002, Uttar Pradesh, India

Corresponding author: Abhay K Pandey, PhD, Professor, Department of Biochemistry, University of Allahabad, University Road, Allahabad (Prayagraj) 211002, Uttar Pradesh, India. akpandey23@rediffmail.com

Abstract

Irritable bowel syndrome (IBS) is a chronic functional disorder which alters gastrointestinal (GI) functions, thus leading to compromised health status. Pathophysiology of IBS is not fully understood, whereas abnormal gut brain axis (GBA) has been identified as a major etiological factor. Recent studies are suggestive for visceral hyper-sensitivity, altered gut motility and dysfunctional autonomous nervous system as the main clinical abnormalities in IBS patients. Bidirectional signalling interactions among these abnormalities are derived through various exogenous and endogenous factors, such as microbiota population and diversity, microbial metabolites, dietary uptake, and psychological abnormalities. Strategic efforts focused to study these interactions including probiotics, antibiotics and fecal transplantations in normal and germfree animals are clearly suggestive for the pivotal role of gut microbiota in IBS etiology. Additionally, neurotransmitters act as communication tools between enteric microbiota and brain functions, where serotonin (5-hydroxytryptamine) plays a key role in pathophysiology of IBS. It regulates GI motility, pain sense and inflammatory responses particular to mucosal and brain activity. In the absence of a better understanding of various interconnected crosstalks in GBA, more scientific efforts are required in the search of novel and targeted therapies for the management of IBS. In this review, we have summarized the gut microbial composition, interconnected signalling pathways and their regulators, available therapeutics, and the gaps needed to fill for a better management of IBS.

Key Words: Irritable bowel syndrome; Microbiota; Gut brain axis; Stress; Serotonin

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Irritable bowel syndrome (IBS) is a prevalent gastrointestinal disorder with a dysregulated gut brain communication. Gut microbiota functional characterization is still underappreciated but their roles have been found to be pivotal. Various microbial species and their metabolites with altered composition and diversity have been found to be specific to IBS. Clinical manipulation of these microbial species improved the symptom profile in IBS patients while the associated mechanisms have been identified for a bidirectional communication between gut microbiota and brain. This in turn seems promising for future treatments specific to microbiota manipulation and targeting various cross-talks for the management of IBS and associated symptoms.

Citation: Singh SV, Ganguly R, Jaiswal K, Yadav AK, Kumar R, Pandey AK. Molecular signalling during cross talk between gut brain axis regulation and progression of irritable bowel syndrome: A comprehensive review. World J Clin Cases 2023; 11(19): 4458-4476

URL: https://www.wjgnet.com/2307-8960/full/v11/i19/4458.htm DOI: https://dx.doi.org/10.12998/wjcc.v11.i19.4458

INTRODUCTION

Human body consist of trillions of microbial cells, majority of which inhabit the gastrointestinal (GI) tract thus forming a microbial colonization with a dynamic ecological environment, commonly known as "microbiota" [1,2]. This microbiota is comprised of approximately 500 transient and indigenous species of bacteria, viruses, fungi and protozoa[3,4]. Among all, bacteria are the most abundant microbial community dominated with the members of Firmicutes and Bacteroidetes phyla[5,6]. In recent years, research focusing intestinal microbiota functional characterization led to the identification of a bidirectional crosstalk between brain and gut microbiota, thus forming a gut-brain axis (GBA)[7,8]. Moreover, GBA includes central nervous system (CNS), enteric nervous system (ENS), hypothalamicpituitary-adrenal (HPA) axis, gut and its microbiota (Figure 1). Interestingly, these components have been identified to be interconnected through various coordinated signalling pathways[8,9]. Abnormalities in these pathways and their regulators have been identified for the etiology of irritable bowel syndrome (IBS). IBS is a common GI bowel disorder featured with altered GI motility, visceral hypersensitivity, post-infection reactivity, small intestinal bacterial overload, carbohydrate malabsorption and intestinal inflammation[10,11]. These abnormalities result in dysbiosis, recurrent abdominal pain and distressed bowel habits.

IBS is affecting 10%-25% of global world population, especially in developed countries with a poorly deciphered pathophysiology^[12]. Considering the predominant symptoms and bowel habits, IBS is further classified into four subtypes: diarrhea-predominant IBS (IBS-D), constipation-predominant IBS (IBS-C), mixed IBS (IBS-M), and unclassified IBS[13,14]. Interestingly, women are more susceptible (1.67 times) than men for IBS. In recent years, IBS symptoms have been predominantly found to be associated with environmental factors such as, diet, enteric microbial communities, host genetics and psychology [15]. However, the ways in which these factors contribute to the etiology of IBS are not fully deciphered. The treatments available for IBS are also not very specific. Generally, dietary fiber supplementation for IBS-C, opioids for IBS-D and abdominal pain with bloating, and low doses of antidepressants are recommended for the management of various symptoms associated with IBS[12,13,15]. However, modulating the gut-brain axis seems to be a promising target for the development of novel therapeutic for IBS[16]. In addition to this, various metabolic disorders have also been identified to be massively regulated by gut microbiota and their metabolites[6,17,18]. Involvement of gut microbiota metabolites in maintaining homeostasis including host immunity, physiological functions (digestion and nutrition) and biosynthesis of vitamins have also been recently validated by various research groups [18,19]. Experimental sets of data intoning functional cruciality of microbiota in IBS advocate their widespread interactions not localized only with intestinal cells and ENS, but also directly with CNS through neuroendocrine and metabolic pathways [19,20]. The ENS composed of semiautonomous effector system is also connected to central autonomic network and modulated via afferent and efferent communications through parasympathetic and sympathetic nerves[21,22].

Ongoing bidirectional brain-gut interactions are significantly influenced by serotonergic [5-hydroxytryptamine (5-HT)] pathway, where serotonin also known as 5-HT is an important neurotransmitter and paracrine signalling molecule^[23,24]. Serotonin is synthesized by enterochromaffin cells (EC) of the gut and by serotonergic neurons in the CNS^[25]. Aberrant 5-HT signalling has been found to be accountable for various GI disorders, including IBS, diarrhea and chronic constipation, and functional dyspepsia[26, 27]. Various components of serotonin signalling, including EC cells count, serotonin level, tryptophan hydroxylase activity, and expression of serotonin-selective reuptake transporters have also been found to be altered in IBS[28]. 5-HT signalling in between 5-HT receptors (on postsynaptic and presynaptic neurons at CNS and intestinal serotonergic neurons) and serotonin transporters (SERT) of various cell



Singh SV et al. Interaction of gut-brain-axis and IBS progression



DOI: 10.12998/wjcc.v11.i19.4458 Copyright ©The Author(s) 2023.

Figure 1 Regulatory cross talk between gut and brain. The central nervous system and enteric nervous system mainly communicate through vagal and autonomic pathways to control various gastrointestinal functions in a bidirectional approach. These signalling cascades are also regulated through mood and cognitive processes and significantly affected by environmental factors and early adverse events. These signalling crosstalks are significantly altered in irritative bowel syndrome (IBS) thus resulting in abnormal gastrointestinal motility, visceral hypersensitivity and bowel movements. Perturbations in the gut microbiome composition and diversity through various sources alter this cross talk, thus resulting in interceptive feedback to brain and causing functional and neuroplastic changes. Microbial metabolites released from various microbial populations have critical role in these crosstalks, thus having therapeutic values for the management of IBS. ANS: Autonomic nervous system; HPA: Hypothalamic-pituitary-adrenal axis; ECC: Enterochromaffin cells; ICC: Interstitial cells of Cajal; SMC: Smooth muscle cells; SCFAs: Short chain fatty acids; ENS: Enteric nervous system.

> types of GI tract is crucial for proper functioning of gut-brain communication [29,30]. Additionally, 5-HT is produced by the chemical conversion of tryptophan to 5-hydroxytryptamine, a reaction catalyzed by enzyme tryptophan hydroxylase (TPH1 in EC cells and TPH2 in neurons)[31]. Stored into vesicles formed through the vesicular monoamine transporter (VMAT; VMAT1 in EC cells and VMAT2 in neurons), 5-HT is further released into the extracellular space where it binds to different serotonin receptors (5-HTR)[32]. Further, it is known to regulate peristaltic, secretory, vasodilatory, vagal and nociceptive reflexes (Figure 2)[26,27,33].

> The present review summarizes recent updates consisting key components of microbiota derived GBA, molecular signalling during cross talk between GBA regulation and in the progression of IBS. Herein, we have also discussed the diagnostic and therapeutic implications of microbiota in the management of IBS and related symptoms.

IBS

IBS is a GI disorder characterized by chronic abdominal pain and altered bowel habits in the absence of any organic cause which significantly affects the quality of life[34]. It is the most diagnosed functional GI disorder accounting for approximately 30 percent of all referrals to gastroenterologists^[35]. The pathophysiology of IBS is the result of aberrated and interconnected signalling networks, especially between gut microbiome and brain[34,35]. However, the associated mechanisms and pathways are still unclear.

Pathophysiology of IBS

Traditionally, IBS was known to be associated with altered GI motility, visceral hypersensitivity and distorted pain perception[34,35]. Studies on IBS indicated the pivotal role of inflammation, alterations in fecal flora, and bacterial overgrowth[36]. IBS is also known as a psychosomatic illness since it is associated with mood disorders with abnormal psychiatric conditions[36,37]. Furthermore, mucosal immune activation, inflammatory cells and elevated inflammatory markers have been recorded in IBS





DOI: 10.12998/wjcc.v11.i19.4458 Copyright ©The Author(s) 2023.

Figure 2 Serotonin mediated bidirectional cross talk between gut microbiota and brain. Serotonin (5-HT) plays a key role in microbiota-gut-brain communications to modulate gastrointestinal (GI) and central nervous system (CNS) functions. Serotonin action is mediated through various signalling mechanisms between 5-HT receptors located in postsynaptic and presynaptic neurons at CNS and intestinal serotonergic neurons, and in different cell types of GI tract. Serotonin is synthesized by enterochromaffin cells (EC) in the gut and serotonergic neurons in the CNS. Microbial associated molecular patterns from microbiota directly affect the serotonergic system, mainly through modulating the activity and expression of serotonin transporter (SERT) and serotonin receptors (5-HTRs), as well as the synthesis of 5-HT in GI tract. Stored into vesicles forms through the vesicular monoamine transporter (VMAT; VMAT1 in EC cells and VMAT2 in neurons), 5-HT is further released into the extracellular space where its binds to different serotonin receptors (5-HTR). At the same place taken up by the neurons, enterocytes or platelets through the SERT limit the 5-HT mediated signalling crosstalk. 5-HT: 5-Hydroxytryptamine; TLR: Toll-like receptors; NLR: NOD-like receptors; SFCA: Short-chain fatty acids; MAMP: Molecular patterns from microbiota; SERT: Serotonin transporter; VMAT: Vesicular monoamine transporter.

patients[34,35]. Various studies associated with the effect of antibiotics in changing the gut microbiota diversity and complexities are co-related with gut microbiota profiles and IBS symptoms[34,38]. Perturbations in GBA have been also proposed as the main mechanisms in the pathophysiology of IBS. In a recent study, corticotrophin-releasing hormone (CRH) was also found to have an important role in IBS and in augmentation of intestinal mucosal inflammation[39-41].

Role of stress in IBS

Studies are suggestive for co-morbidity of stress-related psychiatric illness and IBS, since 50%-60% of IBS patients have been diagnosed with various psychosocial health issues[42,43]. Psychological stress plays an important role in the pathophysiology of IBS, since it critically influences gut-brain axis and its associated metabolism[44]. It is also known to regulate intestinal motility and permeability, visceral hypersensitivity, immune responses, and gut microbiota composition[45,46]. Among the possible mechanisms for this regulation, immune response dependent secretion of various proinflammatory cytokines seems to be the pivotal one[47]. These cytokines activate the HPA and hypothalamic-autonomic nervous system (ANS) axes along with the release of corticotropin releasing factor, adreno-corticotropic hormone and cortisol, and all these subsequently control the gut homeostasis (Figure 3) [48]. It is established that these also alter neurotransmitter release within the enteric nervous system which thereafter affect gut motility, secretion and epithelial permeability *via* tight junction dysregulation[49]. An abnormal progression of monoamine neurotransmitter systems, including 5-HT and noradrenaline have been noticed in many stress related disorders. Furthermore, serotonin is known for its pivotal role in emotional response and in pain management by modulating brain-gut-microbiota axis, which is also significantly affected by early life stress[50].

Singh SV et al. Interaction of gut-brain-axis and IBS progression



DOI: 10.12998/wjcc.v11.i19.4458 Copyright ©The Author(s) 2023.

Figure 3 Impact of stress on irritable bowel syndrome. A bidirectional signalling has been identified between brain and gut, under stress condition in irritable bowel syndrome (IBS). Stress induces the activation of hypothalamic-pituitary-adrenal axis which results in the release of various stress factors such as corticotropin-releasing factor from hypothalamus, thus causing the release of glucocorticoids from adrenal glands. This collectively alters bowel function and stimulates the upregulation of immune system, which directly or indirectly regulates gut function. In addition to this, stress also modulates the tryptophan metabolism by up-regulating kynurenine pathway which generates neurotoxic and neuroprotective metabolites. Enzymes responsible for the degradation of tryptophan are significantly affected by immune pathways (indoleamine-2,3-dioxygenase) and stress levels (tryptophan-2,3-dioxygenase). Under stressed conditions and due to excess availability of kynurenine and its metabolites, the activation of these pathways lead to potential serotonergic deficiency along with altered enteric nervous system and central nervous system functionalities, thus leading to microbial dysbiosis, as a classical symptom of IBS. CRF: Corticotropin-releasing factor; ANS: Autonomic nervous system; CNS: Central nervous system; ACTH: Adrenocorticotropic hormone; IDO: Indoleamine-2,3-dioxygenase; TDO: Tryptophan-2,3dioxygenase.

MICROBIOTA AND GBA

Human GI tract hosts various genera of bacteria that are present on its mucosal surface. The term "microbiota" refers to the ecological system composed of various commensal bacteria in the GI tract, particularly localized on the lining of mucosal surface[51]. The term "microbiome" refers to metagenome of the microbiota and their genetic architecture^[52]. Mucosal surface is the only layer separating host tissues from various germ lines by which host immune system remains protective against various pathogens in GI tract[52,53].

Microbial colonization

Although, majority of microbiota is still uncharacterized, it consists of a very delicate community of commensal organisms that have evolved over millions of years. Among all, four bacterial phyla including gram-positive Firmicutes (Lactobacillus spp.) and Actinobacteria (Bifidobacterium spp.) along with gram-negative Bacteroides and Proteobacteria are the most prevalent[6,54,55]. In addition to bacterial strains, various viruses, protozoa, archae, and fungus also constitute GI microbiota. Multidisciplinary studies were also performed to characterize the gut microbiota, thus revealing the system as a state of symbiosis[56-59]. Homeostasis of host immune system and mucosal barrier are



generally altered by dysbiosis, which in turn promotes the invasion and growth of pathogenic species. Microbiota also regulates gut inflammation with the presence of immune cells in gut wall[60]. Although, the complete sketch of a balanced microbiome is still unknown and each person is assumed to have a distinct microbiota (called as microbiome fingerprint), healthy individuals are generally known to have similar profile and distribution of bacterial phylotypes^[61].

Microbial colonization is a vital early life phenomenon as a part of development of a healthy microbiota which seems to be crucial for the development of GBA[62,63]. Altered microbial colonization in human GI tract has been clearly identified to exhibit adverse health effects in the later years of the host[63]. The colonization process is also regulated with various factors such as birth mode and feeding of babies, since the breastfed and vaginally born babies were reported to be initially colonised by the member of Bifidobacteria, Lactobacilli, and Bacteroides species[64]. As compared to pre-term babies, fullterm babies are reported for rapid maturation of gut microbiome in their first year of life. There is several evidence supporting the existence of a microbiota and foetal meconium, where the colonisation is reported to start even before the delivery, which suggest that these organisms could also colonise the foetal gut[65]. Various bacteria like Staphylococcus, Streptococcus, Bifidobacterium, and Lactobacillus are also reported in human milk[66]. Thus, colonization is significantly affected in neonatal period by human breast milk and the technique of delivery[67]. Moreover, mononuclear cells carrying bacterial population from the mother's gut to the mammary gland through an entero-mammary pathway are also playing pivotal role in gut immunological maturation of the infant[68]. This mammary microbiome also possesses anti-infective, anti-inflammatory, immunomodulatory and metabolic activities which also constitute for microbiota colonization in neonatal gut[69,70].

Role of microbiome in GBA

The human microbiota play fundamental role in host physiology and pathology, and alteration in microbial population (dysbiosis) has been found to be significantly associated with various GI disorders [71]. A balanced interaction between microbiota and its host seems to be beneficial and essential for intestinal health and host metabolism. Under normal conditions, mucosal microbiota have been found to have significant role in digestion of food, vitamins synthesis, angiogenesis, epithelial cell and in development and maturation of the host defense system^[72]. Recently, it has been evident that the intestinal bacterial population significantly affects the CNS physiology and gut inflammation, mainly by a bidirectional communication consisting of various signalling pathways. This communication commonly known as GBA[73], consists of multiple inter-connections including vagus nerve, immune components, and microbial metabolites[74,75].

An enhanced size of microbiota and neuronal development takes place generally in starting five years of an infant, where the neuronal development is mainly configured by maternal microbiota [76]. The effect of gut microbiota on neurodevelopment of an infant was also examined and proved using various experiments on germ-free (GF) or specific pathogen-free mice. These mice were treated with various antibiotics to alter the microbial diversity and thereby to study their importance in gut microbiome[77, 78]. These experiments clearly suggested that various neurological problems appeared in the treated animals due to improper maturation of gut microbiome. Results of these studies are suggestive of exaggerated HPA axis in germ free animals with impaired social behaviors, reduced anxiety and increased motor activity [79,80]. In the same experiment, altered neuronal developments and behaviors were found to be improved when the newborn infants were supplemented with various microbial flora [81]. These altered behavioral phenotypes were also found to deregulate various genes and metabolites involved in HPA axis regulation. Among these, some common regulators are adrenaline, 5-hydroxytryptophan, postsynaptic density protein 95, dopamine and synaptophysin[82].

Scientific reports suggest that production of various neuroactive molecules by gut microbes directly or indirectly play critical role in GBA[83]. Among these neurotransmitters, acetylcholine, γ aminobutyric acid and serotonin are released by various bacterial populations belonging to Bifidobacteria, Lactobacillus, Enterococcus, and Streptococcus species[84]. Interestingly, 90% of total serotonin is produced by gut microbiota, and used for mood disorders and functional regulation of CNS and GI tract[84,85]. Serotonin is known to bind to 5-HT receptors on microglial surface, thus inducing the release of inflammatory cytokines, which in turn play important role in the maintenance of gut inflammation[86]. In addition to 5-HT, tryptophan is also a serotonin precursor and known for influencing microglia activity in gut lumen. Moreover, tryptophan and its derivatives have been found to regulate CNS inflammation, particularly following an aryl hydrocarbon receptor (Ahr) mediated mechanism, which in turn is responsible for microglial activation and transcriptional regulation of astrocytes[87]. The vitality of tryptophan and its metabolism in maintaining CNS homeostasis has been reported recently, where GF animals were found to have augmented 5-HT levels and 5-hydroxyindoleacetic acid in the hippocampus and serum samples, compared to conventionally colonized control mice[87,88]. These findings are suggestive towards the ability of microbiota in maintaining CNS serotonergic neurotransmission, mainly through systemic circulation. These initial findings were also validated by colonizing these animals post-weaning, where they were found to have sufficient levels of tryptophan in peripheral samples and reduced anxiety. Interestingly, this colonization was found ineffective in reversing the CNS neurochemical consequences exhibited by GF animals[89]. In a recent study, one major finding was listed where an intact and diversified microbiota (post birth) was found to



produce quinolinic acid and N-methyl-D-aspartate agonist, as a part of tryptophan metabolism regulated by microglial cells[90]. Both have a critical role in the pathogenesis of several neurological conditions such as Huntington's disease and behavioral disorders[91]. In experimental setup where GF mice were recolonized with bacterial population belonging to *Clostridium tyrobutyricum*, they exhibited colonized mucus layer, and regulated immune and gut barrier homeostasis[92].

Furthermore, Lactobacillus rhamnosus (JB-1) containing probiotics supplementations in previously colonized mice were found to be able to reduce anxiety and depressive behavior^[93]. Recently, researchers also found that microbiota dependent alterations in neural synapses along with fear extinction behavior are not mainly due to altered HPA axis, they are also due to a reduction in various neuroactive metabolites' levels such as phenyl sulfate, pyrocatechol sulfate, and indoxyl sulfate[94,95]. This was further confirmed with the abraded levels of these metabolites in various samples collected from experimental animals, such as fecal sample, serum, and CSF. Collectively, gut microbiome and its associated bacterial populations were found to be pivotal in maintaining the level of various neuroactive metabolites through which they regulate the behavioral patterns in healthy subjects.

Regulatory crosstalk between intestinal microbiome and brain

In humans, ENS is generally referred as the second brain which is located inside the digestive tract wall and shielded by a mucous membrane from gut lumen, having five times higher neurons and ganglia than spinal cord. An autonomous system regulating microbiome and GBA is known to be bidirectional whereas somatic sensory system received signals in pathological conditions causes abnormal feelings like discomfort, nausea, and pain which in turn exhibit bowel associated problems[96]. Herein, ANS transports afferent signals received from gut lumen to CNS through various routes such as enteric, spinal and vagal[96]. In these communications, vagus nerve plays a critical role as it has millions of nerve terminals (80% are afferent). Moreover, the same pathway is also used to send efferent signals from CNS to the gut wall[97]. In case of stress and anxiety, neuroendocrine system in response to stress regulates various vital body functions, including digestion and immunity which are mainly dependent on the HPA axis[98]. For example, generation of corticosteroids in response to environmental stress is also found to be dependent on HPA axis[97]. Collectively, neural and hormonal signals regulate brain and microbiome for further regulation of gut cell activity in controlled conditions.

In addition, signal transducing cells such as enterochromaffine cells (EC) and dendritic cells (DC) produce various neurotransmitters (Figure 2) viz., 5-HT, somatostatin, cholecystokinin and CRH[99-101]. All these are important for signal-based regulation of microbiota and CNS as these directly affect the microbial behavior. EC serve as luminal sensors for observing a great range of bacteria and microbial compounds in the gut[102]. These microbial populations have neurotransmitter receptors, and their activation is a key factor to understand the mode of microbiota functions and composition. A specialized process called as "inter-kingdom signalling" is a well-known regulatory pathway through which bacteria communicate with gut epithelial cells, mainly by using oligopeptides and monoamines which are also known for their neurotransmitter behavior [103,104]. Numerous neurotransmitters are produced by ENS exhibiting critical roles in GBA regulation, where each has a specific signalling pathway. Some neurotransmitters such as 5-HT, somatostatin, dopamine, neuropeptide Y, peptide YY, cholecystokinin, and corticotropin-releasing factor are also necessary for GBA regulation[96,101,104].

EC in GI tract produces majority of body's 5-HT and dopamine. An increased synthesis of 5-HT has been found as a key regulator through which microbiota regulates HPA[97]. Additionally, gut microbiota senses the EC cells mainly via short-chain fatty acids (SCFA) (butyrate and acetate) to promote 5-HT synthesis that regulates GI motility, secretion and immunological responses[96]. Hence, a change in microbiota composition significantly affects the levels of 5-HT and its altered level contributes towards the pathophysiology of IBS. Interestingly, microbiota is also connected with CNS through TLR signalling pathway [105,106]. TLR is principally expressed on immune cells of gut wall and neurons of the ENS. Among microbial populations, gram-negative bacterial membrane component lipopolysaccharide (LPS) is selectively detected by TLR which in turn is responsible for the production of proinflammatory cytokines via NF-KB pathway[106]. In colonic biopsies of IBS-D patients, higher TLR expression has been recorded suggesting microbiota and host immune system interaction [102,104]. Furthermore, smooth muscle dysfunction and paralysis in septic ileus is also known to be significantly influenced by the number of cytokines produced by immune cells, in response to LPS through TLR mediated signalling[106].

INTESTINAL MICROBIOME AND SEROTONERGIC SIGNALLING

Serotonergic system

5-HT is an important neurotransmitter and paracrine signalling molecule in the gut where its major role has been identified in the modulation of gut-brain communication and in functional GI disorders^[21]. In previous studies, altered levels of 5-HT have been reported in various CNS related disorders including anxiety, depression, obsessive compulsive disorder, phobias and in other psychiatric disorders[67,69]. 5-HT modulators including SSRIs and specific 5-HT receptor agonists and antagonists have also been



used to treat various metabolic disorders such as, migraine, nausea, obesity, chronic pain, hypertension, vascular disorders, and sexual dysfunction[107].

5-HT signalling in GI tract motility

Among all the neurotransmitter-based regulations of GBA, 5-HT mediated signalling is primary where 5-HT mainly targets receptor subtypes of seven 5-HT family receptors, having 15 subtypes of recognition sites[85]. Additionally, seven types of 5-HT receptors have a wide range of biological tasks including enhancing mucosal permeability and visceral hypersensitivity, inflammation and immune cell activation, and gut motility [107]. Currently, 7 families of 5-HT receptors have been identified, where 5-HT3 and 5-HT4 receptors are in intestine, presynaptic positions and, in sensory and mesenteric neurons [107]. In clinical practices, medications that are known to target these receptors have been frequently used for the management of IBS complications. Enteric microbiota has also been found to control the expression of 5-HT receptors[107]. 5-HT mainly activates peristaltic reflexes in GI tract (Figure 4) thus causing ascending contractile and descending relaxant limbs. 5-HT has been also found to control segmental motor patterns in small intestine. A study performed in tryptophan hydroxylase deficient (Tph2-/- mice) mice demonstrated the functional aspects of 5-HT in GI motility mainly by attributing with significant reductions in contractile complexes and synaptic transmission due to lower level of 5-HT[108,109]. This is further indicative for the significance of serotonergic neurons, comparing to EC cells for constitutive GI motility. Additionally, ENS dopaminergic neurons were found to be immature in Tph2-/- mice, which are responsible for homeostatic GI movement[110]. Moreover, in vivo and in vitro studies have shown decreased intestinal motility in mice with SERT Ala56 mutation which was restored by 5-HT4 receptor antagonists[111]. A SERT antagonist named 'fluoxetine' was found to improve GI motility in SERT-/- mice, along with a higher level of 5-HT, thus indicating the importance of 5-HT in GI motility, by targeting SERT[112].

Various antibiotic treatments have been found to impede GI motility along with a reduced production of peripheral 5-HT level, which was also confirmed by GF animal studies where a slower GI transit was recorded in these animals compared to control mice having normal gut microbiota[66,112]. In GF mice, treatment with pharmacological blockers of 5-HT4 receptors significantly improved GI transit along with a higher level of luminal 5-HT level [113]. Additionally, 5-hydroxyindole has been produced as a key metabolite of serotonin metabolic pathway (Figure 4), mainly by increasing the colonic motility, especially by activating L-type calcium channels[114]. Recently, in a BTBR (BTBR T+ Itpr3tf/J) mice model of autism spectrum disorder, an altered serotonergic pathway has been found along with downregulation of Tph1 gene, while SERT expression was upregulated along with a reduced level of 5-HT and its producing host (Blautia spp)[115]. A list of various enteric microbiota and their mode of regulating serotonergic pathway are listed in Table 1. Additionally, 5-HT level in combination with gut microbial stimulus can also lift the proportion of M20000 macrophages, which are known to be stimulatory for GI motility particularly in the colonic muscle layer located nearby ENS[117]. Various microbial byproducts such as SCFAs were also found to enhance colonic transit with the release of 5-HT intra-luminally, mainly by targeting GPR43 receptor located on mucosal mast cells[117]. Therefore, these findings clearly suggest that through various mechanisms enteric microbial population controls 5-HT level and associated signalling for the control of GI motility, and dysregulation in this system resulted in abnormal GI movement, which is linked with IBS symptoms[116,117]. It has been widely accepted that IBS patients have altered motor and stool patterns, where enhanced GI motility produce abnormal gut contractions resulting in abdominal pain and discomfort.

Role of gut microbe-mediated 5-HT signalling in visceral pain sensation

Chronic abdominal pain is a crucial aspect of IBS which is also influenced by intestinal gut microbiota [118]. Animal studies are further suggestive for the ability of fecal microbiota to transmit hypersensitivity to colonic distension in rats[119]. These results suggest that abnormal pain perception associated with IBS is also critically derived from gut microbial components[118,119]. Additionally, neurotransmitters derived from gut microbiota have been also found to be decisive for the perception of visceral discomfort. Among the other neurotransmitters, a preferential role of 5-HT has been found to modulate the intestinal pain by activating mesenteric sensory nerve fibers along with the activation of vagal and spinal afferent fibers[120]. Microbial colonization and their commensal activities have been reported to be essential for promoting excitability of the gut sensory neurons, which are responsible for the development of homeostatic pain sensitivity[121]. In GF mice, where a little mucosal inflammation has been found to be associated with visceral hypersensitivity caused by abnormal pain processing in the brain, it can be corrected by fecal microbiota transplantation (FMT) therapies derived from conventional mice[122]. FMTs are also known to stimulate primary nociceptive neurons in the dorsal root ganglia (DRG) and various gut microbial components such as TLR ligands, formyl peptide receptor 1 agonists and SCFAs[122]. These can modulate direct or indirect enhancement of visceral pain sensitivity. Additionally, microbiota derived kynurenic acid, serine proteases and bile acids can reduce pain sensitivity by inactivating DRG neurons or indirectly by releasing opioid-like factors, mainly from the mucosal immune cells[123,124].

Table 1 Types of regulation of serotonergic (5-Hydroxytryptamine) pathway by specific enteric microbiota in gastrointestinal tract							
Microbiota spp.	5-HT pathway	Mechanisms of action and observations	Ref.				
Akkermansia muciniphila (Amuc_1100)	Upregulation	Promote intestinal 5-HT biosynthesis and extracellular availability through TLR2 signalling	[129]				
Akkermansia muciniphila (extracellular vesicles)	Upregulation	Increase expression of the Htr4 gene, and decreases expression of the Htr2B, Htr3B, and Htr7 genes	[129]				
Bacteriodes thetaiotaomicron	Upregulation	Restore 5-HT+ EC cells and shape EC networks in the GI tract of GF mice by producing SCFAs	[130]				
Bifidobacterium dentium	Upregulation	Increase intestinal 5-HT level; expressions of 5-HTra receptors 2a and 4, and SERT by producing acetate	[131]				
Bifidobacterium longum and	Downregulation	Upregulate SERT expression	[132]				
Lactobacillus acidophilus							
Bifidobacterium pseudolongum	Downregulation	Diminish EC cells	[133]				
Clostridium ramosum	Upregulation	Promote 5-HT synthesis in colonic EC cells and program differentiation of intestinal stem progenitors toward a secretory 5-HT-producing lineage	[134]				
Corynebacterium spp., Enterococcus spp., Strepto- coccus spp.	Upregulation	Enable the direct production of 5-HT	[135]				
Escherichia coli Nissle 1917	Upregulation	Enhance 5-HT bioavailability in ileal tissue through interaction with compounds secreted from host tissue	[136]				
Indigenous spore-forming bacteria	Upregulation	Enhance colonic 5-HT pathway by upregulation of Htr4	[137]				
Lactobacillus acidophilus	Down regulation	Upregulate SERT expression	[138]				
Lactobacillus plantarum IS-10506	Upregulation	Increase gut 5-HT production along with brain 5-HTT, neurotrophin, and brain-derived neurotrophic factor	[139]				
Lactobacillus plantarum PS128	Upregulation	Increase 5-HT+ cells in the gut and alter expression levels of Tph1, Chga, Slc6a4, and Htr4	[140]				
Lactobacillus rhamnosus	Down regulation	Upregulate gene and protein level of SERT	[141]				
SadA-expressing Staphylococci, Trichinella spiralis and Campylobacter jejuni (pathogens)	Upregulation	Promote converting 5-HTP into 5-HT; increase EC cell number and reduce SERT expression	[141]				

5-HT: 5-Hydroxytryptamine; 5-HTP: 5-Hydroxytryptophan; EC: Enterochromaffin cell; SERT: Serotonin transporters.

Interestingly, the mode of action of 5-HT mediated signalling depends on the type of 5-HT receptor activation and the release of enteric 5-HT, which has been significantly linked to the intensity of abdominal discomfort in IBS patients[125]. The etiology of IBS also includes visceral hypersensitivity along with the nociceptive process and these are mainly regulated with 5-HT3 receptor activation, where it is mainly expressed on peripheral terminals of spinal afferent nerves and vagal afferent nerve endings in stomach [125]. 5-HT3 receptor has been also identified to influence other neurotransmitters in the brain, where a 5-HT3 receptor antagonist (Ramosetron) has been found to lower the visceral hypersensitivity and modifies GI transit in IBS-D patients [126]. Additionally, 5-HT3 receptor antagonists have demonstrated anti-inflammatory effects particularly to the enhanced permeability of gut and mucosal inflammation reported to be caused by various microbial populations^[125]. Moreover, an impaired intestinal barrier linked to dysfunctional 5-HT metabolism in IBS patients has been observed recently [126]. Collectively, these findings are indicative of the vitality of gut microbiota in 5-HT-mediated pain perception, where visceral hypersensitivity resulted from the dysfunction of serotonergic pathway has been linked to enteric dysbiosis as reported in IBS cases[127,128].

Role of gut-microbe-mediated 5-HT signalling in mucosal inflammation and immune response

Among the various pathways underpinning the IBS pathogenesis, a persistent and low-grade mucosal inflammation has been reported in majority of clinical cases, with aberrant immune cell activation [90-129]. In IBS patients, abdominal pain has been characterized with a significant infiltration of mast cells in the colonic mucosa, which is also accompanied with an augmented level of mucosal 5-HT[28,29]. Although, exact underlying mechanisms and causes of mucosal inflammation are not completely known, dysbiosis of the gut microbiota along with altered serotonergic signalling have been identified as the important contributors for the onset of IBS. Indeed, augmented numbers of mucosal and EC cells have been recorded in various inflammatory diseases, such as inflammatory bowel disease (IBD), where a range of 5-HT receptors are expressed to stimulate intestinal inflammation through serotonergic



Figure 4 Tryptophan derived serotonergic metabolism in gut brain axis cross talk. Upon absorption in the gut, L-tryptophan is metabolized through three pathways, particular to bacterial cells (indole pathway) of gut microbiota and mammalian cells (kynurenine and serotonin pathways). Tryptophan is converted to 5-hydoxytryptophan (5-HTP) by tryptophan hydroxylase and thereafter converted to -5-HT by the enzyme aromatic L-amino acid decarboxylase. In addition to this, tryptophan is also metabolized to indole derivatives and kynurenine, which also perform various biological functions. Tryptophan derived 5-HT regulates various functions in central nervous system (including emotion, cognition, stress, and visceral perception) and in enteric nervous system (gastrointestinal motility and secretion). CRF: Corticotropin-releasing factor; ANS: Autonomic nervous system; CNS: Central nervous system; GI: Gastrointestinal; 5-HT: 5-Hydroxytryptamine; HPA: Hypothalamic-pituitary-adrenal.

> signalling[118]. In colitis, 5-HT directs proinflammatory cytokines generations by stimulating T cells, peritoneal macrophages and splenic DCs in NFkB-dependent manner [130].

> Moreover, enhanced 5-HT availability has been recorded in SERT-deficient animals sensitive to gut mucosal inflammation whereas Tph1-/- mice with a lower level of 5-HT have been found to be resistant to experimental colitis[131]. These results suggest that intestinal inflammation is significantly regulated by enteric 5-HT, mainly by acting as pro-inflammatory immune regulator. Various results derived from a post-inflammatory IBS rat model also reveal the onset of visceral hypersensitivity along with fecal microbial dysbiosis, along with an elevated serum level of 5-HT[132]. Additionally, in recent studies lamina propria mast cells (MC) were found to be significantly co-related with an enhanced and spontaneous release of 5-HT in IBS patients[133]. Interestingly, histamine has also been identified as an important biogenic amine found to have pathophysiological role in IBS, where the exact mechanism is still not fully deciphered.

> Usually, histamine played critical roles in regulating GI motility, gastric acid and mucosal ion secretion[49,133]. Histamine H1 receptors (H1R) are known to be involved in mediating sensorineural signalling and vascular dilatation, where their activation is known to regulate food and water intake and diurnal feeding rhythm. In addition to this, stimulation of histamine H2 receptors (H2R) have been known for the degranulation of MCs and production of various antibodies, T helper (Th) 1 cytokines, and for T-cell proliferation [133,134]. However, evidences are indicative for the overproduction of histamine by MCs that has been known to cause diarrhea, with an increased neuronal secretomotor function. In constipation, histamine also induces altered enteric neuron function resulting in an excessive segmental contractile colonic motor activity [50,135]. Although, evidences suggested that use of various agents targeting the histamine receptors (HRs) have been found to be potential therapeutic option for IBS patients[134].

> In addition to this, upregulated expression of various colonic receptors of 5-HT (5-HT3A/5-HT2B) along with impaired junction proteins has been found in test animals[136]. Additionally, 5-HT3A receptor antagonist administration or FMT derived from the faeces of normal healthy rats, were found to be able to alleviate IBS-like symptoms[136]. Moreover, SCFAs from the gut microbiota have been appeared to have a dual nature directing GI mucosal immunity and inflammation, which are known to be crucial for maintaining gut homeostasis[58]. These cause the over expression of G-protein coupled receptors and induction of regulatory T cells in the gut which in turn increases the integrity of epithelial cell barrier, and thus exhibiting anti-inflammatory effects [136]. In parallel to this, SCFAs are known to cause mucosal inflammation by promoting mucosal 5-HT synthesis (serotonergic pathway) and upregulation of TPH1 transcription[137].



Microbiota mediated serotonergic signalling outside GI tract

Various preclinical and clinical data have demonstrated the critical role of 5-HT derived from gut in glucose and lipid metabolism, and in various metabolic disorders[138,139]. Various antagonist specific to 5-HT receptors have been beneficial in reversing the clinical manifestations received by altered 5-HT metabolism. For example, fluoxetine was used to prevent T. sanguinis population by lowering serum triglyceride levels and changing the expression of lipid metabolism genes[139]. Additionally, the gut was found to be significantly regulated by its microbiota member "Clostridium ramosum" which is found to be associated with lipid transport and storage functions in animals. Furthermore, glucose homeostasis is also controlled by various members of enteric microbiota, mainly with the modulation of EC cell's 5-HT production where this was mainly targeted with the inhibition or genetic depletion of TPH1[140]. Purine metabolism was also found to be regulated with host-microbial metabolic route in IBS patients[141]. These results are clearly indicative for the 5-HT-dependent regulation of various metabolic pathways in the pathogenesis of IBS.

ROLE OF GUT MICROBIOME DERIVED METABOLITES IN IBS

The population of gut microbiota has been reported to be massive along with a modular genome which in turn reported to be benefitting the host and adapted to gut environment. These microbes carry out a variety of tasks, such as xenobiotic metabolism, vitamin production, pathogen defense, and dietary fiber fermentation[115,136,142]. Microbiota derived metabolites are small substances synthesized as intermediate or end products of microbial metabolisms, which are also the principal regulators through which gut microbiota plays an important role in host specific metabolism[136]. These metabolites may be produced directly by bacteria, by alteration of host molecules like bile acids, or through bacterial metabolism of food components. Immunological maturation, immune homeostasis, host energy metabolism, and mucosal integrity maintenance are all identified to be influenced by microbial metabolites mediated endocrine signalling[11,12,14]. IBD's pathophysiology has been principally linked to certain groups of metabolites, such as bile acids, SCFAs, and tryptophan which have been found in a variety of biological tissues, including faeces, urine, serum, liver, and cerebrospinal fluid, thus having significant impact on the physiology of the host[17,136]. Additionally, some metabolites have been also found to be specific to Crohn's disease patients and non-IBD controls, where individuals with IBD were found to have higher amounts of bile acids, amino acids, and sphingolipids[143]. Interestingly, these individuals have lower levels of cholesterols, phenylbenzodioxanes, indoles, tetrapyrroles and longchain fatty acids[144]. SCFAs are derived from carbohydrates and are localized to microbiome, regulating the function of intestinal macrophages, appetite, fat accumulation, intestinal motility and energy metabolism^[145]. Other SCFAs produced by microbial fermentation include acetate, propionate, and butyrate, as well as the gases methane, hydrogen sulphide and some other intermediates [145,146]. Butyrate, along with other SCFAs, inhibits epithelial stem cells and promotes epithelial homeostasis by producing IL-18 through inflammasome activation, which is known to be the main source of energy in colonic epithelial cells[147].

Like SCFAs, medium-chain fatty acid levels have been found to be depleted in IBD. Thus, it is hypothesized that certain fatty acids, such as conjugated linoleic acid might exert some anti-inflammatory effects by activating the peroxisome proliferator-activated receptor- γ (PPAR- γ)[148]. Additionally, bile acids also influence host metabolism and immune system via interacting with receptors, such as the transmembrane G protein-coupled receptor 5 (TGR5) and farnesoid X receptor (FXR)[149]. By activating type 2 iodothyronine deiodinase, TGR5 enhances insulin sensitivity (via GLP1), thus regulating energy expenditure (in muscle and brown adipose tissue), and gall bladder relaxation[149]. Activation of bile acid receptor (FXR) also affects host metabolism through various ways, including decreased lipogenesis, hepatic gluconeogenesis, and liver regeneration, as well as by generating antimicrobial peptides for liver regeneration[150,151]. Constitutional makeup of gut microbiota is also significantly influenced by bile acid composition. Furthermore, activation of small intestine FXR also reduces bacterial translocation and growth[152].

In addition to other metabolic intermediates, tryptophan is a well-known precursor for the synthesis of several metabolites such as serotonin, melatonin, nicotinamide and vitamin B3[153]. Gut is the main site of tryptophan synthesis and metabolism, where it is converted by commensal microbiota into indoles agonists of pregnane X receptor and AhR, which have pivotal roles in mucosal immunity and homeostasis[154,155]. Moreover, dysbiosis in IBD also causes a significant decrease in microbial tryptophan activation, which in turn increases metabolism (Figure 4)[148,149]. Indole also stimulates GLP1 release while other indole derivatives [such as indoleacetic acid, indole-3-acetaldehyde, indole-3aldehyde, indoleacrylic acid (IA) and IPA] are also known to act as agonists for AhR[156]. Additionally, AhR also act as a transcription factor which having developmental and tissue-dependent effects on T cell immunity[157]. Reduced AhR expression has been noted in inflamed mucosal samples from Crohn's disease patients whereas serum samples from these patients were also found to have altered xenobiotic metabolism[158]. Moreover, enhanced catabolism of fatty acids, reduced level of amino acid metabolites including serotonergic and indole derivatives of tryptophan, as well as decreased levels of



Table 2 Various therapies prescribed for the treatment of irritable bowel syndrome along with their possible mode of action					
Therapy	Description	Proposed mechanism(s) of action			
Prebiotics	Ingested compounds targeted to stimulate gut microbiota	Mechanism of action undefined, but may include: Anti-inflammatory effects; inhibition of pathogen adherence; and growth of intestinal mucosal layer			
Probiotics	Ingested microorganisms (e.g., bacteria)	Mechanism of action undefined, but may include: Inhibition of pathogenic microor- ganism colonization; support intestinal barrier integrity and function; production of beneficial micronutrients; and activation and augmentation of the enteric nervous system			
Rifaximin	Nonabsorbable, bile-soluble antibiotic indicated for the treatment of adults with IBS- D	Antibacterial against Gram-positive and Gram-negative bacteria: Modulation of gut- immune signalling; inhibition of bacterial translocation; SIBO eradication (in some patients); causing decreases in GI methane concentrations in combination; and with the antibiotic neomycin (in patients with IBS-C)			
SBI	Prescription medical food for patients with IBS-D	Modulation of gut microbiota: Causing decreases in GI permeability			
SYN-010	Derivative of the HMG-CoA reductase inhibitor lovastatin lactone; currently in development for the treatment of patients with IBS-C	Inhibition of methane production by Methanobrevibacter smithii			
Dietary modification	Variable; one example is the low FODMAP diet	Causing decreases in GI gas production			
		Causing decreases in intra-luminal fluid production			

FODMAP: Fermentable oligo-, di-, monosaccharides and polyols; GI: Gastrointestinal; HMG-CoA: 3-hydroxy-3-methylglutaryl-coenzyme A; IBS-C: Constipation-predominant irritable bowel syndrome; IBS-D: Diarrhea-predominant irritable bowel syndrome; SBI: Serum-derived bovine immunoglobulin; SIBO: Small intestinal bacterial overgrowth.

phenylalanine and histidine metabolite ergothioneine has been also found in these patients [153,155,158].

MICROBIOME TARGETED THERAPEUTICS FOR IBS

Restoring the gut microbiome seems to be promising and therapeutic strategy for the management of IBS with diversified clinical scenarios. Moreover, maintaining the GI microbiome to a level of optimal health with the ideal fecal product is known to be challenging and beneficial to human being. In recent years, FMT has been introduced to yield promising results in various gut disorders such as in Clostridioides difficile infection, where FMT has shown high cure rates, compared to other standard therapies [55,159,160]. This initial lead has been also implanted with microbiome manipulation for the treatment of IBD, ulcerative colitis and Crohn's disease, with a higher rate of cure effectively [161,162]. However, designing FMT formulations as the treatment for IBS is very challenging, since the optimal microbiome composition in healthy people is poorly defined with a higher degree of variability [163]. In addition to this, disrupted microbiota which would be ideally restored via FMT is known to be highly variable.

Interestingly, dysbiosis of several inflammatory markers is known to induce the expansion of facultative anaerobes, where the mechanisms by which microbiota are disrupted remain unclear. At the same time, GI tract is known to be contributory for key determinants of FMT success whereas, some patients that received FMT also developed systemic infections owing to the adverse effects of FMT[61, 63,148]. To avoid the side effects received from FMT application, one more way is to administer autologous FMT (auto-FMT also known as microbiome restoration), where patient sample is screened for various pathogens, banked and then re-administered to the same patient after disrupting microbiome. In this case, restoring the microbiome samples for a long period of time has storage and safety issues, leading to future health problems. Meanwhile, several IBS therapeutics (Table 2) have been developed which are known to target the gut microbiota and target secondary consequences of alterations in the gut microbiota[164].

CONCLUSION

IBS is a prevalent GI disorder where mechanisms regulating the interconnected cross talk between brain and gut microbiota have been found to be altered. Various metabolic pathways and diseases have also been found to be associated with gut microbiota architecture. The pathophysiology of IBS is still not fully deciphered, whereas a complex network of interaction between various genes, metabolic pathways, behavioral events, host immune response and psychosocial factors has been found to be contributory for IBS and its associated symptoms. Interestingly, serotonin (5-HT) as a neurotransmitter secreted by EC cells regulates GI motility, secretion, and sensation, whereas altered 5-HT signalling has



been found contributory in the pathophysiology of IBS. Furthermore, environmental stress has a significant impact on IBS etiology where it significantly regulates the neuroendocrine system and gut functions, mainly through immune system mediated mechanisms. The dietary composition and its intake have a pivotal role in the regulation of IBS, hence usage of quality foods which are gluten free, low fat and FODMAP content, tryptophan and fiber rich may be prominent approach for the management of IBS.

ACKNOWLEDGEMENTS

Shiv Vardan Singh acknowledges University Grants Commission (UGC), New Delhi, India for Dr DS Kothari Fellowship. Risha Ganguly, Kritika Jaiswal and Ramesh Kumar acknowledge financial support from UGC/Council of Scientific and Industrial Research, New Delhi, India in the form of Junior and Senior Research Fellowships. Aditya Kumar Yadav acknowledges financial support from UGC in the form of CRET fellowship. All the authors also acknowledge DST-FIST and UGC-SAP facilities of the Department of Biochemistry, University of Allahabad, Prayagraj, India.

FOOTNOTES

Author contributions: Pandey AK and Singh SV conceptualized the idea; Singh SV, Ganguly R, Jaiswal K, Yadav AK, and Kumar R performed the literature search; Singh SV and Ganguly R wrote the first draft of the manuscript and validated the references; Pandey AK critically reviewed and revised the manuscript; and all authors have read and approved the final manuscript.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is noncommercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: India

ORCID number: Shiv Vardan Singh 0000-0002-6720-5719; Risha Ganguly 0000-0002-6190-8923; Ramesh Kumar 0000-0002-8236-5016; Abhay K Pandey 0000-0002-4774-3085.

S-Editor: Chen YL L-Editor: A P-Editor: Cai YX

REFERENCES

- Grace-Farfaglia P, Frazier H, Iversen MD. Essential Factors for a Healthy Microbiome: A Scoping Review. Int J Environ Res Public Health 2022; 19 [PMID: 35886216 DOI: 10.3390/ijerph19148361]
- León ED, Francino MP. Roles of Secretory Immunoglobulin A in Host-Microbiota Interactions in the Gut Ecosystem. 2 Front Microbiol 2022; 13: 880484 [PMID: 35722300 DOI: 10.3389/fmicb.2022.880484]
- 3 Yan X, Si H, Zhu Y, Li S, Han Y, Liu H, Du R, Pope PB, Qiu Q, Li Z. Integrated multi-omics of the gastrointestinal microbiome and ruminant host reveals metabolic adaptation underlying early life development. Microbiome 2022; 10: 222 [PMID: 36503572 DOI: 10.1186/s40168-022-01396-8]
- He J, Yi L, Hai L, Ming L, Gao W, Ji R. Characterizing the bacterial microbiota in different gastrointestinal tract segments 4 of the Bactrian camel. Sci Rep 2018; 8: 654 [PMID: 29330494 DOI: 10.1038/s41598-017-18298-7]
- Leite G, Pimentel M, Barlow GM, Chang C, Hosseini A, Wang J, Parodi G, Sedighi R, Rezaie A, Mathur R. Age and the 5 aging process significantly alter the small bowel microbiome. Cell Rep 2021; 36: 109765 [PMID: 34592155 DOI: 10.1016/j.celrep.2021.109765]
- Kumar Singh A, Cabral C, Kumar R, Ganguly R, Kumar Rana H, Gupta A, Rosaria Lauro M, Carbone C, Reis F, Pandey 6 AK. Beneficial Effects of Dietary Polyphenols on Gut Microbiota and Strategies to Improve Delivery Efficiency. Nutrients 2019; 11 [PMID: 31540270 DOI: 10.3390/nu11092216]
- Guo P, Lei M, Hu S, Xu Z, Zhou Y, Zhou P, Huang R. Long-term LDR exposure may induce cognitive impairments: A possible association through targeting gut microbiota-gut-brain axis. Ecotoxicol Environ Saf 2023; 249: 114351 [PMID: 36508818 DOI: 10.1016/j.ecoenv.2022.114351]
- Muhammad F, Fan B, Wang R, Ren J, Jia S, Wang L, Chen Z, Liu XA. The Molecular Gut-Brain Axis in Early Brain 8 Development. Int J Mol Sci 2022; 23 [PMID: 36499716 DOI: 10.3390/ijms232315389]



- Dai CL, Liu F, Iqbal K, Gong CX. Gut Microbiota and Immunotherapy for Alzheimer's Disease. Int J Mol Sci 2022; 23 9 [PMID: 36499564 DOI: 10.3390/ijms232315230]
- Kraaij R, Schuurmans IK, Radjabzadeh D, Tiemeier H, Dinan TG, Uitterlinden AG, Hillegers M, Jaddoe VWV, Duijts L, 10 Moll H, Rivadeneira F, Medina-Gomez C, Jansen PW, Cecil CAM. The gut microbiome and child mental health: A population-based study. Brain Behav Immun 2023; 108: 188-196 [PMID: 36494050 DOI: 10.1016/j.bbi.2022.12.006]
- Dothel G, Barbaro MR, Di Vito A, Ravegnini G, Gorini F, Monesmith S, Coschina E, Benuzzi E, Fuschi D, Palombo M, Bonomini F, Morroni F, Hrelia P, Barbara G, Angelini S. New insights into irritable bowel syndrome pathophysiological mechanisms: contribution of epigenetics. J Gastroenterol 2023 [PMID: 37160449 DOI: 10.1007/s00535-023-01997-6]
- Shrestha B, Patel D, Shah H, Hanna KS, Kaur H, Alazzeh MS, Thandavaram A, Channar A, Purohit A, Venugopal S. The 12 Role of Gut-Microbiota in the Pathophysiology and Therapy of Irritable Bowel Syndrome: A Systematic Review. Cureus 2022; 14: e28064 [PMID: 36127988 DOI: 10.7759/cureus.28064]
- Canavan C, West J, Card T. The epidemiology of irritable bowel syndrome. Clin Epidemiol 2014; 6: 71-80 [PMID: 13 24523597 DOI: 10.2147/CLEP.S40245]
- 14 Osadchuk AM, Osadchuk MA, Kvetnoĭ IM. [Irritated bowel syndrome: clinico-morphological types]. Klin Med (Mosk) 2007; 85: 46-50 [PMID: 17523405]
- Sheptulin AA, Vize-Khripunova MA. [NEWS IN ETIOLOGY AND PATHOGENESIS OF IRRITATED BOWEL 15 SYNDROME]. Klin Med (Mosk) 2016; 94: 92-96 [PMID: 27459756]
- Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, Gordon JI. The human microbiome project. Nature 16 2007; 449: 804-810 [PMID: 17943116 DOI: 10.1038/nature06244]
- 17 Hu Y, Chen Z, Xu C, Kan S, Chen D. Disturbances of the Gut Microbiota and Microbiota-Derived Metabolites in Inflammatory Bowel Disease. Nutrients 2022; 14 [PMID: 36501169 DOI: 10.3390/nu14235140]
- Singh AK, Singla RK, Pandey AK. Chlorogenic Acid: A Dietary Phenolic Acid with Promising Pharmacotherapeutic 18 Potential. Curr Med Chem 2023; 30: 3905-3926 [PMID: 35975861 DOI: 10.2174/0929867329666220816154634]
- 19 Camilleri M. Diagnosis and Treatment of Irritable Bowel Syndrome: A Review. JAMA 2021; 325: 865-877 [PMID: 33651094 DOI: 10.1001/jama.2020.22532]
- Basseri RJ, Weitsman S, Barlow GM, Pimentel M. Antibiotics for the treatment of irritable bowel syndrome. 20 Gastroenterol Hepatol (N Y) 2011; 7: 455-493 [PMID: 22298980]
- Wood JD. Enteric nervous system, serotonin, and the irritable bowel syndrome. Curr Opin Gastroenterol 2001; 17: 91-97 21 [PMID: 17031157 DOI: 10.1097/00001574-200101000-00017]
- Holland AM, Bon-Frauches AC, Keszthelyi D, Melotte V, Boesmans W. The enteric nervous system in gastrointestinal 22 disease etiology. Cell Mol Life Sci 2021; 78: 4713-4733 [PMID: 33770200 DOI: 10.1007/s00018-021-03812-y]
- 23 Alcaino C, Knutson KR, Treichel AJ, Yildiz G, Strege PR, Linden DR, Li JH, Leiter AB, Szurszewski JH, Farrugia G, Bevder A. A population of gut epithelial enterochromaffin cells is mechanosensitive and requires Piezo2 to convert force into serotonin release. Proc Natl Acad Sci U S A 2018; 115: E7632-E7641 [PMID: 30037999 DOI: 10.1073/pnas.1804938115]
- Dickson I. Gut mechanosensors: enterochromaffin cells feel the force via PIEZO2. Nat Rev Gastroenterol Hepatol 2018; 24 15: 519 [PMID: 30100606 DOI: 10.1038/s41575-018-0059-9]
- Young SN. How to increase serotonin in the human brain without drugs. J Psychiatry Neurosci 2007; 32: 394-399 [PMID: 25 18043762
- 26 Noddin L, Callahan M, Lacy BE. Irritable bowel syndrome and functional dyspepsia: different diseases or a single disorder with different manifestations? MedGenMed 2005; 7: 17 [PMID: 16369243]
- Gwee KA, Chua AS. Functional dyspepsia and irritable bowel syndrome, are they different entities and does it matter? 27 World J Gastroenterol 2006; 12: 2708-2712 [PMID: 16718757 DOI: 10.3748/wjg.v12.i17.2708]
- Sangkuhl K, Klein TE, Altman RB. Selective serotonin reuptake inhibitors pathway. Pharmacogenet Genomics 2009; 19: 28 907-909 [PMID: 19741567 DOI: 10.1097/FPC.0b013e32833132cb]
- Lukiw WJ, Li W, Bond T, Zhao Y. Facilitation of Gastrointestinal (GI) Tract Microbiome-Derived Lipopolysaccharide 29 (LPS) Entry Into Human Neurons by Amyloid Beta-42 (Aβ42) Peptide. Front Cell Neurosci 2019; 13: 545 [PMID: 31866832 DOI: 10.3389/fncel.2019.00545]
- 30 **Guzel T**, Mirowska-Guzel D. The Role of Serotonin Neurotransmission in Gastrointestinal Tract and Pharmacotherapy. Molecules 2022; 27 [PMID: 35268781 DOI: 10.3390/molecules27051680]
- Maffei ME. 5-Hydroxytryptophan (5-HTP): Natural Occurrence, Analysis, Biosynthesis, Biotechnology, Physiology and 31 Toxicology. Int J Mol Sci 2020; 22 [PMID: 33375373 DOI: 10.3390/ijms22010181]
- Celada P, Puig M, Amargós-Bosch M, Adell A, Artigas F. The therapeutic role of 5-HT1A and 5-HT2A receptors in 32 depression. J Psychiatry Neurosci 2004; 29: 252-265 [PMID: 15309042]
- 33 Saatçi Oğlu Ö, BuketTomruk N. Antidepressant Treatment Strategies in the Perinatal Period with a Focus on SSRI use. NPA 2013; 50: 93-94 [DOI: 10.4274/npa.y7029]
- 34 Shukla R, Ghoshal U, Ranjan P, Ghoshal UC. Expression of Toll-like Receptors, Pro-, and Anti-inflammatory Cytokines in Relation to Gut Microbiota in Irritable Bowel Syndrome: The Evidence for Its Micro-organic Basis. J Neurogastroenterol Motil 2018; 24: 628-642 [PMID: 30347939 DOI: 10.5056/jnm18130]
- Fukudo S. [Irritable Bowel Syndrome, Emotion Regulation, and Gut Microbiota]. Brain Nerve 2016; 68: 607-615 [PMID: 35 27279158 DOI: 10.11477/mf.1416200448]
- Bhattarai Y, Muniz Pedrogo DA, Kashyap PC. Irritable bowel syndrome: a gut microbiota-related disorder? Am J Physiol 36 Gastrointest Liver Physiol 2017; 312: G52-G62 [PMID: 27881403 DOI: 10.1152/ajpgi.00338.2016]
- Bennet SM, Ohman L, Simren M. Gut microbiota as potential orchestrators of irritable bowel syndrome. Gut Liver 2015; 37 9: 318-331 [PMID: 25918261 DOI: 10.5009/gn114344]
- Lee BJ, Bak YT. Irritable bowel syndrome, gut microbiota and probiotics. J Neurogastroenterol Motil 2011; 17: 252-266 38 [PMID: 21860817 DOI: 10.5056/inm.2011.17.3.252]
- 39 Fukudo S. Role of corticotropin-releasing hormone in irritable bowel syndrome and intestinal inflammation. J



Gastroenterol 2007; 42 Suppl 17: 48-51 [PMID: 17238026 DOI: 10.1007/s00535-006-1942-7]

- Sagami Y, Shimada Y, Tayama J, Nomura T, Satake M, Endo Y, Shoji T, Karahashi K, Hongo M, Fukudo S. Effect of a 40 corticotropin releasing hormone receptor antagonist on colonic sensory and motor function in patients with irritable bowel syndrome. Gut 2004; 53: 958-964 [PMID: 15194643 DOI: 10.1136/gut.2003.018911]
- 41 Singh AK, Bishayee A, Pandey AK. Targeting Histone Deacetylases with Natural and Synthetic Agents: An Emerging Anticancer Strategy. Nutrients 2018; 10 [PMID: 29882797 DOI: 10.3390/nu10060731]
- Torii A, Toda G. Management of irritable bowel syndrome. Intern Med 2004; 43: 353-359 [PMID: 15206545 DOI: 42 10.2169/internalmedicine.43.353]
- Levy RL, Cain KC, Jarrett M, Heitkemper MM. The relationship between daily life stress and gastrointestinal symptoms 43 in women with irritable bowel syndrome. J Behav Med 1997; 20: 177-193 [PMID: 9144039 DOI: 10.1023/a:1025582728271]
- Whitehead WE, Crowell MD, Robinson JC, Heller BR, Schuster MM. Effects of stressful life events on bowel symptoms: 44 subjects with irritable bowel syndrome compared with subjects without bowel dysfunction. Gut 1992; 33: 825-830 [PMID: 1624167 DOI: 10.1136/gut.33.6.825]
- Qin HY, Cheng CW, Tang XD, Bian ZX. Impact of psychological stress on irritable bowel syndrome. World J 45 Gastroenterol 2014; 20: 14126-14131 [PMID: 25339801 DOI: 10.3748/wjg.v20.i39.14126]
- Alonso C, Guilarte M, Vicario M, Ramos L, Ramadan Z, Antolín M, Martínez C, Rezzi S, Saperas E, Kochhar S, Santos J, 46 Malagelada JR. Maladaptive intestinal epithelial responses to life stress may predispose healthy women to gut mucosal inflammation. Gastroenterology 2008; 135: 163-172.e1 [PMID: 18455999 DOI: 10.1053/j.gastro.2008.03.036]
- Patacchioli FR, Angelucci L, Dellerba G, Monnazzi P, Leri O. Actual stress, psychopathology and salivary cortisol levels 47 in the irritable bowel syndrome (IBS). J Endocrinol Invest 2001; 24: 173-177 [PMID: 11314746 DOI: 10.1007/BF033438381
- Posserud I, Agerforz P, Ekman R, Björnsson ES, Abrahamsson H, Simrén M. Altered visceral perceptual and 48 neuroendocrine response in patients with irritable bowel syndrome during mental stress. Gut 2004; 53: 1102-1108 [PMID: 15247175 DOI: 10.1136/gut.2003.017962]
- Vanuytsel T, van Wanrooy S, Vanheel H, Vanormelingen C, Verschueren S, Houben E, Salim Rasoel S, Tóth J, Holvoet 49 L, Farré R, Van Oudenhove L, Boeckxstaens G, Verbeke K, Tack J. Psychological stress and corticotropin-releasing hormone increase intestinal permeability in humans by a mast cell-dependent mechanism. Gut 2014; 63: 1293-1299 [PMID: 24153250 DOI: 10.1136/gutjnl-2013-305690]
- Esposito P, Chandler N, Kandere K, Basu S, Jacobson S, Connolly R, Tutor D, Theoharides TC. Corticotropin-releasing 50 hormone and brain mast cells regulate blood-brain-barrier permeability induced by acute stress. J Pharmacol Exp Ther 2002; 303: 1061-1066 [PMID: 12438528 DOI: 10.1124/jpet.102.038497]
- 51 Menees S, Chey W. The gut microbiome and irritable bowel syndrome. F1000Res 2018; 7 [PMID: 30026921 DOI: 10.12688/f1000research.14592.11
- Chong PP, Chin VK, Looi CY, Wong WF, Madhavan P, Yong VC. The Microbiome and Irritable Bowel Syndrome A 52 Review on the Pathophysiology, Current Research and Future Therapy. Front Microbiol 2019; 10: 1136 [PMID: 31244784 DOI: 10.3389/fmicb.2019.01136]
- González-Castro AM, Martínez C, Salvo-Romero E, Fortea M, Pardo-Camacho C, Pérez-Berezo T, Alonso-Cotoner C, 53 Santos J, Vicario M. Mucosal pathobiology and molecular signature of epithelial barrier dysfunction in the small intestine in irritable bowel syndrome. J Gastroenterol Hepatol 2017; 32: 53-63 [PMID: 27087165 DOI: 10.1111/jgh.13417]
- Enck P, Aziz Q, Barbara G, Farmer AD, Fukudo S, Mayer EA, Niesler B, Quigley EM, Rajilić-Stojanović M, Schemann 54 M, Schwille-Kiuntke J, Simren M, Zipfel S, Spiller RC. Irritable bowel syndrome. Nat Rev Dis Primers 2016; 2: 16014 [PMID: 27159638 DOI: 10.1038/nrdp.2016.14]
- Devanarayana NM, Rajindrajith S. Irritable bowel syndrome in children: Current knowledge, challenges and opportunities. World J Gastroenterol 2018; 24: 2211-2235 [PMID: 29881232 DOI: 10.3748/wjg.v24.i21.2211]
- Sinagra E, Morreale GC, Mohammadian G, Fusco G, Guarnotta V, Tomasello G, Cappello F, Rossi F, Amvrosiadis G, 56 Raimondo D. New therapeutic perspectives in irritable bowel syndrome: Targeting low-grade inflammation, immunoneuroendocrine axis, motility, secretion and beyond. World J Gastroenterol 2017; 23: 6593-6627 [PMID: 29085207 DOI: 10.3748/wjg.v23.i36.6593]
- Alnoman A, Badeghiesh AM, Baghlaf HA, Dahan MH. Pregnancy, delivery, and neonatal outcomes among women with 57 irritable bowel syndrome (IBS) an evaluation of over 9 million deliveries. J Matern Fetal Neonatal Med 2022; 35: 5935-5942 [PMID: 33823718 DOI: 10.1080/14767058.2021.1903421]
- Talavera JIR, Parrill AM, Elsayad C, Fogel J, Riggs JC, Peng B. The association between ectopic pregnancy and 58 inflammatory bowel disease, irritable bowel syndrome, and celiac disease: A systematic review. J Obstet Gynaecol Res 2021; 47: 1601-1609 [PMID: 33733538 DOI: 10.1111/jog.14705]
- 59 Ganguly R, Gupta A, Pandey AK. Role of baicalin as a potential therapeutic agent in hepatobiliary and gastrointestinal disorders: A review. World J Gastroenterol 2022; 28: 3047-3062 [PMID: 36051349 DOI: 10.3748/wjg.v28.i26.3047]
- Nabavi-Rad A, Sadeghi A, AsadzadehAghdaei H, Yadegar A, Smith SM, Zali MR. The double-edged sword of probiotic 60 supplementation on gut microbiota structure in Helicobacter pylori management. Gut Microbes 2022; 14: 2108655 [PMID: 35951774 DOI: 10.1080/19490976.2022.2108655]
- Farzaei MH, Singh AK, Kumar R, Croley CR, Pandey AK, Coy-Barrera E, Kumar Patra J, Das G, Kerry RG, Annunziata G, Tenore GC, Khan H, Micucci M, Budriesi R, Momtaz S, Nabavi SM, Bishayee A. Targeting Inflammation by Flavonoids: Novel Therapeutic Strategy for Metabolic Disorders. Int J Mol Sci 2019; 20 [PMID: 31597283 DOI: 10.3390/ijms20194957]
- Cozma-Petruț A, Loghin F, Miere D, Dumitrașcu DL. Diet in irritable bowel syndrome: What to recommend, not what to 62 forbid to patients! World J Gastroenterol 2017; 23: 3771-3783 [PMID: 28638217 DOI: 10.3748/wjg.v23.i21.3771]
- 63 Senn V, Bassler D, Choudhury R, Scholkmann F, Righini-Grunder F, Vuille-Dit-Bile RN, Restin T. Microbial Colonization From the Fetus to Early Childhood-A Comprehensive Review. Front Cell Infect Microbiol 2020; 10: 573735 [PMID: 33194813 DOI: 10.3389/fcimb.2020.573735]



- Cerdó T, García-Santos JA, Rodríguez-Pöhnlein A, García-Ricobaraza M, Nieto-Ruíz A, G Bermúdez M, Campoy C. 64 Impact of Total Parenteral Nutrition on Gut Microbiota in Pediatric Population Suffering Intestinal Disorders. Nutrients 2022; **14** [PMID: 36364953 DOI: 10.3390/nu14214691]
- 65 Takiishi T, Fenero CIM, Câmara NOS. Intestinal barrier and gut microbiota: Shaping our immune responses throughout life. Tissue Barriers 2017; 5: e1373208 [PMID: 28956703 DOI: 10.1080/21688370.2017.1373208]
- Wennerberg J, Sharma S, Nilsson PM, Ohlsson B. A possible association between early life factors and burden of 66 functional bowel symptoms in adulthood. Scand J Prim Health Care 2021; 39: 506-514 [PMID: 34806535 DOI: 10.1080/02813432.2021.2004823
- Chitkara DK, van Tilburg MA, Blois-Martin N, Whitehead WE. Early life risk factors that contribute to irritable bowel 67 syndrome in adults: a systematic review. Am J Gastroenterol 2008; 103: 765-74; quiz 775 [PMID: 18177446 DOI: 10.1111/j.1572-0241.2007.01722.x
- Collado MC, Cernada M, Neu J, Pérez-Martínez G, Gormaz M, Vento M. Factors influencing gastrointestinal tract and 68 microbiota immune interaction in preterm infants. Pediatr Res 2015; 77: 726-731 [PMID: 25760550 DOI: 10.1038/pr.2015.54]
- Cilieborg MS, Boye M, Sangild PT. Bacterial colonization and gut development in preterm neonates. Early Hum Dev 69 2012; 88 Suppl 1: S41-S49 [PMID: 22284985 DOI: 10.1016/j.earlhumdev.2011.12.027]
- Cassir N, Simeoni U, La Scola B. Gut microbiota and the pathogenesis of necrotizing enterocolitis in preterm neonates. 70 Future Microbiol 2016; 11: 273-292 [PMID: 26855351 DOI: 10.2217/fmb.15.136]
- Sproat T, Payne RP, Embleton ND, Berrington J, Hambleton S. T Cells in Preterm Infants and the Influence of Milk Diet. Front Immunol 2020; 11: 1035 [PMID: 32582165 DOI: 10.3389/fimmu.2020.01035]
- Singh P, Alm EJ, Kelley JM, Cheng V, Smith M, Kassam Z, Nee J, Iturrino J, Lembo A. Effect of antibiotic pretreatment on bacterial engraftment after Fecal Microbiota Transplant (FMT) in IBS-D. Gut Microbes 2022; 14: 2020067 [PMID: 35014601 DOI: 10.1080/19490976.2021.2020067]
- Singh P, Lembo A. Emerging Role of the Gut Microbiome in Irritable Bowel Syndrome. Gastroenterol Clin North Am 73 2021; 50: 523-545 [PMID: 34304786 DOI: 10.1016/j.gtc.2021.03.003]
- 74 Zaman S, Lippman SI, Schneper L, Slonim N, Broach JR. Glucose regulates transcription in yeast through a network of signaling pathways. Mol Syst Biol 2009; 5: 245 [PMID: 19225458 DOI: 10.1038/msb.2009.2]
- Amabebe E, Anumba DOC. Female Gut and Genital Tract Microbiota-Induced Crosstalk and Differential Effects of 75 Short-Chain Fatty Acids on Immune Sequelae. Front Immunol 2020; 11: 2184 [PMID: 33013918 DOI: 10.3389/fimmu.2020.02184]
- Hurley E, Mullins D, Barrett MP, O'Shea CA, Kinirons M, Ryan CA, Stanton C, Whelton H, Harris HMB, O'Toole PW. 76 The microbiota of the mother at birth and its influence on the emerging infant oral microbiota from birth to 1 year of age: a cohort study. J Oral Microbiol 2019; 11: 1599652 [PMID: 32128038 DOI: 10.1080/20002297.2019.1599652]
- Tun HM, Bridgman SL, Chari R, Field CJ, Guttman DS, Becker AB, Mandhane PJ, Turvey SE, Subbarao P, Sears MR, Scott JA, Kozyrskyj AL; Canadian Healthy Infant Longitudinal Development (CHILD) Study Investigators. Roles of Birth Mode and Infant Gut Microbiota in Intergenerational Transmission of Overweight and Obesity From Mother to Offspring. JAMA Pediatr 2018; 172: 368-377 [PMID: 29459942 DOI: 10.1001/jamapediatrics.2017.5535]
- De Palma G, Lynch MD, Lu J, Dang VT, Deng Y, Jury J, Umeh G, Miranda PM, Pigrau Pastor M, Sidani S, Pinto-78 Sanchez MI, Philip V, McLean PG, Hagelsieb MG, Surette MG, Bergonzelli GE, Verdu EF, Britz-McKibbin P, Neufeld JD, Collins SM, Bercik P. Transplantation of fecal microbiota from patients with irritable bowel syndrome alters gut function and behavior in recipient mice. Sci Transl Med 2017; 9 [PMID: 28251905 DOI: 10.1126/scitranslmed.aaf6397]
- Craven LJ, Silverman M, Burton JP. Transfer of altered behaviour and irritable bowel syndrome with diarrhea (IBS-D) 79 through fecal microbiota transplant in mouse model indicates need for stricter donor screening criteria. Ann Transl Med 2017; 5: 490 [PMID: 29299452 DOI: 10.21037/atm.2017.10.03]
- Juncadella AC, Moss A. Fecal microbiota transplantation as a possible treatment of irritable bowel syndrome. Ann Transl 80 Med 2017; 5: 506 [PMID: 29299467 DOI: 10.21037/atm.2017.09.13]
- Human Microbiome Project Consortium. Structure, function and diversity of the healthy human microbiome. Nature 81 2012; 486: 207-214 [PMID: 22699609 DOI: 10.1038/nature11234]
- Vlasova AN, Kandasamy S, Chattha KS, Rajashekara G, Saif LJ. Comparison of probiotic lactobacilli and bifidobacteria 82 effects, immune responses and rotavirus vaccines and infection in different host species. Vet Immunol Immunopathol 2016; 172: 72-84 [PMID: 26809484 DOI: 10.1016/j.vetimm.2016.01.003]
- 83 Clapp M, Aurora N, Herrera L, Bhatia M, Wilen E, Wakefield S. Gut microbiota's effect on mental health: The gut-brain axis. Clin Pract 2017; 7: 987 [PMID: 29071061 DOI: 10.4081/cp.2017.987]
- Kolodziejczak M, Béchade C, Gervasi N, Irinopoulou T, Banas SM, Cordier C, Rebsam A, Roumier A, Maroteaux L. 84 Serotonin Modulates Developmental Microglia via 5-HT2B Receptors: Potential Implication during Synaptic Refinement of Retinogeniculate Projections. ACS Chem Neurosci 2015; 6: 1219-1230 [PMID: 25857335 DOI: 10.1021/cn5003489]
- 85 Rothhammer V, Kenison JE, Li Z, Tjon E, Takenaka MC, Chao CC, Alves de Lima K, Borucki DM, Kaye J, Quintana FJ. Aryl Hydrocarbon Receptor Activation in Astrocytes by Laquinimod Ameliorates Autoimmune Inflammation in the CNS. Neurol Neuroimmunol Neuroinflamm 2021; 8 [PMID: 33408169 DOI: 10.1212/NXI.00000000000946]
- Chen Y, Palm F, Lesch KP, Gerlach M, Moessner R, Sommer C. 5-hydroxyindolacetic acid (5-HIAA), a main metabolite 86 of serotonin, is responsible for complete Freund's adjuvant-induced thermal hyperalgesia in mice. Mol Pain 2011; 7: 21 [PMID: 21447193 DOI: 10.1186/1744-8069-7-21]
- Neufeld KM, Kang N, Bienenstock J, Foster JA. Reduced anxiety-like behavior and central neurochemical change in 87 germ-free mice. Neurogastroenterol Motil 2011; 23: 255-264, e119 [PMID: 21054680 DOI: 10.1111/j.1365-2982.2010.01620.x]
- Lugo-Huitrón R, Ugalde Muñiz P, Pineda B, Pedraza-Chaverrí J, Ríos C, Pérez-de la Cruz V. Quinolinic acid: an 88 endogenous neurotoxin with multiple targets. Oxid Med Cell Longev 2013; 2013: 104024 [PMID: 24089628 DOI: 10.1155/2013/104024]
- 89 Alharthi A, Alhazmi S, Alburae N, Bahieldin A. The Human Gut Microbiome as a Potential Factor in Autism Spectrum



Disorder. Int J Mol Sci 2022; 23 [PMID: 35163286 DOI: 10.3390/ijms23031363]

- 90 Chernikova MA, Flores GD, Kilroy E, Labus JS, Mayer EA, Aziz-Zadeh L. The Brain-Gut-Microbiome System: Pathways and Implications for Autism Spectrum Disorder. Nutrients 2021; 13 [PMID: 34960049 DOI: 10.3390/nu13124497
- Rutsch A, Kantsjö JB, Ronchi F. The Gut-Brain Axis: How Microbiota and Host Inflammasome Influence Brain 91 Physiology and Pathology. Front Immunol 2020; 11: 604179 [PMID: 33362788 DOI: 10.3389/fimmu.2020.604179]
- Dobson GP, Letson HL, Biros E, Morris J. Specific pathogen-free (SPF) animal status as a variable in biomedical 92 research: Have we come full circle? EBioMedicine 2019; 41: 42-43 [PMID: 30803932 DOI: 10.1016/j.ebiom.2019.02.038]
- Graham BM, Milad MR. The study of fear extinction: implications for anxiety disorders. Am J Psychiatry 2011; 168: 93 1255-1265 [PMID: 21865528 DOI: 10.1176/appi.ajp.2011.11040557]
- 94 Ahmed H, Leyrolle Q, Koistinen V, Kärkkäinen O, Layé S, Delzenne N, Hanhineva K. Microbiota-derived metabolites as drivers of gut-brain communication. Gut Microbes 2022; 14: 2102878 [PMID: 35903003 DOI: 10.1080/19490976.2022.2102878
- Doroszkiewicz J, Groblewska M, Mroczko B. The Role of Gut Microbiota and Gut-Brain Interplay in Selected Diseases 95 of the Central Nervous System. Int J Mol Sci 2021; 22 [PMID: 34576191 DOI: 10.3390/ijms221810028]
- Rea K, Dinan TG, Cryan JF. The microbiome: A key regulator of stress and neuroinflammation. Neurobiol Stress 2016; 4: 96 23-33 [PMID: 27981187 DOI: 10.1016/j.ynstr.2016.03.001]
- 97 Zubcevic J, Richards EM, Yang T, Kim S, Sumners C, Pepine CJ, Raizada MK. Impaired Autonomic Nervous System-Microbiome Circuit in Hypertension. Circ Res 2019; 125: 104-116 [PMID: 31219753 DOI: 10.1161/CIRCRESAHA.119.313965]
- Stephens MA, Wand G. Stress and the HPA axis: role of glucocorticoids in alcohol dependence. Alcohol Res 2012; 34: 468-483 [PMID: 23584113]
- Lund ML, Egerod KL, Engelstoft MS, Dmytriyeva O, Theodorsson E, Patel BA, Schwartz TW. Enterochromaffin 5-HT 00 cells - A major target for GLP-1 and gut microbial metabolites. Mol Metab 2018; 11: 70-83 [PMID: 29576437 DOI: 10.1016/j.molmet.2018.03.004]
- Reigstad CS, Salmonson CE, Rainey JF 3rd, Szurszewski JH, Linden DR, Sonnenburg JL, Farrugia G, Kashyap PC. Gut 100 microbes promote colonic serotonin production through an effect of short-chain fatty acids on enterochromaffin cells. FASEB J 2015; 29: 1395-1403 [PMID: 25550456 DOI: 10.1096/fj.14-259598]
- 101 Martin AM, Lumsden AL, Young RL, Jessup CF, Spencer NJ, Keating DJ. The nutrient-sensing repertoires of mouse enterochromaffin cells differ between duodenum and colon. Neurogastroenterol Motil 2017; 29 [PMID: 28251760 DOI: 10.1111/nmo.13046
- Jones LA, Sun EW, Martin AM, Keating DJ. The ever-changing roles of serotonin. Int J Biochem Cell Biol 2020; 125: 102 105776 [PMID: 32479926 DOI: 10.1016/j.biocel.2020.105776]
- 103 Li Q, Ren Y, Fu X. Inter-kingdom signaling between gut microbiota and their host. Cell Mol Life Sci 2019; 76: 2383-2389 [PMID: 30911771 DOI: 10.1007/s00018-019-03076-7]
- 104 Gupta A, Singh AK, Kumar R, Jamieson S, Pandey AK, Bishayee A. Neuroprotective Potential of Ellagic Acid: A Critical Review. Adv Nutr 2021; 12: 1211-1238 [PMID: 33693510 DOI: 10.1093/advances/nmab007]
- Fung TC, Olson CA, Hsiao EY. Interactions between the microbiota, immune and nervous systems in health and disease. 105 Nat Neurosci 2017; 20: 145-155 [PMID: 28092661 DOI: 10.1038/nn.4476]
- 106 Cox LM, Weiner HL. Microbiota Signaling Pathways that Influence Neurologic Disease. Neurotherapeutics 2018; 15: 135-145 [PMID: 29340928 DOI: 10.1007/s13311-017-0598-8]
- Pithadia AB, Jain SM. 5-Hydroxytryptamine Receptor Subtypes and their Modulators with Therapeutic Potentials. J Clin 107 Med Res 2009; 1: 72-80 [PMID: 22505971 DOI: 10.4021/jocmr2009.05.1237]
- Jacobsen JP, Medvedev IO, Caron MG. The 5-HT deficiency theory of depression: perspectives from a naturalistic 5-HT 108 deficiency model, the tryptophan hydroxylase 2Arg439His knockin mouse. Philos Trans R Soc Lond B Biol Sci 2012; 367: 2444-2459 [PMID: 22826344 DOI: 10.1098/rstb.2012.0109]
- Zaniewska M, Mosienko V, Bader M, Alenina N. Tph2 Gene Expression Defines Ethanol Drinking Behavior in Mice. 109 Cells 2022; 11 [PMID: 35269497 DOI: 10.3390/cells11050874]
- Koopman N, Katsavelis D, Hove AST, Brul S, Jonge WJ, Seppen J. The Multifaceted Role of Serotonin in Intestinal 110 Homeostasis. Int J Mol Sci 2021; 22 [PMID: 34502396 DOI: 10.3390/ijms22179487]
- Gaster LM, King FD. Serotonin 5-HT3 and 5-HT4 receptor antagonists. Med Res Rev 1997; 17: 163-214 [PMID: 111 9057164 DOI: 10.1002/(sici)1098-1128(199703)17:2<163::aid-med2>3.0.co;2-y]
- Margolis KG, Li Z, Stevanovic K, Saurman V, Israelyan N, Anderson GM, Snyder I, Veenstra-VanderWeele J, Blakely 112 RD, Gershon MD. Serotonin transporter variant drives preventable gastrointestinal abnormalities in development and function. J Clin Invest 2016; 126: 2221-2235 [PMID: 27111230 DOI: 10.1172/JCI84877]
- Blatter LA, McGuigan JA. Estimation of the upper limit of the free magnesium concentration measured with Mg-sensitive 113 microelectrodes in ferret ventricular muscle: (1) use of the Nicolsky-Eisenman equation and (2) in calibrating solutions of the appropriate concentration. Magnesium 1988; 7: 154-165 [DOI: 10.1113/expphysiol.1986.sp003005]
- Waclawiková B, Bullock A, Schwalbe M, Aranzamendi C, Nelemans SA, van Dijk G, El Aidy S. Gut bacteria-derived 5-114 hydroxyindole is a potent stimulant of intestinal motility via its action on L-type calcium channels. PLoS Biol 2021; 19: e3001070 [PMID: 33481771 DOI: 10.1371/journal.pbio.3001070]
- Taniya MA, Chung HJ, Al Mamun A, Alam S, Aziz MA, Emon NU, Islam MM, Hong SS, Podder BR, Ara Mimi A, 115 Aktar Suchi S, Xiao J. Role of Gut Microbiome in Autism Spectrum Disorder and Its Therapeutic Regulation. Front Cell Infect Microbiol 2022; 12: 915701 [PMID: 35937689 DOI: 10.3389/fcimb.2022.915701]
- 116 Wang J, Xu W, Wang R, Cheng R, Tang Z, Zhang M. The outer membrane protein Amuc_1100 of Akkermansia muciniphila promotes intestinal 5-HT biosynthesis and extracellular availability through TLR2 signalling. Food Funct 2021; 12: 3597-3610 [PMID: 33900345 DOI: 10.1039/d1fo00115a]
- Modasia A, Parker A, Jones E, Stentz R, Brion A, Goldson A, Defernez M, Wileman T, Ashley Blackshaw L, Carding 117 SR. Regulation of Enteroendocrine Cell Networks by the Major Human Gut Symbiont Bacteroides thetaiotaomicron. Front



Microbiol 2020; 11: 575595 [PMID: 33240233 DOI: 10.3389/fmicb.2020.575595]

- 118 Engevik MA, Luck B, Visuthranukul C, Ihekweazu FD, Engevik AC, Shi Z, Danhof HA, Chang-Graham AL, Hall A, Endres BT, Haidacher SJ, Horvath TD, Haag AM, Devaraj S, Garey KW, Britton RA, Hyser JM, Shroyer NF, Versalovic J. Human-Derived Bifidobacterium dentium Modulates the Mammalian Serotonergic System and Gut-Brain Axis. Cell Mol Gastroenterol Hepatol 2021; 11: 221-248 [PMID: 32795610 DOI: 10.1016/j.jcmgh.2020.08.002]
- Wheatcroft J, Wakelin D, Smith A, Mahoney CR, Mawe G, Spiller R. Enterochromaffin cell hyperplasia and decreased 119 serotonin transporter in a mouse model of postinfectious bowel dysfunction. Neurogastroenterol Motil 2005; 17: 863-870 [PMID: 16336502 DOI: 10.1111/j.1365-2982.2005.00719.x]
- 120 Tsuruta T, Saito S, Osaki Y, Hamada A, Aoki-Yoshida A, Sonoyama K. Organoids as an exvivo model for studying the serotonin system in the murine small intestine and colon epithelium. Biochem Biophys Res Commun 2016; 474: 161-167 [PMID: 27105910 DOI: 10.1016/j.bbrc.2016.03.165]
- Tatsuoka M, Osaki Y, Ohsaka F, Tsuruta T, Kadota Y, Tochio T, Hino S, Morita T, Sonoyama K. Consumption of 121 indigestible saccharides and administration of Bifidobacterium pseudolongum reduce mucosal serotonin in murine colonic mucosa. Br J Nutr 2022; 127: 513-525 [PMID: 33849681 DOI: 10.1017/S0007114521001306]
- 122 Mandić AD, Woting A, Jaenicke T, Sander A, Sabrowski W, Rolle-Kampcyk U, von Bergen M, Blaut M. Clostridium ramosum regulates enterochromaffin cell development and serotonin release. Sci Rep 2019; 9: 1177 [PMID: 30718836 DOI: 10.1038/s41598-018-38018-z]
- Yano JM, Yu K, Donaldson GP, Shastri GG, Ann P, Ma L, Nagler CR, Ismagilov RF, Mazmanian SK, Hsiao EY. 123 Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis. Cell 2015; 161: 264-276 [PMID: 25860609 DOI: 10.1016/j.cell.2015.02.047]
- Nzakizwanayo J, Dedi C, Standen G, Macfarlane WM, Patel BA, Jones BV. Escherichia coli Nissle 1917 enhances 124 bioavailability of serotonin in gut tissues through modulation of synthesis and clearance. Sci Rep 2015; 5: 17324 [PMID: 26616662 DOI: 10.1038/srep17324]
- Shajib MS, Wang H, Kim JJ, Sunjic I, Ghia JE, Denou E, Collins M, Denburg JA, Khan WI. Interleukin 13 and serotonin: 125 linking the immune and endocrine systems in murine models of intestinal inflammation. PLoS One 2013; 8: e72774 [PMID: 24015275 DOI: 10.1371/journal.pone.0072774]
- Cao YN, Feng LJ, Wang BM, Jiang K, Li S, Xu X, Wang WQ, Zhao JW, Wang YM. Lactobacillus acidophilus and Bifidobacterium longum supernatants upregulate the serotonin transporter expression in intestinal epithelial cells. Saudi J Gastroenterol 2018; 24: 59-66 [PMID: 29451186 DOI: 10.4103/sig.SJG 333 17]
- 127 Ranuh R, Athiyyah AF, Darma A, Risky VP, Riawan W, Surono IS, Sudarmo SM. Effect of the probiotic Lactobacillus plantarum IS-10506 on BDNF and 5HT stimulation: role of intestinal microbiota on the gut-brain axis. Iran J Microbiol 2019; 11: 145-150 [PMID: 31341569]
- Chen CM, Wu CC, Huang CL, Chang MY, Cheng SH, Lin CT, Tsai YC. Lactobacillus plantarum PS128 Promotes 128 Intestinal Motility, Mucin Production, and Serotonin Signaling in Mice. Probiotics Antimicrob Proteins 2022; 14: 535-545 [PMID: 34327633 DOI: 10.1007/s12602-021-09814-3]
- Cao YN, Feng LJ, Liu YY, Jiang K, Zhang MJ, Gu YX, Wang BM, Gao J, Wang ZL, Wang YM. Effect of Lactobacillus 129 rhamnosus GG supernatant on serotonin transporter expression in rats with post-infectious irritable bowel syndrome. World J Gastroenterol 2018; 24: 338-350 [PMID: 29391756 DOI: 10.3748/wjg.v24.i3.338]
- 130 Yip JLK, Balasuriya GK, Spencer SJ, Hill-Yardin EL. The Role of Intestinal Macrophages in Gastrointestinal Homeostasis: Heterogeneity and Implications in Disease. Cell Mol Gastroenterol Hepatol 2021; 12: 1701-1718 [PMID: 34506953 DOI: 10.1016/j.jcmgh.2021.08.021]
- Fukumoto S, Tatewaki M, Yamada T, Fujimiya M, Mantyh C, Voss M, Eubanks S, Harris M, Pappas TN, Takahashi T. 131 Short-chain fatty acids stimulate colonic transit via intraluminal 5-HT release in rats. Am J Physiol Regul Integr Comp Physiol 2003; 284: R1269-R1276 [PMID: 12676748 DOI: 10.1152/ajpregu.00442.2002]
- Mishima Y, Ishihara S. Enteric Microbiota-Mediated Serotonergic Signaling in Pathogenesis of Irritable Bowel 132 Syndrome. Int J Mol Sci 2021; 22 [PMID: 34638577 DOI: 10.3390/ijms221910235]
- Crouzet L, Gaultier E, Del'Homme C, Cartier C, Delmas E, Dapoigny M, Fioramonti J, Bernalier-Donadille A. The 133 hypersensitivity to colonic distension of IBS patients can be transferred to rats through their fecal microbiota. Neurogastroenterol Motil 2013; 25: e272-e282 [PMID: 23433203 DOI: 10.1111/nmo.12103]
- Barbara G, Wang B, Stanghellini V, de Giorgio R, Cremon C, Di Nardo G, Trevisani M, Campi B, Geppetti P, Tonini M, 134 Bunnett NW, Grundy D, Corinaldesi R. Mast cell-dependent excitation of visceral-nociceptive sensory neurons in irritable bowel syndrome. Gastroenterology 2007; 132: 26-37 [PMID: 17241857 DOI: 10.1053/j.gastro.2006.11.039]
- Luczynski P, Tramullas M, Viola M, Shanahan F, Clarke G, O'Mahony S, Dinan TG, Cryan JF. Microbiota regulates visceral pain in the mouse. Elife 2017; 6 [PMID: 28629511 DOI: 10.7554/eLife.25887]
- Bai T, Zhang L, Wang H, Qian W, Song J, Hou X. Fecal Microbiota Transplantation Is Effective in Relieving Visceral 136 Hypersensitivity in a Postinfectious Model. Biomed Res Int 2018; 2018: 3860743 [PMID: 29992140 DOI: 10.1155/2018/3860743
- Ciapala K, Mika J, Rojewska E. The Kynurenine Pathway as a Potential Target for Neuropathic Pain Therapy Design: 137 From Basic Research to Clinical Perspectives. Int J Mol Sci 2021; 22 [PMID: 34681715 DOI: 10.3390/ijms222011055]
- Staats Pires A, Tan VX, Heng B, Guillemin GJ, Latini A. Kynurenine and Tetrahydrobiopterin Pathways Crosstalk in 138 Pain Hypersensitivity. Front Neurosci 2020; 14: 620 [PMID: 32694973 DOI: 10.3389/fnins.2020.00620]
- 139 Mawe GM, Hoffman JM. Serotonin signalling in the gut--functions, dysfunctions and therapeutic targets. Nat Rev Gastroenterol Hepatol 2013; 10: 473-486 [PMID: 23797870 DOI: 10.1038/nrgastro.2013.105]
- Min YW, Rhee PL. The clinical potential of ramosetron in the treatment of irritable bowel syndrome with diarrhea (IBS-140 D). Therap Adv Gastroenterol 2015; 8: 136-142 [PMID: 25949526 DOI: 10.1177/1756283X15572580]
- Machu TK. Therapeutics of 5-HT3 receptor antagonists: current uses and future directions. Pharmacol Ther 2011; 130: 141 338-347 [PMID: 21356241 DOI: 10.1016/j.pharmthera.2011.02.003]
- Carco C, Young W, Gearry RB, Talley NJ, McNabb WC, Roy NC. Increasing Evidence That Irritable Bowel Syndrome 142



and Functional Gastrointestinal Disorders Have a Microbial Pathogenesis. *Front Cell Infect Microbiol* 2020; **10**: 468 [PMID: 33014892 DOI: 10.3389/fcimb.2020.00468]

- 143 Ng QX, Soh AYS, Loke W, Lim DY, Yeo WS. The role of inflammation in irritable bowel syndrome (IBS). J Inflamm Res 2018; 11: 345-349 [PMID: 30288077 DOI: 10.2147/JIR.S174982]
- 144 Khan WI, Ghia JE. Gut hormones: emerging role in immune activation and inflammation. *Clin Exp Immunol* 2010; 161: 19-27 [PMID: 20408856 DOI: 10.1111/j.1365-2249.2010.04150.x]
- 145 Kwon YH, Wang H, Denou E, Ghia JE, Rossi L, Fontes ME, Bernier SP, Shajib MS, Banskota S, Collins SM, Surette MG, Khan WI. Modulation of Gut Microbiota Composition by Serotonin Signaling Influences Intestinal Immune Response and Susceptibility to Colitis. *Cell Mol Gastroenterol Hepatol* 2019; 7: 709-728 [PMID: 30716420 DOI: 10.1016/j.jcmgh.2019.01.004]
- 146 Li N, Ghia JE, Wang H, McClemens J, Cote F, Suehiro Y, Mallet J, Khan WI. Serotonin activates dendritic cell function in the context of gut inflammation. *Am J Pathol* 2011; **178**: 662-671 [PMID: 21281798 DOI: 10.1016/j.ajpath.2010.10.028]
- 147 Lucarini E, Di Pilato V, Parisio C, Micheli L, Toti A, Pacini A, Bartolucci G, Baldi S, Niccolai E, Amedei A, Rossolini GM, Nicoletti C, Cryan JF, O'Mahony SM, Ghelardini C, Di Cesare Mannelli L. Visceral sensitivity modulation by faecal microbiota transplantation: the active role of gut bacteria in pain persistence. *Pain* 2022; 163: 861-877 [PMID: 34393197 DOI: 10.1097/j.pain.0000000002438]
- 148 Morita H, Mochiki E, Takahashi N, Kawamura K, Watanabe A, Sutou T, Ogawa A, Yanai M, Ogata K, Fujii T, Ohno T, Tsutsumi S, Asao T, Kuwano H. Effects of 5-HT2B, 5-HT3 and 5-HT4 receptor antagonists on gastrointestinal motor activity in dogs. *World J Gastroenterol* 2013; 19: 6604-6612 [PMID: 24151388 DOI: 10.3748/wjg.v19.i39.6604]
- 149 Melhem H, Kaya B, Ayata CK, Hruz P, Niess JH. Metabolite-Sensing G Protein-Coupled Receptors Connect the Diet-Microbiota-Metabolites Axis to Inflammatory Bowel Disease. *Cells* 2019; 8 [PMID: 31091682 DOI: 10.3390/cells8050450]
- 150 Kerckhoffs AP, ter Linde JJ, Akkermans LM, Samsom M. SERT and TPH-1 mRNA expression are reduced in irritable bowel syndrome patients regardless of visceral sensitivity state in large intestine. *Am J Physiol Gastrointest Liver Physiol* 2012; **302**: G1053-G1060 [PMID: 22323131 DOI: 10.1152/ajpgi.00153.2011]
- Slifirski G, Król M, Turło J. 5-HT Receptors and the Development of New Antidepressants. Int J Mol Sci 2021; 22 [PMID: 34445721 DOI: 10.3390/ijms22169015]
- 152 Liu N, Sun S, Wang P, Sun Y, Hu Q, Wang X. The Mechanism of Secretion and Metabolism of Gut-Derived 5-Hydroxytryptamine. Int J Mol Sci 2021; 22 [PMID: 34360695 DOI: 10.3390/ijms22157931]
- Fung TC, Vuong HE, Luna CDG, Pronovost GN, Aleksandrova AA, Riley NG, Vavilina A, McGinn J, Rendon T, Forrest LR, Hsiao EY. Intestinal serotonin and fluoxetine exposure modulate bacterial colonization in the gut. *Nat Microbiol* 2019; 4: 2064-2073 [PMID: 31477894 DOI: 10.1038/s41564-019-0540-4]
- Gershon MD. 5-Hydroxytryptamine (serotonin) in the gastrointestinal tract. *CurrOpin Endocrinol Diabetes Obes* 2013;
 20: 14-21 [PMID: 23222853 DOI: 10.1097/MED.0b013e32835bc703]
- 155 Ahluwalia B, Iribarren C, Magnusson MK, Sundin J, Clevers E, Savolainen O, Ross AB, Törnblom H, Simrén M, Öhman L. A Distinct Faecal Microbiota and Metabolite Profile Linked to Bowel Habits in Patients with Irritable Bowel Syndrome. *Cells* 2021; 10 [PMID: 34200772 DOI: 10.3390/cells10061459]
- 156 Dalile B, Van Oudenhove L, Vervliet B, Verbeke K. The role of short-chain fatty acids in microbiota-gut-brain communication. Nat Rev Gastroenterol Hepatol 2019; 16: 461-478 [PMID: 31123355 DOI: 10.1038/s41575-019-0157-3]
- 157 Bromke MA, Krzystek-Korpacka M. Bile Acid Signaling in Inflammatory Bowel Disease. Int J Mol Sci 2021; 22 [PMID: 34445800 DOI: 10.3390/ijms22169096]
- Ma C, Vasu R, Zhang H. The Role of Long-Chain Fatty Acids in Inflammatory Bowel Disease. *Mediators Inflamm* 2019; 2019: 8495913 [PMID: 31780872 DOI: 10.1155/2019/8495913]
- 159 Heimerl S, Moehle C, Zahn A, Boettcher A, Stremmel W, Langmann T, Schmitz G. Alterations in intestinal fatty acid metabolism in inflammatory bowel disease. *Biochim Biophys Acta* 2006; 1762: 341-350 [PMID: 16439103 DOI: 10.1016/j.bbadis.2005.12.006]
- 160 Parada Venegas D, De la Fuente MK, Landskron G, González MJ, Quera R, Dijkstra G, Harmsen HJM, Faber KN, Hermoso MA. Short Chain Fatty Acids (SCFAs)-Mediated Gut Epithelial and Immune Regulation and Its Relevance for Inflammatory Bowel Diseases. *Front Immunol* 2019; 10: 277 [PMID: 30915065 DOI: 10.3389/fimmu.2019.00277]
- 161 Gasaly N, Hermoso MA, Gotteland M. Butyrate and the Fine-Tuning of Colonic Homeostasis: Implication for Inflammatory Bowel Diseases. Int J Mol Sci 2021; 22 [PMID: 33802759 DOI: 10.3390/ijms22063061]
- 162 Feng Z, Long W, Hao B, Ding D, Ma X, Zhao L, Pang X. A human stool-derived Bilophilawadsworthia strain caused systemic inflammation in specific-pathogen-free mice. *Gut Pathog* 2017; 9: 59 [PMID: 29090023 DOI: 10.1186/s13099-017-0208-7]
- 163 Lefever S, Van Den Bossche D, Van Moerkercke W, D'Hondt M, Alegret Pampols MDC, Struyve M, De Bel A, Boudewijns M. Ruminococcus gnavus bacteremia, an uncommon presentation of a common member of the human gut microbiota: case report and literature review. *Acta Clin Belg* 2019; 74: 435-438 [PMID: 31658898 DOI: 10.1080/17843286.2018.1541045]
- 164 Barrett E, Fitzgerald P, Dinan TG, Cryan JF, Ross RP, Quigley EM, Shanahan F, Kiely B, Fitzgerald GF, O'Toole PW, Stanton C. Bifidobacterium breve with α-linolenic acid and linoleic acid alters fatty acid metabolism in the maternal separation model of irritable bowel syndrome. *PLoS One* 2012; 7: e48159 [PMID: 23185248 DOI: 10.1371/journal.pone.0048159]

Zaishideng® WJCC | https://www.wjgnet.com



Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

