

World Journal of *Orthopedics*

World J Orthop 2023 April 18; 14(4): 171-267



REVIEW

- 171 Meniscus tears treatment: The good, the bad and the ugly-patterns classification and practical guide
Simonetta R, Russo A, Palco M, Costa GG, Mariani PP

MINIREVIEWS

- 186 Effects of different pelvic osteotomies on acetabular morphology in developmental dysplasia of hip in children
Wen Z, Wu YY, Kuang GY, Wen J, Lu M
- 197 Background, techniques, applications, current trends, and future directions of minimally invasive endoscopic spine surgery: A review of literature
Tang K, Goldman S, Avrumova F, Lebl DR

ORIGINAL ARTICLE**Basic Study**

- 207 Bridge plating in the setting of radiocarpal instability: Does distal fixation to the second or third metacarpal matter? A cadaveric study
Tabeyo E, Saucedo JM, Srinivasan RC, Shah AR, Karamanos E, Rockwood J, Rodriguez-Merchan EC

Retrospective Cohort Study

- 218 Clinical outcomes of cemented distal femur replacements with all-polyethylene tibial components for oncologic indications
Christ AB, Chung BC, Urness M, Mayer LW, Gettleman BS, Heckmann ND, Menendez LR
- 231 Acute hospital-community hospital care bundle for elderly orthopedic surgery patients: A propensity score-matched economic analysis
Tan IEH, Chok AY, Zhao Y, Chen Y, Koo CH, Aw J, Soh MHT, Foo CH, Ang KA, Tan EJKW, Tan AHC, Au MKH

Retrospective Study

- 240 Knowledge and attitudes of orthopedic surgeons regarding prosthesis joint infection
Aytekin MN, Hasanoglu I, Öztürk R, Tosun N

Observational Study

- 248 Perceived barriers and facilitators of day-case surgery for major foot and ankle procedures? A cross-sectional survey of United Kingdom surgeons
Berry A, Houchen-Wolloff L, Crane N, Townshend D, Clayton R, Mangwani J

CASE REPORT

- 260 Subclinical ankle joint tuberculous arthritis - The role of scintigraphy: A case series
Primadhi RA, Kartamihardja AHS

ABOUT COVER

Editorial Board Member of *World Journal of Orthopedics*, Saubhik Das, MS, Assistant Professor, Department of Orthopaedics, Rajendra Institute of Medical Sciences, Ranchi 834009, Jharkhand, India. drsaubhikk@hotmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Orthopedics* (WJO, *World J Orthop*) is to provide scholars and readers from various fields of orthopedics with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJO mainly publishes articles reporting research results and findings obtained in the field of orthopedics and covering a wide range of topics including arthroscopy, bone trauma, bone tumors, hand and foot surgery, joint surgery, orthopedic trauma, osteoarthritis, osteoporosis, pediatric orthopedics, spinal diseases, spine surgery, and sports medicine.

INDEXING/ABSTRACTING

WJO is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 Journal Citation Indicator (JCI) for WJO as 0.62. The WJO's CiteScore for 2021 is 2.4 and Scopus CiteScore rank 2021: Orthopedics and Sports Medicine is 139/284.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Ying-Yi Yuan; **Production Department Director:** Xiang Li; **Editorial Office Director:** Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Orthopedics

ISSN

ISSN 2218-5836 (online)

LAUNCH DATE

November 18, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Massimiliano Leigheb

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/2218-5836/editorialboard.htm>

PUBLICATION DATE

April 18, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/gerinfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/gerinfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/gerinfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Effects of different pelvic osteotomies on acetabular morphology in developmental dysplasia of hip in children

Zhi Wen, Yu-Yuan Wu, Gao-Yan Kuang, Jie Wen, Min Lu

Specialty type: Pediatrics

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): D

Grade E (Poor): 0

P-Reviewer: Maj MK, Malaysia; Ning B, China

Received: December 29, 2022

Peer-review started: December 29, 2022

First decision: January 9, 2023

Revised: January 19, 2023

Accepted: March 30, 2023

Article in press: March 30, 2023

Published online: April 18, 2023



Zhi Wen, Graduate School, Hunan University of Chinese Medicine, Changsha 410007, Hunan Province, China

Zhi Wen, Gao-Yan Kuang, Min Lu, Department of Joint Orthopedics, The First Hospital of Hunan University of Chinese Medicine, Changsha 410007, Hunan Province, China

Yu-Yuan Wu, Department of Pediatric Orthopedics, Traditional Chinese Medicine Hospital in Huaihua, Huaihua 418000, Hunan Province, China

Jie Wen, Department of Pediatric Orthopedics, Hunan Provincial People's Hospital, The First Affiliated Hospital of Hunan Normal University, Changsha 410013, Hunan Province, China

Corresponding author: Jie Wen, PhD, Associate Professor, Department of Pediatric Orthopedics, Hunan Provincial People's Hospital, The First Affiliated Hospital of Hunan Normal University, No. 61 West Jiefang Road, Changsha 410013, Hunan Province, China.
cashwj@qq.com

Abstract

Developmental dysplasia of hip seriously affects the health of children, and pelvic osteotomy is an important part of surgical treatment. Improving the shape of the acetabulum, preventing or delaying the progression of osteoarthritis is the ultimate goal of pelvic osteotomies. Re-directional osteotomies, reshaping osteotomies and salvage osteotomies are the three most common types of pelvic osteotomy. The influence of different pelvic osteotomy on acetabular morphology is different, and the acetabular morphology after osteotomy is closely related to the prognosis of the patients. But there lacks comparison of acetabular morphology between different pelvic osteotomies, on the basis of retrospective analysis and measurable imaging indicators, this study predicted the acetabular shape after developmental dysplasia of the hip pelvic osteotomy in order to help clinicians make reasonable and correct decisions and improve the planning and performance of pelvic osteotomy.

Key Words: Developmental dysplasia of the hip; Pelvic osteotomy; Acetabular morphology; Re-directional osteotomies; Reshaping osteotomies; Salvage osteotomies

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Developmental dysplasia of hip seriously affects the health of children, and pelvic osteotomy is an important part of surgical treatment. Improving the shape of the acetabulum, preventing or delaying the progression of osteoarthritis is the ultimate goal of pelvic osteotomies. This manuscript reviewed influences of different pelvic osteotomy on acetabular morphology and the acetabular morphology after osteotomy. Accurate prediction of acetabular shape after pelvic osteotomies can help clinicians to make reasonable and correct decisions and improve the planning and performance of pelvic osteotomies.

Citation: Wen Z, Wu YY, Kuang GY, Wen J, Lu M. Effects of different pelvic osteotomies on acetabular morphology in developmental dysplasia of hip in children. *World J Orthop* 2023; 14(4): 186-196

URL: <https://www.wjgnet.com/2218-5836/full/v14/i4/186.htm>

DOI: <https://dx.doi.org/10.5312/wjo.v14.i4.186>

INTRODUCTION

Developmental dysplasia of the hip (DDH) leads to loss of the normal anatomical relationship between the femoral head and acetabulum due to developmental defects of the acetabulum, resulting in abnormal morphology and position of the hip joint[1-3]. DDH is the most common hip deformity in children. The global incidence rate is 1‰–2‰. Surgical correction is still the most challenging problem for pediatric orthopedic surgeons[4,5]. Delayed diagnosis of DDH means that more complex and higher failure rates of treatment are needed, so early diagnosis and treatment are important[6]. The main goal of DDH treatment is to achieve and maintain concentric reduction of the femoral head and acetabulum, to ensure the normal development of the hip joint, and to avoid osteonecrosis of the femoral head and reoperation[7,8]. Surgical treatment of DDH is usually determined according to the age, imaging examination and clinical symptoms of the child. In patients with progressive acetabular dysplasia, pelvic osteotomy is necessary to minimize the risk of hip dislocation in adults[9]. Common surgical procedures for DDH patients include pelvic osteotomy, open reduction, capsular arthroplasty and femoral osteotomy[10]. To restore the anatomical morphology and mechanical structure of the pelvis and ensure normal development and life of children, pelvic osteotomy is almost inevitable[11,12]. Morphology of the acetabulum after pelvic osteotomy is closely related to prognosis of DDH. Many researchers have proposed their own methods of pelvic osteotomy that have their own characteristics, indications and clinical effects. There are three common types of pelvic osteotomy: Redirectional, reshaping and salvage[11]. On the basis of retrospective analysis and measurable imaging indicators, this study predicted the acetabular shape after DDH pelvic osteotomy in order to help clinicians make reasonable and correct decisions and improve the planning and performance of pelvic osteotomy.

NORMAL ACETABULAR MORPHOLOGY IN CHILDREN

The hip joint consists of a round head of the femur and a cup-shaped acetabulum, which forms a unique ball and fossa shape that helps stabilize the joint and allows a dynamic range of rotational motion[13]. The normal acetabular structure consists of the upper ilium, the anterior pubic bone and the posterior ischial bone. In early infants, the acetabulum consists of a cartilage complex composed of Y-type and acetabular cartilage. The acetabular cartilage is located at the outer two thirds of the acetabular fossa, and the Y-type cartilage is located at the junction of the pubic bone, ischium and ilium[14,15]. With the growth and development of infants, the ossification center in acetabular cartilage appears and develops successively; the pubic ossification center develops and forms the anterior wall of the acetabulum; the ossification center of the ischium develops and forms the posterior wall of the acetabulum; and the ossification center of the ilium develops and forms the superior wall of the acetabulum. The cartilage complex of the normal acetabulum fuses at the age of 12–16 years, completely forming the fossa of the joint, and covering the entire femoral head. The stress direction determines whether the growing hip joint tends to concentric growth, and determines the morphology and structure of the acetabulum under stress[16]. The acetabular cup usually faces downward, lateral and anterior, opposite to the upper, medial and lateral direction of the femoral head, forming a stable spherical joint structure of the hip joint[17]. Acetabular cartilage is interrupted in the center of the acetabular fossa and is also known as lunate cartilage because it is similar to a crescent shape. Lunate cartilage is located on the lateral and superior articular surface of the acetabulum, which is thicker and stronger than femoral head cartilage[18]. The glenoid labrum is a fibrocartilage structure that surrounds the outer edge of the bony structure of the acetabulum, which functionally deepens the acetabular cup and improves the stability of the hip joint[19,20].

ACETABULAR MORPHOLOGY DURING DDH

Morphological changes of the acetabulum and proximal femur during DDH includes acetabular cartilage degeneration, acetabular fossa shallower, acetabular roof tilted, acetabulum lost its normal and nearly round shape[21,22]. Subluxation or dislocation of the hip is characterized by widening of teardrops, loss of lateral depression of the acetabulum, elongation of the posterior upper edge of the acetabulum, increase of the ratio of long diameter to short diameter of the acetabulum, and loss of normal concentric circle relationship between the proximal femur and the acetabulum[5,23]. Due to the reduction of the contact area between the acetabulum and femoral head, the load of the acetabulum increases, and the structure of the acetabulum propagates laterally under stress, resulting in excessive anteversion of the femoral neck and varying degrees of hip varus[24,25]. The structural changes of the hip joint also cause compensatory changes in the pelvis and spine, leading to a variety of clinical symptoms. It was believed that dislocation of the hip joint during DDH was caused by insufficient depth and excessive tilt of the acetabulum, which could not accommodate the femoral head. However, recent studies have shown that the development between the acetabulum and femoral head follows the Wolff rule[26]. Accordingly, the arrangement of bone trabeculae is affected by the dynamic distribution of bone load, and the mechanical pressure on the bone is conducive to stimulating bone development. When the acetabulum loses the centripetal pressure from the femoral head, the structural development of the acetabulum is affected, and the depth, width and inclination of the acetabulum become abnormal, and even acetabular cartilage degenerates. With the development of DDH, the anatomical morphology of the acetabulum is completely lost and the hip joint is totally dislocated. In the case of total dislocation of the hip, the normal acetabulum loses the pressure from the femoral head, resulting in hypertrophy of the articular pelvis, and the hypertrophic pelvis adheres to the joint capsule and ligaments, which further affects the reduction of the femoral head.

EVALUATION OF ACETABULAR MORPHOLOGY BY COMPUTED TOMOGRAPHY

Computed tomography (CT) is a widely used method for evaluation of the hip joint. Because acetabular dysplasia is not just a simple malrotation or anterolateral defect, three-dimensional (3D) CT imaging can help doctors to determine the exact shape of the acetabulum[27]. Because of the simplicity and speed of obtaining images, CT has become an important method for the diagnosis of hip joint structural changes, preoperative evaluation and postoperative re-examination[25]. The traditional 3D reconstruction of CT can observe the acetabular morphology of DDH in many directions, accurately measuring the acetabulum, and carrying out further quantitative analysis, so as to better evaluate the pathological changes of DDH and morphology of the acetabulum after osteotomy. Some parameters that cannot be measured by conventional X-ray films can be measured by 3D CT, such as bony acetabular index and anteversion. Therefore, CT and 3D reconstruction scanning have important guiding significance for clinicians. The bone reconstruction model of 3D CT provides an omnidirectional and multiangle observation index for clinicians to observe the pathological changes of the hip joint and the curative effect of surgery, and has value for guiding surgical planning and prediction of postoperative acetabular shape[28]. Li *et al*[29] used computer-aided design for quantitative dynamic analysis of hip joint morphology, and found that it was a better guide for surgical planning and evaluation of treatment results. Some clinical studies have found that the value and safety of using 3D CT to guide pelvic osteotomy is higher than that of ordinary CT[30]. In addition, 3D CT can also be combined with finite element analysis method to carry out biomechanical analysis of acetabular morphology after DDH, which has important guiding significance for the evaluation of surgical efficacy and improvement of surgical methods of DDH[31].

EVALUATION OF ACETABULAR MORPHOLOGY BY X-RAY FILM

X-ray film can dynamically observe the changes of bone structure during child growth and development, which is the most basic imaging method to evaluate the structural changes of the hip joint, and is important for guiding screening and evaluation of acetabular morphology[32-34]. The following are the main parameters in the evaluation of acetabular morphology by X-ray film[35,36]: Degree of acetabular anteversion, central edge angle (CEA), Acetabular index (AI), sharp angle, angle of iliac wall, and acetabular coverage. The normal acetabulum is usually tilted forward, and it is of clinical importance when the angle is increased by $> 7^\circ$ compared with normal[24]. CEA is an important index to describe the shape of the acetabulum[37,38]. It is normal when the CEA is $> 25^\circ$, but abnormal when it is $< 20^\circ$, suggesting that the acetabulum is incomplete. When CEA is between 20° and 25° , it is necessary to closely follow up and observe its changing trends. AI is an objective measurement index of acetabular dysplasia, which can be used to judge the severity and therapeutic effect of DDH. The normal range of AI before age 2 years is 20° - 25° , and AI $> 30^\circ$ after 2 years is considered abnormal[38, 39]. The sharp angle reflects the development of the acetabulum and coverage of the hip joint on the

femoral head. It is used to diagnose and predict the progress of hip dislocation. The normal value is 33°–38°. Dynamic observation is needed when the value is 42°–47°, and abnormality can be diagnosed when it is > 47°[40]. The angle of the iliac wall is the angle between the line from the anterior inferior iliac spine to the outer upper edge of the acetabulum and the Y-shaped cartilage line on both sides of the acetabulum. The normal value of the infant is about 55°. The iliac wall angle increases slowly with infant age, and there is little difference between the normal value and the outlier range, so it is necessary to compare the two sides on the X-ray film[41]. The acetabular coverage rate[42] reflects accommodation of the acetabulum to the femoral head. It is the ratio of the distance from the inner edge of the femoral head to the vertical line of the upper edge of the acetabulum to the diameter of the femoral head, and the normal value is > 75%. X-ray film is a simple, fast and economic examination method, which can quickly diagnose DDH from a large number of pelvic films, effectively reduce the misdiagnosis rate and evaluating the shape of the pelvis and acetabulum after surgery.

PELVIC OSTEOTOMY AND ACETABULAR MORPHOLOGY

The ability of acetabular remodeling has an age limit, and previous studies have shown that the acetabulum can be remodeled under the age of 5 years[43]. Acetabular dysplasia over a certain age needs to be corrected by pelvic osteotomy. Different methods of pelvic osteotomy depend on the age of DDH patients and the degree of deformity[44,45]. Normal development of the acetabulum depends on the femoral head under the concentric circle position, but DDH changes the biomechanical structure of the hip joint, which limits the morphological development of the acetabulum in children, so the purpose of the operation is to restore the concentric reduction of the hip joint[46]. Pelvic osteotomy can improve deformity of the acetabulum, increase the stress area of the hip joint, restore the coverage of acetabular cartilage, and reconstruct the biomechanical relationship of the hip joint[47]. When acetabular dysplasia is associated with subluxation or dislocation, pelvic osteotomy is necessary to minimize the risk of hip dislocation in adults[11]. The older the onset of DDH, the worse the prognosis. It is generally believed that osteotomy has a lot of complications in patients aged > 8 years[48]. The effects of different pelvic osteotomy on acetabular morphology are different, which provides a basis for different patients to formulate specific osteotomy methods[49]. The current imaging technology has been able to measure the relevant indicators of acetabular morphology, but the operations process of different patients can't be total the same, so it is difficult to measure acetabular morphology accurately by different osteotomies. Previous researchers used the lateral CEA to assess acetabular developmental defects, but Daniel *et al*[50] found that acetabular deficiency in borderline hip dysplasia can be underestimated by lateral CEA alone. Some researchers used 3D printing technology to measure the relevant indexes of acetabular morphology after different osteotomies. The real proportional anatomical model can accurately measure the AI before and after surgery, and it provides conditions for optimizing the surgical plan and improving the method. Researchers had described many types of pelvic osteotomy. Different pelvic osteotomies have different advantages and indications, and the shape of the acetabulum after osteotomy is also different. There are three common types of pelvic osteotomy: Redirectional, reshaping and salvage[11]. We will describe the effects of the different types of pelvic osteotomy on the morphology of the acetabulum.

REDIRECTIONAL OSTEOTOMY

Redirectional osteotomy of the pelvis changes the direction of the acetabulum, increases the contact area between the acetabulum and the femoral head, and increases the area of hyaline cartilage in the weight-bearing area of the acetabulum. It is characterized by repositioning the acetabulum to improve the anterolateral acetabular cover without changing the size and shape of the hip, which is an incomplete osteotomy[51]. The common redirectional osteotomies include Salter, periacetabular and triple pelvic osteotomy.

Salter osteotomy

Salter osteotomy is a classic method of pelvic osteotomy, which belongs to the osteotomies that change the direction of the acetabulum. It was first proposed by Salter in 1961[52]. The osteotomy line of Salter is from the anterior inferior iliac spine to the great ischial notch, and the entire acetabulum is turned forward and outward with the pubic symphysis as the fulcrum. It changes the overall orientation of the acetabulum to increase the inclusiveness of the acetabulum to the femoral head, and the shape and volume of the acetabulum remain unchanged[3]. Therefore, Salter osteotomy is suitable for the following[52,53]: (1) Children aged 18 mo to 6 years with unossified pubic symphysis; (2) anterolateral acetabular defect as the main defect; and (3) AI is basically normal. Salter osteotomy requires internal fixation support, so the wedge-shaped bone removed from the ilium is placed into the fracture line and fixed with Kirschner wire to increase stability of the acetabulum after osteotomy. In the Salter

procedure, the rotation angle and distance of the distal iliac bone are not only the difficulties of the operation, but also the key to its effectiveness. Previous studies have shown that Salter osteotomy changes the acetabular center with the movement of the distal bone mass, and the AI decreases yearly. At 3–4 years after the operation, the AI is stable to about 15°, the CEA increases yearly, and returns to the normal range for about 5 years, and the postoperative obturator area is significantly reduced[54].

Periacetabular osteotomy

Periacetabular osteotomy, also known as GANZ osteotomy, is suitable for patients with hip pain and when the femoral head is located in the center of the acetabulum, and the degree of hip deformity and arthritis is low. It was first proposed by Ganz in 1983[55]. The periacetabular osteotomy procedure is mainly to amputate the pubic bone, ischium and ilium, and under the condition of maintaining the continuity of the pelvis, rotate the acetabulum to make the acetabulum cover the femoral head to the best extent[56]. Because periacetabular osteotomy does not destroy the acetabular blood supply and the pelvic ring, it can improve the biomechanical relationship between the acetabulum and femoral head, reduce the probability of postoperative acetabular necrosis, and avoid the occurrence of hip arthritis [45]. Andrew *et al*[57] found that among athletes with symptomatic hip dysplasia who received periacetabular osteotomy, all acetabular parameters were improved, and the return to exercise rate of competitive athletes was > 70%. Periacetabular osteotomy allows the acetabulum to be relocated in multiple directions to correct the deficiency of the anterior coverage of the femoral head, so even severe acetabular deformities can be satisfactorily corrected by accurate rotation and internal displacement of the acetabular center. Because periacetabular osteotomy maintains the integrity of the posterior column of the acetabulum, better initial stability is obtained[55,58]. Through long-term follow-up, researchers have found that in the patients treated with periacetabular osteotomy, the fracture healed well, the lateral coverage of the femoral head, AI angle and Sharp angle were significantly improved, the cartilage area of the acetabular weight-bearing area was increased, and the gait and hip joint function were significantly improved.

Triple pelvic osteotomy

Triple pelvic osteotomy, which includes osteotomy of the ilium, pubis and ischium around the acetabulum, is a complete redirected osteotomy, which solves the problems of limited movement of bone mass and lateralization of the hip joint after osteotomy[59,60]. This complete redirected osteotomy increases the range of motion of the acetabulum by cutting off the pubic bone, ischium and ilium, so that the rotated acetabulum can completely contain the femoral head and achieve concentric reduction of the hip joint. At present, there are three common triple pelvic osteotomy procedures, which are LeCoeur, Steel and Tonis[61]. By amputating the ilium and the superior and inferior branches of the pubis near the symphysis pubis, the LeCoeur procedure limits the rotation of the acetabulum because the position of the amputated pubis and ischium is far from the acetabulum[11]. Steel osteotomy cuts the ischial bone at the ischial tubercle through the posterior approach, and the pubic bone and ilium through the anterior approach. Because the osteotomy is close to the acetabulum, the range of acetabular adjustment is better than that of LeCoeur. The position of the Tonnis osteotomy is closer to the acetabulum than the Steel osteotomy is, so the acetabular rotation is greater[62]. Like other redirected osteotomies, triple pelvic osteotomy is a rotational osteotomy that does not change the size and shape of the acetabulum [63]. The premise of this operation is that the structure of the hip joint is intact, which is mainly suitable for older children with more severe acetabular dysplasia, and it is an unstable osteotomy that requires solid internal and external fixation[59,64]. Triple pelvic osteotomy can improve the CEA, lateral coverage of the femoral head, and stress distribution of acetabular cartilage, and can effectively correct acetabular deformities.

RESHAPING OSTEOTOMIES

Compared with pelvic redirection osteotomies, pelvic reshaping osteotomies are incomplete, which increases the coverage of the acetabulum and femoral head by bending acetabular vertices[65]. Therefore, this kind of operation is also called acetabuloplasty. Because it is an incomplete osteotomy, the position of the bone mass after operation is stable and usually does not require internal fixation. There are three common methods of reshaping osteotomies, namely, Pemberton, Dega and SanDiego. Researchers[49] compared the effects of these three osteotomies on acetabular morphology using 3D-printed models, and found that Pemberton and Dega increased the upper and anterior coverage of the acetabulum, resulting in retroversion of the acetabulum, while SanDiego increased the posterior coverage of the acetabulum, resulting in acetabular anteversion.

Pemberton pelvic osteotomy

Pemberton pelvic osteotomy is a periarticular osteotomy, also known as Pemberton acetabuloplasty, which was first proposed by Pemberton in 1968[66,67]. Pemberton pelvic osteotomy has a wide range of adaptation, and can be used in children under 12 years old, but because it affects development of the

pelvis and pubic symphysis, caution is needed in children under 6 years old. Pemberton pelvic osteotomy takes the Y-type cartilage around the acetabulum as the rotation fulcrum and controls the shape of the acetabulum by adjusting the position of the bone cortex on the posterior side of the ilium. It increases the coverage of the anterior edge of the acetabulum, and significantly increases the coverage of the outer upper edge and posterior edge of the acetabulum[68]. Pemberton pelvic osteotomy can adjust acetabular direction, correct acetabular deformity and increase acetabular depth according to the degree of acetabular defect. Therefore, it can reduce AI, increase the coverage of acetabulum and femoral head, stabilize the hip joint, and reduce deformity of the acetabulum[69,70]. Because the Pemberton procedure can reduce AI, it is suitable for cases where the acetabulum needs to be corrected by $> 15^\circ$ and the acetabulum is shallow and steep.

Degapelvic osteotomy

Dega osteotomy is one of the commonly used osteotomies in the treatment of DDH. It was first proposed and applied in clinic Dega in 1958[71,72]. The Dega procedure is a type of acetabuloplasty to change the shape and inclination of the acetabulum, mainly to increase the lateral coverage of the acetabulum, which can change the volume of the acetabulum without damaging the Y-type cartilage [73]. Dega operation uses the iliac bone of incomplete osteotomy above the Y-type cartilage as a hinge to change the direction and inclination of the acetabulum. Its advantage is that the position of iliac osteotomy can be adjusted according to the location of the acetabular defect, so that acetabular defects in different parts can be corrected[74]. A retrospective analysis[75] showed that AI decreased significantly in patients who underwent Dega acetabuloplasty, and postoperative Sharp angle was also improved, which can improve AI and inclusiveness of the femoral head.

San Diego pelvic osteotomy

SanDiego osteotomy is a modified Dega osteotomy to improve lateral and posterior acetabular coverage [11,76]. The SanDiego operation is an incomplete osteotomy. The medial iliac bone cortex is intact except for the anterior and posterior sides, and the ischial and pubic cortices are also continuous[51]. In iliac osteotomy, the wedge-shaped bone can be used to stretch the osteotomy surface, so this kind of monocortical osteotomy enables doctors to change the shape of the acetabulum by changing the size and position of the bone graft, thus solving the problem of insufficient acetabular coverage[76]. Because San Diego osteotomy mainly increases coverage of the acetabulum, the shape of the acetabulum tilts forward after osteotomy, the volume of the acetabulum decreases, and the depth of the acetabulum increases.

SALVAGE PELVIC OSTEOTOMIES

If DDH is untreated or inadequately treated, it aggravates the deformity of the hip joint, which eventually develops into a nonfunctional joint with pain symptoms[77]. At that time, the change in the bony structure of the hip joint is more serious, the shaping ability of the bone decreases, the shape of the femoral head and acetabulum are difficult to reduce by simple methods, and in severe cases, cartilage injury of the hip joint occurs[78]. At that time, surgeons should pay attention to recovery of hip joint function and perform salvage surgery, so as to delay the progress of arthritis and reduce the probability of osteonecrosis of the femoral head. The common salvage pelvic osteotomies are Chiari and Shelf.

Chiari pelvic osteotomy

Chiari osteotomy, also known as intrapelvic osteotomy, was first proposed and applied by Chiari *et al* [79] in 1974. Chiari osteotomy changes the weight-bearing line of the hip joint and increases the weight-bearing area by moving the position of iliac osteotomy. It is a type of joint capsule plasty, which belongs to single-plane osteotomy[80]. Therefore, Chiari operation is suitable for older children with abnormal acetabular-femoral head index, non concentric reduction of the femoral head, acetabular deformity and lack of plastic potential[81,82]. The Chiari procedure cuts the ilium from front to back from the area between the acetabulum and the anterior inferior iliac spine along the ischial notch, and then the bone mass of the distal end of the ilium is moved inward and upward. In this process, it is necessary to ensure the coverage of the acetabulum to the femoral head and avoid the impact between the femoral head and ilium[51,78]. The coverage of the acetabulum depends on the width of the ilium. In order to achieve satisfactory coverage of the lateral acetabulum, it is usually necessary to move the distal bone completely inward to the width of the ilium. If there is insufficient containment of the anterior side of the femoral head, iliac alar osteotomy can be performed, then coverage on the anterior side of the femoral head was improved. The Chiari procedure distributes the weight-bearing area of the acetabulum by expanding the inclusion of the acetabulum to the femoral head, but the damaged acetabular cartilage and glenoid labrum may still be located in the weight-bearing area after salvage surgery. Therefore, although this operation can better restore the acetabular-femoral head index, it cannot solve the defect in front of the acetabulum, and shortens the transverse diameter of the pelvis, resulting in unsatisfactory surgical results[83].

Table 1 Summation of previous studies with their case number and conclusions

Ref.	Pelvic osteotomies	Number of cases/hip	Conclusion
Sawamura <i>et al</i> [52]	Salter osteotomy	42/50	Shown favorable outcomes with satisfactory at skeletal maturity
Scott <i>et al</i> [53]		58/78	Improve acetabular morphology, reduced the incidence of advanced osteoarthritis and the probability of THA
Tan <i>et al</i> [55]	PAO	?/225	Lead to good deformity correction for both acetabular retroversion and hip dysplasia
Ali <i>et al</i> [56]		314/?	All acetabular parameters were observed to be improved
Dornacher <i>et al</i> [63]	TPO	176/206	Acetabular parameters recovered well, intraoperative fluoroscopy increased the accuracy of the operation
Lyu <i>et al</i> [64]		43/48	Improve parameters of the acetabulum, makes the complex operation safer and more effective
Bhatti <i>et al</i> [67]	PPO	60/82	The hips with Pemberton's acetabuloplasty exhibited better acetabular coverage and progressive development of hips as compared to Salter's osteotomy
Czubak <i>et al</i> [71]	Dega osteotomy	45/52	It is a safe and adequate operation to restore the acetabulum to normal or close to normal
Badrinath <i>et al</i> [76]	San Diego osteotomy	?/45	It is a safe and effective alternative to treat acetabular dysplasia in patients with typical DDH and can solve specific acetabular defects
Ito <i>et al</i> [82]	Chiari pelvic osteotomy	31/32	Improve acetabular index, relieve pain and reduce hip replacement rate
Dammerer <i>et al</i> [81]		12/12	Improve the acetabular coverage of the hip joint and suitable for some specific older children
Yoo <i>et al</i> [84]	Shelf pelvic osteotomy	14/14	Improve the depth and coverage of acetabulum and can stimulate the growth of acetabulum
Benad <i>et al</i> [85]		56/61	Relieve pain symptoms, improve joint function and delay THA in patients

PAO: Periacetabular osteotomy; TPO: Triple pelvic osteotomy; PPO: Pemberton pelvic osteotomy; THA: Total hip arthroplasty; DDH: Developmental dysplasia of the hip.

Shelf pelvic osteotomy

Shelf osteotomy, also known as acetabular extension, is mainly suitable for older children with femoral head and acetabular mismatch and hip dislocation. The anterior, posterior and lateral weight-bearing surface of the acetabulum is enlarged by osteotomy of the ilium and bone grafting at the fracture line, so the acetabular coverage of the femoral head is increased[11]. Shelf operation is a salvage operation, which is suitable for older children who have failed nonoperative treatment of DDH[84]. Shelf acetabuloplasty has a favorable, stimulatory effect on acetabular growth[85]. The operation does not change the direction or biomechanical structure of the acetabulum, but increases the volume, so it can improve stability of the hip joint. After Shelf osteotomy, the central marginal angle of the hip joint increases, the coverage of the acetabulum to the femoral head increases, and the acetabular shaping is improved[86].

CONCLUSION

Pelvic osteotomy is an important method for surgical treatment of DDH. There are three common types of pelvic osteotomy: redirection, reshaping and salvage. The choice of pelvic osteotomy depends on the patient's age, acetabular development and disease severity. No operation is the best, and no operation is suitable for all cases. It is particularly important for pediatric orthopedic surgeons to choose the osteotomy with which they are most familiar and which is most suitable for patients. Different methods of pelvic osteotomy have different effects on acetabular morphology, and the acetabular morphology is closely related to prognosis (Table 1). It is still one of the most challenging problems for pediatric orthopedic surgeons to master the principles of various pelvic osteotomies and to understand the morphological changes in the acetabulum after pelvic osteotomy. At present, the evaluation of acetabular morphology after pelvic osteotomy mainly depends on ordinary X-ray or 3D CT reconstruction, but the measurement of acetabular morphology is not accurate because of the non-replicability of different patients and the individual differences of observers. In the future, the acetabular morphology of different pelvic osteotomy methods will be measured by 3D printing, and the morphological changes in the acetabulum before and after surgery will be accurately measured by real proportional anatomical models, so as to provide a basis for optimizing the surgical plan and improving the osteotomy method.

FOOTNOTES

Author contributions: Wen J conceived and coordinated the study, Wen J and Lu M designed the study, Wen Z wrote the paper; Wu YY did literature review, Kuang GY revised the paper; Wen J and Lu M contribute equally to this study, they share co-corresponding author; All authors reviewed the results and approved the final version of the manuscript.

Supported by Scientific Research Project of Hunan Education Department, No. 21A0054.

Conflict-of-interest statement: All the authors declare that they have no competing interests.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Yu-Yuan Wu 0000-0002-2694-1680; Jie Wen 0000-0002-5734-4678.

S-Editor: Liu JH

L-Editor: A

P-Editor: Zhao S

REFERENCES

- Li L, Wang X, Zhao Q, Wang E, Wang L, Cheng J, Zhang L, Wang B. CX3CR1 polymorphisms associated with an increased risk of developmental dysplasia of the hip in human. *J Orthop Res* 2017; **35**: 377-380 [PMID: 27176135 DOI: 10.1002/jor.23294]
- Yang S, Zusman N, Lieberman E, Goldstein RY. Developmental Dysplasia of the Hip. *Pediatrics* 2019; **143** [PMID: 30587534 DOI: 10.1542/peds.2018-1147]
- Yang G, Zhang Z, Hou X, Luo D, Cheng H, Xiao K, Liu H, Zhang H. A New Parameter of Hip Instability in Developmental Dysplasia of the Hip (DDH): Teardrop Distance. *Front Surg* 2022; **9**: 899960 [PMID: 36034389 DOI: 10.3389/fsurg.2022.899960]
- Sewell MD, Rosendahl K, Eastwood DM. Developmental dysplasia of the hip. *BMJ* 2009; **339**: b4454 [PMID: 19934187 DOI: 10.1136/bmj.b4454]
- Schaeffer EK, Study Group I, Mulpuri K. Developmental dysplasia of the hip: addressing evidence gaps with a multicentre prospective international study. *Med J Aust* 2018; **208**: 359-364 [PMID: 29716513 DOI: 10.5694/mja18.00154]
- von Kries R, Ihme N, Altenhofen L, Niethard FU, Krauspe R, Rückinger S. General ultrasound screening reduces the rate of first operative procedures for developmental dysplasia of the hip: a case-control study. *J Pediatr* 2012; **160**: 271-275 [PMID: 21962602 DOI: 10.1016/j.jpeds.2011.08.037]
- Kotlarsky P, Haber R, Bialik V, Eidelman M. Developmental dysplasia of the hip: What has changed in the last 20 years? *World J Orthop* 2015; **6**: 886-901 [PMID: 26716085 DOI: 10.5312/wjo.v6.i11.886]
- Rees HW. Management of Osteoarthritis of the Hip. *J Am Acad Orthop Surg* 2020; **28**: e288-e291 [PMID: 31800436 DOI: 10.5435/JAAOS-D-19-00416]
- Moraleda L, Bravo C, Forriol F, Albiñana J. Does Orientation of the Femoral Head Affect Acetabular Development? *J Pediatr Orthop* 2019; **39**: 416-421 [PMID: 31393302 DOI: 10.1097/BPO.0000000000000974]
- Chen M, Shang XF. Surgical treatment for young adult hip dysplasia: joint-preserving options. *Int Orthop* 2016; **40**: 891-900 [PMID: 26216530 DOI: 10.1007/s00264-015-2927-z]
- Venkatadass K, Durga Prasad V, Al Ahmadi NMM, Rajasekaran S. Pelvic osteotomies in hip dysplasia: why, when and how? *EFORT Open Rev* 2022; **7**: 153-163 [PMID: 35192506 DOI: 10.1530/EOR-21-0066]
- Pisecky L, Großbötzl G, Stevoska S, Klotz MCM, Haas C, Gotterbarm T, Luger M, Gahleitner M. Short Term Radiological Outcome of Combined Femoral and Ilium Osteotomy in Pelvic Reconstruction of the Child. *Children (Basel)* 2022; **9** [PMID: 35327813 DOI: 10.3390/children9030441]
- Birkenmaier C, Jorysz G, Jansson V, Heimkes B. Normal development of the hip: a geometrical analysis based on planimetric radiography. *J Pediatr Orthop B* 2010; **19**: 1-8 [PMID: 19829156 DOI: 10.1097/BPB.0b013e32832f5aeb]
- Krebs V, Incavo SJ, Shields WH. The anatomy of the acetabulum: what is normal? *Clin Orthop Relat Res* 2009; **467**: 868-875 [PMID: 18648904 DOI: 10.1007/s11999-008-0317-1]
- Harris JD, Lewis BD, Park KJ. Hip Dysplasia. *Clin Sports Med* 2021; **40**: 271-288 [PMID: 33673886 DOI: 10.1016/j.csm.2020.11.004]
- Heimkes B, Wegener V, Birkenmaier C, Ziegler CM. Physiologic and Pathologic Development of the Infantile and Adolescent Hip Joint: Descriptive and Functional Aspects. *Semin Musculoskelet Radiol* 2019; **23**: 477-488 [PMID: 31556083 DOI: 10.1055/s-0039-1693975]
- Nogami R, Kaku N, Shimada T, Tabata T, Tagomori H, Tsumura H. Three-dimensional architecture of the acetabular labrum in the human hip joint. *Med Mol Morphol* 2020; **53**: 21-27 [PMID: 31144043 DOI: 10.1007/s00795-019-00228-3]

- 18 **Müller M**, Rakow A, Wassilew GI, Winkler T, Perka C. Prediction of time to prosthesis implantation as a function of joint anatomy in patients with developmental dysplasia of the hip. *J Orthop Surg Res* 2019; **14**: 471 [PMID: 31888687 DOI: 10.1186/s13018-019-1511-4]
- 19 **Lertwanich P**, Plakseychuk A, Kramer S, Linde-Rosen M, Maeyama A, Fu FH, Smolinski P. Biomechanical evaluation contribution of the acetabular labrum to hip stability. *Knee Surg Sports Traumatol Arthrosc* 2016; **24**: 2338-2345 [PMID: 25749654 DOI: 10.1007/s00167-015-3555-2]
- 20 **Johannsen AM**, Ejnisman L, Behn AW, Shibata K, Thio T, Safran MR. Contributions of the Capsule and Labrum to Hip Mechanics in the Context of Hip Microinstability. *Orthop J Sports Med* 2019; **7**: 2325967119890846 [PMID: 31903404 DOI: 10.1177/2325967119890846]
- 21 **Gala L**, Clohisy JC, Beaulé PE. Hip Dysplasia in the Young Adult. *J Bone Joint Surg Am* 2016; **98**: 63-73 [PMID: 26738905 DOI: 10.2106/JBJS.O.00109]
- 22 **Troelsen A**. Assessment of adult hip dysplasia and the outcome of surgical treatment. *Dan Med J* 2012; **59**: B4450 [PMID: 22677250]
- 23 **Schwend RM**, Shaw BA, Segal LS. Evaluation and treatment of developmental hip dysplasia in the newborn and infant. *Pediatr Clin North Am* 2014; **61**: 1095-1107 [PMID: 25439013 DOI: 10.1016/j.pcl.2014.08.008]
- 24 **Cai Z**, Zhao Q, Li L, Zhang L, Ji S. Can Computed Tomography Accurately Measure Acetabular Anterversion in Developmental Dysplasia of the Hip? *J Pediatr Orthop* 2018; **38**: e180-e185 [PMID: 29356794 DOI: 10.1097/BPO.0000000000001141]
- 25 **Jia J**, Li L, Zhang L, Zhao Q, Liu X. Three dimensional-CT evaluation of femoral neck anteversion, acetabular anteversion and combined anteversion in unilateral DDH in an early walking age group. *Int Orthop* 2012; **36**: 119-124 [PMID: 21858684 DOI: 10.1007/s00264-011-1337-0]
- 26 **Erridge S**, Goh EL, Chidambaram S. Subclinical developmental dysplasia of the hip could predispose to femoroacetabular impingement. *Med Hypotheses* 2017; **102**: 144-145 [PMID: 28478819 DOI: 10.1016/j.mehy.2017.03.028]
- 27 **Ramo BA**, De La Rocha A, Sucato DJ, Jo CH. A New Radiographic Classification System for Developmental Hip Dysplasia is Reliable and Predictive of Successful Closed Reduction and Late Pelvic Osteotomy. *J Pediatr Orthop* 2018; **38**: 16-21 [PMID: 26866641 DOI: 10.1097/BPO.0000000000000733]
- 28 **Frick SL**, Kim SS, Wenger DR. Pre- and postoperative three-dimensional computed tomography analysis of triple innominate osteotomy for hip dysplasia. *J Pediatr Orthop* 2000; **20**: 116-123 [PMID: 10641700]
- 29 **Li H**, Zhang Z, Li C, Liang Z, Liu Z, Li H. Computer-assisted design model to evaluate the outcome of combined osteotomies in Legg-Calvé-Perthes disease. *Front Pediatr* 2022; **10**: 920840 [PMID: 36003490 DOI: 10.3389/fped.2022.920840]
- 30 **van Bosse HJ**, Lee D, Henderson ER, Sala DA, Feldman DS. Pelvic positioning creates error in CT acetabular measurements. *Clin Orthop Relat Res* 2011; **469**: 1683-1691 [PMID: 21365336 DOI: 10.1007/s11999-011-1827-9]
- 31 **Zhao X**, Chosa E, Totoribe K, Deng G. Effect of periacetabular osteotomy for acetabular dysplasia clarified by three-dimensional finite element analysis. *J Orthop Sci* 2010; **15**: 632-640 [PMID: 20953924 DOI: 10.1007/s00776-010-1511-z]
- 32 **Expert Panel on Pediatric Imaging**, Nguyen JC, Dorfman SR, Rigsby CK, Iyer RS, Alazraki AL, Anupindi SA, Bardo DME, Brown BP, Chan SS, Chandra T, Garber MD, Moore MM, Pandya NK, Shet NS, Siegel A, Karmazyn B. ACR Appropriateness Criteria® Developmental Dysplasia of the Hip-Child. *J Am Coll Radiol* 2019; **16**: S94-S103 [PMID: 31054762 DOI: 10.1016/j.jacr.2019.02.014]
- 33 **Babcock DS**, Hernandez RJ, Kushner DC, Cohen HL, Gelfand MJ, McAlister WH, Parker BR, Royal SA, Slovis TL, Smith WL, Strain JD, Strife JL, Tosi L. Developmental dysplasia of the hip. American College of Radiology. ACR Appropriateness Criteria. *Radiology* 2000; **215** Suppl: 819-827 [PMID: 11037505]
- 34 **Nie Y**, Pei F, Shen B, Kang P, Li Z. Implication of acetabular width on the anteroposterior pelvic radiograph of patients with developmental dysplasia of the hip. *J Arthroplasty* 2015; **30**: 489-494 [PMID: 25311162 DOI: 10.1016/j.arth.2014.09.024]
- 35 **Keller MS**, Nijs EL. The role of radiographs and US in developmental dysplasia of the hip: how good are they? *Pediatr Radiol* 2009; **39** Suppl 2: S211-S215 [PMID: 19308388 DOI: 10.1007/s00247-008-1107-3]
- 36 **Sherman B**, Lalonde FD, Schlechter JA. Measuring the Acetabular Index: An Accurate and Reliable Alternative Method of Measurement. *AJR Am J Roentgenol* 2021; **217**: 172-176 [PMID: 33909466 DOI: 10.2214/AJR.20.23358]
- 37 **Li Y**, Guo Y, Shen X, Liu H, Mei H, Xu H, Canavese F; Chinese Multi-center Pediatric Orthopedic Study Group (CMPOS). Radiographic outcome of children older than twenty-four months with developmental dysplasia of the hip treated by closed reduction and spica cast immobilization in human position: a review of fifty-one hips. *Int Orthop* 2019; **43**: 1405-1411 [PMID: 30980114 DOI: 10.1007/s00264-019-04315-z]
- 38 **Li Y**, Guo Y, Li M, Zhou Q, Liu Y, Chen W, Li J, Canavese F, Xu H; Multi-center Pediatric Orthopedic Study Group of China. Acetabular index is the best predictor of late residual acetabular dysplasia after closed reduction in developmental dysplasia of the hip. *Int Orthop* 2018; **42**: 631-640 [PMID: 29285666 DOI: 10.1007/s00264-017-3726-5]
- 39 **Stetzelberger VM**, Leibold CS, Steppacher SD, Schwab JM, Siebenrock KA, Tannast M. The Acetabular Wall Index Is Associated with Long-term Conversion to THA after PAO. *Clin Orthop Relat Res* 2021; **479**: 1052-1065 [PMID: 33605631 DOI: 10.1097/CORR.0000000000001641]
- 40 **Li Q**, Zhong L, Huang H, Liu H, Qin Y, Wang Y, Zhou Z, Yang W, Qin M, Wang J, Zhou T, Wang D, Xu M, Huang Y. Auxiliary diagnosis of developmental dysplasia of the hip by automated detection of Sharp's angle on standardized anteroposterior pelvic radiographs. *Medicine (Baltimore)* 2019; **98**: e18500 [PMID: 31876738 DOI: 10.1097/MD.00000000000018500]
- 41 **Swami VG**, Liu M, Mabee M, Jaremko JL. Normal values and variation of radiographic and CT infant lateral iliac wall angles in normal and dysplastic hips. *Hip Int* 2016; **26**: 602-607 [PMID: 27768217 DOI: 10.5301/hipint.5000393]
- 42 **Alrashdi NZ**, Motl RW, Aguiar EJ, Ryan MK, Perumean-Chaney SE, Ithurburn MP. Mobility-related outcomes for periacetabular osteotomy in persons with acetabular dysplasia: setting the stage for measurement of real-world outcomes. *J Hip Preserv Surg* 2021; **8**: 367-381 [PMID: 35505808 DOI: 10.1093/jhps/hnab086]
- 43 **Miyake T**, Tetsunaga T, Endo H, Yamada K, Sanki T, Fujiwara K, Nakata E, Ozaki T. Predicting acetabular growth in

- developmental dysplasia of the hip following open reduction after walking age. *J Orthop Sci* 2019; **24**: 326-331 [PMID: 30377015 DOI: 10.1016/j.jos.2018.09.015]
- 44 de Courtivron B, Brulefert K, Portet A, Odent T. Residual acetabular dysplasia in congenital hip dysplasia. *Orthop Traumatol Surg Res* 2022; **108**: 103172 [PMID: 34896582 DOI: 10.1016/j.otsr.2021.103172]
 - 45 Ahmad SS, Haertlé M, Konrads C, Derksen A, Windhagen H, Wirries N. The Scientific Evolution of Periacetabular Osteotomy: A Global Review. *J Clin Med* 2022; **11** [PMID: 36294420 DOI: 10.3390/jcm11206099]
 - 46 Vaquero-Picado A, González-Morán G, Garay EG, Moraleda L. Developmental dysplasia of the hip: update of management. *EFORT Open Rev* 2019; **4**: 548-556 [PMID: 31598333 DOI: 10.1302/2058-5241.4.180019]
 - 47 Nepple JJ, Fowler LM, Larson CM. Decision-making in the Borderline Hip. *Sports Med Arthrosc Rev* 2021; **29**: 15-21 [PMID: 33395225 DOI: 10.1097/JSA.0000000000000298]
 - 48 Thomas SR. A review of long-term outcomes for late presenting developmental hip dysplasia. *Bone Joint J* 2015; **97-B**: 729-733 [PMID: 26033050 DOI: 10.1302/0301-620X.97B.35395]
 - 49 Caffrey JP, Jeffords ME, Farnsworth CL, Bomar JD, Upasani VV. Comparison of 3 Pediatric Pelvic Osteotomies for Acetabular Dysplasia Using Patient-specific 3D-printed Models. *J Pediatr Orthop* 2019; **39**: e159-e164 [PMID: 30300278 DOI: 10.1097/BPO.0000000000001271]
 - 50 Dornacher D, Lutz B, Fuchs M, Zippelius T, Reichel H. Acetabular deficiency in borderline hip dysplasia is underestimated by lateral center edge angle alone. *Arch Orthop Trauma Surg* 2022 [PMID: 36271941 DOI: 10.1007/s00402-022-04652-6]
 - 51 Selberg CM, Chidsey B, Skelton A, Mayer S. Pelvic Osteotomies in the Child and Young Adult Hip: Indications and Surgical Technique. *J Am Acad Orthop Surg* 2020; **28**: e230-e237 [PMID: 31714417 DOI: 10.5435/JAAOS-D-19-00223]
 - 52 Sawamura K, Kitoh H, Kaneko H, Kitamura A, Hattori T. Appropriate Surgical Timing of Salter Innominate Osteotomy for Residual Acetabular Dysplasia in Children. *J Pediatr Orthop* 2022; **42**: e971-e975 [PMID: 36040038 DOI: 10.1097/BPO.0000000000002258]
 - 53 Scott EJ, Dolan LA, Weinstein SL. Closed Vs. Open Reduction/Salter Innominate Osteotomy for Developmental Hip Dislocation After Age 18 Months: Comparative Survival at 45-Year Follow-up. *J Bone Joint Surg Am* 2020; **102**: 1351-1357 [PMID: 32769602 DOI: 10.2106/JBJS.19.01278]
 - 54 Sun Y, Zhao L, Yu B, Hu Y, Chang S. [Imaging evaluations of Salter innominate osteotomy for developmental dysplasia of the hip]. *Zhonghua Yi Xue Za Zhi* 2014; **94**: 1080-1082 [PMID: 24851892]
 - 55 Tan SHS, Tan JHI, Lim AKS, Hui JH. Periacetabular osteotomy for acetabular retroversion: A systematic review and meta-analysis. *Orthop Traumatol Surg Res* 2021; **107**: 103078 [PMID: 34583014 DOI: 10.1016/j.otsr.2021.103078]
 - 56 Ali M, Malviya A. Complications and outcome after periacetabular osteotomy - influence of surgical approach. *Hip Int* 2020; **30**: 4-15 [PMID: 31526038 DOI: 10.1177/1120700019871195]
 - 57 Curley AJ, Padmanabhan S, Chishti Z, Parsa A, Jimenez AE, Domb BG. Periacetabular Osteotomy in Athletes With Symptomatic Hip Dysplasia Allows for Participation in Low-, Moderate-, and High-Impact Sports, With Greater Than 70% Return to Sport for Competitive Athletes: A Systematic Review. *Arthroscopy* 2023; **39**: 868-880 [PMID: 36528217 DOI: 10.1016/j.arthro.2022.12.004]
 - 58 Ellsworth BK, Bram JT, Sink EL. Hip Morphology in Periacetabular Osteotomy (PAO) Patients Treated for Developmental Dysplasia of the Hip (DDH) as Infants Compared With Those Without Infant Treatment. *J Pediatr Orthop* 2022; **42**: e565-e569 [PMID: 35667051 DOI: 10.1097/BPO.0000000000002137]
 - 59 Zahedi AR, Lüring C, Janßen D. [Tönnis and Kalchschmidt triple pelvic osteotomy]. *Orthopade* 2016; **45**: 673-677 [PMID: 27385385 DOI: 10.1007/s00132-016-3291-4]
 - 60 Grigoryan G, Korcek L, Eidelman M, Paley D, Nelson S. Direct Lateral Approach for Triple Pelvic Osteotomy. *J Am Acad Orthop Surg* 2020; **28**: e64-e70 [PMID: 31157758 DOI: 10.5435/JAAOS-D-16-00918]
 - 61 Mimura T, Mori K, Kawasaki T, Imai S, Matsusue Y. Triple pelvic osteotomy: Report of our mid-term results and review of literature. *World J Orthop* 2014; **5**: 14-22 [PMID: 24649410 DOI: 10.5312/wjo.v5.i1.14]
 - 62 Tönnis D, Kalchschmidt K, Heinecke A. [Acetabular rotations y triple pelvic osteotomy by the Tönnis method]. *Orthopade* 1998; **27**: 733-742 [PMID: 9871921]
 - 63 Dornacher D, Sgroi M, Freitag T, Reichel H, Lutz B. Acetabular orientation in triple pelvic osteotomy: is intraoperative fluoroscopy reliable? *Arch Orthop Trauma Surg* 2022 [PMID: 35947170 DOI: 10.1007/s00402-022-04568-1]
 - 64 Lyu X, Yang Z, Wang Y, Zhang T, Lu M, Bian Z. Novel Minimally-Invasive Triple Pelvic Osteotomy: JiShuiTan Minimally-Invasive Approach. *J Pediatr Orthop* 2022; **42**: e154-e162 [PMID: 34839311 DOI: 10.1097/BPO.00000000000002019]
 - 65 Gillingham BL, Sanchez AA, Wenger DR. Pelvic osteotomies for the treatment of hip dysplasia in children and young adults. *J Am Acad Orthop Surg* 1999; **7**: 325-337 [PMID: 10504359 DOI: 10.5435/00124635-199909000-00005]
 - 66 Liu KY, Wu KW, Lee CC, Lin SC, Kuo KN, Chang JF, Wang TM. A Long-Term Study of Alignment Correction Following Proximal Femoral Varus Osteotomy and Pemberton Osteotomy in Children With Legg-Calvé-Perthes Disease and Developmental Dysplasia of the Hip. *Front Pediatr* 2022; **10**: 835447 [PMID: 35463899 DOI: 10.3389/fped.2022.835447]
 - 67 Bhatti A, Abbasi I, Naeem Z, Jaffri K, Bhatti MY. A Comparative Study of Salter Versus Pemberton Osteotomy in Open Reduction of Developmental Dysplastic Hips and Clinical Evaluation on Bhatti's Functional Score System. *Cureus* 2021; **13**: e12626 [PMID: 33585115 DOI: 10.7759/cureus.12626]
 - 68 Bulut M, Azboy I, Ozkul E, Karakurt L. Comparison of Iliac and Femoral Autograft Practices in Pemberton Pelvic Osteotomy. *J Pediatr Orthop* 2021; **41**: 46-50 [PMID: 32804869 DOI: 10.1097/BPO.0000000000001665]
 - 69 Yu J, Duan F, Guo W, Wang D, Qin X, Fu G, Chen T. Consistency of Indices Obtained via Hip Medial Ultrasound and Magnetic Resonance Imaging in Reduction and Spica Cast Treatment for Developmental Dysplasia of the Hip. *Ultrasound Med Biol* 2021; **47**: 58-67 [PMID: 33039172 DOI: 10.1016/j.ultrasmedbio.2020.08.010]
 - 70 Wada A, Fujii T, Takamura K, Yanagida H, Taketa M, Nakamura T. Pemberton osteotomy for developmental dysplasia of the hip in older children. *J Pediatr Orthop* 2003; **23**: 508-513 [PMID: 12826952]

- 71 **Czubak J**, Kowalik K, Kawalec A, Kwiatkowska M. Dega pelvic osteotomy: indications, results and complications. *J Child Orthop* 2018; **12**: 342-348 [PMID: 30154924 DOI: 10.1302/1863-2548.12.180091]
- 72 **Krieg AH**, Hefti F. [Acetabuloplasty - The Dega and Pemberton technique]. *Orthopade* 2016; **45**: 653-658 [PMID: 27385388 DOI: 10.1007/s00132-016-3295-0]
- 73 **Jóźwiak M**, Grzegorzewski A, Napiontek M, Woźniak Ł. The significance of the Dega osteotomy. *J Pediatr Orthop B* 2023; **32**: 207-210 [PMID: 36125907 DOI: 10.1097/BPB.0000000000001005]
- 74 **Woźniak Ł**, Idzior M, Jóźwiak M. Dega Transiliac Pelvic Osteotomy: the History of Surgical Technique Development. *Ortop Traumatol Rehabil* 2020; **22**: 455-464 [PMID: 33506802 DOI: 10.5604/01.3001.0014.6057]
- 75 **Wells J**, Millis M, Kim YJ, Bulat E, Miller P, Matheney T. Survivorship of the Bernese Periacetabular Osteotomy: What Factors are Associated with Long-term Failure? *Clin Orthop Relat Res* 2017; **475**: 396-405 [PMID: 27172819 DOI: 10.1007/s11999-016-4887-z]
- 76 **Badrinath R**, Bomar JD, Wenger DR, Mubarak SJ, Upasani VV. Comparing the Pemberton osteotomy and modified San Diego acetabuloplasty in developmental dysplasia of the hip. *J Child Orthop* 2019; **13**: 172-179 [PMID: 30996742 DOI: 10.1302/1863-2548.13.190004]
- 77 **Wyles CC**, Vargas JS, Heidenreich MJ, Mara KC, Peters CL, Clohisy JC, Trousdale RT, Sierra RJ. Natural History of the Dysplastic Hip Following Modern Periacetabular Osteotomy. *J Bone Joint Surg Am* 2019; **101**: 932-938 [PMID: 31094985 DOI: 10.2106/JBJS.18.00983]
- 78 **Zhen Y**, Yin C, Tan S, Yuan Q, Zhu L, Wang X. Retrospective analysis of the radiographic indicators for peri-acetabular osteotomy of developmental dysplasia in children. *Int Orthop* 2016; **40**: 341-345 [PMID: 26614106 DOI: 10.1007/s00264-015-3038-6]
- 79 **Chiari K**. Medial displacement osteotomy of the pelvis. *Clin Orthop Relat Res* 1974; 55-71 [PMID: 4817245 DOI: 10.1097/00003086-197401000-00008]
- 80 **Migaud H**, Chantelot C, Giraud F, Fontaine C, Duquennoy A. Long-term survivorship of hip shelf arthroplasty and Chiari osteotomy in adults. *Clin Orthop Relat Res* 2004; 81-86 [PMID: 15043097 DOI: 10.1097/00003086-200401000-00014]
- 81 **Dammerer D**, Braito M, Ferlic P, Kaufmann G, Kosiol J, Biedermann R. Long-term clinical and radiological outcome in patients with severe Legg-Calvé-Perthes disease after Chiari pelvic osteotomy: a mean of 14 years follow-up. *Hip Int* 2022; **32**: 807-812 [PMID: 33566697 DOI: 10.1177/1120700020988150]
- 82 **Ito H**, Matsuno T, Minami A. Chiari pelvic osteotomy for advanced osteoarthritis in patients with hip dysplasia. *J Bone Joint Surg Am* 2005; **87** Suppl 1: 213-225 [PMID: 16140795 DOI: 10.2106/JBJS.E.00204]
- 83 **Siebenrock KA**, Anwander H, Zurmühle CA, Tannast M, Slongo T, Steppacher SD. Head reduction osteotomy with additional containment surgery improves sphericity and containment and reduces pain in Legg-Calvé-Perthes disease. *Clin Orthop Relat Res* 2015; **473**: 1274-1283 [PMID: 25384430 DOI: 10.1007/s11999-014-4048-1]
- 84 **Yoo WJ**, Moon HJ, Cho TJ, Choi IH. Does shelf acetabuloplasty influence acetabular growth and remodeling? *Clin Orthop Relat Res* 2012; **470**: 2411-2420 [PMID: 22048866 DOI: 10.1007/s11999-011-2163-9]
- 85 **Benad K**, Martinot P, Dartus J, Girard J, Putman S, Migaud H. Influence of shelf acetabuloplasty on the outcomes of total hip arthroplasty in hips with dysplasia: a case-control study. *Int Orthop* 2022; **46**: 989-997 [PMID: 35113185 DOI: 10.1007/s00264-022-05322-3]
- 86 **Grzegorzewski A**, Synder M, Kmiec K, Krajewski K, Polguj M, Sibiński M. Shelf acetabuloplasty in the treatment of severe Legg-Calvé-Perthes disease: good outcomes at midterm follow-up. *Biomed Res Int* 2013; **2013**: 859483 [PMID: 24377097 DOI: 10.1155/2013/859483]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

