

World Journal of *Clinical Cases*

World J Clin Cases 2023 May 6; 11(13): 2855-3113



Contents

Thrice Monthly Volume 11 Number 13 May 6, 2023

OPINION REVIEW

- 2855 Long-term implications of fetal growth restriction
D'Agostin M, Di Sipio Morgia C, Vento G, Nobile S

REVIEW

- 2864 Appraisal of gastric stump carcinoma and current state of affairs
Shukla A, Kalayarasan R, Gnanasekaran S, Pottakkat B
- 2874 Burden of severe infections due to carbapenem-resistant pathogens in intensive care unit
Pace MC, Corrente A, Passavanti MB, Sansone P, Petrou S, Leone S, Fiore M

MINIREVIEWS

- 2890 Individualized diabetes care: Lessons from the real-world experience
Khor XY, Pappachan JM, Jeeyavudeen MS
- 2903 Clinical management of dural defects: A review
Dong RP, Zhang Q, Yang LL, Cheng XL, Zhao JW
- 2916 Potential impact of music interventions in managing diabetic conditions
Eseadi C, Amedu AN
- 2925 Implications of obesity and adiposopathy on respiratory infections; focus on emerging challenges
Lempesis IG, Georgakopoulou VE

ORIGINAL ARTICLE

Case Control Study

- 2934 Association of C-reactive protein and complement factor H gene polymorphisms with risk of lupus nephritis in Chinese population
Li QY, Lv JM, Liu XL, Li HY, Yu F

Retrospective Study

- 2945 Comparison of the application value of transvaginal ultrasound and transabdominal ultrasound in the diagnosis of ectopic pregnancy
Hu HJ, Sun J, Feng R, Yu L

Observational Study

- 2956 Assessment of knowledge, cultural beliefs, and behavior regarding medication safety among residents in Harbin, China
Liu XT, Wang N, Zhu LQ, Wu YB

SYSTEMATIC REVIEWS

- 2966** Palliative oral care in terminal cancer patients: Integrated review
Silva ARP, Bodanezi AV, Chrun ES, Lisboa ML, de Camargo AR, Munhoz EA

META-ANALYSIS

- 2981** Effect of preoperative inspiratory muscle training on postoperative outcomes in patients undergoing cardiac surgery: A systematic review and meta-analysis
Wang J, Wang YQ, Shi J, Yu PM, Guo YQ
- 2992** Efficacy and safety of intravenous tranexamic acid in total shoulder arthroplasty: A meta-analysis
Deng HM

CASE REPORT

- 3002** Awake laparoscopic cholecystectomy: A case report and review of literature
Mazzone C, Sofia M, Sarv  I, Litrico G, Di Stefano AML, La Greca G, Latteri S
- 3010** Bilateral malignant glaucoma with bullous keratopathy: A case report
Ma YB, Dang YL
- 3017** Finger compartment syndrome due to a high-pressure washer injury: A case report
Choi JH, Choi SY, Hwang JH, Kim KS, Lee SY
- 3022** Primary dedifferentiated chondrosarcoma of the lung with a 4-year history of breast cancer: A case report
Wen H, Gong FJ, Xi JM
- 3029** Importance of proper ventilator support and pulmonary rehabilitation in obese patients with heart failure: Two case reports
Lim EH, Park SH, Won YH
- 3038** Multiple flexor tendon ruptures due to osteochondroma of the hamate: A case report
Kwon TY, Lee YK
- 3045** Fractional flow reserve measured *via* left internal mammary artery after coronary artery bypass grafting: Two case reports
Zhang LY, Gan YR, Wang YZ, Xie DX, Kou ZK, Kou XQ, Zhang YL, Li B, Mao R, Liang TX, Xie J, Jin JJ, Yang JM
- 3052** Uterine artery embolization combined with percutaneous microwave ablation for the treatment of prolapsed uterine submucosal leiomyoma: A case report
Zhang HL, Yu SY, Cao CW, Zhu JE, Li JX, Sun LP, Xu HX
- 3062** Metachronous urothelial carcinoma in the renal pelvis, bladder, and urethra: A case report
Zhang JQ, Duan Y, Wang K, Zhang XL, Jiang KH
- 3070** Unusual phenomenon-“polyp” arising from a diverticulum: A case report
Liew JJJ, Lim WS, Koh FH

- 3076** Idiopathic steno-occlusive disease with bilateral internal carotid artery occlusion: A Case Report
Hamed SA, Yousef HA
- 3086** Solitary acral persistent papular mucinosis nodule: A case report and summary of eight Korean cases
Park YJ, Shin HY, Choi WK, Lee AY, Lee SH, Hong JS
- 3092** Eosinophilic fasciitis difficult to differentiate from scleroderma: A case report
Lan TY, Wang ZH, Kong WP, Wang JP, Zhang N, Jin DE, Luo J, Tao QW, Yan ZR
- 3099** Misdiagnosis of scalp angiosarcoma: A case report
Yan ZH, li ZL, Chen XW, Lian YW, Liu LX, Duan HY
- 3105** Discrepancy among microsatellite instability detection methodologies in non-colorectal cancer: Report of 3 cases
Şenocak Taşçı E, Yıldız İ, Erdamar S, Özer L

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Jina Yun, MD, PhD, Assistant Professor, Division of Hematology-Oncology, Department of Internal Medicine, Soonchunhyang University Bucheon Hospital, Soonchunhyang University School of Medicine, Bucheon 14584, South Korea. 19983233@schmc.ac.kr

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

May 6, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Burden of severe infections due to carbapenem-resistant pathogens in intensive care unit

Maria Caterina Pace, Antonio Corrente, Maria Beatrice Passavanti, Pasquale Sansone, Stephen Petrou, Sebastiano Leone, Marco Fiore

Specialty type: Critical care medicine

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C
Grade D (Fair): D
Grade E (Poor): 0

P-Reviewer: Hu H, China; Jain N

Received: December 29, 2022

Peer-review started: December 29, 2022

First decision: January 17, 2023

Revised: February 17, 2023

Accepted: April 4, 2023

Article in press: April 4, 2023

Published online: May 6, 2023



Maria Caterina Pace, Antonio Corrente, Maria Beatrice Passavanti, Pasquale Sansone, Marco Fiore, Department of Women, Child and General and Specialized Surgery, University of Campania "Luigi Vanvitelli", Naples 80138, Italy

Stephen Petrou, Department of Emergency Medicine, University of California San Francisco, San Francisco, CA 94143, United States

Sebastiano Leone, Division of Infectious Diseases, "San Giuseppe Moscati" Hospital, Avellino 83100, Italy

Corresponding author: Marco Fiore, MD, Doctor, Department of Women, Child and General and Specialized Surgery, University of Campania "Luigi Vanvitelli", Piazza Luigi Miraglia, 2, Naples 80138, Italy. marco.fiore@unicampania.it

Abstract

Intensive care units (ICU) for various reasons, including the increasing age of admitted patients, comorbidities, and increasingly complex surgical procedures (e.g., transplants), have become "the epicenter" of nosocomial infections, these are characterized by the presence of multidrug-resistant organisms (MDROs) as the cause of infection. Therefore, the perfect match of fragile patients and MDROs, as the cause of infection, makes ICU mortality very high. Furthermore, carbapenems were considered for years as last-resort antibiotics for the treatment of infections caused by MDROs; unfortunately, nowadays carbapenem resistance, mainly among Gram-negative pathogens, is a matter of the highest concern for worldwide public health. This comprehensive review aims to outline the problem from the intensivist's perspective, focusing on the new definition and epidemiology of the most common carbapenem-resistant MDROs (*Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacteriales*) to emphasize the importance of the problem that must be permeating clinicians dealing with these diseases.

Key Words: Antimicrobial resistance; Multidrug-resistant; PDR; Carbapenem-resistance; Multidisciplinary critical care; Intensive care unit

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Intensive care units for various reasons have become "the epicenter" of nosocomial infections due to multidrug-resistant organisms: a perfect combination of critically ill patients and multidrug-resistant organisms, as the cause of infection, makes these patients' mortality very high. This comprehensive review aims to outline the problem from the clinician's perspective, focusing on the new definition and epidemiology of the most common multidrug-resistant organisms that are *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacterales* to emphasize the importance of the problem.

Citation: Pace MC, Corrente A, Passavanti MB, Sansone P, Petrou S, Leone S, Fiore M. Burden of severe infections due to carbapenem-resistant pathogens in intensive care unit. *World J Clin Cases* 2023; 11(13): 2874-2889

URL: <https://www.wjgnet.com/2307-8960/full/v11/i13/2874.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i13.2874>

INTRODUCTION

Carbapenem resistance is such an important public health issue worldwide[1,2] that the 2017 World Health Organization (WHO) global priority list of pathogens ranks carbapenem-resistant *Enterobacteriaceae* (CRE), carbapenem-resistant *Pseudomonas aeruginosa* (CRPA), and carbapenem-resistant *Acinetobacter baumannii* (CRAB) in the highest priority category (*i.e.*, Critical)[3]. Infections sustained by these bacteria lead to longer lengths of stay, increased healthcare costs, and higher mortality[4-6], especially in patients admitted to the intensive care unit (ICU)[7]. Many studies demonstrated the link between carbapenem use and carbapenem resistance[8-10]. This has even greater clinical relevance when we consider that the rise in the consumption rate of carbapenems was 45% worldwide[11]. Carbapenems are the third most widely used class of antibiotics worldwide for community-acquired infections in ICU (10.7%) and the first class for hospital-acquired infections (HAI) (21.5%)[12]. This comprehensive review aims to analyze from the perspective of worldwide epidemiology the global burden of severe infections supported by carbapenems-resistant germs in the ICU setting.

LITERATURE SEARCH

To review the published clinical data on the epidemiology of carbapenem resistance in the ICU setting, a systematic search of the biomedical literature was conducted. Medline (*via* PubMed) was searched, limited from 2012 to 2022, for articles using the following terms: [(carbapenem or imipenem or meropenem or doripenem or ertapenem) and (resistance or resistant or susceptible or susceptibility)] or (carbapenemase). The result of this search was combined with three separate searches for "*Pseudomonas aeruginosa*", "*Acinetobacter baumannii*" and "*Enterobacterales* or *Enterobacteriaceae*". The retrieved studies were scheduled from the geographical area of origin in the five continents: "Africa", "America", "Asia", "Europe", "and Australia".

DEFINITIONS

Carbapenem-resistant Gram-negative bacteria (GNBs), namely, CRE (*e.g.*, *Klebsiella pneumoniae*, *Escherichia coli*), CRAB and CRPA, are a matter of national and international concern as they are an emerging cause of HAI that pose a significant threat to public health. The term 'CROS' is used as a generic term that refers to all of these GNBs[13]. Centers for disease control and prevention (CDC) define CRE as multidrug-resistant organisms that are resistant to at least one of the carbapenem antibiotics (ertapenem, meropenem, doripenem, or imipenem) or produce a carbapenemase. CRE is a phenotypic definition (*i.e.*, based on the organism susceptibility pattern). A lot of different mechanisms (*i.e.*, genotypes) can result in carbapenem resistance, for example, the production of enzymes that break down carbapenems and related antimicrobials making them ineffective: CRE that produce carbapenemases are called carbapenemase-producing CRE (CP-CRE); therefore, CP-CRE are a subset of all CRE (approximately 30% of CRE carry a carbapenemase), carbapenemase genes are often on mobile genetic elements, which can be easily shared between bacteria, leading to the rapid spread of resistance. Carbapenemases are classified by ambler into three classes - A, B and D (class C includes enzymes that hydrolyze primarily cephalosporins[14]) based on their central catalytic domain and substrate preference[15]. Class A [*e.g.*, *Klebsiella pneumoniae* carbapenemase (KPC), imipenem-hydrolyzing β -lactamase and *Serratia marcescens* enzyme] and D [oxacillin carbapenemase/oxacillinase (OXA)] carbapenemases have serine residues in their active sites and hence are called serine-proteases, while

Class B [New Delhi metallo- β -lactamase (NDM), Verona integron-encoded metallo- β -lactamase (VIM) and imipenemase metallo- β -lactamase (IMP)] enzymes are metallo- β -lactamases with zinc in the active site[16]. The five carbapenemases most frequently identified in CRE are KPC, which was the first carbapenemase identified in the United States (US) in 2001, the NDM, VIM, oxacillinase-48 (OXA-48-type), and IMP[17]. The European committee on antimicrobial susceptibility testing defined the meropenem breakpoints for *Escherichia coli* and *Klebsiella pneumoniae* as $S \leq 2$ mg/L and $R > 8$ mg/L; the corresponding breakpoints for ertapenem are $S \leq 0.5$ mg/L and $R > 0.5$ mg/L. Isolates with meropenem minimum inhibitory concentration (MIC) > 2 mg/L and/or ertapenem MIC > 0.5 mg/L are considered resistant and should be investigated for carbapenem resistance mechanisms. This approach will not identify all *Escherichia coli* and *klebsiella pneumoniae* isolates but will detect most isolates with clinically significant carbapenem non-susceptibility. As the CDC also the European CDC encourages proceeding with the detection of carbapenemase production in carbapenem non-susceptible isolates with MIC values above the susceptible breakpoint[18].

EPIDEMIOLOGY

To monitor antibiotic resistance and plan contrast strategies, the different continents established epidemiological surveillance networks: European antimicrobial resistance surveillance network and central Asian and eastern European surveillance of antimicrobial resistance in Europe and Asia while the national healthcare safety network at the CDC in the US. They documented that multidrug-resistant organisms (MDROs) have become much more prevalent during the last decade[19-21]. CDC estimates that each year in the US, at least 2.8 million people get an antibiotic-resistant infection, and more than 35000 people die. The estimated national cost to treat infections caused by six MDROs identified in the last CDC report and frequently found in healthcare can be substantial—more than \$4.6 billion annually [22]. In a report conducted for “the review on antimicrobial resistance (AMR)”, commissioned in July 2014 by the United Kingdom prime minister, it is predicted that the toll of global antimicrobial resistance will be 10 million deaths per year and up to \$100 trillion lost to the global economy by 2050 [23]. In a survey promoted by the European society of intensive care medicine, 12.4% of ICU physicians reported that they had, during the preceding six months, at least one patient with an infection caused by a bacterium resistant to all or almost all antibiotics available in their ICU[24]. An international multicenter study concluded that 19% of patients admitted to the ICU for more than 24 hours acquired an infection, with rates ranging between 2.3% and 49.2% depending on the hospital unit[25]. The most common ICU-acquired infections are pneumonia, surgical site infection, gastrointestinal infection, urinary tract infection (UTI) and bloodstream infection (BSI)[26]. In a large surveillance report from 183 US hospitals, 84% of BSI were related to the use of a central line catheter, 39% of pneumonia cases were ventilator-associated pneumonia and 68% of UTIs were related to urinary catheters[27]. According to the Gram staining results, bacteria can be classified into 2 categories: GNBs and Gram-positive bacteria (GPBs). Infections caused by multidrug-resistant GNBs are more frequent than multidrug-resistant GPBs, compared to the past. In a large prevalence study on infected ICU patients with isolates from 75 countries, 62% were GNBs, 47% were GPBs and 19% were fungal[28]. Many acronyms help clinicians remember the most prevalent germs: ESKAPE organisms identify a group of highly resistant germs that 'escape' to β -lactam antibiotics and consist of *Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter spp.*, *Pseudomonas aeruginosa*, and *Enterobacter spp.*[29,30]. ESKAPE organisms represent the 6 most common MDROs of HAI[31]. However, since it was pointed out that this acronym excluded other enteric GNBs including *Escherichia coli*, it was modified into ESKAPE+C where “c” refers to *Clostridium difficile*, an important nosocomial pathogen that may easily acquire an MDROs phenotype and “e” refers *Enterobacteriaceae* covering all enteric GNBs including *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus spp.* and *Enterobacter spp.*[32]. In Europe and other areas, of particular concern is the rapid spread of resistance mediated by extended-spectrum β -lactamases (ESBLs), especially in *Klebsiella pneumoniae*. ESBLs organisms are usually resistant to multiple antimicrobials, including third-generation and fourth-generation cephalosporins and aztreonam[33]. Sader and colleagues in their large cross-national research study reported that among *Escherichia coli* isolates from the ICUs, 13.7% were ESBLs producers while ESBLs-*klebsiella spp.* were 17.2%[34]. Another antibiotic class that over time increased the resistance of *Escherichia coli* is that of fluoroquinolones, usually considered active in this species[35, 36]. Resistance of *Pseudomonas aeruginosa* to fluoroquinolones and imipenem has increased rapidly; above 10% of *Pseudomonas aeruginosa* are now resistant to multiple antibiotics classes such as cephalosporins, carbapenems, aminoglycosides and fluoroquinolones[33]. The increased use of carbapenems, which are among the most effective classes of antibiotics active against MDROs contributed to the emergence of CRE or CRAB[37,38]: Up to 25% of *Acinetobacter baumannii* isolates are CRAB[33]. The CRAB prevalence in Europe seems to be higher in south-eastern Europe, with the highest prevalence in Romania (86.5% meropenem 94.6% imipenem resistance)[39]. In the American continent, there seems to be a north-south gradient with all isolated *Acinetobacter baumannii* resistant to carbapenems in Uruguay[40], and practically absent in Canada[41]. More contained data come from Asia with China which seems to have the greatest number of CRAB. As for the African continent, there

are few studies on the prevalence of carbapenem resistance[42,43]; in a study conducted in Uganda, the prevalence of CRAB is 81.25% [44]. Table 1 and Figure 1 report the worldwide prevalence of meropenem-resistant *Acinetobacter baumannii*; we decided to use meropenem as a benchmark to determine the occurrence of carbapenem resistance, to make tables and figures easier to read because *in vitro* studies involving isolates from ICU patients indicate that meropenem is more active against most GNBs than other comparators (including imipenem)[45]. More contained data concern the CRPA: In Europe, the data are more varied with very variable resistance, also between homogeneous nations in terms of geography, economy, and social progress; for example, in the Netherlands, the prevalence is 8.3%-17% [46] while in Germany it is 66.7% [47]. In North America the prevalence does not seem to exceed the two-fifths of the isolates, on the contrary, in a study conducted in Costa Rica, these exceeded four-fifths [48]. In Asia, the highest prevalence is in Korea with 92.9% of the BSI isolated from a burn ICU [49]. In Africa, the prevalence varies from about half of the isolates to almost all, as in Uganda with 88.8% of the CRPA [44] (Table 2). Figure 2 reports the worldwide prevalence of meropenem-resistant *Pseudomonas aeruginosa*.

CRE account for approximately 20%-70% of *Enterobacteriales* isolated in Europe [50,51], in North America, they remain almost non-existent in Canada [41], with a prevalence similar to the European one in the US [52-55]. In Asia data are very varied with a prevalence in China of 56.6%-76.7% of carbapenem-resistant *Klebsiella pneumoniae* (CRKP) [56,57]. From studies conducted in the African continent, Tunisia seems to be the country with the highest prevalence with a percentage of 85.2% of CRKP [58]. In Table 3 we reported the worldwide prevalence of meropenem-resistant *Enterobacteriales*, and in Figure 3 is shown the worldwide prevalence of CRKP which is the most common CRE.

RISK FACTORS

Many risk factors can contribute to the genesis of antimicrobial resistance. They can be categorized as host, environmental, human, and protective barrier integrity factors [109]. Host risk factors include advanced age, organ and bone marrow transplant, end-stage renal disease in dialysis, intra-abdominal surgical procedures, cancer chemotherapy, immunosuppressive disease or therapy [26,110-112]. Prior use of antibiotics (90 days), prolonged antimicrobial usage and hospitalization (more than 5 days), use of indwelling catheters, long mechanical ventilation and residence in nursing homes and long-term care facilities are other important risk factors [110,112,113]. Numerous drugs used in ICU can be a risk factor predisposing patients to infections such as pneumonia (*e.g.*, sedatives and muscle relaxants because they can reduce the cough and swallow reflexes) or gastrointestinal infections (*e.g.*, proton pump inhibitors for stress ulcer prophylaxis because they disrupt the normal non-pathogenic bacterial flora) [110]. In this category, an important independent risk factor is previous MDROs infection or MDROs colonization. If the latter case occurs the probability of developing an infection is high [113]. Considering that some microorganisms can survive on surfaces, environmental is a category of risk factors, very dangerous for the genesis of antimicrobial resistance. It includes poor cleaning and disinfection of environmental surfaces as well as medical devices used for patient care (*e.g.*, stethoscopes, thermometers, suction apparatus) that so became a source or reservoir to disseminate germs to other patients [114]. Among environmental risk factors, colonization pressure is of great importance. First described by Bonten for vancomycin-resistant *Enterococci* [115], and later for other bacteria as well [116-118], it is a critical parameter in the epidemiology of MDROs defined as the proportion of patients colonized with a microorganism in a given geographic area for a specified period [119]. It can be used to estimate the probability of cross-contamination [118], which is in turn an important indicator of poor hygiene especially when there is a clonal relationship of isolates [120]. In their study, Arvaniti *et al* [121] found that out of the total number of patients admitted to their ICU, 5.7% were already colonized at the hospitalization and of these 15.7% acquired *Acinetobacter spp.* during their ICU stay.

The main physical barriers of our body are the skin and mucosa membranes. They represent the first defensive bulwark against infections in general and therefore also for those supported by MDROs. Damage or interruption of their integrity using invasive devices in the ICU increases the risk of infections. In a recent meta-analysis by Hui Ang and Xuan, it was found that male gender (OR 1.40, 95%CI: 1.09, 1.80), having an operative procedure (OR 1.31, 95%CI: 1.10, 1.56), a central venous catheter (OR 1.22, 95%CI: 1.01, 1.48), mechanical ventilation (OR 1.25, 95%CI: 1.07, 1.46), previous antibiotic therapy (OR 1.66, 95%CI: 1.41, 1.96), length of ICU stay (weighted mean difference 8.18, 95%CI: 0.27, 16.10) were the identified risk factors associated with MDROs infections in ICU [122].

CURRENT AND FUTURE STRATEGIES AGAINST ANTIMICROBIAL RESISTANCE IN ICU

Infection prevention strategies can be divided into vertical or horizontal approaches [123,125]. Both go to integrate themselves into complex and various strategies to prevent MDROs infections. Vertical approaches involve the reduction of the risk of colonization, infection and transmission from high-risk pathogens or a specific group of them (*e.g.*, *Clostridium difficile*, multidrug-resistant GNBs, and others)

Table 1 Worldwide prevalence of Meropenem-resistant *Acinetobacter Baumannii*, n %

Continent	Country	Prevalence	Site of infection	Ref.
Africa	Uganda	81.25	Mix	[44]
America	Brazil	22.8-94.2	Mix	[42,59]
	Canada	4.4	Mix	[41]
	French Guiana	16.2	Mix	[60]
	Mexico	56.6	Mix	[61]
	Uruguay	100	Mix	[40]
	United States	61.2-74.2	Mix	[26]
	China	76.7-91.8	Mix	[42,43]
Asia	India	65.2	VAP	[62]
	Indonesia	16.7-68	Mix	[50,63]
	Iran	53.8-94.5	Mix	[64,65]
	Jordan	88.2	Mix	[66]
	Kazakhstan	44.4	Mix	[67]
	Korea	55.8-91.8	Mix	[52,53]
	Saudi Arabia	6.2-52.6	Mix	[68,69]
	Taiwan	50.7	Mix	[70]
	Thailand	40.5-69	Mix	[71,72]
	Vietnam	84	VAP	[73]
Europe	Germany	43 ¹	Mix	[74,75]
	Greece	58.9	Mix	[76]
	Italy	70	VAP	[77]
	Lithuania	30	VAP	[78]
	Poland	74.9-92.3	Mix	[54,55]
	Romania	86.5	Mix	[39]
	Russia	38-67.5	Mix	[79,80]
	Serbia	82	HAC	[81]
		85.3	VAP	
	Spain	86.05	Mix	[82]
	Switzerland	37	Mix	[51]

¹Authors used carbapenems other than Meropenem or do not specify the carbapenem tested.

VAP: Ventilator-associated pneumonia; HAC: Hospital-acquired condition; Mix: More than one infection site or aggregated data about them.

[124]. For this reason, they are valuable tools in controlling and managing an outbreak[123,124]. Vertical approaches are centered on the use of active surveillance testing to detect patients who are MDROs carriers (*i.e.*, asymptomatic colonizers) and separate them from patients who are not colonized with that specific pathogen. This is because asymptomatic colonizers can spread the microorganism contaminating the environment and devices and favoring transmission through direct and indirect contact [124]. Examples of active surveillance testing are a rectal culture for CRE. Vertical strategies include also contact precaution and targeted decolonization (TD) for specific pathogens. TD has some limitations: the different decolonization strategies reduce the diffusion of a single specific target organism and not all-important organisms, such as multidrug-resistant GNBs and VRE, have options for decolonization[126]. Horizontal infection prevention strategies aim to reduce the risk of infections sustained by a broad spectrum of pathogens[124]. They include standard precautions (such as hand hygiene and use of personal protective equipment) and antimicrobial stewardship (AS). It should be noted that some interventions falling within the vertical approach, such as the use of gloves with or without gowns or the decolonization of the skin, can be applied to all patients (*i.e.*, in a horizontal approach), not just those

Table 2 Worldwide prevalence of Meropenem-resistant *Pseudomonas Aeruginosa*, n %

Continent	Country	Prevalence	Site of infection	Ref.
Africa	Egypt	41.82-78	Mix	[83,84]
	Lebanon	42.9	Mix	[83]
	Libya	46	Mix	[83]
	Tunisia	53.7 ¹	Mix	[83]
	Uganda	88.8	Mix	[83]
America	Brazil	22.9-51.8	Mix	[85,86]
	Canada	18.3	Mix	[41]
	Costa Rica	91.3	Ns	[48]
	United States	12.9-43.3	Mix	[87,88]
Asia	Indonesia	12.4-38.1	Mix	[89,90]
	Indonesia	12.4-38.1	Mix	[89,90]
	Iran	25	BSI	[91]
	Korea	50-92.9	BSI	[49,92]
	Qatar	85.7	Mix	[93]
	Saudi Arabia	52.5	Mix	[83]
	Taiwan	22.5	Mix	[94]
	United Arab Emirates	7.7	Mix	[83]
	Turkey	46.7	Mix	[95]
Europe	Germany	61-66.7	Mix	[47,75]
	Netherlands	8.3-17	Mix	[46]
	Serbia	65.1	HAC	[81]
		70.2	VAP	
	Switzerland	27	Mix	[51]

¹Authors used carbapenems other than Meropenem or do not specify the carbapenem tested.

BSI: Bloodstream infections; VAP: Ventilator-associated pneumonia; HAC: Hospital-acquired condition; NS: Not specified; Mix: More than one infection site or aggregated data about them.

with a specific pathogen. According to the CDC and the WHO, hand hygiene remains the simplest and most important practice in infection control. In May 2009 the WHO drew up a simple and precise infographic (called "The 5 moments of hand hygiene") for hand hygiene or the transition from one patient to the next, to prevent cross-transmission[127]. Despite the evidence showing the effectiveness of hand hygiene in preventing infections and efforts to increase compliance rate, it remains low at between 40% and 60% [128,129]. AS is a set of strategies used to improve the use of antibiotics and limit the onset of resistance. It is centred on a systematic approach in multidisciplinary teams[130,131].

An AS programme should provide for: (1) The systematic search for causal agents by carrying out targeted crop surveys; the use of molecular biology tests can also enable important data to be obtained quickly; (2) Limiting the use of broad-spectrum drugs and reducing the duration of empirical therapy through de-escalation strategies[132], with timely replacement of these drugs with other narrow-spectrum drugs; (3) Base therapies on pharmacokinetic and pharmacodynamic criteria adapted to the conditions of critical patients and any changes in the volume of distribution, metabolism, and elimination of drugs; and (4) Optimization of therapy (*i.e.*, adequate dosage, optimal mode of administration for the shortest possible time).

About AS it is important to note that data suggest that 30% to 60% of antibiotics prescribed in ICU are unnecessary, inappropriate, or suboptimal[133]. One of the possible reasons for this is the widespread belief that once the diagnosis of infection is made it is necessary to immediately start the antibiotic therapy with broad-spectrum drugs as each delay is associated with a worsening of the patient's outcome. This is true in infections with a rapid evolution (*e.g.*, Meningitis) or for patients hemodynamically unstable. However, data suggest that in patients with infection but stable, a limited delay in the start of antibiotic therapy allowing the execution of targeted cultures would allow a more appropriate

Table 3 Worldwide prevalence of Meropenem-resistant *Enterobacteriales*, n %

Continent	Country	Pathogen	Prevalence	Site of infection	Ref.
Africa	Egypt	<i>Enterobacter cloacae</i> ; <i>Escherichia coli</i> ; <i>Klebsiella pneumoniae</i>	43.5; 27.1; 53.7	Mix	[96]
	Morocco	<i>Enterobacteriales</i>	2.6	Mix	[97]
	South Africa	<i>Enterobacter spp.</i> Other; <i>Klebsiella spp.</i>	18; 6; 18	Mix	[98]
	Tunisia	<i>Enterobacter aerogenes</i> ; <i>Enterobacter cloacae</i> ; <i>Escherichia coli</i> ; <i>K. Pneumoniae</i> ; <i>Providencia Stuarthii</i>	0.9; 9.8; 2.9; 85.2; 0.9	Mix	[58]
America	Argentina	<i>Enterobacteriales</i>	2.8	BSI	[99]
	Canada	<i>Enterobacter cloacae</i> ; <i>Escherichia coli</i> ; <i>K. pneumoniae</i> ; <i>S marcescens</i>	0.8; 0.1; 0.2; 0.5	Mix	[41]
	United States	<i>Citrobacter spp.</i> ; <i>Enterobacter aerogenes</i> ; <i>Enterobacter cloacae</i> ; <i>Escherichia coli</i> ; <i>Klebsiella oxytoca</i> ; <i>Klebsiella pneumoniae</i>	4; 6; 42; 14; 4; 30	Mix	[87,88,100,101]
Asia	China	<i>Escherichia coli</i> ; <i>Klebsiella Pneumoniae</i>	11.9; 57-76.7	Mix	[56,57,102]
	India	<i>Klebsiella spp.</i>	54	Mix	[103]
	Iran	<i>Klebsiella Pneumoniae</i>	25.3	Mix	[104]
	Korea	<i>Enterobacteriales</i>	31.1	BSI	[49]
	Pakistan	<i>Klebsiella Pneumoniae</i>	72	Mix	[97]
	Turkey	<i>Klebsiella Pneumoniae</i>	44.7-67.47	Mix	[105]
Europe	France	<i>Enterobacteriales</i>	72.8	Mix	[106]
	Germany	<i>Escherichia coli</i> ; <i>Klebsiella Pneumoniae</i>	3; 13	Mix	[75]
	Greece	<i>Klebsiella pneumoniae</i>	74	NS	[107]
	Russia	<i>Escherichia coli</i> ; <i>Klebsiella spp.</i> ; <i>Proteus spp.</i>	3; 16; 29	NS	[108]
	Serbia	<i>Enterobacter spp.</i>	36.4/35.9	HAC/VAP	[81]
		<i>Klebsiella pneumoniae</i>	50/56.8		
		<i>Proteus mirabilis</i>	40/39.5		
	Spain	<i>Enterobacteriales</i>	30.3	NS	[84]
	Switzerland	<i>Enterobacter spp.</i> ; <i>Escherichia coli</i> ; <i>Klebsiella pneumoniae</i>	77; 8; 11	Mix	[50]

BSI: Bloodstream infections; VAP: Ventilator-associated pneumonia; HAC: Hospital-acquired condition; NS: Not specified; Mix: More than one infection site or aggregated data about them.

treatment and an improvement of the outcome[134]. It seems to be essential to identify protocols for the quickest identification of the germ causing the infection[135], in order not to use combination therapies whose efficacy on MDROs is not always the most effective[136,137]. A paradigmatic case seems to be the use of colistin in combination, which is the most common use in clinical practice[138], but randomized studies have not shown any benefits even in strains resistant to retrospectively identified as colistin-resistant[139]. Environmental cleaning and disinfection are other essential horizontal strategies for the control of infections and especially the prevention of cross-contamination[109]. It is important that in every hospital there is a systematic protocol for environmental cleaning and disinfection. It should address regular daily high-touch areas frequently exposed to human contact and emphasize adequate disinfection of the discharged patient's room as a terminal cleaning practice[140].

Currently, antibiotics are still the first therapeutic weapon for patients with MDROs infection in ICU [141]. Despite government efforts and incentives for pharmacological research of new molecules, few antimicrobial agents remain effective against MDROs that are available in clinical practice. New antimicrobial agents recently approved or in advanced phases of clinical development including the new beta-lactam and beta-lactamase inhibitor combinations (ceftolozane/tazobactam, ceftazidime/avibactam, meropenem/vaborbactam, imipenem/cilastatin/relebactam, aztreonam/avibactam), siderophore cephalosporins (cefiderocol), aminoglycosides (plazomicin) and tetracyclines (eravacycline)[142]. Numerous incentives have been provided to encourage researchers to work on alternative strategies to reverse the resistance trend. There are numerous alternative therapeutic weapons to antimicrobials in the study that could be used in the future[141]. Our microbiota remains an important ally in the battle against MDROs infections. Therefore, it must remain unaltered. Two therapeutic options are currently being investigated to remove the antibiotic residues active in the colonic space where the highest




Figure 1 Worldwide prevalence of Meropenem-resistant *Acinetobacter Baumannii*.




Figure 2 Worldwide prevalence of Meropenem-resistant *Pseudomonas Aeruginosa*.

concentrations of intestinal bacteria are found. The first is the use of an engineered, broad-spectrum beta-lactamase that aims at decaying any beta-lactamase in the gut. The second is colon-delivered active charcoal, which aims to adsorb free colonic compounds[141]. Phage therapy is another therapeutic alternative with an interest in the future. A serious advantage of phages over antibiotics is that is highly specific. For this, they can be a perfect weapon to decontaminate MDROs from the gastrointestinal tract, as only MDROs strains would be targeted while commensal strains would be spared[141]. Like phage another specific future possibility against MDROs infection is antibodies. To overcome the issue of immune reaction against monoclonal antibodies, they are now humanized. Examples of antibodies that are being developed in this context target virulence factors: Alpha-toxin of *Staphylococcus aureus*, the type III secretion system of *Pseudomonas aeruginosa*, and the toxin B of *Clostridium difficile*[141]. In addition, a vaccine against multidrug-resistant *Acinetobacter baumannii* is also under investigation at the preclinical stage[141].

A Specific carbapenem-resistant and carbapenemase-producing Organism Prevention Program for Public Health and Healthcare is recently uploaded by the California Department of Public Health; it is clearly articulated ten different points: (1) Laboratory Identification (implement the updated laboratory breakpoints for carbapenems and *Enterobacteriales*); (2) Surveillance (ensure that the laboratory rapidly notifies infection prevention and clinical staff when a patient with carbapenem resistance is identified);




Figure 3 Worldwide prevalence of Meropenem-resistant *Klebsiella pneumonia*.

(3) Colonization Testing (perform CRE colonization testing upon ICU admission of high-risk patients); (4) Infection Control Measures (place patients infected or colonized with CRE in a single room whenever possible, and implement Standard and Contact precautions); (5) Adherence Monitoring (use infection control assessment and adherence monitoring tools); (6) Environmental Cleaning (Ensure thorough daily and terminal environmental cleaning. Focus on high-touch surfaces or any shared reusable medical equipment); (7) Interfacility Communication (Communicate CRE status to the receiving facility ahead of time to ensure appropriate care is maintained when transferring a patient); (8) AS (Implement strategies to limit the use of broad-spectrum antimicrobial agents and an antimicrobial stewardship program); (9) Regional Prevention (Participate in regional efforts to prevent the spread of drug-resistant infections); and (10) Reporting (Report CPO cases through CalREDIE electronic laboratory reporting[143]).

CONCLUSION

Antimicrobial resistance remains a huge public health problem on a global scale whose weight has a huge cost in terms of health expenditure and human lives. At present, antimicrobial agents remain the only causal therapeutic strategy available. Thanks to the efforts of research, in the future, we could use new therapeutic weapons as alternatives or even superior to antimicrobial agents[141]. At present, it is important to preserve the effectiveness of the last molecules put on the market, through a systematic implementation of strategies to minimize or prevent risk factors (first the pressure selection) and the spread of MDROs. For this purpose, *in primis*, the knowledge of local epidemiology and the creation of antimicrobial programs and diagnostic stewardship are mandatory to ensure the appropriateness of antimicrobial therapies. The WHO Global Action Plan on antimicrobial resistance gives strategic objectives, one of which is to strengthen knowledge through surveillance to cover the gaps in knowledge on the incidence, prevalence, and range of antimicrobial resistance across different geographical regions[144]. In our review, it is evident that there are huge differences in the epidemiology of different nations and that in most of the geographical regions, there are no data. Finally, a multidisciplinary approach including intensivists, microbiologists, pharmacists, and infectious disease specialists should play a key role to optimize antimicrobial treatment and minimizing inappropriate use of antibiotics in an era of limited pharmacological options[142]. To our knowledge, this is the first comprehensive review of the global burden of severe infections due to carbapenem-resistant pathogens focusing on ICU, as well as an evaluation of the limited availability of data. Previous reports focused on the overall antimicrobial resistance aggregating data from different inpatient wards and not exclusively from ICU[145].

FOOTNOTES

Author contributions: Corrente A and Marco F designed the study and performed the research; Pace MC, Passavanti

MB, Sansone P and Leone S supervised the manuscript; Petrou S provided critical reviews and revised the Language; Corrente A and Fiore M wrote the manuscript.

Conflict-of-interest statement: The authors have nothing to disclose.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Italy

ORCID number: Maria Caterina Pace 0000-0002-9352-4780; Antonio Corrente 0000-0002-8490-3148; Maria Beatrice Passavanti 0000-0002-9659-0847; Pasquale Sansone 0000-0003-0873-3586; Stephen Petrou 0000-0001-9627-5444; Sebastiano Leone 0000-0001-7852-4101; Marco Fiore 0000-0001-7263-0229.

S-Editor: Ma YJ

L-Editor: A

P-Editor: Zhao S

REFERENCES

- Hussein K, Raz-Pasteur A, Finkelstein R, Neuberger A, Shachor-Meyouhas Y, Oren I, Kassis I. Impact of carbapenem resistance on the outcome of patients' hospital-acquired bacteraemia caused by *Klebsiella pneumoniae*. *J Hosp Infect* 2013; **83**: 307-313 [PMID: 23313086 DOI: 10.1016/j.jhin.2012.10.012]
- Centers for Disease Control and Prevention (CDC). Vital signs: carbapenem-resistant Enterobacteriaceae. *MMWR Morb Mortal Wkly Rep* 2013; **62**: 165-170 [PMID: 23466435]
- World Health Organization. WHO publishes list of bacteria for which new antibiotics are urgently needed. Available from: <https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-new-antibiotics-are-urgently-needed>
- Bartsch SM, McKinnell JA, Mueller LE, Miller LG, Gohil SK, Huang SS, Lee BY. Potential economic burden of carbapenem-resistant Enterobacteriaceae (CRE) in the United States. *Clin Microbiol Infect* 2017; **23**: 48.e9-48.e16 [PMID: 27642178 DOI: 10.1016/j.cmi.2016.09.003]
- Ben-David D, Kordevani R, Keller N, Tal I, Marzel A, Gal-Mor O, Maor Y, Rahav G. Outcome of carbapenem resistant *Klebsiella pneumoniae* bloodstream infections. *Clin Microbiol Infect* 2012; **18**: 54-60 [PMID: 21722257 DOI: 10.1111/j.1469-0691.2011.03478.x]
- Patel G, Huprikar S, Factor SH, Jenkins SG, Calfee DP. Outcomes of carbapenem-resistant *Klebsiella pneumoniae* infection and the impact of antimicrobial and adjunctive therapies. *Infect Control Hosp Epidemiol* 2008; **29**: 1099-1106 [PMID: 18973455 DOI: 10.1086/592412]
- van Loon K, Voor In 't Holt AF, Vos MC. A Systematic Review and Meta-analyses of the Clinical Epidemiology of Carbapenem-Resistant Enterobacteriaceae. *Antimicrob Agents Chemother* 2018; **62** [PMID: 29038269 DOI: 10.1128/AAC.01730-17]
- Coppy M, Jeanne-Leroyer C, Noize P, Dumartin C, Boyer A, Bertrand X, Dubois V, Rogues AM. Antibiotics associated with acquisition of carbapenem-resistant *Pseudomonas aeruginosa* in ICUs: a multicentre nested case-control study. *J Antimicrob Chemother* 2019; **74**: 503-510 [PMID: 30376042 DOI: 10.1093/jac/dky427]
- Woerther PL, Lepeule R, Burdet C, Decousser JW, Ruppé É, Barbier F. Carbapenems and alternative β -lactams for the treatment of infections due to extended-spectrum β -lactamase-producing Enterobacteriaceae: What impact on intestinal colonisation resistance? *Int J Antimicrob Agents* 2018; **52**: 762-770 [PMID: 30176355 DOI: 10.1016/j.ijantimicag.2018.08.026]
- Raman G, Avendano EE, Chan J, Merchant S, Puzniak L. Risk factors for hospitalized patients with resistant or multidrug-resistant *Pseudomonas aeruginosa* infections: a systematic review and meta-analysis. *Antimicrob Resist Infect Control* 2018; **7**: 79 [PMID: 29997889 DOI: 10.1186/s13756-018-0370-9]
- Van Boeckel TP, Gandra S, Ashok A, Caudron Q, Grenfell BT, Levin SA, Laxminarayan R. Global antibiotic consumption 2000 to 2010: an analysis of national pharmaceutical sales data. *Lancet Infect Dis* 2014; **14**: 742-750 [PMID: 25022435 DOI: 10.1016/S1473-3099(14)70780-7]
- Versporten A, Zarb P, Caniaux I, Gros MF, Drapier N, Miller M, Jarlier V, Nathwani D, Goossens H. Global-PPS network. Antimicrobial consumption and resistance in adult hospital inpatients in 53 countries: results of an internet-based global point prevalence survey. *Lancet Glob Health* 2018; **6**: e619-e629 [PMID: 29681513 DOI: 10.1016/s2214-109x(18)30186-4]
- World Health Organization. Implementation manual to prevent and control the spread of carbapenem-resistant organisms at the national and health care facility level. Geneva, 201. Accessed September 30, 2022. Available from: <https://www.who.int/publications/i/item/WHO-UHC-SDS-2019-6>
- Bush K, Jacoby GA. Updated functional classification of beta-lactamases. *Antimicrob Agents Chemother* 2010; **54**: 969-976 [PMID: 19995920 DOI: 10.1128/AAC.01009-09]
- Ambler RP. The structure of beta-lactamases. *Philos Trans R Soc Lond B Biol Sci* 1980; **289**: 321-331 [PMID: 6109327 DOI: 10.1098/rstb.1980.0049]

- 16 Nordmann P, Poirel L. Epidemiology and Diagnostics of Carbapenem Resistance in Gram-negative Bacteria. *Clin Infect Dis* 2019; **69**: S521-S528 [PMID: 31724045 DOI: 10.1093/cid/ciz824]
- 17 CRE Technical Information. Healthcare-Associated Infections (HAIs). Centres for Disease Control and Prevention (CDC). Accessed September 30, 2022. Available from: <https://www.cdc.gov/hai/organisms/cre/technical-info.html>
- 18 European Centre for Disease Prevention and Control. Expert consensus protocol on carbapenem resistance detection and characterisation for the survey of carbapenem- and/or colistin-resistant Enterobacteriaceae – Version 3.0. Stockholm: ECDC; 2019. Accessed September 30, 2022. Available from: <https://www.ecdc.europa.eu/sites/default/files/documents/expert-consensus-protocol-carbapenem-resistance.pdf>
- 19 European Centre for Disease Prevention and Control. Antimicrobial resistance in the EU/EEA (EARS-Net) - Annual Epidemiological Report 2019. Accessed September 30, 2022. Available from: <https://www.ecdc.europa.eu/en/publications-data/surveillance-antimicrobial-resistance-europe-2019>
- 20 World Health Organization. Regional Office for E. Central Asian and European surveillance of antimicrobial resistance: annual report 2020. Accessed September 30, 2022. Available from: <https://apps.who.int/iris/handle/10665/345873>
- 21 Weiner-Lastinger LM, Abner S, Edwards JR, Kallen AJ, Karlsson M, Magill SS, Pollock D, See I, Soe MM, Walters MS, Dudeck MA. Antimicrobial-resistant pathogens associated with adult healthcare-associated infections: Summary of data reported to the National Healthcare Safety Network, 2015-2017. *Infect Control Hosp Epidemiol* 2020; **41**: 1-18 [PMID: 31767041 DOI: 10.1017/ice.2019.296]
- 22 Centers for Disease Control and Prevention. About Antimicrobial Resistance. Accessed September 30, 2022. Available from: <https://www.cdc.gov/drugresistance/about.html>
- 23 Review on Antimicrobial Resistance. Tackling drug-resistant infections globally. Accessed September 30, 2022. Available from: <https://amr-review.org/>
- 24 Lepape A, Jean A, De Waele J, Friggeri A, Savey A, Vanhems P, Gustin MP, Monnet DL, Garnacho-Montero J, Kohlenberg A. European intensive care physicians' experience of infections due to antibiotic-resistant bacteria. *Antimicrob Resist Infect Control* 2020; **9**: 1 [PMID: 31908772 DOI: 10.1186/s13756-019-0662-8]
- 25 Alberti C, Brun-Buisson C, Burchardi H, Martin C, Goodman S, Artigas A, Sicignano A, Palazzo M, Moreno R, Boulmé R, Lepage E, Le Gall R. Epidemiology of sepsis and infection in ICU patients from an international multicentre cohort study. *Intensive Care Med* 2002; **28**: 108-121 [PMID: 11907653 DOI: 10.1007/s00134-001-1143-z]
- 26 MacVane SH. Antimicrobial Resistance in the Intensive Care Unit: A Focus on Gram-Negative Bacterial Infections. *J Intensive Care Med* 2017; **32**: 25-37 [PMID: 26772199 DOI: 10.1177/0885066615619895]
- 27 Magill SS, Edwards JR, Bamberg W, Beldavs ZG, Dumyati G, Kainer MA, Lynfield R, Maloney M, McAllister-Hollod L, Nadle J, Ray SM, Thompson DL, Wilson LE, Fridkin SK; Emerging Infections Program Healthcare-Associated Infections and Antimicrobial Use Prevalence Survey Team. Multistate point-prevalence survey of health care-associated infections. *N Engl J Med* 2014; **370**: 1198-1208 [PMID: 24670166 DOI: 10.1056/NEJMoa1306801]
- 28 Vincent JL, Rello J, Marshall J, Silva E, Anzueto A, Martin CD, Moreno R, Lipman J, Gomersall C, Sakr Y, Reinhart K; EPIC II Group of Investigators. International study of the prevalence and outcomes of infection in intensive care units. *JAMA* 2009; **302**: 2323-2329 [PMID: 19952319 DOI: 10.1001/jama.2009.1754]
- 29 Guervil DJ, Chau T. Trends in multidrug-resistant gram-negative bacilli and the role of prolonged β -lactam infusion in the intensive care unit. *Crit Care Nurs Q* 2013; **36**: 345-355 [PMID: 24002425 DOI: 10.1097/CNQ.0b013e3182a10d2f]
- 30 Iovene MR, Pota V, Galdiero M, Corvino G, Di Lella FM, Stelitano D, Passavanti MB, Pace MC, Alfieri A, Di Franco S, Aurilio C, Sansone P, Niyas VKM, Fiore M. First Italian outbreak of VIM-producing *Serratia marcescens* in an adult polyvalent intensive care unit, August-October 2018: A case report and literature review. *World J Clin Cases* 2019; **7**: 3535-3548 [PMID: 31750335 DOI: 10.12998/wjcc.v7.i21.3535]
- 31 Rice LB. Federal funding for the study of antimicrobial resistance in nosocomial pathogens: no ESCAPE. *J Infect Dis* 2008; **197**: 1079-1081 [PMID: 18419525 DOI: 10.1086/533452]
- 32 Peterson LR. Bad bugs, no drugs: no ESCAPE revisited. *Clin Infect Dis* 2009; **49**: 992-993 [PMID: 19694542 DOI: 10.1086/605539]
- 33 Fish DN, Ohlinger MJ. Antimicrobial resistance: factors and outcomes. *Crit Care Clin* 2006; **22**: 291-311, vii [PMID: 16678001 DOI: 10.1016/j.ccc.2006.02.006]
- 34 Sader HS, Farrell DJ, Flamm RK, Jones RN. Antimicrobial susceptibility of Gram-negative organisms isolated from patients hospitalised with pneumonia in US and European hospitals: results from the SENTRY Antimicrobial Surveillance Program, 2009-2012. *Int J Antimicrob Agents* 2014; **43**: 328-334 [PMID: 24630306 DOI: 10.1016/j.ijantimicag.2014.01.007]
- 35 Robicsek A, Jacoby GA, Hooper DC. The worldwide emergence of plasmid-mediated quinolone resistance. *Lancet Infect Dis* 2006; **6**: 629-640 [PMID: 17008172 DOI: 10.1016/S1473-3099(06)70599-0]
- 36 Jacoby GA. Mechanisms of resistance to quinolones. *Clin Infect Dis* 2005; **41** Suppl 2: S120-S126 [PMID: 15942878 DOI: 10.1086/428052]
- 37 Fiore M, Di Franco S, Alfieri A, Passavanti MB, Pace MC, Petrou S, Martora F, Leone S. Spontaneous bacterial peritonitis due to carbapenemase-producing Enterobacteriaceae: Etiology and antibiotic treatment. *World J Hepatol* 2020; **12**: 1136-1147 [PMID: 33442443 DOI: 10.4254/wjh.v12.i12.1136]
- 38 Baditoui L, Axente C, Lungeanu D, Muntean D, Horhat F, Moldovan R, Hogeia E, Bedreag O, Sandesc D, Licker M. Intensive care antibiotic consumption and resistance patterns: a cross-correlation analysis. *Ann Clin Microbiol Antimicrob* 2017; **16**: 71 [PMID: 29132352 DOI: 10.1186/s12941-017-0251-8]
- 39 Lăzureanu V, Poroşnicu M, Gândac C, Moişil T, Bădiţoiu L, Laza R, Musta V, Crişan A, Marinescu AR. Infection with *Acinetobacter baumannii* in an intensive care unit in the Western part of Romania. *BMC Infect Dis* 2016; **16** Suppl 1: 95 [PMID: 27169538 DOI: 10.1186/s12879-016-1399-0]
- 40 Bado I, Papa-Ezdra R, Delgado-Blas JF, Gaudio M, Gutiérrez C, Cordeiro NF, García-Fulgueiras V, Araújo Pirez L, Seija V, Medina JC, Rieppi G, Gonzalez-Zorn B, Vignoli R. Molecular Characterization of Carbapenem-Resistant *Acinetobacter*

- baumannii in the Intensive Care Unit of Uruguay's University Hospital Identifies the First *rmtC* Gene in the Species. *Microb Drug Resist* 2018; **24**: 1012-1019 [PMID: 29920143 DOI: 10.1089/mdr.2017.0300]
- 41 **Denisuik AJ**, Garbutt LA, Golden AR, Adam HJ, Baxter M, Nichol KA, Lagacé-Wiens P, Walkty AJ, Karlowsky JA, Hoban DJ, Mulvey MR, Zhanel GG. Antimicrobial-resistant pathogens in Canadian ICUs: results of the CANWARD 2007 to 2016 study. *J Antimicrob Chemother* 2019; **74**: 645-653 [PMID: 30500898 DOI: 10.1093/jac/dky477]
 - 42 **Fiorentin Vandresen D**, Carolina Lucio L, Shigueyasu Yamada R, Paula Vieira A, Ani Caovilla Follador F, Pitt Benedetti V, Pereira Bento Casaril K, Viviane Buzanello Martins C, Welter Wendt G, Elize Defante Ferreto L. Associated factors of *Acinetobacter baumannii* complex in hospitalized patients: A case-control study. *J Infect Dev Ctries* 2021; **15**: 73-80 [PMID: 33571148 DOI: 10.3855/jidc.13525]
 - 43 **Chen Y**, Yang Y, Liu L, Qiu G, Han X, Tian S, Zhao J, Chen F, Grundmann H, Li H, Sun J, Han L. High prevalence and clonal dissemination of OXA-72-producing *Acinetobacter baumannii* in a Chinese hospital: a cross sectional study. *BMC Infect Dis* 2018; **18**: 491 [PMID: 30268099 DOI: 10.1186/s12879-018-3359-3]
 - 44 **Aruhukumama D**, Najjuka CF, Kajumbula H, Okee M, Mboowa G, Sserwadda I, Mayanja R, Joloba ML, Kateete DP. bla(VIM)- and bla(OXA)-mediated carbapenem resistance among *Acinetobacter baumannii* and *Pseudomonas aeruginosa* isolates from the Mulago hospital intensive care unit in Kampala, Uganda. *BMC Infect Dis* 2019; **19**: 853 [PMID: 31619192 DOI: 10.1186/s12879-019-4510-5]
 - 45 **Hurst M**, Lamb HM. Meropenem: a review of its use in patients in intensive care. *Drugs* 2000; **59**: 653-680 [PMID: 10776838 DOI: 10.2165/00003495-200059030-00016]
 - 46 **Croughs PD**, Klaassen CHW, van Rosmalen J, Maghddid DM, Boers SA, Hays JP, Goessens WHF; Dutch Antibiotic Resistance Surveillance Group. Unexpected mechanisms of resistance in Dutch *Pseudomonas aeruginosa* isolates collected during 14 years of surveillance. *Int J Antimicrob Agents* 2018; **52**: 407-410 [PMID: 29772393 DOI: 10.1016/j.ijantimicag.2018.05.009]
 - 47 **Schäfer E**, Malecki M, Tellez-Castillo CJ, Pfennigwerth N, Marlinghaus L, Higgins PG, Mattner F, Wendel AF. Molecular surveillance of carbapenemase-producing *Pseudomonas aeruginosa* at three medical centres in Cologne, Germany. *Antimicrob Resist Infect Control* 2019; **8**: 208 [PMID: 31893042 DOI: 10.1186/s13756-019-0665-5]
 - 48 **Toval F**, Guzmán-Marte A, Madriz V, Somogyi T, Rodríguez C, García F. Predominance of carbapenem-resistant *Pseudomonas aeruginosa* isolates carrying blaIMP and blaVIM metallo-β-lactamases in a major hospital in Costa Rica. *J Med Microbiol* 2015; **64**: 37-43 [PMID: 25355933 DOI: 10.1099/jmm.0.081802-0]
 - 49 **Park JJ**, Seo YB, Choi YK, Kym D, Lee J. Changes in the prevalence of causative pathogens isolated from severe burn patients from 2012 to 2017. *Burns* 2020; **46**: 695-701 [PMID: 31630835 DOI: 10.1016/j.burns.2019.09.008]
 - 50 **Anggraini D**, Kemal RA, Hadi U, Kuntaman K. The susceptibility pattern and distribution of blaOXA-23 genes of clinical isolate *Acinetobacter baumannii* in a tertiary hospital, Indonesia. *J Infect Dev Ctries* 2022; **16**: 821-826 [PMID: 35656953 DOI: 10.3855/jidc.15902]
 - 51 **Barnsteiner S**, Baty F, Albrich WC, Babouee Flury B, Gasser M, Plüss-Suard C, Schlegel M, Kronenberg A, Kohler P; Swiss Centre for Antibiotic Resistance (ANRESIS). Antimicrobial resistance and antibiotic consumption in intensive care units, Switzerland, 2009 to 2018. *Euro Surveill* 2021; **26** [PMID: 34794535 DOI: 10.2807/1560-7917.ES.2021.26.46.2001537]
 - 52 **Liu Q**, Li W, Du X, Zhong T, Tang Y, Feng Y, Tao C, Xie Y. Risk and Prognostic Factors for Multidrug-Resistant *Acinetobacter Baumannii* Complex Bacteremia: A Retrospective Study in a Tertiary Hospital of West China. *PLoS One* 2015; **10**: e0130701 [PMID: 26083415 DOI: 10.1371/journal.pone.0130701]
 - 53 **Chiang CH**, Pan SC, Yang TS, Matsuda K, Kim HB, Choi YH, Hori S, Wang JT, Sheng WH, Chen YC, Chang FY, Chang SC. Healthcare-associated infections in intensive care units in Taiwan, South Korea, and Japan: recent trends based on national surveillance reports. *Antimicrob Resist Infect Control* 2018; **7**: 129 [PMID: 30455867 DOI: 10.1186/s13756-018-0422-1]
 - 54 **Ziólkowski G**, Pawłowska I, Krawczyk L, Wojkowska-Mach J. Antibiotic consumption versus the prevalence of multidrug-resistant *Acinetobacter baumannii* and *Clostridium difficile* infections at an ICU from 2014-2015. *J Infect Public Health* 2018; **11**: 626-630 [PMID: 29548877 DOI: 10.1016/j.jiph.2018.02.003]
 - 55 **Litwin A**, Fedorowicz O, Duszynska W. Characteristics of Microbial Factors of Healthcare-Associated Infections Including Multidrug-Resistant Pathogens and Antibiotic Consumption at the University Intensive Care Unit in Poland in the Years 2011-2018. *Int J Environ Res Public Health* 2020; **17** [PMID: 32977435 DOI: 10.3390/ijerph17196943]
 - 56 **Chang H**, Wei J, Zhou W, Yan X, Cao X, Zuo L, Chen S, Yao K, Huang R, Chen Y, Wu C. Risk factors and mortality for patients with Bloodstream infections of *Klebsiella pneumoniae* during 2014-2018: Clinical impact of carbapenem resistance in a large tertiary hospital of China. *J Infect Public Health* 2020; **13**: 784-790 [PMID: 31843651 DOI: 10.1016/j.jiph.2019.11.014]
 - 57 **Zhang WX**, Chen HY, Chen C, Chen JH, Wan FS, Li LX, Chen M, Zhang J. Resistance Phenotype and Molecular Epidemiology of Carbapenem-Resistant *Klebsiella pneumoniae* Isolates in Shanghai. *Microb Drug Resist* 2021; **27**: 1312-1318 [PMID: 34297609 DOI: 10.1089/mdr.2020.0390]
 - 58 **Ben Helal R**, Dziri R, Chedly M, Klibi N, Barguelli F, El Asli MS, Ben Moussa M. Occurrence and Characterization of Carbapenemase-Producing Enterobacteriaceae in a Tunisian Hospital. *Microb Drug Resist* 2018; **24**: 1361-1367 [PMID: 29596032 DOI: 10.1089/mdr.2018.0013]
 - 59 **Kurihara MNL**, Sales RO, Silva KED, Silva GD, Mansano MCT, Mahmoud FF, Simionatto S. High lethality rate of carbapenem-resistant *Acinetobacter baumannii* in Intensive Care Units of a Brazilian hospital: An epidemiologic surveillance study. *Rev Soc Bras Med Trop* 2022; **55**: e05292021 [PMID: 35522809 DOI: 10.1590/0037-8682-0529-2021]
 - 60 **Mahamat A**, Bertrand X, Moreau B, Hommel D, Couppie P, Simonnet C, Kallel H, Demar M, Djossou F, Nacher M. Clinical epidemiology and resistance mechanisms of carbapenem-resistant *Acinetobacter baumannii*, French Guiana, 2008-2014. *Int J Antimicrob Agents* 2016; **48**: 51-55 [PMID: 27236843 DOI: 10.1016/j.ijantimicag.2016.03.006]
 - 61 **Uc-Cachón AH**, Gracida-Osorno C, Luna-Chi IG, Jiménez-Guillermo JG, Molina-Salinas GM. High Prevalence of Antimicrobial Resistance Among Gram-Negative Isolated Bacilli in Intensive Care Units at a Tertiary Care Hospital in Yucatán Mexico. *Medicina (Kaunas)* 2019; **55** [PMID: 31540314 DOI: 10.3390/medicina55090588]

- 62 **Kumari M**, Verma S, Venkatesh V, Gupta P, Tripathi P, Agarwal A, Siddiqui SS, Arshad Z, Prakash V. Emergence of blaNDM-1 and blaVIM producing Gram-negative bacilli in ventilator-associated pneumonia at AMR Surveillance Regional Reference Laboratory in India. *PLoS One* 2021; **16**: e0256308 [PMID: [34495985](#) DOI: [10.1371/journal.pone.0256308](#)]
- 63 **Saharman YR**, Karuniawati A, Sedono R, Aditjaningsih D, Sudarmono P, Goessens WHF, Klaassen CHW, Verbrugh HA, Severin JA. Endemic carbapenem-nonsusceptible *Acinetobacter baumannii*-calcoaceticus complex in intensive care units of the national referral hospital in Jakarta, Indonesia. *Antimicrob Resist Infect Control* 2018; **7**: 5 [PMID: [29344351](#) DOI: [10.1186/s13756-017-0296-7](#)]
- 64 **Pourajam S**, Kalantari E, Talebzadeh H, Mellali H, Sami R, Soltaninejad F, Amra B, Sajadi M, Alenaseri M, Kalantari F, Solgi H. Secondary Bacterial Infection and Clinical Characteristics in Patients With COVID-19 Admitted to Two Intensive Care Units of an Academic Hospital in Iran During the First Wave of the Pandemic. *Front Cell Infect Microbiol* 2022; **12**: 784130 [PMID: [35281440](#) DOI: [10.3389/fcimb.2022.784130](#)]
- 65 **Ranjbar R**, Farahani A. Study of genetic diversity, biofilm formation, and detection of Carbapenemase, MBL, ESBL, and tetracycline resistance genes in multidrug-resistant *Acinetobacter baumannii* isolated from burn wound infections in Iran. *Antimicrob Resist Infect Control* 2019; **8**: 172 [PMID: [31719975](#) DOI: [10.1186/s13756-019-0612-5](#)]
- 66 **Nazer LH**, Kharabsheh A, Rimawi D, Mubarak S, Hawari F. Characteristics and Outcomes of *Acinetobacter baumannii* Infections in Critically Ill Patients with Cancer: A Matched Case-Control Study. *Microb Drug Resist* 2015; **21**: 556-561 [PMID: [26430944](#) DOI: [10.1089/mdr.2015.0032](#)]
- 67 **Viderman D**, Brotfain E, Khamzina Y, Kapanova G, Zhumadilov A, Poddighe D. Bacterial resistance in the intensive care unit of developing countries: Report from a tertiary hospital in Kazakhstan. *J Glob Antimicrob Resist* 2019; **17**: 35-38 [PMID: [30448518](#) DOI: [10.1016/j.jgar.2018.11.010](#)]
- 68 **Aljindan R**, Bukharie H, Alomar A, Abdalhamid B. Prevalence of digestive tract colonization of carbapenem-resistant *Acinetobacter baumannii* in hospitals in Saudi Arabia. *J Med Microbiol* 2015; **64**: 400-406 [PMID: [25657302](#) DOI: [10.1099/jmm.0.000033](#)]
- 69 **Al-Hamad A**, Pal T, Leskafi H, Abbas H, Hejles H, Alsubikhy F, Darwish D, Ghazawi A, Sonnevend A. Molecular characterization of clinical and environmental carbapenem resistant *Acinetobacter baumannii* isolates in a hospital of the Eastern Region of Saudi Arabia. *J Infect Public Health* 2020; **13**: 632-636 [PMID: [31551188](#) DOI: [10.1016/j.jiph.2019.08.013](#)]
- 70 **Chen YP**, Liang CC, Chang R, Kuo CM, Hung CH, Liao TN, Liao CS. Detection and Colonization of Multidrug Resistant Organisms in a Regional Teaching Hospital of Taiwan. *Int J Environ Res Public Health* 2019; **16** [PMID: [30925662](#) DOI: [10.3390/ijerph16071104](#)]
- 71 **Chusri S**, Chongsuvivatwong V, Rivera JI, Silpapojakul K, Singkhamanan K, McNeil E, Doi Y. Molecular epidemiology and spatiotemporal analysis of hospital-acquired *Acinetobacter baumannii* infection in a tertiary care hospital in southern Thailand. *J Hosp Infect* 2017; **95**: 53-58 [PMID: [27865634](#) DOI: [10.1016/j.jhin.2016.10.003](#)]
- 72 **Kiddee A**, Assawatheptawee K, Na-Udom A, Treebupachatsakul P, Wangteeraprasert A, Walsh TR, Niumsup PR. Risk Factors for Gastrointestinal Colonization and Acquisition of Carbapenem-Resistant Gram-Negative Bacteria among Patients in Intensive Care Units in Thailand. *Antimicrob Agents Chemother* 2018; **62** [PMID: [29891594](#) DOI: [10.1128/AAC.00341-18](#)]
- 73 **Le Minh V**, Thi Khanh Nhu N, Vinh Phat V, Thompson C, Huong Lan NP, Thieu Nga TV, Thanh Tam PT, Tuyen HT, Hoang Nhu TD, Van Hao N, Thi Loan H, Minh Yen L, Parry CM, Trung Nghia HD, Campbell JI, Hien TT, Thwaites L, Thwaites G, Van Vinh Chau N, Baker S. In vitro activity of colistin in antimicrobial combination against carbapenem-resistant *Acinetobacter baumannii* isolated from patients with ventilator-associated pneumonia in Vietnam. *J Med Microbiol* 2015; **64**: 1162-1169 [PMID: [26297024](#) DOI: [10.1099/jmm.0.000137](#)]
- 74 **Wendel AF**, Malecki M, Otchwemah R, Tellez-Castillo CJ, Sakka SG, Mattner F. One-year molecular surveillance of carbapenem-susceptible *A. baumannii* on a German intensive care unit: diversity or clonality. *Antimicrob Resist Infect Control* 2018; **7**: 145 [PMID: [30505434](#) DOI: [10.1186/s13756-018-0436-8](#)]
- 75 **Maechler F**, Peña Diaz LA, Schröder C, Geffers C, Behnke M, Gastmeier P. Prevalence of carbapenem-resistant organisms and other Gram-negative MDRO in German ICUs: first results from the national nosocomial infection surveillance system (KISS). *Infection* 2015; **43**: 163-168 [PMID: [25395161](#) DOI: [10.1007/s15010-014-0701-6](#)]
- 76 **Karampatakis T**, Tsergouli K, Politi L, Diamantopoulou G, Iosifidis E, Antachopoulos C, Karyoti A, Mouloudi E, Tsakris A, Roilides E. Molecular Epidemiology of Endemic Carbapenem-Resistant Gram-Negative Bacteria in an Intensive Care Unit. *Microb Drug Resist* 2019; **25**: 712-716 [PMID: [30589601](#) DOI: [10.1089/mdr.2018.0266](#)]
- 77 **Delle Rose D**, Pezzotti P, Fortunato E, Sordillo P, Gini S, Boros S, Meledandri M, Gallo MT, Prignano G, Caccese R, D'Ambrosio M, Citterio G, Rocco M, Leonardi F, Natoli S, Fontana C, Favaro M, Celeste MG, Franci T, Testore GP, Andreoni M, Sarmati L. Clinical predictors and microbiology of ventilator-associated pneumonia in the intensive care unit: a retrospective analysis in six Italian hospitals. *Eur J Clin Microbiol Infect Dis* 2016; **35**: 1531-1539 [PMID: [27272120](#) DOI: [10.1007/s10096-016-2694-9](#)]
- 78 **Čiginskienė A**, Dambrasienė A, Rello J, Adukauskienė D. Ventilator-Associated Pneumonia due to Drug-Resistant *Acinetobacter baumannii*: Risk Factors and Mortality Relation with Resistance Profiles, and Independent Predictors of In-Hospital Mortality. *Medicina (Kaunas)* 2019; **55** [PMID: [30781896](#) DOI: [10.3390/medicina55020049](#)]
- 79 **Mayanskiy N**, Chebotar I, Alyabieva N, Kryzhanovskaya O, Savinova T, Turenok A, Bocharova Y, Lazareva A, Polikarpova S, Karaseva O. Emergence of the Uncommon Clone ST944/ST78 Carrying bla(OXA-40-like) and bla(CTX-M-like) Genes Among Carbapenem-Nonsusceptible *Acinetobacter baumannii* in Moscow, Russia. *Microb Drug Resist* 2017; **23**: 864-870 [PMID: [28437227](#) DOI: [10.1089/mdr.2016.0302](#)]
- 80 **Ershova K**, Savin I, Kurdyumova N, Wong D, Danilov G, Shifrin M, Alexandrova I, Sokolova E, Fursova N, Zelman V, Ershova O. Implementing an infection control and prevention program decreases the incidence of healthcare-associated infections and antibiotic resistance in a Russian neuro-ICU. *Antimicrob Resist Infect Control* 2018; **7**: 94 [PMID: [30083313](#) DOI: [10.1186/s13756-018-0383-4](#)]
- 81 **Djordjevic ZM**, Folc MM, Jankovic SM. Distribution and antibiotic susceptibility of pathogens isolated from adults with

- hospital-acquired and ventilator-associated pneumonia in intensive care unit. *J Infect Public Health* 2017; **10**: 740-744 [PMID: 28189513 DOI: 10.1016/j.jiph.2016.11.016]
- 82 **Rodríguez-Lucas C**, Rodicio MR, Vázquez X, Escudero D, Quindós B, Alaguero M, Fernández J. Extensively drug-resistant *Acinetobacter baumannii* carrying bla(OXA-23-like) and armA in a hospital after an intervention in the intensive care unit which ended a long-standing endemicity. *Eur J Clin Microbiol Infect Dis* 2021; **40**: 385-389 [PMID: 32808109 DOI: 10.1007/s10096-020-04009-0]
 - 83 **Al-Orphaly M**, Hadi HA, Eltayeb FK, Al-Hail H, Samuel BG, Sultan AA, Skariah S. Epidemiology of Multidrug-Resistant *Pseudomonas aeruginosa* in the Middle East and North Africa Region. *mSphere* 2021; **6** [PMID: 34011686 DOI: 10.1128/mSphere.00202-21]
 - 84 **Elbaradei A**, Sayedahmed MS, El-Sawaf G, Shawky SM. Screening of mcr-1 among Gram-Negative Bacteria from Different Clinical Samples from ICU Patients in Alexandria, Egypt: One-Year Study. *Pol J Microbiol* 2022; **71**: 83-90 [PMID: 35635164 DOI: 10.33073/pjm-2022-011]
 - 85 **de Almeida Silva KCF**, Calomino MA, Deutsch G, de Castilho SR, de Paula GR, Esper LMR, Teixeira LA. Molecular characterization of multidrug-resistant (MDR) *Pseudomonas aeruginosa* isolated in a burn center. *Burns* 2017; **43**: 137-143 [PMID: 27595453 DOI: 10.1016/j.burns.2016.07.002]
 - 86 **Rodrigues YC**, Furlaneto IP, Maciel AHP, Quaresma AJPG, de Matos ECO, Conceição ML, Vieira MCDS, Brabo GLDC, Sarges EDSNF, Lima LNGC, Lima KVB. High prevalence of atypical virulotype and genetically diverse background among *Pseudomonas aeruginosa* isolates from a referral hospital in the Brazilian Amazon. *PLoS One* 2020; **15**: e0238741 [PMID: 32911510 DOI: 10.1371/journal.pone.0238741]
 - 87 **Jarrell AS**, Kruer RM, Berescu LD, Pronovost PJ, Trivedi JB. Factors associated with in-hospital mortality among critically ill surgical patients with multidrug-resistant Gram-negative infections. *J Crit Care* 2018; **43**: 321-326 [PMID: 29239831 DOI: 10.1016/j.jcrc.2017.10.035]
 - 88 **Sader HS**, Mendes RE, Streit JM, Carvalhaes CG, Castanheira M. Antimicrobial susceptibility of Gram-negative bacteria from intensive care unit and non-intensive care unit patients from United States hospitals (2018-2020). *Diagn Microbiol Infect Dis* 2022; **102**: 115557 [PMID: 34673293 DOI: 10.1016/j.diagmicrobio.2021.115557]
 - 89 **Pelegrin AC**, Saharman YR, Griffon A, Palmieri M, Mirande C, Karuniawati A, Sedono R, Aditjaningsih D, Goessens WHF, van Belkum A, Verbrugh HA, Klaassen CHW, Severin JA. High-Risk International Clones of Carbapenem-Nonsusceptible *Pseudomonas aeruginosa* Endemic to Indonesian Intensive Care Units: Impact of a Multifaceted Infection Control Intervention Analyzed at the Genomic Level. *mBio* 2019; **10** [PMID: 31719179 DOI: 10.1128/mBio.02384-19]
 - 90 **Saharman YR**, Pelegrin AC, Karuniawati A, Sedono R, Aditjaningsih D, Goessens WHF, Klaassen CHW, van Belkum A, Mirande C, Verbrugh HA, Severin JA. Epidemiology and characterisation of carbapenem-non-susceptible *Pseudomonas aeruginosa* in a large intensive care unit in Jakarta, Indonesia. *Int J Antimicrob Agents* 2019; **54**: 655-660 [PMID: 31398483 DOI: 10.1016/j.ijantimicag.2019.08.003]
 - 91 **Emami A**, Pirbonyeh N, Keshavarzi A, Bazargani A, Hassanpour S, Javanmardi F. Evaluating the Saliva of Burn ICU Patients for Resistant Infections Harbor Metallo- β -Lactamase Genes. *J Burn Care Res* 2020; **41**: 647-651 [PMID: 31930340 DOI: 10.1093/jbcr/iraa007]
 - 92 **Kang JS**, Moon C, Mun SJ, Lee JE, Lee SO, Lee S, Lee SH. Antimicrobial Susceptibility Trends and Risk Factors for Antimicrobial Resistance in *Pseudomonas aeruginosa* Bacteremia: 12-Year Experience in a Tertiary Hospital in Korea. *J Korean Med Sci* 2021; **36**: e273 [PMID: 34751008 DOI: 10.3346/jkms.2021.36.e273]
 - 93 **Baiou A**, Elbuzidi AA, Bakdach D, Zaout A, Alarbi KM, Bintaher AA, Ali MMB, Elarabi AM, Ali GAM, Daghfal J, Almaslamani MA, Ibrahim ASS, Alkhal A, Omrani AS. Clinical characteristics and risk factors for the isolation of multi-drug-resistant Gram-negative bacteria from critically ill patients with COVID-19. *J Hosp Infect* 2021; **110**: 165-171 [PMID: 33561503 DOI: 10.1016/j.jhin.2021.01.027]
 - 94 **Lin KY**, Lauderdale TL, Wang JT, Chang SC. Carbapenem-resistant *Pseudomonas aeruginosa* in Taiwan: Prevalence, risk factors, and impact on outcome of infections. *J Microbiol Immunol Infect* 2016; **49**: 52-59 [PMID: 24662016 DOI: 10.1016/j.jmii.2014.01.005]
 - 95 **Acar A**, Karaahmetoğlu G, Akalın H, Altay AF. Pooled prevalence and trends of antimicrobial resistance in *Pseudomonas aeruginosa* clinical isolates over the past 10 years in Turkey: A meta-analysis. *J Glob Antimicrob Resist* 2019; **18**: 64-70 [PMID: 30753904 DOI: 10.1016/j.jgar.2019.01.032]
 - 96 **Koth S**, Lyman M, Ismail G, Abd El Fattah M, Girgis SA, Etman A, Hafez S, El-Kholy J, Zaki MES, Rashed HG, Khalil GM, Sayyoub O, Talaat M. Epidemiology of Carbapenem-resistant Enterobacteriaceae in Egyptian intensive care units using National Healthcare-associated Infections Surveillance Data, 2011-2017. *Antimicrob Resist Infect Control* 2020; **9**: 2 [PMID: 31911830 DOI: 10.1186/s13756-019-0639-7]
 - 97 **Chabah M**, Chems M, Zerouali K, Alloula O, Lehlmi M, Habzi A, Benomar S. Healthcare-associated infections due to carbapenemase-producing Enterobacteriaceae: Bacteriological profile and risk factors. *Med Mal Infect* 2016; **46**: 157-162 [PMID: 26897308 DOI: 10.1016/j.medmal.2015.12.015]
 - 98 **Law T**, Chibabhai V, Nana T. Analysis and comparison of cumulative antibiograms for the Charlotte Maxeke Johannesburg Academic Hospital adult intensive care and high-care units, 2013 and 2017. *S Afr Med J* 2019; **110**: 55-64 [PMID: 31865944 DOI: 10.7196/SAMJ.2019.v110i1.13841]
 - 99 **Lipari FG**, Hernández D, Vilaró M, Caeiro JP, Saka HA. [Clinical, epidemiological and microbiological characterization of bacteremia produced by carbapenem-resistant enterobacteria in a university hospital in Córdoba, Argentina]. *Rev Chilena Infectol* 2020; **37**: 362-370 [PMID: 33399656 DOI: 10.4067/S0716-10182020000400362]
 - 100 **Johnson JK**, Robinson GL, Pineles LL, Ajao AO, Zhao L, Albrecht JS, Harris AD, Thom KA, Furuno JP. Carbapenem MICs in *Escherichia coli* and *Klebsiella* Species Producing Extended-Spectrum β -Lactamases in Critical Care Patients from 2001 to 2009. *Antimicrob Agents Chemother* 2017; **61** [PMID: 28167543 DOI: 10.1128/AAC.01718-16]
 - 101 **Tamma PD**, Kazmi A, Bergman Y, Goodman KE, Ekunseitan E, Amoah J, Simner PJ. The Likelihood of Developing a Carbapenem-Resistant Enterobacteriaceae Infection during a Hospital Stay. *Antimicrob Agents Chemother* 2019; **63** [PMID: 31138574 DOI: 10.1128/AAC.00757-19]
 - 102 **Li Y**, Sun QL, Shen Y, Zhang Y, Yang JW, Shu LB, Zhou HW, Wang Y, Wang B, Zhang R, Wang S, Shen Z. Rapid

- Increase in Prevalence of Carbapenem-Resistant Enterobacteriaceae (CRE) and Emergence of Colistin Resistance Gene *mer-1* in CRE in a Hospital in Henan, China. *J Clin Microbiol* 2018; **56** [PMID: 29386265 DOI: 10.1128/JCM.01932-17]
- 103 **Sodhi K**, Mittal V, Arya M, Kumar M, Phillips A, Kajla B. Pattern of colistin resistance in Klebsiella isolates in an Intensive Care Unit of a tertiary care hospital in India. *J Infect Public Health* 2020; **13**: 1018-1021 [PMID: 31818712 DOI: 10.1016/j.jiph.2019.10.013]
- 104 **Darabi N**, Motazakker M, Khalkhali HR, Yousefi S. A multicenter study of β -lactamase-producing Klebsiella pneumoniae isolated from university teaching hospitals of Urmia, Iran. *J Infect Dev Ctries* 2019; **13**: 690-697 [PMID: 32069252 DOI: 10.3855/jidc.9985]
- 105 **Candevir Ulu A**, Güven Gökmen T, Kibar F, Kurtaran B, Önen C, Kuşçu F, İnal AS, Kömür S, Yaman A, Aksu HSZ, Taşova Y. Molecular epidemiology of carbapenem-resistant Klebsiella pneumoniae at a Turkish centre: Is the increase of resistance a threat for Europe? *J Glob Antimicrob Resist* 2017; **11**: 10-16 [PMID: 28743652 DOI: 10.1016/j.jgar.2017.06.012]
- 106 **David B**, Moussiegt A, Dinh A, Bouchand F, Matt M, Senard O, Deconinck L, Espinasse F, Lawrence C, Fortineau N, Saleh-Mghir A, Caballero S, Escut L, Salomon J. Germs of thrones - spontaneous decolonization of Carbapenem-Resistant Enterobacteriaceae (CRE) and Vancomycin-Resistant Enterococci (VRE) in Western Europe: is this myth or reality? *Antimicrob Resist Infect Control* 2018; **7**: 100 [PMID: 30123500 DOI: 10.1186/s13756-018-0390-5]
- 107 **Karampatakis T**, Antachopoulos C, Iosifidis E, Tsakris A, Roilides E. Molecular epidemiology of carbapenem-resistant Klebsiella pneumoniae in Greece. *Future Microbiol* 2016; **11**: 809-823 [PMID: 27206024 DOI: 10.2217/fmb-2016-0042]
- 108 **Partina I**, Kalinogorskaya O, Kojima S, Gostev V, Volkova M, Ageevets V, Lobzin Y, Sidorenko S. Surveillance of antimicrobial susceptibility of Enterobacteriaceae pathogens isolated from intensive care units and surgical units in Russia. *Jpn J Antibiot* 2016; **69**: 41-51 [PMID: 27290829]
- 109 **Riley MM**. The Rising Problem of Multidrug-Resistant Organisms in Intensive Care Units. *Crit Care Nurse* 2019; **39**: 48-55 [PMID: 31371367 DOI: 10.4037/ccn2019773]
- 110 **Brusselsaers N**, Vogelaers D, Blot S. The rising problem of antimicrobial resistance in the intensive care unit. *Ann Intensive Care* 2011; **1**: 47 [PMID: 22112929 DOI: 10.1186/2110-5820-1-47]
- 111 **Umscheid CA**, Agarwal RK, Brennan PJ; Healthcare Infection Control Practices Advisory Committee. Updating the guideline development methodology of the Healthcare Infection Control Practices Advisory Committee (HICPAC). *Am J Infect Control* 2010; **38**: 264-273 [PMID: 20116133 DOI: 10.1016/j.ajic.2009.12.005]
- 112 **Djibré M**, Fedun S, Le Guen P, Vimont S, Hafiani M, Fulgencio JP, Parrot A, Denis M, Fartoukh M. Universal versus targeted additional contact precautions for multidrug-resistant organism carriage for patients admitted to an intensive care unit. *Am J Infect Control* 2017; **45**: 728-734 [PMID: 28285725 DOI: 10.1016/j.ajic.2017.02.001]
- 113 **Martín-Loeches I**, Diaz E, Vallés J. Risks for multidrug-resistant pathogens in the ICU. *Curr Opin Crit Care* 2014; **20**: 516-524 [PMID: 25188366 DOI: 10.1097/MCC.0000000000000124]
- 114 **Kanamori H**, Parobek CM, Juliano JJ, van Duin D, Cairns BA, Weber DJ, Rutala WA. A Prolonged Outbreak of KPC-3-Producing Enterobacter cloacae and Klebsiella pneumoniae Driven by Multiple Mechanisms of Resistance Transmission at a Large Academic Burn Center. *Antimicrob Agents Chemother* 2017; **61** [PMID: 27919898 DOI: 10.1128/AAC.01516-16]
- 115 **Bonten MJ**, Slaughter S, Ambergen AW, Hayden MK, van Voorhis J, Nathan C, Weinstein RA. The role of "colonization pressure" in the spread of vancomycin-resistant enterococci: an important infection control variable. *Arch Intern Med* 1998; **158**: 1127-1132 [PMID: 9605785 DOI: 10.1001/archinte.158.10.1127]
- 116 **Merrill J**, Santoli F, Appéré de Vecchi C, Tran B, De Jonghe B, Outin H. "Colonization pressure" and risk of acquisition of methicillin-resistant Staphylococcus aureus in a medical intensive care unit. *Infect Control Hosp Epidemiol* 2000; **21**: 718-723 [PMID: 11089656 DOI: 10.1086/501721]
- 117 **Harris AD**, Johnson JK, Thom KA, Morgan DJ, McGregor JC, Ajao AO, Moore AC, Comer AC, Furuno JP. Risk factors for development of intestinal colonization with imipenem-resistant Pseudomonas aeruginosa in the intensive care unit setting. *Infect Control Hosp Epidemiol* 2011; **32**: 719-722 [PMID: 21666406 DOI: 10.1086/660763]
- 118 **Ajao AO**, Harris AD, Roghmann MC, Johnson JK, Zhan M, McGregor JC, Furuno JP. Systematic review of measurement and adjustment for colonization pressure in studies of methicillin-resistant Staphylococcus aureus, vancomycin-resistant enterococci, and clostridium difficile acquisition. *Infect Control Hosp Epidemiol* 2011; **32**: 481-489 [PMID: 21515979 DOI: 10.1086/659403]
- 119 **Bonten MJ**, Gaillard CA, Johanson WG Jr, van Tiel FH, Smeets HG, van der Geest S, Stobberingh EE. Colonization in patients receiving and not receiving topical antimicrobial prophylaxis. *Am J Respir Crit Care Med* 1994; **150**: 1332-1340 [PMID: 7952561 DOI: 10.1164/ajrcm.150.5.7952561]
- 120 **Bonten MJ**. Colonization pressure: a critical parameter in the epidemiology of antibiotic-resistant bacteria. *Crit Care* 2012; **16**: 142 [PMID: 22849650 DOI: 10.1186/cc11417]
- 121 **Arvaniti K**, Lathyris D, Ruimy R, Haidich AB, Koulourida V, Nikolaidis P, Matamis D, Miyakis S. The importance of colonization pressure in multiresistant Acinetobacter baumannii acquisition in a Greek intensive care unit. *Crit Care* 2012; **16**: R102 [PMID: 22694969 DOI: 10.1186/cc11383]
- 122 **Ang H**, Sun X. Risk factors for multidrug-resistant Gram-negative bacteria infection in intensive care units: A meta-analysis. *Int J Nurs Pract* 2018; **24**: e12644 [PMID: 29575345 DOI: 10.1111/ijn.12644]
- 123 **Popovich KJ**. Another success story for horizontal infection control strategies--which one? *Crit Care Med* 2014; **42**: 2292-2293 [PMID: 25226120 DOI: 10.1097/CCM.0000000000000566]
- 124 **Septimus E**, Weinstein RA, Perl TM, Goldmann DA, Yokoe DS. Approaches for preventing healthcare-associated infections: go long or go wide? *Infect Control Hosp Epidemiol* 2014; **35**: 797-801 [PMID: 24915206 DOI: 10.1086/676535]
- 125 **Wenzel RP**, Edmond MB. Infection control: the case for horizontal rather than vertical interventional programs. *Int J Infect Dis* 2010; **14** Suppl 4: S3-S5 [PMID: 20851010 DOI: 10.1016/j.ijid.2010.05.002]
- 126 **Calfee DP**, Salgado CD, Milstone AM, Harris AD, Kuhar DT, Moody J, Aureden K, Huang SS, Maragakis LL, Yokoe DS; Society for Healthcare Epidemiology of America. Strategies to prevent methicillin-resistant Staphylococcus aureus transmission and infection in acute care hospitals: 2014 update. *Infect Control Hosp Epidemiol* 2014; **35**: 772-796 [PMID:

- 24915205 DOI: [10.1086/676534](https://doi.org/10.1086/676534)]
- 127 **World Health Organization.** WHO guidelines on hand hygiene in health care. Accessed September 30, 2022. Available from: <https://www.who.int/publications/i/item/9789241597906>
 - 128 **Erasmus V, Daha TJ, Brug H, Richardus JH, Behrendt MD, Vos MC, van Beeck EF.** Systematic review of studies on compliance with hand hygiene guidelines in hospital care. *Infect Control Hosp Epidemiol* 2010; **31**: 283-294 [PMID: [20088678](https://pubmed.ncbi.nlm.nih.gov/20088678/) DOI: [10.1086/650451](https://doi.org/10.1086/650451)]
 - 129 **Kowitt B, Jefferson J, Mermel LA.** Factors associated with hand hygiene compliance at a tertiary care teaching hospital. *Infect Control Hosp Epidemiol* 2013; **34**: 1146-1152 [PMID: [24113597](https://pubmed.ncbi.nlm.nih.gov/24113597/) DOI: [10.1086/673465](https://doi.org/10.1086/673465)]
 - 130 **Luyt CE, Bréchet N, Trouillet JL, Chastre J.** Antibiotic stewardship in the intensive care unit. *Crit Care* 2014; **18**: 480 [PMID: [25405992](https://pubmed.ncbi.nlm.nih.gov/25405992/) DOI: [10.1186/s13054-014-0480-6](https://doi.org/10.1186/s13054-014-0480-6)]
 - 131 **Onorato L, Macera M, Calò F, Monari C, Russo F, Iovene MR, Signoriello G, Annibale R, Pace MC, Aurilio C, Gaeta GB, Coppola N.** The effect of an antimicrobial stewardship programme in two intensive care units of a teaching hospital: an interrupted time series analysis. *Clin Microbiol Infect* 2020; **26**: 782.e1-782.e6 [PMID: [31678230](https://pubmed.ncbi.nlm.nih.gov/31678230/) DOI: [10.1016/j.cmi.2019.10.021](https://doi.org/10.1016/j.cmi.2019.10.021)]
 - 132 **Tabah A, Bassetti M, Kollef MH, Zahar JR, Paiva JA, Timsit JF, Roberts JA, Schouten J, Giamarellou H, Rello J, De Waele J, Shorr AF, Leone M, Poulakou G, Depuydt P, Garnacho-Montero J.** Antimicrobial de-escalation in critically ill patients: a position statement from a task force of the European Society of Intensive Care Medicine (ESICM) and European Society of Clinical Microbiology and Infectious Diseases (ESCMID) Critically Ill Patients Study Group (ESGCI). *Intensive Care Med* 2020; **46**: 245-265 [PMID: [31781835](https://pubmed.ncbi.nlm.nih.gov/31781835/) DOI: [10.1007/s00134-019-05866-w](https://doi.org/10.1007/s00134-019-05866-w)]
 - 133 **Ventola CL.** The antibiotic resistance crisis: part 1: causes and threats. *P T* 2015; **40**: 277-283 [PMID: [25859123](https://pubmed.ncbi.nlm.nih.gov/25859123/)]
 - 134 **Hranjec T, Rosenberger LH, Swenson B, Metzger R, Flohr TR, Politano AD, Riccio LM, Popovsky KA, Sawyer RG.** Aggressive versus conservative initiation of antimicrobial treatment in critically ill surgical patients with suspected intensive-care-unit-acquired infection: a quasi-experimental, before and after observational cohort study. *Lancet Infect Dis* 2012; **12**: 774-780 [PMID: [22951600](https://pubmed.ncbi.nlm.nih.gov/22951600/) DOI: [10.1016/S1473-3099\(12\)70151-2](https://doi.org/10.1016/S1473-3099(12)70151-2)]
 - 135 **Fiore M, Maraolo AE, Leone S, Gentile I, Cuomo A, Schiavone V, Bimonte S, Pace MC, Cascella M.** Spontaneous peritonitis in critically ill cirrhotic patients: a diagnostic algorithm for clinicians and future perspectives. *Ther Clin Risk Manag* 2017; **13**: 1409-1414 [PMID: [29081656](https://pubmed.ncbi.nlm.nih.gov/29081656/) DOI: [10.2147/TCRM.S144262](https://doi.org/10.2147/TCRM.S144262)]
 - 136 **Fiore M, Alfieri A, Di Franco S, Pace MC, Simeon V, Ingoglia G, Cortegiani A.** Ceftazidime-Avibactam Combination Therapy Compared to Ceftazidime-Avibactam Monotherapy for the Treatment of Severe Infections Due to Carbapenem-Resistant Pathogens: A Systematic Review and Network Meta-Analysis. *Antibiotics (Basel)* 2020; **9** [PMID: [32645986](https://pubmed.ncbi.nlm.nih.gov/32645986/) DOI: [10.3390/antibiotics9070388](https://doi.org/10.3390/antibiotics9070388)]
 - 137 **Fiore M, Corrente A, Pace MC, Alfieri A, Simeon V, Ippolito M, Giaratano A, Cortegiani A.** Ceftolozane-Tazobactam Combination Therapy Compared to Ceftolozane-Tazobactam Monotherapy for the Treatment of Severe Infections: A Systematic Review and Meta-Analysis. *Antibiotics (Basel)* 2021; **10** [PMID: [33467508](https://pubmed.ncbi.nlm.nih.gov/33467508/) DOI: [10.3390/antibiotics10010079](https://doi.org/10.3390/antibiotics10010079)]
 - 138 **Giacobbe DR, Saffioti C, Losito AR, Rinaldi M, Aurilio C, Bolla C, Boni S, Borgia G, Carannante N, Cassola G, Ceccarelli G, Corcione S, Dalla Gasperina D, De Rosa FG, Dentone C, Di Bella S, Di Lauria N, Feasi M, Fiore M, Fossati S, Franceschini E, Gori A, Granata G, Grignolo S, Grossi PA, Guadagnino G, Lagi F, Maraolo AE, Marinò V, Mazzitelli M, Mularoni A, Oliva A, Pace MC, Parisini A, Patti F, Petrosillo N, Pota V, Raffaelli F, Rossi M, Santoro A, Tascini C, Torti C, Trecarichi EM, Venditti M, Viale P, Signori A, Bassetti M, Del Bono V, Giannella M, Mikulska M, Tumbarello M, Viscoli C; SITA GIOVANI (Young Investigators Group of the Società Italiana Terapia Antinfettiva) and the COLI-CROSS Study Group.** Use of colistin in adult patients: A cross-sectional study. *J Glob Antimicrob Resist* 2020; **20**: 43-49 [PMID: [31207379](https://pubmed.ncbi.nlm.nih.gov/31207379/) DOI: [10.1016/j.jgar.2019.06.009](https://doi.org/10.1016/j.jgar.2019.06.009)]
 - 139 **Dickstein Y, Lellouche J, Ben Dalak Amar M, Schwartz D, Nutman A, Daitch V, Yahav D, Leibovici L, Skiada A, Antoniadou A, Daikos GL, Andini R, Zampino R, Durante-Mangoni E, Mouton JW, Friberg LE, Dishon Benattar Y, Bitterman R, Neuberger A, Carmeli Y, Paul M; AIDA Study Group.** Treatment Outcomes of Colistin- and Carbapenem-resistant *Acinetobacter baumannii* Infections: An Exploratory Subgroup Analysis of a Randomized Clinical Trial. *Clin Infect Dis* 2019; **69**: 769-776 [PMID: [30462182](https://pubmed.ncbi.nlm.nih.gov/30462182/) DOI: [10.1093/cid/ciy988](https://doi.org/10.1093/cid/ciy988)]
 - 140 **Gupta R, Hannon E, Huprikar S, Bassily-Marcus A, Manasia A, Oropello J, Kohli-Seth R.** Getting to zero: Reduction in the incidence of multidrug-resistant organism infections using an integrated infection control protocol in an intensive care unit. *Am J Infect Control* 2016; **44**: 1695-1697 [PMID: [27575774](https://pubmed.ncbi.nlm.nih.gov/27575774/) DOI: [10.1016/j.ajic.2016.07.003](https://doi.org/10.1016/j.ajic.2016.07.003)]
 - 141 **Bassetti M, Poulakou G, Ruppe E, Bouza E, Van Hal SJ, Brink A.** Antimicrobial resistance in the next 30 years, humankind, bugs and drugs: a visionary approach. *Intensive Care Med* 2017; **43**: 1464-1475 [PMID: [28733718](https://pubmed.ncbi.nlm.nih.gov/28733718/) DOI: [10.1007/s00134-017-4878-x](https://doi.org/10.1007/s00134-017-4878-x)]
 - 142 **Leone S, Damiani G, Pezone I, Kelly ME, Cascella M, Alfieri A, Pace MC, Fiore M.** New antimicrobial options for the management of complicated intra-abdominal infections. *Eur J Clin Microbiol Infect Dis* 2019; **38**: 819-827 [PMID: [30903538](https://pubmed.ncbi.nlm.nih.gov/30903538/) DOI: [10.1007/s10096-019-03533-y](https://doi.org/10.1007/s10096-019-03533-y)]
 - 143 **Healthcare-Associated Infections (Hai) Program.** Carbapenem-resistant and Carbapenemase-producing Organisms for Public Health and Healthcare Providers. Accessed September 30, 2022. Available from: https://www.cdph.ca.gov/Programs/CHCQ/HAI/Pages/CRE_InfectionPreventionStrategies.aspx
 - 144 **World Health Organization.** Global action plan on antimicrobial resistance. Accessed February 1, 2023. Available from: <https://www.who.int/publications/i/item/9789241509763>
 - 145 **Antimicrobial Resistance Collaborators.** Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet* 2022; **399**: 629-655 [PMID: [35065702](https://pubmed.ncbi.nlm.nih.gov/35065702/) DOI: [10.1016/S0140-6736\(21\)02724-0](https://doi.org/10.1016/S0140-6736(21)02724-0)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

