

World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2023 August 27; 15(8): 1559-1840



Contents

Monthly Volume 15 Number 8 August 27, 2023

MINIREVIEWS

- 1559 Impact of tumour rupture risk on the oncological rationale for the surgical treatment choice of gastrointestinal stromal tumours
Peparini N
- 1564 Prevention and treatment of hepatic encephalopathy during the perioperative period of transjugular intrahepatic portosystemic shunt
Wang LJ, Yao X, Qi Q, Qin JP
- 1574 Vascular complications of chronic pancreatitis and its management
Walia D, Saraya A, Gunjan D
- 1591 Historical changes in surgical strategy and complication management for hepatic cystic echinococcosis
A JD, Chai JP, Jia SL, A XR

ORIGINAL ARTICLE

Basic Study

- 1600 High spindle and kinetochore-associated complex subunit-3 expression predicts poor prognosis and correlates with adverse immune infiltration in hepatocellular carcinoma
Zheng LL, Wang YR, Liu ZR, Wang ZH, Tao CC, Xiao YG, Zhang K, Wu AK, Li HY, Wu JX, Xiao T, Rong WQ

Case Control Study

- 1615 Post-transplant biliary complications using liver grafts from deceased donors older than 70 years: Retrospective case-control study
Jimenez-Romero C, Justo-Alonso I, del Pozo-Elso P, Marcacuzco-Quinto A, Martín-Arriscado-Arroba C, Manrique-Municio A, Calvo-Pulido J, García-Sesma A, San Román R, Caso-Maestro O
- 1629 Goldilocks principle of minimally invasive surgery for gastric subepithelial tumors
Chang WJ, Tsao LC, Yen HH, Yang CW, Chang HC, Kor CT, Wu SC, Lin KH

Retrospective Cohort Study

- 1641 Prognosis after splenectomy plus pericardial devascularization *vs* transjugular intrahepatic portosystemic shunt for esophagogastric variceal bleeding
Qi WL, Wen J, Wen TF, Peng W, Zhang XY, Shen JY, Li X, Li C
- 1652 Initial suction drainage decreases severe postoperative complications after pancreatic trauma: A cohort study
Li KW, Wang K, Hu YP, Yang C, Deng YX, Wang XY, Liu YX, Li WQ, Ding WW

Retrospective Study

- 1663** Radiation therapy prior to a pancreaticoduodenectomy for adenocarcinoma is associated with longer operative times and higher blood loss
Aploks K, Kim M, Stroeve S, Ostapenko A, Sim YB, Sooriyakumar A, Rahimi-Ardabili A, Seshadri R, Dong XD
- 1673** Prognostic significance of preoperative lymphocyte to monocyte ratio in patients with signet ring gastric cancer
Liu HL, Feng X, Tang MM, Zhou HY, Peng H, Ge J, Liu T
- 1684** Clinical efficacy of total laparoscopic splenectomy for portal hypertension and its influence on hepatic hemodynamics and liver function
Qi RZ, Li ZW, Chang ZY, Chang WH, Zhao WL, Pang C, Zhang Y, Hu XL, Liang F
- 1693** Accurate resection of hilar cholangiocarcinoma using eOrganmap 3D reconstruction and full quantization technique
Cui DP, Fan S, Guo YX, Zhao QW, Qiao YX, Fei JD
- 1703** Regional differences in islet amyloid deposition in the residual pancreas with new-onset diabetes secondary to pancreatic ductal adenocarcinoma
Wang R, Liu Y, Liang Y, Zhou L, Chen MJ, Liu XB, Tan CL, Chen YH
- 1712** Risk factors and their interactive effects on severe acute pancreatitis complicated with acute gastrointestinal injury
Chen JH, Zhang MF, Du WC, Zhang YA
- 1719** Effects of ultrasound monitoring of gastric residual volume on feeding complications, caloric intake and prognosis of patients with severe mechanical ventilation
Xu XY, Xue HP, Yuan MJ, Jin YR, Huang CX
- 1728** Enhanced recovery nursing and mental health education on postoperative recovery and mental health of laparoscopic liver resection
Li DX, Ye W, Yang YL, Zhang L, Qian XJ, Jiang PH
- 1739** Changing trends in gastric and colorectal cancer among surgical patients over 85 years old: A multicenter retrospective study, 2001–2021
Chen K, Li M, Xu R, Zheng PP, Chen MD, Zhu L, Wang WB, Wang ZG

Observational Study

- 1751** Knowledge, attitude, and practice of monitoring early gastric cancer after endoscopic submucosal dissection
Yang XY, Wang C, Hong YP, Zhu TT, Qian LJ, Hu YB, Teng LH, Ding J
- 1761** Anti-reflux effects of a novel esophagogastric asymmetric anastomosis technique after laparoscopic proximal gastrectomy
Pang LQ, Zhang J, Shi F, Pang C, Zhang CW, Liu YL, Zhao Y, Qian Y, Li XW, Kong D, Wu SN, Zhou JF, Xie CX, Chen S
- 1774** Prognostic scores in primary biliary cholangitis patients with advanced disease
Feng J, Xu JM, Fu HY, Xie N, Bao WM, Tang YM

SYSTEMATIC REVIEWS

- 1784** Maternal choledochal cysts in pregnancy: A systematic review of case reports and case series
Augustin G, Romic I, Miličić I, Mikuš M, Herman M
- 1799** Intraoperative pancreas stump perfusion assessment during pancreaticoduodenectomy: A systematic scoping review
Robertson FP, Spiers HVM, Lim WB, Loveday B, Roberts K, Pandanaboyana S
- 1808** Comparison between upfront surgery and neoadjuvant chemotherapy in patients with locally advanced gastric cancer: A systematic review
Fiflis S, Papakonstantinou M, Giakoustidis A, Christodoulidis G, Louri E, Papadopoulos VN, Giakoustidis D

CASE REPORT

- 1819** Long-term survival of patients with hepatocellular carcinoma with hepatic, pulmonary, peritoneal and rare colon metastasis: A case report
Gong YQ, Lu TL, Chen CW
- 1825** Donor hepatic artery reconstruction based on human embryology: A case report
Zhang HZ, Lu JH, Shi ZY, Guo YR, Shao WH, Meng FX, Zhang R, Zhang AH, Xu J
- 1831** Outpatient hybrid endoscopic submucosal dissection with SOUTEN for early gastric cancer, followed by endoscopic suturing of the mucosal defect: A case report
Ito R, Miwa K, Matano Y

LETTER TO THE EDITOR

- 1838** Is endoscopic mucosal resection-precutting superior to conventional methods for removing sessile colorectal polyps?
Yang QY, Zhao Q, Hu JW

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Raja Kalayarasan, MS, DNB, MCh, FRCS (Ed), Additional Professor & Head, Department of Surgical Gastroenterology, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Puducherry 605006, India. kalayarasanraja@yahoo.com

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (WJGS, *World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

INDEXING/ABSTRACTING

The WJGS is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGS as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Rui-Rui Wu; Production Department Director: Xiang Li; Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

August 27, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Historical changes in surgical strategy and complication management for hepatic cystic echinococcosis

Ji-De A, Jin-Ping Chai, Sheng-Long Jia, Xiang-Ren A

Specialty type: Infectious diseases

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B, B

Grade C (Good): C, C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Amante MF, Argentina; El-Nakeep S, Egypt; Mikulic D, Croatia

Received: March 30, 2023

Peer-review started: March 30, 2023

First decision: May 12, 2023

Revised: May 17, 2023

Accepted: June 21, 2023

Article in press: June 21, 2023

Published online: August 27, 2023



Ji-De A, Department of Hepatic Hydatidosis, Qinghai Provincial People's Hospital, Xining 810007, Qinghai Province, China

Jin-Ping Chai, Department of Internal Medicine-Cardiovascular, Qinghai Provincial People's Hospital, Xining 810007, Qinghai Province, China

Sheng-Long Jia, Department of General Surgery, Huangzhong Hospital of Traditional Chinese Medicine, Xining 810007, Qinghai Province, China

Xiang-Ren A, Department of Clinical Laboratory, Qinghai Provincial People's Hospital, Xining 810007, Qinghai Province, China

Corresponding author: Xiang-Ren A, MBBS, Chief Physician, Doctor, Department of Clinical Laboratory, Qinghai Provincial People's Hospital, No. 2 Gonghe Road, Chengdong District, Xining 810007, Qinghai Province, China. xiangrena_001@163.com

Abstract

Echinococcosis is a zoonotic parasitic disease caused by *Escherichia* larvae. It frequently involves the liver (70%-75%), followed by the lungs (15%-20%), and occasionally the brain, heart, spleen, bone, and other organs. The main pathogenic forms of human echinococcosis currently include cystic echinococcosis (CE) and alveolar echinococcosis (AE). CE is globally distributed, while the distribution of AE is generally restricted to the northern hemisphere. In China, CE accounts for 75% of all echinococcosis cases. With rapid advances in surgical techniques in recent decades, the surgical strategy for CE has changed, especially with the continuous improvement of surgical methods and the expansion of surgical contraindications. To further understand the changes in surgical treatment strategies for hepatic CE, we interpreted and analyzed the existing literature addressing the surgical treatment of hepatic CE both domestically and abroad and briefly summarized them in chronological order. This review aims to provide a deeper understanding of the progress in the surgical treatment of hepatic CE to provide clearer avenues for its clinical diagnosis and treatment.

Key Words: Echinococcosis; Hepatic cystic; Internal capsule; External capsule; Complication; Surgical excision

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Many articles have been published on the surgical methods and postoperative complications of hepatic cystic echinococcosis (CE), especially regarding the relationship between surgical methods and patient prognosis. However, few studies have examined the historical changes in surgical methods for hepatic CE. Therefore, this review discusses the main surgical methods and complications in the treatment of hepatic CE over time to provide readers with a deeper understanding of the surgical treatment of this disease.

Citation: A JD, Chai JP, Jia SL, A XR. Historical changes in surgical strategy and complication management for hepatic cystic echinococcosis. *World J Gastrointest Surg* 2023; 15(8): 1591-1599

URL: <https://www.wjgnet.com/1948-9366/full/v15/i8/1591.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v15.i8.1591>

INTRODUCTION

Echinococcosis, also known as hydatid disease, is a parasitic disease caused by the echinococcosis tapeworm larvae. The liver is the primary organ involved in echinococcosis, with the brain and other organs occasionally involved[1,2]. To date, at least five species of larvae have been reported to cause echinococcosis, including *Echinococcus granulosus*, *Echinococcus multilocularis*, *Echinococcus oligarthrus*, *Echinococcus vogeli*, and *Echinococcus shiquicus*. However, two main pathogens cause echinococcosis in humans: *Echinococcus granulosus* and *Echinococcus multilocularis*[3-5]. Cystic echinococcosis (CE), caused by *Echinococcus granulosus*, is globally distributed, while alveolar echinococcosis (AE), caused by *Echinococcus multilocularis*, is restricted to the northern hemisphere[6]. The distributions of CE and AE in China are slightly different from the global distribution of echinococcosis, where CE accounts for approximately 75% of all echinococcosis cases[7]. With the rapid advances in surgical techniques in recent decades, the surgical treatment strategy for CE has changed, especially with the continuous improvement of surgical methods and the expansion of surgical contraindications[8]. Original internal capsule excision is a widely used classical procedure because of its safety, practicality, and simplicity. However, owing to the high incidence of postoperative complications, especially residual cavity infection after long-term catheterization, which results in significant physical and psychological trauma, the procedure has been gradually abandoned[9]. Subsequently, various modified methods based on this procedure have evolved, such as internal capsule removal combined with omental tamponade and internal capsule removal combined with external capsule subtotal resection. Although these procedures have addressed many complications caused by partial residual cavity retention, they have not resolved persistent problems associated with internal capsule removal[10]. With the promotion of precision surgery, radical surgical procedures such as external capsule exfoliation and regular or irregular hepatectomy have gradually been accepted by most surgeons in the field[11]. These procedures significantly reduce the incidence of residual cavity complications associated with internal capsule exfoliation. However, radical surgery has stringent requirements for operators and surgical conditions; therefore, it is difficult to popularize this technique. Moreover, forced radical surgery can lead to fatal complications[12,13]. Specialists originally intended to design an individualized surgical plan from the perspective of patient and surgical safety. This article reviews surgical treatment strategies for hepatic CE and the historical evolution of postoperative complications (Table 1).


INITIAL SURGICAL TREATMENT FOR HEPATIC CE

Excision of the internal capsule

A case of successful internal capsule removal for liver CE was first described in 1871[14]. Excision of the internal capsule was the only surgery performed for CE during this period and is still used as a classical surgical approach (Figure 1). Subsequently, some investigators have proposed principles for CE surgery, specifically clearance of the internal capsule, prevention of the extravasation of cyst fluid, elimination of the residual cavity of the external capsule, and prevention of infection[15]. Excision of the internal capsule has the advantages of minimal surgical trauma, safety, and reliability and is a simple operation; however, each procedure has its own drawbacks. Indeed, many common complications can occur after internal capsule removal, such as *in situ* recurrence of the residual cavity, disseminated implantation, biliary fistula, residual cavity infection, and other refractory complications[16]. Some researchers have reported that the recurrence rates of liver CE after internal capsule exocytosis and biliary fistula are as high as 40% and 20%, respectively, and the incidence of residual cavity infection is as high as 65%[17-20]. The main reason for CE recurrence is rupture of the cyst during the operation or failure to completely eliminate the protoscolex when the inner CE cyst is removed, and the protoscolex left in the residual cavity continues to survive and ultimately develops into CE in the outer cavity[21]. The reason for intraperitoneal implantation is that the CE cyst fluid extravasates into the abdomen and pelvis during the operation, contaminating other abdominal organs, and a large number of larvae adhere to the peritoneal surface and survive. The main cause of postoperative biliary fistula is prolonged compression of the bile duct by a CE cyst, leading to intrahepatic bile duct distortion, atrophy, and even distal ischemic necrosis. In addition, owing to the corrosive effect of bile and the dramatic increase in luminal pressure, the external capsule and biliary tract are prone to form internal fistulas. The premise of a postoperative residual cavity infection is the existence of a residual cavity. Some patients experience

Table 1 Main surgical methods for hepatic cystic echinococcosis and their complications in different periods

Period (year)	Surgical methods	Major complications
Initial stage		
1871	Excision of the internal capsule	Recurrence <i>in situ</i> , abdominal implantation, biliary fistula, and residual cavity infection
Mature stage		
1965	Complete removal of the external capsule and hepatectomy	Intraoperative hemorrhage, cyst rupture, and liver dysfunction
1985	Percutaneous puncture and catheter drainage	Allergic reaction, abdominal implantation, residual cavity infection, biliary fistula, and recurrence <i>in situ</i>
2002	Modified excision of the internal capsule	Abdominal implantation and biliary fistula
2010	Internal capsule removal combined with external capsule subtotal resection	Recurrence <i>in situ</i> , disseminated implantation, and biliary fistula
Current stage		
1992	Laparoscopic cyst excision	Intraoperative hemorrhage and cyst rupture
1994	Local ablation	Damage to surrounding tissues and organs. Allergic reaction, cyst rupture, abdominal implantation, and recurrence <i>in situ</i>
2002	Liver transplantation	Organ shortage, postoperative complications, and high cost



DOI: 10.4240/wjgs.v15.i8.1591 Copyright ©The Author(s) 2023.

Figure 1 Excision of the internal capsule. A: After the incision of the peritoneum, the outer capsule of cystic echinococcosis (CE) was ruptured, and the inner capsule was intact; B: The gauze soaked with iodophor was used to isolate the surrounding organs and the intact ascus was removed carefully; C: The CE cyst was large in size and dense in adhesion with the diaphragm. Therefore, we chose to perform the internal capsule enucleation and found an internal fistula in the bile duct of the hydatid cyst, and the contents of the cyst were stained with bile; D: The aspirator aspirated the contents of the capsule, the capsule was repeatedly rinsed with iodophor and wiped with gauze, and a rubber drainage tube was placed.

preoperative cystic cavity infection, or the cyst wall is thick and calcified, and the residual cavity does not heal for a prolonged period. Another major cause of residual cavity infection is the presence of an internal biliary fistula on the cyst wall. When the pressure in the cyst sharply decreases, bile enters the residual cavity from the fistula and causes infection.

MATURE STAGE OF SURGICAL TREATMENT FOR HEPATIC CE

Percutaneous puncture and catheter drainage of CE cysts

Advances and developments in modern medical science have led to a more comprehensive and in-depth understanding of CE, especially when a breakthrough was achieved in the puncture treatment of CE, in which the puncture diagnosis and treatment were no longer considered a “taboo”. In 1985, Livraghi *et al*[22] performed an ultrasound-guided diagnostic puncture for hepatic CE. In the same year, Mueller first reported the success of fine-needle puncture in the treatment of human hepatic CE[23]. In 1990, researchers reported ultrasound-guided puncture sclerotherapy for hepatic CE cysts in a large number of patients. This procedure is mainly aimed at patients with simple cyst-type CE who cannot tolerate surgery or for whom the effect of albendazole is not ideal. It has the advantages of less trauma, safety, effectiveness, low cost, and a short hospital stay[24]. Investigators domestically and abroad have reported that percutaneous tube puncture under ultrasound guidance combined with albendazole for the treatment of CE achieved curative effects similar to those of surgery[25]. Ultrasound-guided percutaneous puncture combined with drug therapy for CE is mainly used for cases of liver and abdominal wall adhesions caused by multiple operations and is a simple and less invasive method. Thousands of cases have been reported, both domestically and internationally. The rate of anaphylaxis and shock of puncture and extravasation of cyst fluid leading to CE implantation and dissemination is not higher than that of traditional surgical methods; as such, it is included in the World Health Organization (WHO) guidelines for the diagnosis and treatment of CE. However, some studies have reported that puncture treatment for CE is also associated with problems, including allergic reactions and cholangiosclerosis caused by head-extinguishing agents such as ethanol and formalin, residual cavity infection, biliary fistula, and recurrence *in situ*[26]. Some investigators have reported a complication rate of approximately 17%, a CE cyst infection rate of 2.6%, and an *in situ* recurrence rate as high as 40%[27-29]. To address this problem, researchers designed a special puncture instrument. Puncture and aspiration of the sac contents have been reported to have positive effects. However, the complication rate remained high after follow-up; therefore, this procedure was abandoned in China[30].

Modified excision of the internal capsule


At the beginning of the 20th century, Wen Hao first proposed modified enucleation of the internal capsule for the treatment of hepatic CE. The goal of this operation is to further reduce the incidence of high residual cavity complications after traditional internal capsule removal. A further improved surgical method is performed based on traditional internal capsule removal[31] to completely remove the internal capsule, remove the external capsule as far as possible, convert the large residual cavity into a small residual cavity, suture and repair the biliary fistula, manage the residual cavity with omental tamponade, which can reduce CE recurrence, abdominal spread, and the occurrence of refractory biliary fistula and residual cavity fluid infection. This surgery is suitable for patients with multiple cysts in whom the external capsule cannot be completely removed. Studies have reported that the incidence rates of biliary fistula and residual cavity infections were 17.56% and 2.70%, respectively. There were no recurrences or deaths, and the cure rate was reported to be 100%[10].

Total cystectomy

In 1965, French researchers proposed the concept of radical surgery for the complete removal of the CE external capsule. Peng *et al*[32,33] reported, for the first time in 1999, the complete excision of a hepatic CE cyst inside the outer membrane, which was based on the liver resection technique, combined with experience in the surgical treatment of hepatic CE. The main technical point and radical purpose of this procedure are to completely remove the CE cyst along the potential gap between the external capsule of the CE and the normal liver tissue (Figure 2). Compared with traditional internal capsule extraction, this operation significantly reduces postoperative bile leakage, residual cavity infection, and recurrence. It is an ideal new surgical method for radical treatment, which has been recognized by the WHO and gradually popularized [34,35]. This procedure is suitable for patients with single or external cysts that can be completely resected. Complete excision of the external capsule can completely remove the lesion; however, this operation requires a high degree of knowledge and proficient surgical skills of the surgeon. Otherwise, the cyst could easily rupture during surgery, resulting in serious intraoperative complications. Some studies have reported that the incidence of biliary fistula complications after complete exfoliation of the external capsule is 2.34%, and the incidence of postoperative liver dysfunction is 3.03%. No cases of postoperative recurrence, residual cavity infection, implantation spread, or death occurred, and the cure rate was 100%[36]. However, owing to the expansive growth of CE cysts, some important vessels in the liver are compressed, deformed, and/or displaced, and an internal fistula may form. Although there is a potential gap between the external capsule and normal liver tissue, these vessels and bile ducts are likely to be destroyed during the removal of the external capsule. Furthermore, intraoperative hemorrhage is significantly increased compared with the removal of the internal capsule and subtotal removal of the external capsule[37,38].


Partial hepatectomy

In 2007, researchers reported that partial hepatectomy was used in France to treat hepatic CE as early as 1965[37]. With advances in liver surgery, hepatectomy has become one of the main radical treatments for hepatic CE[39]. Birnbaum *et al* [40] proposed that active hepatectomy has advantages over conservative internal capsule removal in the treatment of CE. However, not all CE cases should be treated by liver resection because regular liver resection removes part of the normal liver tissue at the same time as the resection of the lesion, especially when the residual liver volume is small; therefore, hepatectomy should be more conservative (Figure 3). This procedure is mainly suitable for patients with multiple and/or large lesions, thick and calcified CE cyst walls, or complications due to intracystic infection[32,41].



DOI: 10.4240/wjgs.v15.i8.1591 Copyright ©The Author(s) 2023.

Figure 2 Total cystectomy. A: The cystic echinococcosis (CE) cyst was located behind the gallbladder, and the lesion size was small; B: The gallbladder and CE cyst were completely removed; C: CE cyst was located in the S5 and S6 segments of the liver, with a certain distance from the first and second hilum hepatitis; D: The CE cyst was completely removed along the edge of the external capsule, and the remaining liver section was definitely hemostatic.



DOI: 10.4240/wjgs.v15.i8.1591 Copyright ©The Author(s) 2023.

Figure 3 Partial hepatectomy. A: Cystic echinococcosis (CE) occupied the entire S3 segment of the liver, so segmentectomy was performed; B: Some liver tissue was observed on the surface of CE.

Internal capsule removal combined with external capsule subtotal resection

In 2010, researchers described the treatment of hepatic CE using internal capsule removal combined with subtotal resection[42]. This procedure combines the advantages of traditional endocystectomy and complete removal of the external capsule, particularly in patients with complex CE or in whom total cystectomy is difficult. Moreover, patients with CE cysts near the main hilar vessels and biliary tract achieved satisfactory results. The primary technical aspects include conventional internal capsule extraction followed by external capsule stripping along the potential gap of the CE external capsule, which maximally preserves the normal liver parenchyma and protects the surrounding hepatic duct system. Simultaneously, separation in this space also reduces bleeding without blocking blood flow into the liver, leaving a residue in the outer capsule wall near important blood vessels and the hilus. It aims to remove the outer capsule wall to the greatest extent possible, reduce the residual cavity volume, or eliminate the residual cavity, thereby significantly reducing postoperative residual cavity infections and biliary fistula complications. However, Zhang *et al*[43] found that, although subtotal external capsule resection removes part of the external capsule, it does not avoid the risk of a residual

internal capsule. For example, during surgery, the internal capsule is routinely removed, and there is the possibility of dissemination and implantation of the cephalic segment during the removal of the internal capsule. In addition, compared with internal capsule resection, subtotal external capsule resection also requires the removal of part of the external capsule, which prolongs the operation and increases the surgical risk and incidence of postoperative complications.

CURRENT STATUS OF SURGICAL TREATMENT FOR HEPATIC CE

Laparoscopic CE cyst excision

With advances and developments in minimally invasive techniques, particularly laparoscopic surgery, to a higher level, the indications for surgery continue to expand and extend to the treatment of various diseases. The first case of laparoscopic liver CE internal capsule excision was reported[44]. In the ensuing ten years, several liver surgery centers in China have successively performed total laparoscopic complete cystectomies and laparoscopic hepatectomies to treat hepatic CE. Furthermore, laparoscopic hepatectomy is safe, reliable, minimally invasive, and has a rapid postoperative recovery [45,46]. However, the technical requirements for laparoscopic hepatectomy are high, the technological maturity curve is long, and cases must be carefully selected. The procedure needs to be performed in experienced endoscopic centers; therefore, it is difficult to promote wide-range adoption and a low recurrence rate in the short term. Moreover, surgical indications should be strictly controlled, such as those without a history of abdominal surgery, and cases with CE cysts located in the superficial part of the liver, and clear exposure of the surgical field[47]. Additionally, the three principles of CE surgery should be strictly observed during the procedure: No cystic fluid overflow, a fully inactivated original segment, and field isolation and protection. Attention should be paid to inactivating the extravasated cystic fluid to prevent infection and implant recurrence; therefore, the choice of indication must be emphasized.

Local ablation therapy for CE

With the popularization of minimally invasive concepts in recent years, local lesion ablation, a newly emerging minimally invasive technique, has been recognized by the medical community for the treatment of diseases, especially small lesions [48]. After more than 20 years of development, many domestic and foreign echinococcosis diagnosis and treatment centers have attempted to apply this technology to the treatment of CE or AE because of its simplicity, minimal trauma, low incidence of postoperative complications, and rapid recovery[49-52]. The echinococcosis diagnosis and treatment center of Qinghai Provincial People's Hospital successfully performed radiofrequency ablation for liver CE disease in more than 40 cases. In addition, we found that patients with a smaller hepatic echinococcosis diameter had a shorter operative duration, less postoperative liver damage, and faster recovery of liver function. The operative duration for a single lesion is shorter than that for multiple lesions, the postoperative liver damage is less severe, and the short term clinical effect is good. Radiofrequency ablation has the advantages of being a simple operation that results in less trauma, rapid patient recovery, and fewer complications, and it can, to a certain extent, reduce the economic burden on patients. In addition, in patients with advanced age, poor pulmonary function, or cirrhosis who do not meet the indications for surgery or cannot undergo surgery, it can replace part of the traditional open surgery. However, there are also many uncertain factors in radiofrequency ablation, such as the severity of damage to the surrounding tissues and organs, which cannot be verified. Therefore, to reduce the damage to organs, tissues, and cells caused by radiofrequency ablation puncture, it is necessary to strictly observe operative specifications, accurate preoperative positioning, and careful intraoperative ultrasound guidance, and control the temperature and ablation time during radiofrequency ablation. Reducing the number of ablation attempts with a radiofrequency needle, repeated irregular insertion of the radiofrequency needle, and avoiding careless operations can effectively reduce damage to intrahepatic bile ducts, tissues, and cells.

Liver transplantation for hepatic CE

Hepatic CE lesions compress the second hilum, resulting in severe liver congestion. Traditional treatments cannot effectively improve liver function, hypersplenism, and portal hypertension or relieve abdominal fluid[9]. Allograft liver transplantation is performed when there is a risk of liver failure after autologous transplantation. However, liver transplantation has some shortcomings, such as organ shortage, postoperative complications, and high costs. Therefore, liver transplantation in patients with hepatic CE is rarely reported globally, and there are no records of liver and biliary disease treatment centers in the authors' location.

Continuous improvements in interventional therapeutic technologies, such as laparoscopy, and their application in clinical practice have ushered in significant changes and breakthroughs in the diagnosis, treatment, and postoperative management of hepatic CE. The boundary between minimally invasive and traditional surgeries is constantly being broken, especially as laparoscopic surgery progresses to higher levels. The application of imaging techniques, such as computed tomography angiography, magnetic resonance cholangiopancreatography, and three-dimensional reconstruction, can reveal the relationship between CE lesions, liver vessels, and the biliary tract and provide detailed preoperative imaging for precision surgery for CE. Despite these remarkable achievements, problems persist in the diagnosis and treatment of hepatic CE. Accurate surgical prediction, postoperative evaluation, and the diagnosis and treatment of postoperative complications require further investigation.

MANAGEMENT OF COMPLICATIONS

Albendazole should be administered for a prolonged period after CE recurrence. Reoperation should be considered for patients with poor drug efficacy to prolong and improve their quality of life. Among patients who experience a postoperative biliary fistula, only unobstructed drainage is needed if the size of the biliary fistula is small. If stubborn biliary leakage occurs, which requires long-term catheterization or repeated debridement, and the liver is severely damaged, partial hepatectomy or reoperation can be performed. The leak is sutured, and biliary duct decompression or endoscopic duodenal papillary incision is made for biliary duct decompression, and the biliary leak can be closed by itself [21]. For patients with postoperative residual cavity infection, the first step is smooth drainage, followed by reasonable use of antibiotics to control the infection. If the infection persists, repeat surgical debridement or resection of the part of the liver with bile leakage should be considered to achieve recovery. To prevent recurrence *in situ* or peritoneal implantation, intraoperative protection and isolation measures should be taken, such as using hypertonic saline or iodophor gauze to isolate the tissues and organs around the lesion during the operation, regularly taking albendazole after the operation, and follow-up work to achieve early detection and treatment as far as possible.

CONCLUSION

Compared with 200 years ago, there is a deeper understanding of the pathogenesis of CE, and medical diagnosis and treatment technologies have improved dramatically. However, key treatment principles for hepatic CE remain unchanged. Complete resection of the lesion and prevention of CE recurrence and implantation are lifelong pursuits of CE specialists. However, our team believes that because we cannot guarantee the integrity of the cyst, we should shift the focus of treatment to prevent the inactivation of the head segment, such as pre-inactivation of the head segment and solidification of the cyst fluid, which can not only reduce the implantation and recurrence rates of CE but also improve the safety of surgery. We hope that there will be qualitative breakthroughs in the surgical treatment of hepatic CE.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the helpful suggestions of the reviewers, who have improved the manuscript.

FOOTNOTES

Author contributions: A JD and Chai JP conceptualized the study, collected the data, designed the review, and drafted the original manuscript; Jia SL was responsible for the investigation and supervision; and A XR supervised and reviewed the final manuscript submitted for publication.

Supported by the Qinghai Province Talent Action Plan of Kunlun; National Natural Science Foundation of China, No. 82260412; Basic Research Project of Qinghai Province, No. 2020-wjzdx-27; Qinghai Province 2022 Innovation Platform Construction Special Project, No. 2022-ZJ-T01; and High-end Innovative Talent Project of Kunlun Talents of Qinghai Province in 2021.

Conflict-of-interest statement: The authors declare that they have no conflict of interest to disclose.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Ji-De A 0000-0003-4478-1972; Jin-Ping Chai 0000-0001-8873-1323; Xiang-Ren A 0000-0002-0305-996X.

S-Editor: Yan JP

L-Editor: Wang TQ

P-Editor: Yan JP

REFERENCES

- 1 Keong B, Wilkie B, Sutherland T, Fox A. Hepatic cystic echinococcosis in Australia: an update on diagnosis and management. *ANZ J Surg* 2018; **88**: 26-31 [PMID: 29024292 DOI: 10.1111/ans.14117]
- 2 Zhang RQ, Chen XH, Wen H. Improved experimental model of hepatic cystic hydatid disease resembling natural infection route with stable growing dynamics and immune reaction. *World J Gastroenterol* 2017; **23**: 7989-7999 [PMID: 29259374 DOI: 10.3748/wjg.v23.i45.7989]

- 3 Nunnari G, Pinzone MR, Gruttadauria S, Celesia BM, Madeddu G, Malaguarnera G, Pavone P, Cappellani A, Cacopardo B. Hepatic echinococcosis: clinical and therapeutic aspects. *World J Gastroenterol* 2012; **18**: 1448-1458 [PMID: 22509076 DOI: 10.3748/wjg.v18.i13.1448]
- 4 Xing ZK, Wang EQ, Liao Y, Gao FY, Zhao XY, Tan XW, Yu XF, Wu XW. Comparison of fibrosis in the surrounding tissues of hepatic echinococcosis granulosa and echinococcosis multilocularis in mice at different stages. *Zhongguo Renshou Gonghuanbing Xuebao* 2021; **37**: 1008-1016 [DOI: 10.3969/j.issn.1002-2694.2021.00.140]
- 5 Guo M, Ailiyali A, Tuerganaili A, Shao YM. Establishment and preliminary application of cloud platform system for hydatid diseases. *Zhongguo Renshou Gonghuanbing Xuebao* 2019; **35**: 970-972
- 6 Cao DP, Zhang YG, Jiang BP, Li JH, Pang MQ, Chen G. The effects of hydatid cyst and alveolar hydatid infection on lymphocyte subsets in Balb/c mice. *Zhongguo Renshou Gonghuanbing Xuebao* 2021; **37**: 421-425
- 7 Gandhiraman K, Balakrishnan R, Ramamoorthy R, Rajeshwari R. Primary Peritoneal Hydatid Cyst Presenting as Ovarian Cyst Torsion: A Rare Case Report. *J Clin Diagn Res* 2015; **9**: QD07-QD08 [PMID: 26436004 DOI: 10.7860/JCDR/2015/14324.6397]
- 8 Bayrak M, Altintas Y. Current approaches in the surgical treatment of liver hydatid disease: single center experience. *BMC Surg* 2019; **19**: 95 [PMID: 31315619 DOI: 10.1186/s12893-019-0553-1]
- 9 Chinese College of Surgeons Chinese Committee for Hydatidology. Expert consensus on the diagnosis and treatment of hepatic cystic and alveolar echinococcosis (2019 edition). *Zhonghua Xiaohua Waikē Zazhi* 2019; **18**: 711-721
- 10 Abulikemu W. The Application of Individualized Therapy in The Surgical treatment of hepatic Hydatid Disease and Its Clinical Significance. Xinjiang Medical University, 2013
- 11 Goja S, Saha SK, Yadav SK, Tiwari A, Soin AS. Surgical approaches to hepatic hydatidosis ranging from partial cystectomy to liver transplantation. *Ann Hepatobiliary Pancreat Surg* 2018; **22**: 208-215 [PMID: 30215042 DOI: 10.14701/ahbps.2018.22.3.208]
- 12 Chen ZY. Surgical treatment of complex hepatic echinococcosis. *Zhongguo Puwai Jichu Yu Linchuang Zazhi* 2017; **24**: 785-787 [DOI: 10.7507/1007-9424.201705115]
- 13 Li MH, Yang ZQ, Yan WT. Surgical treatment of complex hepatic cystic echinococcosis. *Zhongguo Puwai Jichu Yu Linchuang Zazhi* 2020; **27**: 4-6 [DOI: 10.7507/1007-9424.201911079]
- 14 Wen H, Shao YM, Zhao JM, Tuergan A, Li HT, Maimaitijiang AK. Clinical analysis of surgical treatment for human hepatic cystic and alveolar echinococcosis. *Zhonghua Xiaohua Waikē Zazhi* 2007; **6**: 13-18 [DOI: 10.3760/cma.j.issn.1673-9752.2007.01.010]
- 15 Chen J, Wen H. Advances in diagnosis and treatment of hepatic echinococcosis. *Dongnan Daxue Xuebao (Yixueban)* 2018; **37**: 929-934 [DOI: 10.3969/j.issn.1671-6264.2018.05.038]
- 16 Gou P, Wang ZX, Hu CL, Fu SQ, Fan HN, Wang HJ. Retrospective analysis of treatment therapies of alveolar, cystic echinococcosis and their postoperative complications in western area of China. *Shiyong Linchuang Yiyao Zazhi* 2019; **23**: 33-37 [DOI: 10.7619/jcmp.201903010]
- 17 Yan XK, Luo RQ, Shi XP. Laparoscopic treatment of hepatic echinococcosis: report of 18 cases. *Zhongguo Puwai Jichu Yu Linchuang Zazhi* 2012; **19**: 318
- 18 Zhao YY. Prevention and management of complications of retained cavity of postoperative hydatidosis (a report of 173 patients). *Lanzhou Daxue Xuebao (Yixueban)* 2005; **31**: 24-26 [DOI: 10.3969/j.issn.1000-2812.2005.01.007]
- 19 Yisimayili A, Maimaitiuerxun T, Xu QL, Tang RH, Keli M. Cause and treatment of postoperative residual biliary fistula in hepatic cystic echinococcosis. *Gandanyi Waikē Zazhi* 2018; **30**: 498-500, 509 [DOI: 10.11952/j.issn.1007-1954.2018.06.014]
- 20 Tang MJ, Yang JY. Management of accidental rupture of hydatid cyst during 3D laparoscopic cystic hepatic hydatid resection in plateau area. *Zhongguo Weichuan Waikē Zazhi* 2020; **26**: 272-273, 281 [DOI: 10.3969/j.issn.1009-6604.2020.03.020]
- 21 Surmelioglu A, Ozer I, Reyhan E, Dalgic T, Ozdemir Y, Ulas M, Bostanci EB, Akoglu M. Risk Factors for Development of Biliary Complications after Surgery for Solitary Liver Hydatid Cyst. *Am Surg* 2017; **83**: 30-35 [PMID: 28234119]
- 22 Livraghi T, Bosoni A, Giordano F, Lai N, Vettori C. Diagnosis of hydatid cyst by percutaneous aspiration: value of electrolyte determinations. *J Clin Ultrasound* 1985; **13**: 333-337 [PMID: 3924964 DOI: 10.1002/jcu.1870130507]
- 23 Morris DL. Hepatic echinococcal cyst: successful percutaneous drainage. *Radiology* 1986; **159**: 567 [PMID: 3961197 DOI: 10.1148/radiology.159.2.3961197]
- 24 Zeng J, Ye JW, Zhang J, Li J. Advancement on Non-surgical Treatment for the Hepatic Cystic Echinococcosis. *Xiandai Shengwu Yixue Jinzhan* 2018; **18**: 3790-3794 [DOI: 10.13241/j.cnki.pmb.2018.19.042]
- 25 Khuroo MS. Percutaneous Drainage in Hepatic Hydatidosis-The PAIR Technique: Concept, Technique, and Results. *J Clin Exp Hepatol* 2021; **11**: 592-602 [PMID: 34511821 DOI: 10.1016/j.jceh.2021.05.005]
- 26 Giorgio A, De Luca M, Gatti P, Ciraci E, Montesarchio L, Santoro B, Di Sarno A, Coppola C, Giorgio V. Treatment of Hydatid Liver Cyst With Double Percutaneous Aspiration and Ethanol Injection Under Ultrasound Guidance: 6.5-Year Median Follow-up Analysis. *Cardiovasc Intervent Radiol* 2021; **44**: 1214-1222 [PMID: 33987694 DOI: 10.1007/s00270-021-02839-9]
- 27 Gargouri M, Ben Amor N, Ben Chehida F, Hammou A, Gharbi HA, Ben Cheikh M, Kchouk H, Ayachi K, Golvan JY. Percutaneous treatment of hydatid cysts (Echinococcus granulosa). *Cardiovasc Intervent Radiol* 1990; **13**: 169-173 [PMID: 2121344 DOI: 10.1007/BF02575469]
- 28 Muhammedoglu B, Pircanoglu EM, Piskin E, Torun S, Karadağ M, Topuz S, Köktaş S. Treatment of Hepatic Hydatid Disease: Role of Surgery, ERCP, and Percutaneous Drainage: A Retrospective Study. *Surg Laparosc Endosc Percutan Tech* 2020; **31**: 313-320 [PMID: 32956333 DOI: 10.1097/SLE.0000000000000861]
- 29 Yan S, Li M, Tan XL. Interventional ultrasound treatment of 120 cases of hepatic hydatid cyst. *Zhonghua Chaosheng Yixue Zazhi* 1995; **11**: 399-400
- 30 Wen H, Luan MX, Yang WG. Study on standard classification and clinical significance for hepatic echinococcosis. *Xinjiang Yike Daxue Xuebao* 2002; **25**: 129-130 [DOI: 10.3969/j.issn.1009-5551.2002.02.005]
- 31 Tuergan A, Shao YM, Jiang TM, Dai JB, Ran B, Wen H. Diagnosis and treatment of ruptured hepatic cystic echinococcosis: report of 109 cases. *Zhonghua Xiaohua Waikē Zazhi* 2011; **10**: 293-296 [DOI: 10.3760/cma.j.issn.1673-9752.2011.04.015]
- 32 Peng XY, Zhang SJ, Niu JH, Wu XW, Pan HZ, Mo LT, Chen Y, Yang HQ. Total subadventitial cystectomy for the treatment of 30 patients with hepatic hydatid cyst. *Zhonghua Xiaohua Waikē Zazhi* 2002; **17**: 529-530 [DOI: 10.3760/j.issn.1007-631X.2002.09.005]
- 33 Peng XY. New perspectives on surgical treatment of hepatic echinococcosis. *Zhongguo Shiyong Waikē Zazhi* 2003; **23**: 651-653 [DOI: 10.3321/j.issn.1005-2208.2003.11.007]
- 34 Botezatu C, Mastalier B, Patrascu T. Hepatic hydatid cyst - diagnose and treatment algorithm. *J Med Life* 2018; **11**: 203-209 [PMID: 30364592 DOI: 10.25122/jml-2018-0045]

- 35 **Al-Saeedi M**, Ramouz A, Khajeh E, El Rafidi A, Ghamarnejad O, Shafiei S, Ali-Hasan-Al-Saegh S, Probst P, Stojkovic M, Weber TF, Hoffmann K, Mehrabi A. Endocystectomy as a conservative surgical treatment for hepatic cystic echinococcosis: A systematic review with single-arm meta-analysis. *PLoS Negl Trop Dis* 2021; **15**: e0009365 [PMID: 33979343 DOI: 10.1371/journal.pntd.0009365]
- 36 **Gollackner B**, Längle F, Auer H, Maier A, Mittlböck M, Agstner I, Karner J, Langer F, Aspöck H, Loidolt H, Rockenschaub S, Steininger R. Radical surgical therapy of abdominal cystic hydatid disease: factors of recurrence. *World J Surg* 2000; **24**: 717-721 [PMID: 10773125 DOI: 10.1007/s002689910115]
- 37 **Mekeel KL**, Hemming AW. Combined resection of the liver and the inferior vena cava for hydatid disease. *J Gastrointest Surg* 2007; **11**: 1741-1743 [PMID: 17874275 DOI: 10.1007/s11605-007-0292-y]
- 38 **Tuergan A**, Shao YM, Zhao JM, Zhang JH, Li HT, Wen H. Operative technique and indication of total cystectomy for treatment of cystic echinococcosis: a report of 64 cases. *Zhonghua Gandan Waiké Zazhi* 2007; **13**: 247-249 [DOI: 10.3760/cma.j.issn.1007-8118.2007.04.011]
- 39 **Losada Morales H**, Burgos San Juan L, Silva Abarca J, Muñoz Castro C. Experience with the surgical treatment of hepatic hydatidosis: case series with follow-up. *World J Gastroenterol* 2010; **16**: 3305-3309 [PMID: 20614487 DOI: 10.3748/wjg.v16.i26.3305]
- 40 **Birnbaum DJ**, Hardwigsen J, Barbier L, Bouchiba N, Le Treut YP. Is hepatic resection the best treatment for hydatid cyst? *J Gastrointest Surg* 2012; **16**: 2086-2093 [PMID: 22903365 DOI: 10.1007/s11605-012-1993-4]
- 41 **Marom G**, Khoury T, Gazla SA, Merhav H, Padawer D, Benson AA, Zamir G, Luques L, Safadi R, Khalaileh A. Operative treatment of hepatic hydatid cysts: A single center experience. *Asian J Surg* 2019; **42**: 702-707 [PMID: 30446425 DOI: 10.1016/j.asjsur.2018.09.013]
- 42 **Deo KB**, Kumar R, Tiwari G, Kumar H, Verma GR, Singh H. Surgical management of hepatic hydatid cysts - conservative versus radical surgery. *HPB (Oxford)* 2020; **22**: 1457-1462 [PMID: 32229090 DOI: 10.1016/j.hpb.2020.03.003]
- 43 **Zhang HZ**, Chen ZY, Yan LN, Zeng Y, Li B, Wen TF, Wang WT, Wu H, Yang JY. A Single-Center Experience on Surgical Treatment of Hepatic Echinococcosis. *Zhongguo Puwai Jichu Yu Linchuang Zazhi* 2015; **22**: 144-148 [DOI: 10.7507/1007-9424.20150040]
- 44 **Tan JZ**. Laparoscopic resection of hepatic hydatid cyst: report of 10 cases. *Zhonghua Yixue Zazhi* 1994; **74**: 208
- 45 **Abudusaimi A**, Wang SP, Ye EF, He J, Li K, Liu D, Azi G, Zhou L, Li SX, Keli M. Treating experience of laparoscopic surgery for hepatic cystic echinococcosis in children. *Zhonghua Ganzang Waiké Shoushuxue Dianzi Zazhi* 2019; **8**: 54-57 [DOI: 10.3877/cma.j.issn.2095-3232.2019.01.013]
- 46 **Tan Q**, Mu QM, Yang J, He W. Comparative study of clinical effect of three different laparoscopic surgical treatment on hepatic hydatidosis. *Zhonghua Puwaike Shoushuxue Zazhi* 2020; **14**: 186-188 [DOI: 10.3877/cma.j.issn.1674-3946.2020.02.023]
- 47 **Tekin A**, Küçükartallar T, Kartal A, Kaynak A, Ozer S, Tavli S, Belviranlı M, Sahin M, Yol S, Aksoy F, Tekin S, Vatansev C, Erikoglu M. Clinical and surgical profile and follow up of patients with liver hydatid cyst from an endemic region. *J Gastrointest Liver Dis* 2008; **17**: 33-37 [PMID: 18392241]
- 48 **Cairang Y**, Zhang L, Ren B, Ren L, Hou L, Wang H, Zhou Y, Zhang Q, Shao J, Fan H. Efficacy and safety of ultrasound-guided percutaneous microwave ablation for the treatment of hepatic alveolar echinococcosis: A preliminary study. *Medicine (Baltimore)* 2017; **96**: e7137 [PMID: 28682866 DOI: 10.1097/MD.00000000000007137]
- 49 **Lamonaca V**, Virga A, Minervini MI, Di Stefano R, Provenzano A, Tagliareni P, Fleres G, Luca A, Vizzini G, Palazzo U, Gridelli B. Cystic echinococcosis of the liver and lung treated by radiofrequency thermal ablation: an ex-vivo pilot experimental study in animal models. *World J Gastroenterol* 2009; **15**: 3232-3239 [PMID: 19598298 DOI: 10.3748/wjg.15.3232]
- 50 **Sarıçık B**, Kartal A, Esen H, Demircili ME. The Use of Radiofrequency Thermal Ablation Method in The Treatment of Hepatic Hydatid Cysts: Ex vivo Sheep Study. *Türkiye Parazitol Derg* 2019; **43**: 10-15 [PMID: 30938125 DOI: 10.4274/tpd.galenos.2019.5461]
- 51 **Guo JQ**, Hua GY, Zhang XM, Ye FY, Qi LN, Yang R. Preliminary clinical study of ultrasound-guided percutaneous liver puncture microwave ablation of hepatic alveolar echinococcosis. *Zhonghua Chaosheng Yingxiangxue Zazhi* 2017; **26**: 911-913 [DOI: 10.3760/cma.j.issn.1004-4477.2017.10.021]
- 52 **Botsa E**, Thanou I, Nikas I, Thanos L. Treatment of Hepatic Hydatid Cyst in a 7-Year-Old Boy Using a New Type of Radiofrequency Ablation Electrode. *Am J Case Rep* 2017; **18**: 953-958 [PMID: 28871077 DOI: 10.12659/ajcr.904432]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

