

World Journal of *Transplantation*

World J Transplant 2023 September 18; 13(5): 208-289



EVIDENCE REVIEW

- 208 Simultaneous pancreas-kidney transplantation for end-stage renal failure in type 1 diabetes mellitus: Current perspectives
Nagendra L, Fernandez CJ, Pappachan JM

REVIEW

- 221 Evolution of human kidney allograft pathology diagnostics through 30 years of the Banff classification process
Mubarak M, Raza A, Rashid R, Shakeel S

MINIREVIEWS

- 239 Sodium-glucose cotransporter-2 inhibitor use in kidney transplant recipients
Ramakrishnan P, Garg N, Pabich S, Mandelbrot DA, Swanson KJ
- 250 Renal allograft procurement from living unrelated donors in Iran: What falls under the eclipse
Taheri S
- 254 Impact of tacrolimus intra-patient variability in adverse outcomes after organ transplantation
Morais MC, Soares ME, Costa G, Guerra L, Vaz N, Codes L, Bittencourt PL

ORIGINAL ARTICLE**Case Control Study**

- 264 Invasive aspergillosis in liver transplant recipients, an infectious complication with low incidence but significant mortality
Farahani A, Ghiasvand F, Davoudi S, Ahmadinejad Z

Retrospective Cohort Study

- 276 Reasons and effects of the decline of willing related potential living kidney donors
Gadelkareem RA, Abdelgawad AM, Mohammed N, Reda A, Azoz NM, Zarzour MA, Hammouda HM, Khalil M

ABOUT COVER

Peer Reviewer of *World Journal of Transplantation*, Abbas Ghazanfar, MBBS, MRCS(Eng), FCPS, FHEA, FEBS, FRCS, FACS, Consultant Transplant Surgeon, Clinical Lead, Department of Renal and Transplant Surgery, St Georges University Hospitals NHS Foundation Trust, London SW17 0QT, United Kingdom. a.ghazanfar@nhs.net

AIMS AND SCOPE

The primary aim of *World Journal of Transplantation* (WJT, *World J Transplant*) is to provide scholars and readers from various fields of transplantation with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJT mainly publishes articles reporting research results obtained in the field of transplantation and covering a wide range of topics including bone transplantation, brain tissue transplantation, corneal transplantation, descemet stripping endothelial keratoplasty, fetal tissue transplantation, heart transplantation, kidney transplantation, liver transplantation, lung transplantation, pancreas transplantation, skin transplantation, etc.

INDEXING/ABSTRACTING

The WJT is now abstracted and indexed in PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The WJT's CiteScore for 2022 is 2.8 and Scopus CiteScore rank 2022: Transplantation is 23/51.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yan-Liang Zhang; **Production Department Director:** Xu Guo; **Editorial Office Director:** Jia-Ping Yan.

NAME OF JOURNAL

World Journal of Transplantation

ISSN

ISSN 2220-3230 (online)

LAUNCH DATE

December 24, 2011

FREQUENCY

Quarterly

EDITORS-IN-CHIEF

Maurizio Salvadori, Sami Akbulut, Vassilios Papalois, Atul C Mehta

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2220-3230/editorialboard.htm>

PUBLICATION DATE

September 18, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/gerinfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/gerinfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/gerinfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Impact of tacrolimus intra-patient variability in adverse outcomes after organ transplantation

Maria Clara Morais, Maria Eduarda Soares, Gabriela Costa, Laura Guerra, Nayana Vaz, Liana Codes, Paulo Lisboa Bittencourt

Specialty type: Transplantation

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): D

Grade E (Poor): 0

P-Reviewer: Ali A, Iraq; Tanaka H, Japan

Received: June 28, 2023

Peer-review started: June 28, 2023

First decision: July 19, 2023

Revised: July 31, 2023

Accepted: August 11, 2023

Article in press: August 11, 2023

Published online: September 18, 2023



Maria Clara Morais, Gabriela Costa, Laura Guerra, Nayana Vaz, Liana Codes, Paulo Lisboa Bittencourt, School of Medicine, Bahiana School of Medicine and Public Health, Salvador 40290-000, Bahia, Brazil

Maria Eduarda Soares, School of Medicine, Federal University of Bahia, Salvador 40110-100, Bahia, Brazil

Nayana Vaz, Liana Codes, Paulo Lisboa Bittencourt, Unit of Gastroenterology and Hepatology, Portuguese Hospital, Salvador 40130-030, Bahia, Brazil

Corresponding author: Paulo Lisboa Bittencourt, MD, PhD, Chief Doctor, Unit of Gastroenterology and Hepatology, Portuguese Hospital, 914 Av. Princesa Isabel, Barra, Salvador 40130-030, Bahia, Brazil. plbbr@uol.com.br

Abstract

Tacrolimus (Tac) is currently the most common calcineurin-inhibitor employed in solid organ transplantation. High intra-patient variability (IPV) of Tac (Tac IPV) has been associated with an increased risk of immune-mediated rejection and poor outcomes after kidney transplantation. Few data are available concerning the impact of high Tac IPV in non-kidney transplants. However, even in kidney transplantation, there is still a controversy whether high Tac IPV is indeed detrimental in respect to graft and/or patient survival. This may be due to different methods employed to evaluate IPV and distinct time frames adopted to assess graft and patient survival in those reports published up to now in the literature. Little is also known about the influence of high Tac IPV in the development of other untoward adverse events, update of the current knowledge regarding the impact of Tac IPV in different outcomes following kidney, liver, heart, lung, and pancreas transplantation to better evaluate its use in clinical practice.

Key Words: Tacrolimus; Intra-patient variability; Rejection; Organ transplantation; Graft survival; Outcomes

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Tacrolimus is widely used after solid organ transplantation. High intra-patient variability of tacrolimus (Tac IPV) has been associated with poor graft and patient survival. This review summarizes current evidence regarding the impact of high Tac IPV in several outcomes after kidney, liver, heart, lung and pancreas transplantation.

Citation: Morais MC, Soares ME, Costa G, Guerra L, Vaz N, Codes L, Bittencourt PL. Impact of tacrolimus intra-patient variability in adverse outcomes after organ transplantation. *World J Transplant* 2023; 13(5): 254-263

URL: <https://www.wjgnet.com/2220-3230/full/v13/i5/254.htm>

DOI: <https://dx.doi.org/10.5500/wjt.v13.i5.254>

INTRODUCTION

Tacrolimus (Tac) is nowadays the most common immunosuppressive drug employed in solid organ transplantation and has replaced cyclosporin as the most common calcineurin inhibitor used worldwide in immunosuppressive regimens[1, 2]. Tac exhibits a peculiar pharmacokinetic profile with large inter-patient and intra-patient variability (IPV) of whole blood drug levels over time, even when doses remain unchanged. This is usually ascribed to overlapping factors, such as ethnicity, pharmacogenomics, food-drug and drug-drug interactions, non-adherence, enhanced Tac absorption or impaired drug excretion due to either diarrhea or cholestasis, assays variability for Tac levels determinations or even alternate use of Tac compounds or its generic formulations[3-5]. Due to its narrow therapeutic window, therapeutic drug monitoring of trough levels (Cmin) is required to attain target levels of immunosuppression over time, as well as avoidance of undesired Tac side effects such as infections, neurotoxicity and nephrotoxicity, which are usually related to higher drug exposure[1].

High Tac IPV or time in the therapeutic range (TTR) of Tac has been associated with an increased risk of immune-mediated rejection and poor outcomes after kidney transplantation (KT)[5,6]. Few data are available concerning the impact of high Tac IPV in non-kidney transplants[7]. However, even in KT, there is still a controversy whether high Tac IPV is indeed detrimental in respect to graft and/or patient survival[5,6]. This may be due to different methods employed to evaluate IPV and distinct time frames adopted to assess graft and patient survival in the reports published up to now. Little is also known about the influence of high Tac IPV in the development of other harmful adverse events associated with immunosuppression, such as infections, chronic kidney disease, type 2 diabetes, metabolic syndrome and *de novo* or recurrent cancer. The purpose of this review is to provide an update of current knowledge regarding the impact of Tac IPV in different outcomes following KT, liver (LT), heart (HT), lung, kidney/pancreas and bone marrow (BMT) transplantation to better evaluate its use in clinical practice.

STUDY SELECTION

To identify and select studies for this review, research was made in the MedLine/PubMed database, using the following terms: Tacrolimus; intra-patient; variability; transplant; transplantation; rejection; and graft loss. These terms were obtained from Medical Subject Headings (MeSH), using the Booleans "AND" and "OR", from various search algorithms in PubMed.

Search 1: ("tacrolimus"[MeSH Terms] OR "tacrolimus"[All Fields]) AND "intra"[All Fields] AND ("patient s"[All Fields] OR "patients"[MeSH Terms] OR "patients"[All Fields] OR "patient"[All Fields] OR "patients s"[All Fields]) AND ("variabilities"[All Fields] OR "variability"[All Fields] OR "variable"[All Fields] OR "variable s"[All Fields] OR "variables"[All Fields] OR "variably"[All Fields]) AND (y_10[Filter]). Search 2: ((((((tacrolimus[Title/Abstract]) AND (variability[Title/Abstract])) OR (inpatient variability[Title/Abstract])) AND (transplant[Title/Abstract])) OR (transplantation[Title/Abstract])) AND (rejection[Title/Abstract])) OR (graft loss[Title/Abstract])).

Subsequently, an active manual search of studies was carried out through carefully selected articles in gray literature.

The research was completed on May 17th, 2023. A total of 43 articles were included in the present study, 376 studies were found in PubMed database, and after careful reading of abstracts, 15 were included in our final review (references No. 3, 7, 24, 31, 33, 37, 38, 57, 60, 62, 65-67, 69, and 70). The excluded works did not satisfactorily meet the theme proposed for the present work. Furthermore, 28 studies different from the ones previously selected in PubMed database were identified in our active manual research in gray literature, all of which were included in this review.

METHODS USED TO ASSESS TAC IPV

Intra-patient variability of Tac over time has been calculated using different methods, particularly[5-7]: Mean levels and standard deviation (SD) of Tac whole-blood Cmin levels, also expressed as the medication level variability index (MLVI).

Mean absolute deviation (MAD), based on the formula $MAD (\%) = \{[(X_{mean} - X_1) + (X_{mean} - X_2) \dots + (X_{mean} - X_n)]/n$ of Cmin results}, where X is the Tac Cmin level.

Coefficient of variation (CV), calculated according to the formula $CV (\%) = \sigma/\mu$, where σ is the standard deviation, and μ is the mean Tac Cmin levels of all available samples of the individual. CV may be corrected or not by the corresponding Tac dose (C0/D) or defined as time-weighted coefficient of variability using time-weighted standard deviation divided by the mean drug levels.

Tac TTR, calculated by the Rosendaal method[8]. Standard deviation or MLVI are expressed as numbers, classes, or dichotomized intervals, whereas CV and TTR are expressed as percentages, tertiles and dichotomized intervals usually at the median split. There are no universally accepted recommended target levels for each one of those parameters used to assess Tac IPV. It is important to highlight that target levels of Tac Cmin usually vary over time according to the type of organ transplant, donor and recipient risk factors for rejection, comorbidity, occurrence of side effects and local immunosuppression protocols. The TTR may also vary sharply according to adopted Tac Cmin thresholds used by distinct centers in different time frames after organ transplantation[5-7].

FACTORS INFLUENCING TAC IPV

Interpatient variability of Tac has been attributed to interindividual pharmacokinetics variability which may be induced by several factors including drug-food and drug-drug interactions, concurrent clinical events such as diarrhea, cholestasis or liver dysfunction, ethnicity and pharmacogenetics[3,9-13]. In this regard, polymorphisms in CYP3A5, CYP3A4, and SLC and ABC transporter encoding genes have been shown to influence the area under the curve of tacrolimus, leading either to rejection or even toxicity in transplant recipients[4,13].

Tac IPV, on the other hand, has been additionally ascribed to non-adherence after organ transplantation, pre-analytical and analytical variables to measure Cmin in different commercially available biochemical assays and administration of different Tac formulations, including generic substitutions[3,5-7]. However, non-adherence is a common reason reported in medical literature to explain the observed deleterious effect of Tac IPV on patient and graft survival in organ transplantation by some[14-19], but not all authors[20,21].

Several approaches have been proposed to assess IPV, with most studies being retrospective, with different methodologies, considering pediatric and adult populations, with no organ-specific approach. Despite these limitations, IPV provides a sign that patients are at risk.

TAC IPV IN HEART TRANSPLANTATION

Four studies evaluating the influence of Tac IPV on HT outcomes were performed in adult patients using Tac CV[22-24] and Tac TTR[25], and in pediatric subjects using Tac SD/MLVI[26,27].

Concerning pediatric LT, Pollock-Barziv *et al*[26] associated Tac IPV with late rejection as well as worse patient and graft survival, but it is worth to mention that few heart transplant recipients were included in this study. Sirota *et al*[27] found similar results for rejection, but the authors also linked Tac SD/MLVI, particularly when higher than 3, to cardiac allograft vasculopathy and patient survival. In adults, Gueta *et al*[22] described an association of Tac IPV in the first year after HT with rejection after 12 but not between three and 12 mo after HT. On the other hand, Shuker *et al*[23] found no increase in the frequency of either early or late acute rejection or even cardiac allograft vasculopathy in heart transplant recipients. Both authors ascribed the lack of association of Tac IPV with early acute rejection[20] and overall rejection[23] to a higher immunosuppression exposure frequently observed in those heart transplant recipients. González-Vílchez *et al*[24] studied the largest cohort up to now. The authors found an effect of Tac IPV, assessed between four to 12 mo after HT, on the frequency of rejection after one year. Using Tac TTR, Baker *et al*[25] found no effect of Tac IPV on acute rejection within the first 30 d after HT. None of those studies in adults reported other outcomes such as mortality or Tac-related adverse events in association with Tac IPV[22-25]. The main findings related to HT are depicted in Table 1[22-27].

TAC IPV IN LUNG TRANSPLANTATION

Four studies have investigated the impact of Tac IPV in the frequency of acute and chronic rejection or chronic lung allograft dysfunction (CLAD) after lung transplantation in adults using Tac SD/MLVI[28], Tac CV[29] and Tac TTR[29-31]. Gallagher *et al*[28] demonstrated that higher Tac SD/MLVI between 6-12 mo after lung transplantation was associated with development of CLAD with an adverse impact on survival. Acute rejection, on the other hand was not associated with Tac SD/MLVI[28]. Ensor *et al*[30] evaluated retrospectively the role of Tac TTR in 292 Lung transplant recipients the development of acute and chronic rejection. Tac TTR was measured in the first year after lung transplantation. The authors observed a lower likelihood for acute rejection and CLAD whenever Tac TTR was increased by 10%. Lower rates of infection and mortality was also linked to high Tac TTR. Other authors failed to demonstrate higher frequency of acute rejection after lung transplantation when Tac IPV was assessed by either Tac CV or Tac TTR in the first 6 mo after surgery[29]. On the other hand, more recently Japanese investigators disclosed an association of acute rejection with Tac TTR calculated in the first six month after lung transplantation[31]. The main findings related to lung transplantation are depicted in Table 2.

Table 1 Tacrolimus intra-patient variability in heart transplantation: Main findings

Heart transplantation				
Ref.	Sample size	Donor type	Tac-IPV, assessment	Outcome
Gueta <i>et al</i> [22], 2018	72	Deceased	CV	High trough level variability is associated with higher rates of graft rejection, and trough level variability during the first year is associated with increased risk of rejection after HT
Shuker <i>et al</i> [23], 2018	86	Deceased	MAD	A high IPV was not associated with the development and progression of cardiac allograft vasculopathy or development of acute cellular rejection
González-Vílchez <i>et al</i> [24], 2022	1581	Deceased	CV	IPV levels had limited influence on mid-term outcomes in heart transplant, however high IPV may predispose to rejection in initially stable patients
Baker <i>et al</i> [25], 2019	67	Deceased	TTR	Higher TTR was not associated with a lower rate of Acute Cellular Rejection within the first 30 d after heart transplant
Pollock-Barziv <i>et al</i> [26], 2010	144	Deceased	SD	Associated Tac IPV with late rejection as well as worse patient and graft survival, but it is worth to mention that few heart transplant recipients were included in this study
Sirota <i>et al</i> [27], 2021	118	Deceased	SD	SD ≥ 3 is associated with increased risk of poor outcomes

CV: Coefficient of variability; HT: Heart transplant; MAD: Mean absolute deviation; TTR: Time in therapeutic range; SD: Standard deviation.

Table 2 Tacrolimus intra-patient variability in lung transplantation: main findings

Lung transplantation				
Ref.	Sample size	Donor type	Tac-IPV assessment	Outcome
Gallagher <i>et al</i> [28], 2015	110	Non specified	SD	Patients with highly variable trough tacrolimus levels in the second half of the first post-transplant year will likely have similar variability in the second year and are at high risk for subsequent chronic lung allograft dysfunction and death
Kao <i>et al</i> [29], 2021	157	Non specified	CV and TTR	The results suggest that tacrolimus TTR, time in therapeutic range, and variability are not related to the presence of ACR in LTRs
Ensor <i>et al</i> [30], 2018	292	Non specified	TTR	Tacrolimus TTR was predictive of clinical outcomes of ACR, CLAD, infection, and death in lung transplant recipients at 1 yr in this investigation after adjusting for potential confounders
Katada <i>et al</i> [31], 2022	90	Living and deceased	TTR	A lower tacrolimus TTR is a predictor of late acute rejection

SD: Standard deviation; TTR: Time in therapeutic range; ACR: Acute cellular rejection; LTR: Lung transplant rejection; CLAD: Chronic lung allograft dysfunction.

TAC IPV IN LIVER TRANSPLANTATION

Impact of Tac IPV on different outcomes after LT, including allograft rejection, postoperative complications and survival was evaluated in pediatric[26,32-35] and adult[36-42] transplant recipients using SD/MLVI[26,32-34], CV[35-41], and TTR[42].

Most studies performed in children associated higher SD/MLVI to biopsy-proven acute rejection (BPAR) at least six months after LT[42-34]. Defrancq *et al*[35] found similar results in another cohort of children using Tac CV one year after surgery. The authors reported an association between high Tac CV and BPAR and correlated Tac CV with albumin and bilirubin levels in different time frames after LT as well as with missed outpatient consultations, possibly reflecting immunosuppression adherence[33]. Decreased survival was also related to high Tac IPV in just one of those reports including few patients submitted to LT[26].

Up to now, seven reports were published in the medical literature concerning the effect of Tac IPV on post-LT outcomes in adults. In this regard, Christina *et al*[36] disclosed an association between Tac SD/MLVI and BPAR in patients submitted to LT. Similar findings were reported by Del Bello *et al*[37] who reported higher Tac CV measured just after LT discharge with BPAR, as well as with the occurrence of *de novo* anti-donor specific antibodies (*dn*DSA). No impact was seen in patient survival[37]. Van der Veer *et al*[39] evaluated the influence of Tac CV measured between six to 18 mo after LT in adult subjects. The authors were unable to disclose any association between Tac CV and immune mediated graft injury. Rayar *et al*[38] investigated the influence of Tac CV measured in the first 30 d after LT on postoperative outcomes. The authors found an increased frequency of acute kidney injury as well as cardiovascular and

Table 3 Tacrolimus intra-patient variability and liver transplantation: Main findings

Liver transplantation				
Ref.	Sample size	Donor	Tac-IPV assessment	Outcome
Lieber <i>et al</i> [18], 2013	988	Not specified	SD	Non-adherence among liver transplant recipients is associated with increased risk of graft failure
Stuber <i>et al</i> [19], 2008	96	Not specified	SD	The SD has utility of monitoring routine tac blood levels in pediatric recipients for detecting non-adherence prior to clinical rejection
Venkat <i>et al</i> [32], 2008	101	Not specified	SD	Variations in tac blood levels is associated with an increased risk of late allograft rejection in pediatric recipients
Shemesh <i>et al</i> [33], 2017	400	Both living and deceased donor	SD; MLVI	MLVI predicts late acute rejection in pediatric liver transplantation recipients
de Oliveira <i>et al</i> [34], 2017	50	Both living and deceased donor	SD; MLVI	MLVI may be a nice indicator of the risk of medication non-adherence in childhood
Defrancq <i>et al</i> [35], 2019	41	Both living and deceased donor	CV	High Tac IPV may be associated with adverse patient outcomes. Also, there is some impact of biological factors on IPV and therapy adherence
Christina <i>et al</i> [36], 2014	150	Not specified	SD; MLVI	The MLVI is associated with and can predict rejection, possibly related to non-adherence in adult recipients
Del Bello <i>et al</i> [37], 2018	116	Deceased donor only	CV	Tac IPV could be useful to identify patients with a greater risk of graft rejection and pf developing <i>de novo</i> DSA after liver transplantation
Rayar <i>et al</i> [38], 2018	812	Deceased donor only	CV	High CV of Tac concentrations was found to be predictive of Tac-related toxicity and poorer survival
van der Veer <i>et al</i> [39], 2019	326	Both living and deceased donor	CV	High IPV in Tac exposure beyond 6 mo after liver transplantation was not associated with immune-mediated graft injury
Dopazo <i>et al</i> [40], 2022	140	Deceased donor only	CV	High IPV between the third and sixth months appears to be an early and independent predictor of poorer liver transplant outcomes
Kim <i>et al</i> [41], 2022	636	Both living and deceased donor	CV	High Tac IPV was associated with increased risks of overall mortality and HCC recurrence in liver transplantation recipients with HCC

SD: Standard deviation; CV: Coefficient of variation; MLVI: Medication level variability index; DSA: Donor-specific antibodies; HCC: Hepatocellular carcinoma.

neurologic complications in patients with higher Tac CV. Most importantly, shortened graft and patient survival rates were also associated with Tac IPV[38]. Other authors investigated whether Tac CV, assessed between three to six months after LT, could be associated with worse graft and patient long-term outcomes[40]. In this study, lower long-term survival and poorer renal function were similarly observed in those subjects with high Tac CV. Another group of investigators also linked higher Tac IPV using CV with hepatocellular carcinoma (HCC) recurrence after LT[41]. Only one study up to now assessed the impact of Tac IPV after LT employing TTR[42]. The authors found that lower TTR in those subjects, irrespectively of Tac CV, was associated with a higher risk for *dn*DSA and long-term death-censored graft loss. The main findings related to LT are depicted in Table 3.

TAC IPV IN KIDNEY TRANSPLANTATION

Most of the studies published thus far demonstrated an adverse impact of Tac IPV on KT outcomes (Table 4)[43-68]. Most of them were performed in adult recipients using MLVI/SD[44], MAD[45-48,62,66], CV[49-60,63,64,67-68], TTR[61] or other methods[65]. In this regard, several authors have associated higher Tac IPV, usually measured more than 6 mo after KT, with BPAR[44-46], long-term graft loss or dysfunction[47-55,58,59,68] as well as with lower survival[50,51]. In some, but not all reports[62], higher Tac IPV was associated with the development of *dn*DSA[53,67], chronic active antibody mediated rejection[47] and chronic histological lesions in kidney grafts[54,56]. Most authors attributed those Tac IPV-related outcomes to non-adherence[43] and some[57-59] but not others[60] to genetic predisposition. Two other reports compared TTR[61] and a novel Tac variability score (TVS)[64] to conventional IPV measures and found that both were more reliable in their ability to predict worse outcomes after KT.

Concerning pediatric transplant recipients, Pollock-Barziv *et al*[26] were one of the first authors to show a significant association of higher Tac IPV and late graft rejection or loss. Similar findings were subsequently reported by other investigators who reported and association of higher Tac IPV with BPAR and graft loss beyond one year in children and adolescents submitted to KT[66-68].

Table 4 Tacrolimus intra-patient variability and kidney transplantation: Main findings

Kidney transplantation				
Ref.	Sample size	Donor type	Tac-IPV assessment	Outcomes
Borra <i>et al</i> [47], 2010	297	Both living and deceased donor	MAD	Significant relationship between high Tac-IPV and long-term graft failure
Ro <i>et al</i> [45], 2012	249	Both living and deceased donor	MAD	TAC IPV had a significant impact on rejection-free survival. The effect was influenced by CYP3A5 polymorphism
Sapir-Pichhadze <i>et al</i> [44], 2014	356	Both living and deceased donor	MLVI/SD	Increased time-dependent TAC SD may be an independent risk factor for adverse kidney transplant outcomes
O'Regan <i>et al</i> [49], 2016	394	Both living and deceased donor	CV	Inferior renal allograft survival was observed in recipients with higher Tac-IPV
Rodrigo <i>et al</i> [53], 2016	310	Deceased donor only	CV	Tacrolimus level variability is a strong risk factor for dnDSA development and death-censored graft loss
Whalen <i>et al</i> [46], 2017	376	Both living and deceased donor	MAD	Highly variable tacrolimus levels predict worse outcomes postrenal transplantation
Shuker <i>et al</i> [48], 2016	808	Both living and deceased donor	MAD	A high tacrolimus IPV is an independent risk factor for adverse kidney transplant outcomes that can be used as an easy monitoring tool to help identify high-risk RTRs
Vanhove <i>et al</i> [56], 2016	220	Both living and deceased donor	CV	High IPV is related to accelerated progression of chronic histologic lesions before any evidence of renal dysfunction
Rozen-Zvi <i>et al</i> [51], 2017	803	Both living and deceased donor	CV	The combination of high CV and exposure to low drug levels might identify high-risk patients in the early post-transplantation period
Goodall <i>et al</i> [50], 2017	688	Both living and deceased donor	CV	High tacrolimus IPV and clinic nonattendance are associated with inferior allograft survival
Sablik <i>et al</i> [62], 2018	248	Both living and deceased donor	MAD	A high Tac IPV per se does not predispose to the development of chronic active antibody mediated rejection (c-aABMR) but is associated with inferior graft survival once c-aABMR is diagnosed
Seibert <i>et al</i> [57], 2018	1472	Both living and deceased donor	CV	High variability of TAC dose increases risk of acute rejection. High variability of TAC trough increases risk of graft failure
Mo <i>et al</i> [54], 2019	671	Both living and deceased donor	CV	High IPV of Tac is associated with early deterioration of chronic histologic lesions as well as poorer long-term outcomes
Song <i>et al</i> [61], 2019	1241	Living donor only	TTR	Increasing the TTR of tacrolimus in the first year was associated with improved long-term outcomes in living kidney transplants, and TTR may be a novel valuable strategy to monitor tacrolimus exposure
Süsal <i>et al</i> [55], 2019	6638	Deceased donor only	CV	Even in patients with good outcome during the first 3 post-transplant years, a high IPV was associated with inferior graft survival, indicating that a fluctuating tacrolimus trough level at years 1, 2 and 3 post-transplant is a major problem in kidney transplantation
Rahamimov <i>et al</i> [52], 2019	878	Both living and deceased donor	CV	Monitoring CV can help detect the high-risk patients
Gold <i>et al</i> [66], 2020	1419	Deceased donor only	MAD	A more intense and less variable exposure to tacrolimus could improve graft survival strongly in patients with high TAC IPV
Stefanović <i>et al</i> [58], 2020	104	Both living and deceased donor	CV	Combined assessment of tacrolimus IPV and tacrolimus C0/D may categorize patients towards risk of graft deterioration in the long-term post-transplantation period
Stefanović <i>et al</i> [59], 2021	103	Both living and deceased donor	CV	Simultaneous assessment of Tac IPV, C0/D, and CYP3A5 genotype may identify patients at risk of deterioration of graft function in the long-term post-transplantation period
Kim <i>et al</i> [65], 2021	1080	Both living and deceased donor	CV	High tacrolimus IPV significantly increases the risk of graft failure and antibody mediated rejection in patients with high immunological risk
Park <i>et al</i> [63], 2021	1143	Both living and deceased donor	CV	TAC-IPV can significantly affect allograft outcomes even with a high mean TAC-C0
Yin <i>et al</i> [64], 2022	1343	Living donor only	CV	Tac variability score is a novel measure of Tac IPV with higher correlation with graft survival and more convenience in clinical use than CV after kidney transplantation
Baghai Arassi <i>et al</i> [67], 2022	48	Both living and deceased donor	CV	High Tac IPV is associated with an increased risk of dnDSA development and rejection episodes > year 1 posttransplant even in patients with low immunological risk profile

Nuchjumroon <i>et al</i> 188 [60], 2022	Both living and deceased donor	CV	No evidence that the CYP3A5 polymorphisms significantly influence tacrolimus IPV during the 6 to 12 mo after kidney transplantation
---	--------------------------------	----	---

SD: Standard deviation; CV: Coefficient of variation; MLVI: Medication level variability index; DSA: Donor-specific antibodies; TTR: Time in therapeutic range.

TAC IPV IN KIDNEY AND PANCREAS, AND BONE MARROW TRANSPLANTATION

Two studies have evaluated the impact of Tac IPV in the frequency of *dn*DSA, BPAR and graft loss in adult kidney and pancreas recipients[69,70] and in the occurrence of graft *vs* host disease after bone marrow transplantation[70]. The first report failed to disclose an association of Tac IPV with BPAR in adult recipients of kidney and pancreas transplants, but the main purpose of the study was to compare Tac IPV and graft function in groups of patients receiving two different Tac formulations[69]. Davis *et al*[38] on the other hand, demonstrated that Tac TTR was associated to a very high risk of *dn*DSA and a 4-fold risk of graft loss by five years, independently of Tac CV.

In respect to bone marrow transplant recipients, Marco *et al*[70] correlated high Tac IPV, measured by CV, with the occurrence of acute graft *vs* host disease in the first month after BMT (Table 5).

CONCLUSION

Altogether, available data up to now suggest that Tac IPV, possibly due to non-adherence and/or genetic, pharmacologic, or clinically significant factors, is associated with adverse outcomes after organ transplantation, particularly KT. The heterogeneity observed in the results obtained in the reports thus far are probably due to the retrospective design of most studies, the distinct methods used to assess Tac IPV, utilization of different immunosuppression protocols, distinct observation time frames and endpoints. Refinement or combination of different scores may improve usage of Tac IPV in clinical practice in the future.

Table 5 Tacrolimus intra-patient variability in pancreas and bone marrow transplantation: Main findings

Kidney and pancreas, and bone marrow transplant				
Ref.	Sample size	Donor type	Tac-IPV Assessment	Outcome
Torabi <i>et al</i> [69], 2020	39	Both living and deceased donor	CV	The once daily LCPT dosing may facilitate medication adherence and result in improved long-term outcomes
Marco <i>et al</i> [70], 2022	128	Living donor only	CV	Determination of Tac IPV soon after alloHSCT could be useful in identifying greater risks of aGVHD

CV: Coefficient of variation; LCPT: LCP-tacrolimus; alloHSCT: Allogeneic stem cell transplantation; aGVHD: Acute graft-versus-host disease.

FOOTNOTES

Author contributions: Morais MC, Codes L, and Bittencourt PL contributed to concept and design; Morais MC, Costa G, Guerra L, Vaz N, Codes L, and Bittencourt PL contributed to drafting of the manuscript; and all authors have read and approved the final version of the manuscript.

Conflict-of-interest statement: Authors declare no conflict of interests for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Brazil

ORCID number: Maria Clara Morais 0000-0003-3752-584X; Maria Eduarda Soares 0009-0009-3762-8161; Gabriela Costa 0009-0009-3041-8436; Laura Guerra 0009-0000-4145-9019; Nayana Vaz 0000-0002-6766-0164; Liana Codes 0000-0001-5178-8705; Paulo Lisboa Bittencourt 0000-0003-0883-4870.

S-Editor: Chen YL

L-Editor: A

P-Editor: Chen YL

REFERENCES

- 1 **Ong SC**, Gaston RS. Thirty Years of Tacrolimus in Clinical Practice. *Transplantation* 2021; **105**: 484-495 [PMID: [32541562](#) DOI: [10.1097/TP.0000000000003350](#)]
- 2 **Staatz CE**, Tett SE. Clinical pharmacokinetics and pharmacodynamics of tacrolimus in solid organ transplantation. *Clin Pharmacokinet* 2004; **43**: 623-653 [PMID: [15244495](#) DOI: [10.2165/00003088-200443100-00001](#)]
- 3 **Shuker N**, van Gelder T, Hesselink DA. Intra-patient variability in tacrolimus exposure: causes, consequences for clinical management. *Transplant Rev (Orlando)* 2015; **29**: 78-84 [PMID: [25687818](#) DOI: [10.1016/j.trre.2015.01.002](#)]
- 4 **Brunet M**, Pastor-Anglada M. Insights into the Pharmacogenetics of Tacrolimus Pharmacokinetics and Pharmacodynamics. *Pharmaceutics* 2022; **14** [PMID: [36145503](#) DOI: [10.3390/pharmaceutics14091755](#)]
- 5 **Kuypers DRJ**. Inpatient Variability of Tacrolimus Exposure in Solid Organ Transplantation: A Novel Marker for Clinical Outcome. *Clin Pharmacol Ther* 2020; **107**: 347-358 [PMID: [31449663](#) DOI: [10.1002/cpt.1618](#)]
- 6 **Schumacher L**, Leino AD, Park JM. Tacrolimus inpatient variability in solid organ transplantation: A multiorgan perspective. *Pharmacotherapy* 2021; **41**: 103-118 [PMID: [33131078](#) DOI: [10.1002/phar.2480](#)]
- 7 **Coste G**, Lemaître F. The Role of Intra-Patient Variability of Tacrolimus Drug Concentrations in Solid Organ Transplantation: A Focus on Liver, Heart, Lung and Pancreas. *Pharmaceutics* 2022; **14** [PMID: [35214111](#) DOI: [10.3390/pharmaceutics14020379](#)]
- 8 **Rosendaal FR**, Cannegieter SC, van der Meer FJ, Briët E. A method to determine the optimal intensity of oral anticoagulant therapy. *Thromb Haemost* 1993; **69**: 236-239 [PMID: [8470047](#)]
- 9 **Giza P**, Fieck R, Dwulit T, Chudek J, Woźniak I, Więcek A, Kolonko A. Number of Regularly Prescribed Drugs and Inpatient Tacrolimus Trough Levels Variability in Stable Kidney Transplant Recipients. *J Clin Med* 2020; **9** [PMID: [32575525](#) DOI: [10.3390/jcm9061926](#)]
- 10 **Jacobo-Cabral CO**, García-Roca P, Romero-Tejeda EM, Reyes H, Medeiros M, Castañeda-Hernández G, Trocóniz IF. Population pharmacokinetic analysis of tacrolimus in Mexican paediatric renal transplant patients: role of CYP3A5 genotype and formulation. *Br J Clin Pharmacol* 2015; **80**: 630-641 [PMID: [25846845](#) DOI: [10.1111/bcp.12649](#)]
- 11 **Taber DJ**, Su Z, Fleming JN, McGillicuddy JW, Posadas-Salas MA, Treiber FA, Dubay D, Srinivas TR, Mauldin PD, Moran WP, Baliga PK. Tacrolimus Trough Concentration Variability and Disparities in African American Kidney Transplantation. *Transplantation* 2017; **101**: 2931-2938 [PMID: [28658199](#) DOI: [10.1097/TP.0000000000001840](#)]
- 12 **Campagne O**, Mager DE, Tornatore KM. Population Pharmacokinetics of Tacrolimus in Transplant Recipients: What Did We Learn About Sources of Interindividual Variabilities? *J Clin Pharmacol* 2019; **59**: 309-325 [PMID: [30371942](#) DOI: [10.1002/jcph.1325](#)]
- 13 **Tron C**, Lemaître F, Verstuyft C, Petitcollin A, Verdier MC, Bellissant E. Pharmacogenetics of Membrane Transporters of Tacrolimus in Solid Organ Transplantation. *Clin Pharmacokinet* 2019; **58**: 593-613 [PMID: [30415459](#) DOI: [10.1007/s40262-018-0717-7](#)]
- 14 **Shemesh E**, Fine RN. Is calculating the standard deviation of tacrolimus blood levels the new gold standard for evaluating non-adherence to medications in transplant recipients? *Pediatr Transplant* 2010; **14**: 940-943 [PMID: [20887400](#) DOI: [10.1111/j.1399-3046.2010.01396.x](#)]
- 15 **Herblum J**, Dacouris N, Huang M, Zaltzman J, Prasad GVR, Nash M, Chen L. Retrospective Analysis of Tacrolimus Inpatient Variability as a Measure of Medication Adherence. *Can J Kidney Health Dis* 2021; **8**: 20543581211021742 [PMID: [34188946](#) DOI: [10.1177/20543581211021742](#)]
- 16 **Gokoel SRM**, Zwart TC, Moes DJAR, van der Boog PJM, de Fijter JW. No Apparent Influence of Nonadherence on Tacrolimus Inpatient Variability in Stable Kidney Transplant Recipients. *Ther Drug Monit* 2020; **42**: 702-709 [PMID: [32941396](#) DOI: [10.1097/FTD.0000000000000772](#)]
- 17 **Kostalova B**, Mala-Ladova K, Sulkova SD, Denhaerynck K, De Geest S, Maly J. Comparison of different methods to assess tacrolimus concentration intra-patient variability as potential marker of medication non-adherence. *Front Pharmacol* 2022; **13**: 973564 [PMID: [36313323](#) DOI: [10.3389/fphar.2022.973564](#)]
- 18 **Lieber SR**, Volk ML. Non-adherence and graft failure in adult liver transplant recipients. *Dig Dis Sci* 2013; **58**: 824-834 [PMID: [23053889](#) DOI: [10.1007/s10620-012-2412-0](#)]
- 19 **Stuber ML**, Shemesh E, Seacord D, Washington J 3rd, Hellemann G, McDiarmid S. Evaluating non-adherence to immunosuppressant medications in pediatric liver transplant recipients. *Pediatr Transplant* 2008; **12**: 284-288 [PMID: [18331387](#) DOI: [10.1111/j.1399-3046.2008.00923.x](#)]
- 20 **Leino AD**, King EC, Jiang W, Vinks AA, Klawitter J, Christians U, Woodle ES, Alloway RR, Rohan JM. Assessment of tacrolimus inpatient variability in stable adherent transplant recipients: Establishing baseline values. *Am J Transplant* 2019; **19**: 1410-1420 [PMID: [30506623](#) DOI: [10.1111/ajt.15199](#)]
- 21 **Ko H**, Kim HK, Chung C, Han A, Min SK, Ha J, Min S. Association between medication adherence and inpatient variability in tacrolimus concentration among stable kidney transplant recipients. *Sci Rep* 2021; **11**: 5397 [PMID: [33686160](#) DOI: [10.1038/s41598-021-84868-5](#)]
- 22 **Gueta I**, Markovits N, Yarden-Bilavsky H, Raichlin E, Freimark D, Lavee J, Loebstein R, Peled Y. High tacrolimus trough level variability is associated with rejections after heart transplant. *Am J Transplant* 2018; **18**: 2571-2578 [PMID: [29989311](#) DOI: [10.1111/ajt.15016](#)]
- 23 **Shuker N**, Bouamar R, Hesselink DA, van Gelder T, Caliskan K, Manintveld OC, Balk AH, Constantinescu AA. Inpatient Variability in Tacrolimus Exposure Does Not Predict The Development of Cardiac Allograft Vasculopathy After Heart Transplant. *Exp Clin Transplant* 2018; **16**: 326-332 [PMID: [28969528](#) DOI: [10.6002/ect.2016.0366](#)]
- 24 **González-Vilchez F**, Crespo-Leiro MG, Delgado-Jiménez J, Pérez-Villa F, Segovia-Cubero J, Díaz-Molina B, Mirabet-Pérez S, Arizón Del Prado JM, Blasco-Peiró T, Martínez-Sellés M, Almenar-Bonet L, Garrido-Bravo I, Rábago G, Vázquez de Prada JA. Impact of inpatient blood level variability of calcineurin inhibitors on heart transplant outcomes. *Rev Esp Cardiol (Engl Ed)* 2022; **75**: 129-140 [PMID: [33744197](#) DOI: [10.1016/j.rec.2021.02.001](#)]
- 25 **Baker WL**, Steiger S, Martin S, Patel N, Radojevic J, Darsaklis K, O'Bara L, Kutzler H, Dougherty J, Feingold A, Hammond J, Fusco D, Gluck JA. Association Between Time-in-Therapeutic Tacrolimus Range and Early Rejection After Heart Transplant. *Pharmacotherapy* 2019; **39**: 609-613 [PMID: [30892740](#) DOI: [10.1002/phar.2262](#)]
- 26 **Pollock-Barziv SM**, Finkelstein Y, Manlihot C, Dipchand AI, Hebert D, Ng VL, Solomon M, McCrindle BW, Grant D. Variability in tacrolimus blood levels increases the risk of late rejection and graft loss after solid organ transplantation in older children. *Pediatr Transplant* 2010; **14**: 968-975 [PMID: [21040278](#) DOI: [10.1111/j.1399-3046.2010.01409.x](#)]
- 27 **Sirota M**, Heyrend C, Ou Z, Masotti S, Griffiths E, Molina K. Impact of tacrolimus variability on pediatric heart transplant outcomes. *Pediatr Transplant* 2021; **25**: e14043 [PMID: [34390091](#) DOI: [10.1111/petr.14043](#)]

- 28 **Gallagher HM**, Sarwar G, Tse T, Sladden TM, Hii E, Yerkovich ST, Hopkins PM, Chambers DC. Erratic tacrolimus exposure, assessed using the standard deviation of trough blood levels, predicts chronic lung allograft dysfunction and survival. *J Heart Lung Transplant* 2015; **34**: 1442-1448 [PMID: 26186804 DOI: 10.1016/j.healun.2015.05.028]
- 29 **Kao CC**, Segraves J, Parulekar AD. Tacrolimus monitoring parameters are not associated with acute cellular rejection following lung transplantation. *Eur J Clin Pharmacol* 2021; **77**: 63-69 [PMID: 32803287 DOI: 10.1007/s00228-020-02976-z]
- 30 **Ensor CR**, Iasella CJ, Harrigan KM, Morrell MR, Moore CA, Shigemura N, Zeevi A, McDyer JF, Venkataramanan R. Increasing tacrolimus time-in-therapeutic range is associated with superior one-year outcomes in lung transplant recipients. *Am J Transplant* 2018; **18**: 1527-1533 [PMID: 29513387 DOI: 10.1111/ajt.14723]
- 31 **Katada Y**, Nakagawa S, Itoharu K, Suzuki T, Kato R, Endo H, Sugimoto M, Yonezawa A, Nakagawa T, Ohsumi A, Nakajima D, Date H, Terada T. Association between time in therapeutic range of tacrolimus blood concentration and acute rejection within the first three months after lung transplantation. *J Pharm Health Care Sci* 2022; **8**: 25 [PMID: 36180948 DOI: 10.1186/s40780-022-00256-9]
- 32 **Venkat VL**, Nick TG, Wang Y, Bucuvalas JC. An objective measure to identify pediatric liver transplant recipients at risk for late allograft rejection related to non-adherence. *Pediatr Transplant* 2008; **12**: 67-72 [PMID: 18186891 DOI: 10.1111/j.1399-3046.2007.00794.x]
- 33 **Shemesh E**, Bucuvalas JC, Anand R, Mazariegos GV, Alonso EM, Venick RS, Reyes-Mugica M, Annunziato RA, Shneider BL. The Medication Level Variability Index (MLVI) Predicts Poor Liver Transplant Outcomes: A Prospective Multi-Site Study. *Am J Transplant* 2017; **17**: 2668-2678 [PMID: 28321975 DOI: 10.1111/ajt.14276]
- 34 **de Oliveira JTP**, Kielling CO, da Silva AB, Stefani J, Witkowski MC, Smidt CR, Mariano da Rocha CR, Hirakata VN, Grossini MDG, Zanotelli ML, Gonçalves Vieira SM. Variability index of tacrolimus serum levels in pediatric liver transplant recipients younger than 12 years: Non-adherence or risk of non-adherence? *Pediatr Transplant* 2017; **21** [PMID: 29034612 DOI: 10.1111/petr.13058]
- 35 **Defrancq C**, De Wilde N, Raes A, Van Biervliet S, Vande Velde S, Van Winckel M, De Bruyne R, Prytula A. Intra-patient variability in tacrolimus exposure in pediatric liver transplant recipients: Evolution, risk factors, and impact on patient outcomes. *Pediatr Transplant* 2019; **23**: e13388 [PMID: 30916883 DOI: 10.1111/petr.13388]
- 36 **Christina S**, Annunziato RA, Schiano TD, Anand R, Vaidya S, Chuang K, Zack Y, Florman S, Shneider BL, Shemesh E. Medication level variability index predicts rejection, possibly due to nonadherence, in adult liver transplant recipients. *Liver Transpl* 2014; **20**: 1168-1177 [PMID: 24931127 DOI: 10.1002/lt.23930]
- 37 **Del Bello A**, Congy-Jolivet N, Danjoux M, Muscari F, Lavayssière L, Esposito L, Hebral AL, Bellière J, Kamar N. High tacrolimus intra-patient variability is associated with graft rejection, and *de novo* donor-specific antibodies occurrence after liver transplantation. *World J Gastroenterol* 2018; **24**: 1795-1802 [PMID: 29713132 DOI: 10.3748/wjg.v24.i16.1795]
- 38 **Rayar M**, Tron C, Jézéquel C, Beaupaire JM, Petitcollin A, Housset-Debry P, Camus C, Verdier MC, Dehlawi A, Lakéhal M, Desfourneaux V, Meunier B, Sulpice L, Bellissant E, Boudjema K, Lemaître F. High Inpatient Variability of Tacrolimus Exposure in the Early Period After Liver Transplantation Is Associated With Poorer Outcomes. *Transplantation* 2018; **102**: e108-e114 [PMID: 29315140 DOI: 10.1097/TP.0000000000002052]
- 39 **van der Veer MAA**, Nangrahy N, Hesselink DA, Erler NS, Metselaar HJ, van Gelder T, Darwish Murad S. High Inpatient Variability in Tacrolimus Exposure Is Not Associated With Immune-mediated Graft Injury After Liver Transplantation. *Transplantation* 2019; **103**: 2329-2337 [PMID: 30801539 DOI: 10.1097/TP.0000000000002680]
- 40 **Dopazo C**, Bilbao I, García S, Gómez-Gavara C, Caralt M, Campos-Varela I, Castells L, Hidalgo E, Moreso F, Montoro B, Charco R. High inpatient variability of tacrolimus exposure associated with poorer outcomes in liver transplantation. *Clin Transl Sci* 2022; **15**: 1544-1555 [PMID: 35373449 DOI: 10.1111/cts.13276]
- 41 **Kim HJ**, Lee J, Lee JG, Joo DJ, Kim MS. Clinical association between tacrolimus intra-patient variability and liver transplantation outcomes in patients with and without hepatocellular carcinoma. *Sci Rep* 2022; **12**: 16169 [PMID: 36171260 DOI: 10.1038/s41598-022-20636-3]
- 42 **Davis S**, Gralla J, Klem P, Stites E, Wiseman A, Cooper JE. Tacrolimus Inpatient Variability, Time in Therapeutic Range, and Risk of De Novo Donor-Specific Antibodies. *Transplantation* 2020; **104**: 881-887 [PMID: 32224815 DOI: 10.1097/TP.0000000000002913]
- 43 **Gonzales HM**, McGillicuddy JW, Rohan V, Chandler JL, Nadig SN, Dubay DA, Taber DJ. A comprehensive review of the impact of tacrolimus inpatient variability on clinical outcomes in kidney transplantation. *Am J Transplant* 2020; **20**: 1969-1983 [PMID: 32406604 DOI: 10.1111/ajt.16002]
- 44 **Sapir-Pichhadze R**, Wang Y, Famure O, Li Y, Kim SJ. Time-dependent variability in tacrolimus trough blood levels is a risk factor for late kidney transplant failure. *Kidney Int* 2014; **85**: 1404-1411 [PMID: 24336032 DOI: 10.1038/ki.2013.465]
- 45 **Ro H**, Min SI, Yang J, Moon KC, Kim YS, Kim SJ, Ahn C, Ha J. Impact of tacrolimus intraindividual variability and CYP3A5 genetic polymorphism on acute rejection in kidney transplantation. *Ther Drug Monit* 2012; **34**: 680-685 [PMID: 23149441 DOI: 10.1097/FTD.0b013e3182731809]
- 46 **Whalen HR**, Glen JA, Harkins V, Stevens KK, Jardine AG, Geddes CC, Clancy MJ. High Inpatient Tacrolimus Variability Is Associated With Worse Outcomes in Renal Transplantation Using a Low-Dose Tacrolimus Immunosuppressive Regime. *Transplantation* 2017; **101**: 430-436 [PMID: 26950724 DOI: 10.1097/TP.0000000000001129]
- 47 **Borra LC**, Roodnat JJ, Kal JA, Mathot RA, Weimar W, van Gelder T. High within-patient variability in the clearance of tacrolimus is a risk factor for poor long-term outcome after kidney transplantation. *Nephrol Dial Transplant* 2010; **25**: 2757-2763 [PMID: 20190242 DOI: 10.1093/ndt/gfq096]
- 48 **Shuker N**, Shuker L, van Rosmalen J, Roodnat JJ, Borra LC, Weimar W, Hesselink DA, van Gelder T. A high inpatient variability in tacrolimus exposure is associated with poor long-term outcome of kidney transplantation. *Transpl Int* 2016; **29**: 1158-1167 [PMID: 27188932 DOI: 10.1111/tri.12798]
- 49 **O'Regan JA**, Canney M, Connaughton DM, O'Kelly P, Williams Y, Collier G, deFreitas DG, O'Seaghdha CM, Conlon PJ. Tacrolimus trough-level variability predicts long-term allograft survival following kidney transplantation. *J Nephrol* 2016; **29**: 269-276 [PMID: 26374111 DOI: 10.1007/s40620-015-0230-0]
- 50 **Goodall DL**, Willicombe M, McLean AG, Taube D. High Inpatient Variability of Tacrolimus Levels and Outpatient Clinic Nonattendance Are Associated With Inferior Outcomes in Renal Transplant Patients. *Transplant Direct* 2017; **3**: e192 [PMID: 28795143 DOI: 10.1097/TXD.0000000000000710]
- 51 **Rozen-Zvi B**, Schneider S, Lichtenberg S, Green H, Cohen O, Gaft U, Chagnac A, Mor E, Rahamimov R. Association of the combination of time-weighted variability of tacrolimus blood level and exposure to low drug levels with graft survival after kidney transplantation. *Nephrol Dial Transplant* 2017; **32**: 393-399 [PMID: 28025383 DOI: 10.1093/ndt/gfw394]
- 52 **Rahamimov R**, Tifti-Orbach H, Zingerman B, Green H, Schneider S, Chagnac A, Mor E, Fox BD, Rozen-Zvi B. Reduction of exposure to

- tacrolimus trough level variability is associated with better graft survival after kidney transplantation. *Eur J Clin Pharmacol* 2019; **75**: 951-958 [PMID: 30762079 DOI: 10.1007/s00228-019-02643-y]
- 53 **Rodrigo E**, Segundo DS, Fernández-Fresnedo G, López-Hoyos M, Benito A, Ruiz JC, de Cos MA, Arias M. Within-Patient Variability in Tacrolimus Blood Levels Predicts Kidney Graft Loss and Donor-Specific Antibody Development. *Transplantation* 2016; **100**: 2479-2485 [PMID: 26703349 DOI: 10.1097/TP.0000000000001040]
 - 54 **Mo H**, Kim SY, Min S, Han A, Ahn S, Min SK, Lee H, Ahn C, Kim Y, Ha J. Association of Inpatient Variability of Tacrolimus Concentration With Early Deterioration of Chronic Histologic Lesions in Kidney Transplantation. *Transplant Direct* 2019; **5**: e455 [PMID: 31321291 DOI: 10.1097/TXD.0000000000000899]
 - 55 **Süsal C**, Döhler B. Late intra-patient tacrolimus trough level variability as a major problem in kidney transplantation: A Collaborative Transplant Study Report. *Am J Transplant* 2019; **19**: 2805-2813 [PMID: 30859672 DOI: 10.1111/ajt.15346]
 - 56 **Vanhove T**, Vermeulen T, Annaert P, Lerut E, Kuypers DRJ. High Inpatient Variability of Tacrolimus Concentrations Predicts Accelerated Progression of Chronic Histologic Lesions in Renal Recipients. *Am J Transplant* 2016; **16**: 2954-2963 [PMID: 27013142 DOI: 10.1111/ajt.13803]
 - 57 **Seibert SR**, Schladt DP, Wu B, Guan W, Dorr C, Remmel RP, Matas AJ, Mannon RB, Israni AK, Oetting WS, Jacobson PA. Tacrolimus trough and dose intra-patient variability and CYP3A5 genotype: Effects on acute rejection and graft failure in European American and African American kidney transplant recipients. *Clin Transplant* 2018; **32**: e13424 [PMID: 30318646 DOI: 10.1111/ctr.13424]
 - 58 **Stefanović NZ**, Veličković-Radovanović RM, Danković KS, Mitić BP, Paunović GJ, Cvetković MB, Cvetković TP. Combined Effect of Inter- and Inpatient Variability in Tacrolimus Exposure on Graft Impairment Within a 3-Year Period Following Kidney Transplantation: A Single-Center Experience. *Eur J Drug Metab Pharmacokinet* 2020; **45**: 749-760 [PMID: 32886348 DOI: 10.1007/s13318-020-00644-2]
 - 59 **Stefanović N**, Veličković-Radovanović R, Danković K, Pavlović I, Catić-Dordević A, Bašić J, Despotović M, Jevtović-Stoimenov T, Mitić B, Cvetković T. Effect of the Interrelation between CYP3A5 Genotype, Concentration/Dose Ratio and Inpatient Variability of Tacrolimus on Kidney Graft Function: Monte Carlo Simulation Approach. *Pharmaceutics* 2021; **13** [PMID: 34834385 DOI: 10.3390/pharmaceutics13111970]
 - 60 **Nuchjumboon A**, Vacharavivad S, Singhan W, Poosonthornsri M, Chanchaoentana W, Udomkarnjananun S, Townamchai N, Avihingsanon Y, Praditpornsilpa K, Eiam-Ong S. Comparison of Tacrolimus Intra-Patient Variability during 6-12 Months after Kidney Transplantation between CYP3A5 Expressers and Nonexpressers. *J Clin Med* 2022; **11** [PMID: 36362548 DOI: 10.3390/jcm11216320]
 - 61 **Song T**, Yin S, Jiang Y, Huang Z, Liu J, Wang Z, Li L, Zeng J, Fan Y, Wang X, Li X, Lin T. Increasing Time in Therapeutic Range of Tacrolimus in the First Year Predicts Better Outcomes in Living-Donor Kidney Transplantation. *Front Immunol* 2019; **10**: 2912 [PMID: 31921171 DOI: 10.3389/fimmu.2019.02912]
 - 62 **Sablik KA**, Claassen-van Groningen MC, Hesselink DA, van Gelder T, Betjes MGH. Tacrolimus intra-patient variability is not associated with chronic active antibody mediated rejection. *PLoS One* 2018; **13**: e0196552 [PMID: 29746495 DOI: 10.1371/journal.pone.0196552]
 - 63 **Park Y**, Lee H, Eum SH, Kim HD, Ko EJ, Yang CW, Chung BH. Inpatient Variability in Tacrolimus Trough Levels Over 2 Years Affects Long-Term Allograft Outcomes of Kidney Transplantation. *Front Immunol* 2021; **12**: 746013 [PMID: 34659243 DOI: 10.3389/fimmu.2021.746013]
 - 64 **Yin S**, Wang X, Huang Z, Fan Y, Song T, Lin T. Tacrolimus variability score outperforms coefficient of variation in predicting clinical outcomes of living kidney transplantation. *Br J Clin Pharmacol* 2022; **88**: 75-83 [PMID: 33899267 DOI: 10.1111/bcp.14876]
 - 65 **Kim EJ**, Kim SJ, Huh KH, Kim BS, Kim MS, Kim SI, Kim YS, Lee J. Clinical significance of tacrolimus intra-patient variability on kidney transplant outcomes according to pre-transplant immunological risk. *Sci Rep* 2021; **11**: 12114 [PMID: 34108576 DOI: 10.1038/s41598-021-91630-4]
 - 66 **Gold A**, Tönshoff B, Döhler B, Süsal C. Association of graft survival with tacrolimus exposure and late intra-patient tacrolimus variability in pediatric and young adult renal transplant recipients-an international CTS registry analysis. *Transpl Int* 2020; **33**: 1681-1692 [PMID: 32881096 DOI: 10.1111/tri.13726]
 - 67 **Baghai Arassi M**, Gauche L, Schmidt J, Höcker B, Rieger S, Süsal C, Tönshoff B, Fichtner A. Association of intraindividual tacrolimus variability with *de novo* donor-specific HLA antibody development and allograft rejection in pediatric kidney transplant recipients with low immunological risk. *Pediatr Nephrol* 2022; **37**: 2503-2514 [PMID: 35166920 DOI: 10.1007/s00467-022-05426-3]
 - 68 **Pizzo HP**, Ettenger RB, Gjertson DW, Reed EF, Zhang J, Gritsch HA, Tsai EW. Sirolimus and tacrolimus coefficient of variation is associated with rejection, donor-specific antibodies, and nonadherence. *Pediatr Nephrol* 2016; **31**: 2345-2352 [PMID: 27286686 DOI: 10.1007/s00467-016-3422-5]
 - 69 **Torabi J**, Konicki A, Rocca JP, Ajaimy M, Campbell A, Azzi Y, Pynadath C, Liriano-Ward L, Akalin E, Kinkhabwala M, Graham JA. The use of LCP-Tacrolimus (Envarsus XR) in simultaneous pancreas and kidney (SPK) transplant recipients. *Am J Surg* 2020; **219**: 583-586 [PMID: 32122660 DOI: 10.1016/j.amjsurg.2020.02.027]
 - 70 **Marco DN**, Salas MQ, Gutiérrez-García G, Monge I, Riu G, Carcelero E, Roma JR, Llobet N, Arcarons J, Suárez-Lledó M, Martínez N, Pedraza A, Domenech A, Rosiñol L, Fernández-Avilés F, Urbano-Ispizua Á, Rovira M, Brunet M, Martínez C. Impact of Early Inpatient Variability of Tacrolimus Concentrations on the Risk of Graft-Versus-Host Disease after Allogeneic Stem Cell Transplantation Using High-Dose Post-Transplant Cyclophosphamide. *Pharmaceutics (Basel)* 2022; **15** [PMID: 36558980 DOI: 10.3390/ph15121529]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

