


World Journal of *Methodology*

World J Methodol 2023 December 20; 13(5): 373-509



EDITORIAL

- 373 Challenges and limitations of synthetic minority oversampling techniques in machine learning
Alkhalwaldeh IM, Albalkhi I, Naswhan AJ
- 379 Current protocol to achieve dental movement acceleration and pain control with Photo-biomodulation
Dominguez A
- 384 New evidence-based practice: Artificial intelligence as a barrier breaker
Ferreira RM

OPINION REVIEW

- 390 Evidence-based literature review: De-duplication a cornerstone for quality
Hammer B, Virgili E, Bilotta F

REVIEW

- 399 Crohn's disease and clinical management today: How it does?
da Silva Júnior RT, Apolonio JS, de Souza Nascimento JO, da Costa BT, Malheiro LH, Silva Luz M, de Carvalho LS, da Silva Santos C, Freire de Melo F

MINIREVIEWS

- 414 Using national census data to facilitate healthcare research
Colwill M, Poullis A
- 419 Machine learning and deep neural network-based learning in osteoarthritis knee
Ratna HVK, Jeyaraman M, Jeyaraman N, Nallakumarasamy A, Sharma S, Khanna M, Gupta A
- 426 Synoptic review on existing and potential sources for bias in dental research methodology with methods on their prevention and remedies
Agrawal AA, Prakash N, Almagbol M, Alobaid M, Alqarni A, Altamni H

ORIGINAL ARTICLE**Retrospective Study**

- 439 Assessing the readability of online information about jones fracture
Al-Kharouf KFK, Khan FI, Robertson GA
- 446 Impact of COVID-19 lockdown on hospital admissions for epistaxis in Germany
Hoenle A, Wagner M, Lorenz S, Steinhart H

- 456 Effect of vaccination status on CORADS and computed tomography severity score in hospitalized COVID-19 patients: A retrospective study

Binay UD, Karavaş E, Karakeçili F, Barkay O, Aydın S, Şenbil DC

Observational Study

- 466 Study on good clinical practices among researchers in a tertiary healthcare institute in India

Harshita H, Panda PK

- 475 Inflammatory bowel disease among first generation immigrants in Israel: A nationwide epi-Israeli Inflammatory Bowel Disease Research Nucleus study

Stulman M, Focht G, Loewenberg Weisband Y, Greenfeld S, Ben Tov A, Ledderman N, Matz E, Paltiel O, Odes S, Dotan I, Benchimol EI, Turner D

Basic Study

- 484 Sequential extraction of RNA, DNA and protein from cultured cells of the same group

Cui YY

- 492 Urine exosome mRNA-based test for monitoring kidney allograft rejection: Effects of sample transportation and storage, and interference substances

McFaul M, Ventura C, Evans S, Dundar H, Rumpler MJ, McCloskey C, Lowe D, Vlassov AV

CASE REPORT

- 502 Successful hip revision surgery following refracture of a modern femoral stem using a cortical window osteotomy technique: A case report and review of literature

Lucero CM, Luco JB, Garcia-Mansilla A, Slullitel PA, Zanotti G, Comba F, Buttaro MA

ABOUT COVER

Editorial Board Member of *World Journal of Methodology*, Rodrigo Valenzuela, PhD, Associated Professor, Department of Nutrition, Faculty of Medicine, University of Chile, Independence Av. 1027, Santiago 8380000, Chile. rvalenzuelab@med.uchile.cl

AIMS AND SCOPE

The primary aim of *World Journal of Methodology* (*WJM, World J Methodol*) is to provide scholars and readers from various fields of methodology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJM mainly publishes articles reporting research results obtained in the field of methodology and covering a wide range of topics including breath tests, cardiac imaging techniques, clinical laboratory techniques, diagnostic self-evaluation, cardiovascular diagnostic techniques, digestive system diagnostic techniques, endocrine diagnostic techniques, neurological diagnostic techniques, obstetrical and gynecological diagnostic techniques, ophthalmological diagnostic techniques, otological diagnostic techniques, radioisotope diagnostic techniques, respiratory system diagnostic techniques, surgical diagnostic techniques, *etc.*

INDEXING/ABSTRACTING

The *WJM* is now abstracted and indexed in PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Ji-Hong Liu*.

NAME OF JOURNAL

World Journal of Methodology

ISSN

ISSN 2222-0682 (online)

LAUNCH DATE

September 26, 2011

FREQUENCY

Quarterly

EDITORS-IN-CHIEF

Timotius Ivan Hariyanto

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/2222-0682/editorialboard.htm>

PUBLICATION DATE

December 20, 2023

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/gerinfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/gerinfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

<https://www.wjgnet.com/bpg/gerinfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/gerinfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Machine learning and deep neural network-based learning in osteoarthritis knee

Harish V K Ratna, Madhan Jeyaraman, Naveen Jeyaraman, Arulkumar Nallakumarasamy, Shilpa Sharma, Manish Khanna, Ashim Gupta

Specialty type: Orthopedics

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): D
Grade E (Poor): 0

P-Reviewer: Ferreira RM, Portugal;
Wang T, China; Hariyanto TI,
Indonesia

Received: July 16, 2023

Peer-review started: July 16, 2023

First decision: September 13, 2023

Revised: September 14, 2023

Accepted: September 28, 2023

Article in press: September 28, 2023

Published online: December 20,
2023



Harish V K Ratna, Department of Orthopaedics, Rathimed Speciality Hospital, Chennai 600040, Tamil Nadu, India

Madhan Jeyaraman, Naveen Jeyaraman, Arulkumar Nallakumarasamy, Department of Orthopaedics, ACS Medical College and Hospital, Dr MGR Educational and Research Institute, Chennai 600077, Tamil Nadu, India

Madhan Jeyaraman, Ashim Gupta, Department of Orthopaedics, South Texas Orthopaedic Research Institute, Laredo, TX 78045, United States

Shilpa Sharma, Department of Paediatric Surgery, All India Institute of Medical Sciences, New Delhi 110029, India

Manish Khanna, Department of Orthopaedics, Autonomous State Medical College, Ayodhya 224133, Uttar Pradesh, India

Ashim Gupta, Department of Regenerative Medicine, Regenerative Orthopaedics, Noida 201301, Uttar Pradesh, India

Ashim Gupta, Department of Regenerative Medicine, Future Biologics, Lawrenceville, GA 30043, United States

Ashim Gupta, Department of Regenerative Medicine, BioIntegarte, Lawrenceville, GA 30043, United States

Corresponding author: Madhan Jeyaraman, MS, PhD, Assistant Professor, Research Associate, Department of Orthopaedics, ACS Medical College and Hospital, Dr MGR Educational and Research Institute, Chennai 600077, Tamil Nadu, India. madhanjeyaraman@gmail.com

Abstract

Osteoarthritis (OA) of the knee joint is considered the commonest musculoskeletal condition leading to marked disability for patients residing in various regions around the globe. Application of machine learning (ML) in doing research regarding OA has brought about various clinical advances viz, OA being diagnosed at preliminary stages, prediction of chances of development of OA among the population, discovering various phenotypes of OA, calculating the severity in OA structure and also discovering people with slow and fast progression of disease pathology, *etc.* Various publications are available regarding machine

learning methods for the early detection of osteoarthritis. The key features are detected by morphology, molecular architecture, and electrical and mechanical functions. In addition, this particular technique was utilized to assess non-interfering, non-ionizing, and in-vivo techniques using magnetic resonance imaging. ML is being utilized in OA, chiefly with the formulation of large cohorts viz, the OA Initiative, a cohort observational study, the Multi-centre Osteoarthritis Study, an observational, prospective longitudinal study and the Cohort Hip & Cohort Knee, an observational cohort prospective study of both hip and knee OA. Though ML has various contributions and enhancing applications, it remains an imminent field with high potential, also with its limitations. Many more studies are to be carried out to find more about the link between machine learning and knee osteoarthritis, which would help in the improvement of making decisions clinically, and expedite the necessary interventions.

Key Words: Osteoarthritis; Knee; Artificial intelligence; Machine learning; Deep neural network

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Application of machine learning in research has various clinical advances viz, osteoarthritis (OA) knee being diagnosed at preliminary stages, prediction of development of OA, discovering various phenotypes. Large cohorts have been formulated viz, the OA Initiative, the Multi-centre Osteoarthritis Study and the Cohort Hip & Cohort Knee. Many studies are awaited to find about the link between ML and knee OA, which would improve making decisions clinically, and expedite the necessary interventions.

Citation: Ratna HVK, Jeyaraman M, Jeyaraman N, Nallakumarasamy A, Sharma S, Khanna M, Gupta A. Machine learning and deep neural network-based learning in osteoarthritis knee. *World J Methodol* 2023; 13(5): 419-425

URL: <https://www.wjgnet.com/2222-0682/full/v13/i5/419.htm>

DOI: <https://dx.doi.org/10.5662/wjm.v13.i5.419>

INTRODUCTION

Osteoarthritis (OA) of the knee joint is considered the commonest musculoskeletal condition leading to marked disability for patients residing in various regions around the globe[1,2]. OA is a degenerative disease, still, there is no clear knowledge regarding the factors that are responsible for the progression of the disease pathology. In the late stages of the disease, the only option for treatment is total knee arthroplasty, which is not affordable by many, is very much invasive, and highly degrades the individuals' quality of living.

A breakthrough was brought about by artificial intelligence (AI), specifically by machine learning (ML) in the field of medical care, mainly in subspecialties like rheumatology, mainly OA[3-6]. Machine learning includes Supervised, Semi-supervised, Unsupervised, and Reinforcement learning[7]. Deep learning is a subtype of ML that depends on various layers of a network consisting of a neuron architecture, that allows a model to improvise itself and train on its own, which in turn will lead to great accuracy levels by extracting greater level features from the given data[8-10].

Application of ML in doing research regarding OA has brought about various clinical advances viz, OA being diagnosed at preliminary stages, prediction of chances of development of OA among the population, discovering various phenotypes of OA, calculating the severity in OA structure and also discovering people with slow and fast progression of disease pathology, *etc.* The present models for the prediction of the progression of OA have been based on the collaboration of texture descriptors separated from the trabecular bone, Kellgren Lawrence (KL) grade, and clinical and anthropometric data[11-14]. Methods of Machine learning are divided into supervised and unsupervised (Figure 1). In supervised analysis, the results are established and the data are tagged. Whereas, in unsupervised analysis, the results are not established and the data are not tagged. In addition to these, two new categories were added viz, semi-supervised & reinforcement learning, in which the outcomes are only partially established[7].

Models of semi-supervised learning comprise a mixture of data that are tagged & untagged and are predicated on weak monitoring, with limited tagged data that are used to gain information and for the monitoring of untagged data. On the other hand, reinforcement learning is an ML prototype in which learning happens collectively by a sequence of hit-and-miss trials to increase the result achieved after every hit and thereby increase the learning behavior. Amongst the above-mentioned methods, supervised ML methods are most frequently utilized in the health care systems and medical field[15].

ML is being utilized in OA, chiefly with the formulation of large cohorts viz, the OA Initiative (OAI)[16], a cohort observational study, the Multi-centre Osteoarthritis Study[17], an observational, prospective longitudinal study, and the Cohort Hip & Cohort Knee[18], an observational cohort prospective study of both hip and knee OA. Though ML has various contributions and enhancing applications, it remains an imminent field with high potential, also with its limitations.




Figure 1 Various algorithms used in machine learning.

LITERATURE REVIEW

Various publications are available regarding machine learning methods for the early detection of osteoarthritis. The detection of osteoarthritis of the knee in its early stages and the exact & persistent assessment needed to detect changes in cartilage were discussed by Mohd Hani *et al*[19]. The key features are detected by morphology, molecular architecture, and electrical and mechanical functions. In addition, this particular technique was utilized to assess non-interfering, non-ionizing, and in-vivo techniques using magnetic resonance imaging (MRI).

Nelson *et al*[20] elaborated on the various causative factors of Osteoarthritis of the knee viz age, mass, and problems to the joint because of the flexing and kneeling activities. The data was attained by the authors from the OAI to assess the advancement of the knee. Dual Echo Steady State MRI was utilized to scrutinize the various images that are available and to choose the part of comparison and later on, perform a computerized differentiation from the data that is available. The modus operandi and design utilized is the Active Appearance Model. The CDI values that are acquired, are fed into machine learning methods like support vector machines (SVM), artificial neural network (ANN) & Random Forest to achieve precise results.

In a study conducted by Kwon *et al*[21], the radiographic maps were used & applied the model of SVM grouping was to authorize the deep learning model. For the multi-classification system, the deep learning network that was used was the Inceptio-ResNet-v2. Calculations of various parameters viz sensitivity, precision, gait analysis, X-ray findings, and F1-values were separated and skilled with property extraction and depth learning features extraction respectively and the training & conditioning of the SVM model was done by utilizing the training set and the compiled data was utilized in the diagnosis of OA. They found an ambiguous learning technique based on radiographic findings, blue-print-dependent scrutiny of data & analysis of functions. Following the inferences of this study improved the accuracy of the diagnosis of OA of the knee.

Wahyuningrum *et al*[22] utilized Long Short-Term Memory (LSTM) for extraction of the features to inquire about the pre-processing through a convolution neural network (CNN). Their main concerns were on the process of cross-validation, of which 2/3 and 1/3 were taken as training and testing data respectively. They performed the analysis by cropping the data of knee images into 400 × 100 pixels which were then arranged as piled augmented images and in turn were introduced in the manner of sequences, as input into the CNN, because LSTM should be performed sequentially. Three convolution filters followed the stride of Max pooling, and 1000 softmax channels into the LSTM processed the entire data from CNN. They have proposed that the LSTM technique is based on the KL grade and is more precise than deep learning.

An analysis of the cases was performed by Galván-Tejada *et al*[23] on the grounds of the data available from the OAI analysis. In this study they utilized two kinds of radiological grades, the first one being a quantitative analysis score and the second being a semi-quantitative analysis score, with the results of radiological images analyzed by two different sets of radiologists. To measure the variables related to pain in the future, A single variable logistic regression test was conducted. They found that the formation of bone osteophytes acts as the predictor of joint pain in the early stages and there's no correlation between pain in the subsequent period and a decrease in joint space. By the findings of the study, they have found that by using plain radiographs the reasons for potential OA knee can be predicted early. They have demonstrated the correlation between the discriminant X-ray of retrogression and the knee pain that developed.

Due to the difficulties faced in diagnosing knee Osteoarthritis, Christodoulou *et al*[24] brought about the concept of deep neural networks for the categorization of issues faced in diagnosis. In this study, to solve the problems in diagnosing osteoarthritis, taking into consideration the variable health factors that cause osteoarthritis, they have introduced a deep learning network to solve the problems faced in classifying knee OA, as a new and efficient machine learning method. The efficiency of the new method was proven by making discrete subgroups of the treated patients, from the data available and by designing a category for the diagnosis of OA knee. Deep neural networks serve as a successful technique in tackling the several problems of machine learning technique viz classification of images, predilection and disease prognosis, *etc.*

Magnetic resonance imaging was utilized by Du *et al*[25] to find the biomedical information concealed in it to diagnose osteoarthritis. Four techniques of machine learning were utilized to assess the osteoarthritis progression and to figure out the transformation in grades according to Kellgren and Lawrence's (KL) classification system, narrowing of joint space in

both medial or lateral compartments. The 36-dimensional elements were divided into two separate 18-dimensional elements with each 18-dimensional element to be individually examined by a four-graded technique for analyzing the various changes of both lateral and medial compartments. In an attempt to narrow down the regions and increase the performance, the analysis by PCA was utilized. By this technique, the machine learning methods for analysis were used to project the increase in KL grades, and medial and lateral compartment joint space reduction, and helped in delineating the pathological progression of knee OA. For the analysis purpose of PCA, the CDI information was compiled from one of the 36-dimensional elements on the 3D knee model of the MR tibiofemoral compartment.

In another study conducted by Kawathekar *et al*[26], they used geometrical parameters from the radiographs of the knee (anteroposterior view), by calculating the distance between the femur and tibia in active form models. Other approaches like local binary arrangement, which is being utilized in most of the pattern recognition tools are also used in the diagnosis of OA knee.

MACHINE LEARNING TECHNIQUES

The typical programming consists of an Algorithm, that includes a couple of rules that require pre-processing data (input) that is utilized to calculate a form of processed data (output) (Figure 2).

The algorithms of machine learning (Figure 3) can learn from the available data and are utilized for the automatic calculation of the set of rules. To perform such approaches, three components[27,28] are essential viz, pre-processed input data, Output data that the algorithm is supposed to predict, and A comparative tool to verify the predicted output performance.

The working mechanism includes entering both the data into the pipeline, which in turn will learn on its own to transform one data into the other data form. Machine learning techniques can be utilized effectively when there is no feasibility or possibility for defining an algorithm by manual techniques and when there's adequate data that applies to the training method.

Various subtypes of machine learning are available like supervised and unsupervised methods. Supervised methods can be utilized when there is a definite idea about the output data and Unsupervised methods can be utilized to discover anonymous data patterns. For instance, Supervised-learning methods encompass linear array regression, gradient boosting and ANN, *etc.*

ARTIFICIAL NEURAL NETWORKS

In the recent past, AI has been used graciously in the field of medicine. AI is nothing but a division of software engineering that includes algorithm training to reflect the human learning process. The design of the human brain has been the basis of various machine learning methods like the ANN. The ANN consists of a congruence of neurons that are interrelated. The ground structure of the neurons, can transform their inner structure or initiate a process, based on the information that is being fed to them and in turn results in a yield that depends upon both the input provides and the present execution.

CONVOLUTION NEURAL NETWORKS (CNN)

CNN is utilized widely in various modalities like an assertion of images and videos, recommendation systems, and handling of basic languages. Convolutions are vastly used in processing images[29,30], this being one of the reasons behind its introduction to deep learning methods to perform visual tasks. CNN facilitates the advantage of knowing about the local patterns that are concealed in the data given unlike the other systems, which treat the input data variables globally. CNN belong to a special subgroup of ANN, which uses a system known as a convolution in a minimum of at most one of their steps. Yann LeCunn introduced CNN first in 1990 though it was not that popular at that time[31].

Convolutions involve the mathematical assessment of two different functions that are assertions of real values. In imaging modalities, various terminologies are used to describe the input and kernel, which indicates the first and second functions respectively. The output data that is attained is termed the feature map. CNN models date back to one among the historical deep neural network models, which have various layers that are hidden between the general ones, and this, in turn, helps in knowing more information regarding the features that are concealed in the images that are provided as the input. A second type of layer that is often utilized in CNN models conducts pooling undertaking. Due to this pooling, there is a reduction in spatial resolution and the most appropriate features are only retained, which is solely necessary for maintaining a feasible network size.

OBJECT CLASS DETECTION

In today's treatment modalities, technologies of machine vision are utilized mostly for the identification of surgical materials among the available objects. The features of medical image inquiry have helped doctors in various ways viz




Figure 2 Typical programming algorithm.




Figure 3 Machine learning algorithm.

enhancing the diagnostic capabilities of pediatricians, outcomes which are object-guided, treatment-oriented, and extensive evaluation of various body organs like evaluation of clinical images of those in a state of vulnerability. The availability of advanced technological systems enhances the necessity for identifying and diagnosing various diseases like osteoarthritis, but there are various hurdles to the analysis of its probable location.

Nowadays various algorithms are available which are being used in health care to solve these hurdles. Numerous numbers of various technological advancements are happening in the day-to-day period, which focuses on the diagnosis and recognition of many other disease conditions. The traditional object analyzing methods can be utilized in a broad spectrum of imaging methods viz Radiography, Ultrasonography, Computed Tomographic images, and MRI.

There is no standard method of ML which is acceptable in order to apply them for real-time application[32]. Future research should concentrate on making algorithms that can detect OA in the stages of less severity. The lacunae in the real-time application can be fulfilled by utilizing data in larger numbers and multimodality data to make the ML models more accurate and increase the credibility simultaneously[9].

CONCLUSION

The future of medical science relies upon developing new methods of detecting osteoarthritis in its early stages, which would help in providing joint-preserving treatment modalities to patients. Machine learning techniques would develop a new revolution of augmented radiological methods. Machine learning can explore a wide range of designs, to identify the interaction and multiresolution models, which can easily predict the dynamics of the system to find the predisposing factors. Many more studies are to be carried out to find more about the link between machine learning and knee osteoarthritis, which would help in the improvement of making decisions clinically, and expedite the necessary interventions.

FOOTNOTES

Author contributions: Jeyaraman M contributed to conceptualization; Ratna HVK, Jeyaraman N, and Nallakumarasamy A contributed to initial draft; Ratna HVK and Jeyaraman M contributed to final draft; Sharma S, Khanna M, and Gupta A contributed to supervision; All authors accepted to publish the current version of the manuscript.

Conflict-of-interest statement: Authors declare no conflicts of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: India

ORCID number: Harish V K Ratna 0000-0002-9144-0612; Madhan Jeyaraman 0000-0002-9045-9493; Naveen Jeyaraman 0000-0002-4362-3326; Arulkumar Nallakumarasamy 0000-0002-2445-2883; Shilpa Sharma 0000-0001-8695-8372; Manish Khanna 0000-0002-2890-869X; Ashim Gupta 0000-0003-1224-2755.

S-Editor: Liu JH

L-Editor: A

P-Editor: Zhang YL

REFERENCES

- 1 **Vina ER**, Kwok CK. Epidemiology of osteoarthritis: literature update. *Curr Opin Rheumatol* 2018; **30**: 160-167 [PMID: 29227353 DOI: 10.1097/BOR.0000000000000479]
- 2 **Arden N**, Nevitt MC. Osteoarthritis: epidemiology. *Best Pract Res Clin Rheumatol* 2006; **20**: 3-25 [PMID: 16483904 DOI: 10.1016/j.berh.2005.09.007]
- 3 **Jeyaraman M**, Nallakumarasamy A, Jeyaraman N. Industry 5.0 in Orthopaedics. *Indian J Orthop* 2022; **56**: 1694-1702 [PMID: 36187596 DOI: 10.1007/s43465-022-00712-6]
- 4 **Pandit A**, Radstake TRDJ. Machine learning in rheumatology approaches the clinic. *Nat Rev Rheumatol* 2020; **16**: 69-70 [PMID: 31908355 DOI: 10.1038/s41584-019-0361-0]
- 5 **Bohr A**, Memarzadeh K. The rise of artificial intelligence in healthcare applications. *Artificial Intelligence in Healthcare* 2020; 25-60 [DOI: 10.1016/B978-0-12-818438-7.00002-2]
- 6 **Kokkoti C**, Moustakidis S, Papageorgiou E, Giakas G, Tsaopoulos DE. Machine learning in knee osteoarthritis: A review. *Osteoarthr Cartil Open* 2020; **2**: 100069 [PMID: 36474688 DOI: 10.1016/j.ocarto.2020.100069]
- 7 **Sarker IH**. Machine Learning: Algorithms, Real-World Applications and Research Directions. *SN Comput Sci* 2021; **2**: 160 [PMID: 33778771 DOI: 10.1007/s42979-021-00592-x]
- 8 **Yick HTV**, Chan PK, Wen C, Fung WC, Yan CH, Chiu KY. Artificial intelligence reshapes current understanding and management of osteoarthritis: A narrative review. *Journal of Orthopaedics, Trauma and Rehabilitation* 2022; **29**: 22104917221082315 [DOI: 10.1177/22104917221082315]
- 9 **Binvignat M**, Podoia V, Butte AJ, Louati K, Klatzmann D, Berenbaum F, Mariotti-Ferrandiz E, Sellam J. Use of machine learning in osteoarthritis research: a systematic literature review. *RMD Open* 2022; **8** [PMID: 35296530 DOI: 10.1136/rmdopen-2021-001998]
- 10 **Podoia V**. Machine Learning and Artificial Intelligence. *Osteoarthritis and Cartilage* 2020; **28**: S16 [DOI: 10.1016/j.joca.2020.02.010]
- 11 **Janvier T**, Jennane R, Valery A, Harrar K, Delplanque M, Lelong C, Loeuille D, Toumi H, Lespessailles E. Subchondral tibial bone texture analysis predicts knee osteoarthritis progression: data from the Osteoarthritis Initiative: Tibial bone texture & knee OA progression. *Osteoarthritis Cartilage* 2017; **25**: 259-266 [PMID: 27742531 DOI: 10.1016/j.joca.2016.10.005]
- 12 **Kerkhof HJ**, Bierma-Zeinstra SM, Arden NK, Metrustry S, Castano-Betancourt M, Hart DJ, Hofman A, Rivadeneira F, Oei EH, Spector TD, Uitterlinden AG, Janssens AC, Valdes AM, van Meurs JB. Prediction model for knee osteoarthritis incidence, including clinical, genetic and biochemical risk factors. *Ann Rheum Dis* 2014; **73**: 2116-2121 [PMID: 23962456 DOI: 10.1136/annrheumdis-2013-203620]
- 13 **Janvier T**, Jennane R, Toumi H, Lespessailles E. Subchondral tibial bone texture predicts the incidence of radiographic knee osteoarthritis: data from the Osteoarthritis Initiative. *Osteoarthritis Cartilage* 2017; **25**: 2047-2054 [PMID: 28935435 DOI: 10.1016/j.joca.2017.09.004]
- 14 **Kraus VB**, Feng S, Wang S, White S, Ainslie M, Brett A, Holmes A, Charles HC. Trabecular morphometry by fractal signature analysis is a novel marker of osteoarthritis progression. *Arthritis Rheum* 2009; **60**: 3711-3722 [PMID: 19950282 DOI: 10.1002/art.25012]
- 15 **Lo Vercio L**, Amador K, Bannister JJ, Crites S, Gutierrez A, MacDonald ME, Moore J, Mouches P, Rajashekar D, Schimert S, Subbanna N, Tuladhar A, Wang N, Wilms M, Winder A, Forkert ND. Supervised machine learning tools: a tutorial for clinicians. *J Neural Eng* 2020; **17** [PMID: 33036008 DOI: 10.1088/1741-2552/abbf2]
- 16 **Peterfy CG**, Schneider E, Nevitt M. The osteoarthritis initiative: report on the design rationale for the magnetic resonance imaging protocol for the knee. *Osteoarthritis Cartilage* 2008; **16**: 1433-1441 [PMID: 18786841 DOI: 10.1016/j.joca.2008.06.016]
- 17 **Segal NA**, Nevitt MC, Gross KD, Hietpas J, Glass NA, Lewis CE, Torner JC. The Multicenter Osteoarthritis Study: opportunities for rehabilitation research. *PM R* 2013; **5**: 647-654 [PMID: 23953013 DOI: 10.1016/j.pmrj.2013.04.014]
- 18 **Wesseling J**, Boers M, Viergever MA, Hilberdink WK, Lafeber FP, Dekker J, Bijlsma JW. Cohort Profile: Cohort Hip and Cohort Knee (CHECK) study. *Int J Epidemiol* 2016; **45**: 36-44 [PMID: 25172137 DOI: 10.1093/ije/dyu177]
- 19 **Mohd Hani AF**, Malik AS, Kumar D, Kamil R, Razak R, Kiflie A. Features and modalities for assessing early knee osteoarthritis. In: Proceedings of the 2011 International Conference on Electrical Engineering and Informatics 2011: 1-6 [DOI: 10.1109/ICEEI.2011.6021631]
- 20 **Nelson AE**, Fang F, Arbeeve L, Cleveland RJ, Schwartz TA, Callahan LF, Marron JS, Loeser RF. A machine learning approach to knee osteoarthritis phenotyping: data from the FNIH Biomarkers Consortium. *Osteoarthritis Cartilage* 2019; **27**: 994-1001 [PMID: 31002938 DOI: 10.1016/j.joca.2018.12.027]
- 21 **Kwon SB**, Han H-S, Lee MC, Kim HC, Ku Y, Ro DH. Machine Learning-Based Automatic Classification of Knee Osteoarthritis Severity Using Gait Data and Radiographic Images. *IEEE Access* 2020; **8**: 120597-120603 [DOI: 10.1109/ACCESS.2020.3006335]
- 22 **Wahyuningrum RT**, Anifah L, Eddy Purnama IK, Hery Purnomo M. A New Approach to Classify Knee Osteoarthritis Severity from Radiographic Images based on CNN-LSTM Method. In: 2019 IEEE 10th International Conference on Awareness Science and Technology (iCAST). 2019: 1-6 [DOI: 10.1109/ICAwST.2019.8923284]
- 23 **Galván-Tejada JI**, Treviño V, Celaya-Padilla JM, Tamez-Peña JG. Knee Osteoarthritis pain prediction from X-ray imaging: Data from Osteoarthritis Initiative. In: 2014 International Conference on Electronics, Communications and Computers (CONIELECOMP). 2014: 194-199 [DOI: 10.1109/CONIELECOMP.2014.6808590]
- 24 **Christodoulou E**, Moustakidis S, Papandrianos N, Tsaopoulos D, Papageorgiou E. Exploring deep learning capabilities in knee osteoarthritis case study for classification. In: 2019 10th International Conference on Information, Intelligence, Systems and Applications (IISA). 2019: 1-6 [DOI: 10.1109/IISA.2019.8900714]
- 25 **Du Y**, Almajalid R, Shan J, Zhang M. A Novel Method to Predict Knee Osteoarthritis Progression on MRI Using Machine Learning Methods. *IEEE Trans Nanobioscience* 2018; **17**: 228-236 [PMID: 29994316 DOI: 10.1109/TNB.2018.2840082]
- 26 **Kawathekar PP**, Karande KJ. Severity analysis of Osteoarthritis of knee joint from X-ray images: A Literature review. In: 2014 International Conference on Signal Propagation and Computer Technology (ICSPCT 2014).2014: 648-652 [DOI: 10.1109/ICSPCT.2014.6885008]

- 27 **Shamir L**, Ling SM, Scott W, Hochberg M, Ferrucci L, Goldberg IG. Early detection of radiographic knee osteoarthritis using computer-aided analysis. *Osteoarthritis Cartilage* 2009; **17**: 1307-1312 [PMID: 19426848 DOI: 10.1016/j.joca.2009.04.010]
- 28 **Krizhevsky A**, Sutskever I, Hinton GE. ImageNet classification with deep convolutional neural networks. *Commun ACM* 2017; **60**: 84-90 [DOI: 10.1145/3065386]
- 29 **Mahum R**, Rehman SU, Meraj T, Rauf HT, Irtaza A, El-Sherbeeney AM, El-Meligy MA. A Novel Hybrid Approach Based on Deep CNN Features to Detect Knee Osteoarthritis. *Sensors (Basel)* 2021; **21** [PMID: 34577402 DOI: 10.3390/s21186189]
- 30 **Upadhyay A**, Sawant O, Choudhary P. Detection of Knee Osteoarthritis Stages Using Convolutional Neural Network. *SN COMPUT SCI* 2023; **4**: 257 [DOI: 10.1007/s42979-022-01644-6]
- 31 **Yalçın OG**. The Brief History of Convolutional Neural Networks. Medium. 2021. Available from: <https://towardsdatascience.com/the-brief-history-of-convolutional-neural-networks-45afa1046f7f>
- 32 **Nwanosike EM**, Conway BR, Merchant HA, Hasan SS. Potential applications and performance of machine learning techniques and algorithms in clinical practice: A systematic review. *Int J Med Inform* 2022; **159**: 104679 [PMID: 34990939 DOI: 10.1016/j.ijmedinf.2021.104679]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

