

World Journal of *Clinical Cases*

World J Clin Cases 2023 October 6; 11(28): 6670-6973



MINIREVIEWS

- 6670** Neurotransmitters regulate β cells insulin secretion: A neglected factor
Kong CC, Cheng JD, Wang W

ORIGINAL ARTICLE**Case Control Study**

- 6680** Factors influencing the surveillance of re-emerging intracranial infections in elective neurosurgical patients: A single-center retrospective study
Wang JL, Wu XW, Wang SN, Liu X, Xiao B, Wang Y, Yu J

Retrospective Study

- 6688** Clinical value of chemiluminescence method for detection of antinuclear antibody profiles
Xiang HY, Xiang XY, Ten TB, Ding X, Liu YW, Luo CH
- 6698** Value of ultrasound guided biopsy combined with Xpert *Mycobacterium tuberculosis*/resistance to rifampin assay in the diagnosis of chest wall tuberculosis
Yan QH, Chi JY, Zhang L, Xue F, Cui J, Kong HL
- 6707** Research on the intelligent internet nursing model based on the child respiratory and asthma control test scale for asthma management of preschool children
Pei CF, Zhang L, Xu XY, Qin Z, Liang HM
- 6715** Effects of different doses of long-acting growth hormone in treating children with growth hormone deficiency
Xia W, Wang T, Pan JY
- 6725** Efficacy and anti-inflammatory analysis of glucocorticoid, antihistamine and leukotriene receptor antagonist in the treatment of allergic rhinitis
Qiu C, Feng D
- 6733** Subchondral fatigue fracture of the femoral head in young military recruits: Potential risk factors
Yang JZ, Chen P, Chen BH, Zhao B
- 6744** Anemia status of infants and young children aged six to thirty-six months in Ma'an Shan City: A retrospective study
Wang XM, Wang QY, Huang J
- Observational Study**
- 6754** Impact of coronary artery bypass grafting surgery on the chorioretinal biomicroscopic characteristics
Shahriari M, Nikkhah H, Mahjoob MP, Behnaz N, Barkhordari S, Cheraqpour K

Prospective Study

- 6763** Effects of humanized nursing care on negative emotions and complications in patients undergoing hysteromyoma surgery
Liu L, Xiao YH, Zhou XH

Randomized Controlled Trial

- 6774** Randomized controlled trial on the efficacy and safety of autologous serum eye drops in dry eye syndrome
Zheng N, Zhu SQ

SYSTEMATIC REVIEWS

- 6782** Primary adrenal Ewing sarcoma: A systematic review of the literature
Manatakis DK, Tsouknidas I, Mylonakis E, Tasis NP, Antonopoulou MI, Acheimastos V, Mastoropoulou A, Korkolis DP

CASE REPORT

- 6792** Pulmonary artery aneurysm protruding into the bronchus as an endobronchial mass: A case report
Li M, Zhu WY, Wu RR, Wang L, Mo MT, Liu SN, Zhu DY, Luo Z
- 6797** Rare rectal gastrointestinal stromal tumor case: A case report and review of the literature
Dong RX, Wang C, Zhou H, Yin HQ, Liu Y, Liang HT, Pan YB, Wang JW, Cao YQ
- 6806** Bilateral retinal nerve fiber layer thickness reduction in a 9-year-old myopic boy suffering from unilateral optic neuritis: A case report
Zhao FF, Yao SQ, Wang Y, Li TP, Yang JF, Pang CP, Cen LP
- 6812** Application of negative pressure wound therapy after skin grafting in the treatment of skin cancer: A case report
Huang GS, Xu KC
- 6817** Diagnosis and treatment of McCune-Albright syndrome: A case report
Lin X, Feng NY, Lei YJ
- 6823** Paraneoplastic myopathy-related rhabdomyolysis and pancreatic cancer: A case report and review of the literature
Costantini A, Moletta L, Pierobon ES, Serafini S, Valmasoni M, Sperti C
- 6831** Multi-organ hereditary hemorrhagic telangiectasia: A case report
Chen YL, Jiang HY, Li DP, Lin J, Chen Y, Xu LL, Gao H
- 6841** Hyperprogression after anti-programmed death-1 therapy in a patient with urothelial bladder carcinoma: A case report
Yang HY, Du YX, Hou YJ, Lu DR, Xue P
- 6850** Effectiveness of antidepressant repetitive transcranial magnetic stimulation in a patient with refractory psychogenic dysphagia: A case report and review of literature
Woo CG, Kim JH, Lee JH, Kim HJ

- 6857** Entrapment neuropathy of common peroneal nerve by fabella: A case report
Lin JC, Tsai MH, Lin WP, Kuan TS, Lien WC
- 6864** Importance of accurate diagnosis of congenital agenesis of the gallbladder from atypical gallbladder stone presentations: A case report
Sun HJ, Ge F, Si Y, Wang Z, Sun HB
- 6871** Dorsal approach for isolated volar fracture-dislocation of the base of the second metacarpal: A case report
Kurozumi T, Saito M, Odachi K, Masui F
- 6877** Rotationplasty type BIIIb as an effective alternative to limb salvage procedure in adults: Two case reports
Chen ZX, Guo XW, Hong HS, Zhang C, Xie W, Sha M, Ding ZQ
- 6889** Primary cutaneous anaplastic large cell lymphoma with over-expressed Ki-67 transitioning into systemic anaplastic large cell lymphoma: A case report
Mu HX, Tang XQ
- 6895** Confusing finding of quantitative fluorescent polymerase chain reaction analysis in invasive prenatal genetic diagnosis: A case report
Chen C, Tang T, Song QL, He YJ, Cai Y
- 6902** Testicular mixed germ cell tumor: A case report
Xiao QF, Li J, Tang B, Zhu YQ
- 6908** Leukemic transformation during anti-tuberculosis treatment in aplastic anemia-paroxysmal nocturnal hemoglobinuria syndrome: A case report and review of literature
Xiu NN, Yang XD, Xu J, Ju B, Sun XY, Zhao XC
- 6920** Pancreatic arteriovenous malformation treated with transcatheter arterial embolization: Two case reports and review of literature
Shin SH, Cho CK, Yu SY
- 6931** Cecal duplication cyst in an infant presenting as shock: A case report
Kim SM, Lee SH, Park GY, Kim SS, Lee CG, Jin SJ
- 6938** Pulmonary reversed halo cycles and consolidations after immunotherapy: A case report
Suo H, Shi YJ, Huang ZD, Xu K, Huang H
- 6943** Unusual case of emphysematous cystitis mimicking intestinal perforation: A case report
Kang HY, Lee DS, Lee D
- 6949** Malignant proliferative ependymoma of the neck with lymph node metastasis: A case report
Wang K, Wen JZ, Zhou SX, Ye LF, Fang C, Chen Y, Wang HX, Luo X
- 6955** Wandering spleen torsion with portal vein thrombosis: A case report
Zhu XY, Ji DX, Shi WZ, Fu YW, Zhang DK

- 6961** Intracranial infection and sepsis in infants caused by *Salmonella derby*: A case report
Yu JL, Jiang LL, Dong R, Liu SY
- 6967** Large gastric hamartomatous inverted polyp accompanied by advanced gastric cancer: A case report
Park G, Kim J, Lee SH, Kim Y

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Hao Wang, MD, PhD, Associate Professor, Department of Emergency Medicine, John Peter Smith Health Network, Texas Christian University and University of North Texas Health Science Center, School of Medicine, Fort Worth, TX 76104, United States. hwang@ies.healthcare

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xu Guo; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

October 6, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Retrospective Study

Efficacy and anti-inflammatory analysis of glucocorticoid, antihistamine and leukotriene receptor antagonist in the treatment of allergic rhinitis

Chen Qiu, Dai Feng

Specialty type: Medicine, research and experimental**Provenance and peer review:** Unsolicited article; Externally peer reviewed.**Peer-review model:** Single blind**Peer-review report's scientific quality classification**Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0**P-Reviewer:** Autorino R, United States; Dobrzanska J, Poland**Received:** August 1, 2023**Peer-review started:** August 1, 2023**First decision:** August 16, 2023**Revised:** August 29, 2023**Accepted:** September 5, 2023**Article in press:** September 5, 2023**Published online:** October 6, 2023**Chen Qiu**, Department of Pharmacy, Hongshan District Health Service Center, Wuhan 430000, Hubei Province, China**Dai Feng**, Department of Ear-Nose-Throat, Wuchang Hospital Affiliated to Wuhan University of Science and Technology, Wuhan 430063, Hubei Province, China**Corresponding author:** Chen Qiu, MD, Doctor, Department of Pharmacy, Hongshan District Health Service Center, No. 70 Nanhu Road, Hongshan District, Wuhan 430000, Hubei Province, China. qc07200@163.com

Abstract

BACKGROUND

There are many adverse reactions in the treatment of allergic rhinitis (AR) mainly with conventional drugs. Leukotriene receptor antagonists, glucocorticoids and nasal antihistamines can all be used as first-line drugs for AR, but the clinical effects of the three drugs are not clear.

AIM

To examine the impact of glucocorticoids, antihistamines, and leukotriene receptor antagonists on individuals diagnosed with AR, specifically focusing on their influence on serum inflammatory indexes.

METHODS

The present retrospective study focused on the clinical data of 80 patients diagnosed and treated for AR at our hospital between May 2019 and May 2021. The participants were categorized into the control group and the observation group. The control group received leukotriene receptor antagonists, while the observation group was administered glucocorticoids and antihistamines. Conducted an observation and comparison of the symptoms, physical sign scores, adverse reactions, and effects on serum inflammatory indexes in two distinct groups of patients, both before and after treatment.

RESULTS

Subsequent to treatment, the nasal itching score, sneeze score, runny nose score, nasal congestion score, and physical signs score exhibited notable discrepancies ($P < 0.05$), with the observation group demonstrating superior outcomes compared

to the control group ($P < 0.05$). The interleukin (IL)-6, IL-10, tumor necrosis factor- α , Soluble Intercellular Adhesion Molecule-1, Leukotriene D4 after treatment were significantly different and the observation group is better than the control group, which is statistically significant ($P < 0.05$). Following the intervention, the incidence of adverse reactions in the observation group, including symptoms such as nasal dryness, discomfort in the throat, bitter taste in the mouth, and minor erosion of the nasal mucosa, was found to be 7.5%. This rate was significantly lower compared to the control group, which reported an incidence of 27.5%. The difference between the two groups was statistically significant ($P < 0.05$).

CONCLUSION

Glucocorticoids and antihistamines have obvious therapeutic effects, reduce serum inflammatory index levels, relieve symptoms and signs of patients, and promote patients' recovery, which can provide a reference for clinical treatment of AR.

Key Words: Glucocorticoid; Histamine; Leukotriene receptor antagonist; Allergic rhinitis; Serum inflammatory index

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Glucocorticoid, antihistamine and leukotriene receptor antagonist are good therapeutic agents for allergic rhinitis (AR) at present, which can avoid the adverse reactions caused by conventional drug treatment. The purpose of this paper is to explore the synergistic effect of these three drugs in the treatment of AR by combining them. The results show that glucocorticoid and antihistamine have obvious therapeutic effects and are worth popularizing widely in clinic.

Citation: Qiu C, Feng D. Efficacy and anti-inflammatory analysis of glucocorticoid, antihistamine and leukotriene receptor antagonist in the treatment of allergic rhinitis. *World J Clin Cases* 2023; 11(28): 6725-6732

URL: <https://www.wjgnet.com/2307-8960/full/v11/i28/6725.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i28.6725>

INTRODUCTION

Allergic rhinitis (AR) is a kind of allergic disease triggered by exogenous environmental factors, which leads to the immune system response, and it is one of the common diseases of respiratory system. It not only seriously endangers people's health but also easily induces a variety of diseases. AR primarily arises from the inhalation of allergens, which subsequently triggers the release of inflammatory mediators, leading to inflammation of the nasal mucosa[1]. Simultaneously, the etiology of the ailment is intricately associated with physical fitness, genetics, immunity, and additional factors. The primary clinical presentations of AR predominantly encompass nasal congestion, rhinorrhea, nasal pruritus, sneezing, and olfactory impairment[2]. In addition, it may also cause or be complicated by sinusitis, nasal polyps, pharyngitis, otitis media, tracheal and bronchitis, asthma and allergic conjunctivitis and other diseases, which seriously affect people's sleep, study, work and quality of life[3]. AR is primarily managed pharmacologically; however, the utilization of drugs is associated with safety concerns and an increased incidence of adverse reactions[4]. The current treatment methods for AR mainly include nasal glucocorticoid spray, oral antihistamine or nasal spray, intranasal anticholinergic, drug nasal spray, oral decongestant or nose drops, submucosal injection of inferior turbinate and surgery, vidian nerve surgery and nasal concussion treatment, etc[5]. A large number of animal experiments and clinical studies have confirmed that leukotriene receptor antagonists can inhibit the accumulation of inflammatory cells in the airway, inhibit the release of inflammatory mediators, antagonize airway inflammation, reduce airway hyperresponsiveness, and improve lung function in patients with asthma, Relieve clinical symptoms[6]. At present, it is believed that both nasal glucocorticoids and nasal antihistamines can be used as first-line drugs for AR, which can effectively alleviate nasal congestion, runny nose, sneezing and other symptoms[7]. Some scholars also believe that the combined use of nasal glucocorticoids and nasal antihistamines in the treatment of AR is expected to further improve the efficacy, but there is still a lack of sufficient evidence-based medicine[8]. Based on this, we have explored the effects of glucocorticoids, antihistamines, and leukotriene receptor antagonists on patients with AR and their effects on serum inflammatory indicators.

MATERIALS AND METHODS

General information

Before patients are included, patients are fully informed of the project, process and possible risks of the trial, and all patients are signed with informed consent forms and kept on file, and their identity records, basic information and medical records are included and kept on file. The retrospective research object of this study selected 80 patients with AR

who visited our hospital from May 2019 to May 2021. According to the experimental method, they were divided into control group (40 cases) and observation group (40 cases). There was no statistical difference in sex ratio, age distribution and other basic information between the two groups (Table 1).

Inclusion and exclusion criteria

Inclusion criteria: (1) All patients in this study met the "Expert Consensus on Allergic Rhinitis Subcutaneous Immunotherapy"[9] diagnostic criteria for AR; (2) Sneezing, clear watery mucus, nasal congestion, nasal itching and other symptoms appeared more than 2 items (including Item 2), or accompanied by conjunctival congestion and itching around the eyes, and the daily duration of symptoms exceeds 1 h or more; and (3) Pale and edema of the nasal mucosa, watery nasal discharge, positive skin prick test, positive serum specific immunoglobulin E (IgE) test, and consistent with skin prick test results are common. **Exclusion criteria:** (1) The subject has obvious systemic metabolic disease, or the researcher thinks that it may interfere with the evaluation of the research results or affect the safety of life. The subject has received sugar through oral, nasal or systemic routes within the past 3 mo. Corticosteroids, antihistamines, leukotriene receptor inhibitors, various decongestants or theophylline drugs; (2) Combined with serious heart, cerebrovascular, diabetes, hyperthyroidism, connective tissue diseases and hematopoietic diseases or other diseases or The disease is in the acute stage, pregnant or breast-feeding women; and (3) Thorax is obviously deformed, or has lung organic disease, or has other diseases that cannot be applied with, and has received specific immunotherapy within the past 1 year.

Method

Before the start of treatment, the doctor should introduce the uniform and correct method of nasal spray to the patient, and demonstrate on the spot: After waking up in the morning and before going to bed at night, after clearing the nasal secretions, face down, spray nozzle up or slightly outward, and hold in the left hand. Bottle, spray the right nasal cavity, hold the bottle in the right hand, spray the left nasal cavity, once a day in the morning and evening, 2 sprays on each nostril, each spray 50 g, all patients are prohibited from using other treatment drugs and methods during the treatment. Among them, the control group was given a leukotriene receptor antagonist, that is, loratadine tablets (Hainan Xinzhongzheng Pharmaceutical Co., Ltd., H20050628) were administered orally, 10 mg/time, once daily. Montelukast sodium (Hangzhou Merck Pharmaceutical Co., Ltd., J20070058) was taken orally, 10 mg/time, once/d, for 4 wk of treatment. The observation group was treated with glucocorticoids and antihistamines, that is, azelastine hydrochloride nasal spray and budesonide nasal spray were used for nasal spraying. The interval between the two nasal sprays was about 5 min. After waking up in the morning and before going to bed at night, after cleaning up nasal secretions, face down, spray nozzle upward or slightly outward, hold the bottle in the left hand to spray the right nasal cavity, and hold the bottle in the right hand to spray the left nasal cavity, once a day in the morning and evening, on each side Spray 50 g in the nostril, and the treatment time is 8 wk.

Follow-up and observation indicators

Clinical symptoms: Score the four typical symptoms of AR, namely sneezing, runny nose, nasal congestion, and nasal itching, each with a score of 1 to 3. Sneezing score 1 is divided into the number of consecutive sneezes between 3-9, score 2 is divided into the number of consecutive sneezes between 10-14, and score 3 is divided into the number of consecutive sneezes ≥ 15 ; the tear score is calculated based on the number of blows per day, 1 is the number of blows ≤ 4 times, 2 is the number of blows is between 5-9 times, and 3 is the number of blows ≥ 10 ; the nasal congestion score is 1 occasionally nasal congestion, 2 points between occasional nasal congestion and mouth breathing almost all day, 3 points almost all-day mouth breathing; nasal itching score 1 points to intermittent nasal itching, 2 points to sense of ant walking, but tolerable, 3 is divided into ant behavior, unbearable. Sign score: 3 is divided into inferior turbinate and nasal base, nasal septum only, no middle turbinate, or middle turbinate mucosal polypoid change, polyp formation; divided into inferior turbinate and nasal septum (or nasal base) only, inferior turbinate and There is still a small gap between the base of the nose (or septum); 1 is divided into mild swelling of the inferior turbinate, and the septum and middle turbinate are still visible. According to the scoring standard in the "Diagnostic Standards for Eye, Ear, Nose, and Throat Diseases"[10], the severity of symptoms of patients with AR is divided into 4 grades. 0 points: Asymptomatic; 1 point: Intermittent nasal itching, sneezing 3 to 9 times, occasional nasal congestion, the number of nose blows ≤ 4 times a day; 2 points: Obvious symptoms, nasal itching with a sense of ants, tolerable, 10 to 14 sneezes once, with obvious nasal congestion, 5 to 9 times a day to blow the nose; 3 points: Severe nasal itching and ant-like sensation, unbearable, 1 sneeze ≥ 15 , severe nasal congestion, almost all day breathe through your mouth, and blow your nose ≥ 10 times a day. Calculate the curative effect according to the following formula: Integral reduction index = (pre-treatment integral-post-treatment integral)/pre-treatment integral $\times 100\%$. Significantly effective: Integral decline index $\geq 50\%$; effective: Integral decline index $\geq 20\%$; invalid: Integral decline index $< 20\%$. Total effective rate = (marked effect + effective)/total number of cases $\times 100\%$.

Statistical analysis

The data included in this study were statistically analyzed by SPSS 23.0. Shapiro-wilk method was used to test, and the test level was $\alpha = 0.05$. The statistical method of mean standard deviation describes the data that conforms to the normal distribution, and the statistical method of M(QR) describes the data that does not conform to the normal distribution. *T*-test and χ^2 test of independent samples or paired samples are used to count the differences between groups or within groups. Integer or percentage (%) describes the counting data.

Table 1 Comparison of general information between the two groups (n, mean ± SD)				
Group	Gender (Male/Female)	Average age (yr)	Course of disease (wk)	Body mass index (kg/m ²)
Control group (40)	18/22	5.70 ± 1.32	5.6 (3.7-7.9)	27.78 ± 2.32
Observation group (40)	17/23	5.69 ± 1.66	5.5 (3.1-7.7)	27.62 ± 2.66
χ ² /t value	0.051	0.03	0.04	0.02
P value	0.822	0.976	0.965	0.984

RESULTS

Comparison of general information

There was no significant difference between the two groups in baseline data such as sex ratio, mean age, course of disease and body mass index ($P > 0.05$) (Table 1).

Comparison of symptoms and signs scores

Before treatment, there was no significant difference in symptoms and signs between the two groups ($P > 0.05$). After treatment, there were significant differences in nasal itching, sneezing, runny nose, nasal congestion and signs, and the curative effect of the observation group was better than that of the control group. In the control group, the comparison was statistically significant ($P < 0.05$) (Table 2).

Comparison of serum inflammatory indexes

The comparison of serum inflammatory indexes between the two groups before treatment was not statistically significant ($P > 0.05$). However, the levels of interleukin (IL)-6, IL-10, tumor necrosis factor-alpha (TNF-α), Soluble Intercellular Adhesion Molecule-1 (sICAM-1) and Leukotriene D4 (LTD4)-4 were significantly different after treatment, and the observation group was superior to the control group with statistical significance ($P < 0.05$) (Table 3).

Comparison of adverse reactions

After treatment, the adverse reaction rate of the observation group such as dryness of the nose, dry throat discomfort, bitter mouth, and slight erosion of the nasal mucosa was 7.5%, which was significantly lower than the control, 27.5%, which was statistically significant ($P < 0.05$) (Table 4).

DISCUSSION

AR is a series of nasal symptoms caused by IgE-mediated inflammation of the nasal mucosa after allergic individuals are exposed to allergens. The cause of the disease is believed to be the inhalation of allergens and the combination of platelets and eosinophils. The specific IgE binding on cells, mast cells and other cells causes these cells to release inflammatory mediators, which ultimately leads to the inflammatory process of the nasal mucosa[11]. Montelukast sodium is a selective leukotriene receptor antagonist, which can not only block the binding of leukotrienes and leukotriene receptors, so that it can not exert inflammatory effects, but also can inhibit copeptin growth factor Promoting maturation of basophils and eosinophils[12]. It further reduces eosinophils in the airway and surrounding blood, reduces inflammation, and the drug has no obvious adverse effects on vital organs or systems[13].

The results of this study show that after treatment, the degree of nasal itching, sneezing, runny nose, nasal congestion and signs have obviously improved, with statistical differences, and the observation group is better than the control group, indicating that the effect of glucocorticoid and antihistamine treatment is obvious, and the symptoms and signs of the patients are alleviated. The reasons are as follows: Glucocorticoids are the first-line drugs for the treatment of bronchial asthma, as well as the main treatment drugs for AR, and are currently the most effective anti-allergic inflammatory drugs[14]. From clinical observations, long-term regular inhalation of glucocorticoids can effectively control asthma symptoms, improve patients' mental state and quality of life, and reduce asthma mortality[15]. Its main mechanism of action is: Inhibit the synthesis of cytokines by lymphocytes and other immune active cells; Inhibit the chemotaxis and activation of E, reduce the synthesis of leukotrienes and prostaglandins, inhibit the metabolism of arachidonic acid, and promote the contraction of small blood vessels, Reduce blood vessel leakage, improve the responsiveness of respiratory smooth muscle 2 receptors, increase the synthesis of receptors on the cell membrane, etc[16]. Nasal glucocorticoids can effectively improve a series of nasal symptoms induced by AR, and the combined effect is better than antihistamines. Antihistamines are the first choice for patients with AR. They can quickly relieve the symptoms of runny nose, sneezing, nasal itching and dry eyes caused by histamine release[17]. H1 receptor blockers have a positive effect on the control of asthma. They can not only antagonize H1 receptors and reduce receptor sensitivity, but also inhibit the production of histamine stimulated I-8 and reduce the production of inflammatory mediators such as leukotrienes. And release, it has an effect on both the immediate-onset phase and the delayed-onset phase allergy in patients with asthma and AR[18].

Table 2 Comparison of the symptoms and signs scores of the two groups of patients before and after treatment (mean $\pm$ SD)

Group		Symptom score				Physical sign score
		Nasal itching score	Sneeze score	Runny nose score	Nasal congestion score	
Control group	Before therapy	2.14 $\pm$ 0.18	2.08 $\pm$ 0.32	1.84 $\pm$ 0.30	1.79 $\pm$ 0.22	2.13 $\pm$ 0.27
	After treatment	1.76 $\pm$ 0.63 ^a	1.45 $\pm$ 0.26 ^a	1.36 $\pm$ 0.06 ^a	1.37 $\pm$ 0.31 ^a	1.44 $\pm$ 0.23 ^a
Observation group	Before therapy	2.04 $\pm$ 0.61	2.03 $\pm$ 0.10	1.88 $\pm$ 0.80	1.82 $\pm$ 0.39	2.25 $\pm$ 0.44
	After treatment	1.32 $\pm$ 0.31 ^{a,b}	1.09 $\pm$ 0.26 ^{a,b}	0.92 $\pm$ 0.24 ^{a,b}	1.21 $\pm$ 0.31 ^{a,b}	1.17 $\pm$ 0.37 ^{a,b}

^a $P < 0.05$ vs before treatment.^b $P < 0.05$ vs the control group.**Table 3 Comparison of serum inflammatory indexes between the two groups (mean $\pm$ SD)**

Group		IL-6 (ng/L)	TNF- α (pg/L)	IL-10 (ng/L)	sICAM-1 (μ g/L)	LTD4 (mg/kg)
Control group	Before therapy	11.78 $\pm$ 2.32	55.34 $\pm$ 7.40	26.51 $\pm$ 6.32	40.78 $\pm$ 7.32	517.94 $\pm$ 23.40
	After treatment	6.62 $\pm$ 1.66	22.76 $\pm$ 6.64	16.55 $\pm$ 6.11	32.62 $\pm$ 5.66	206.96 $\pm$ 19.24
Observation group	Before therapy	11.78 $\pm$ 2.32	55.28 $\pm$ 7.80	26.81 $\pm$ 4.39	40.93 $\pm$ 7.27	517.75 $\pm$ 10.44
	After treatment	4.62 $\pm$ 0.66	19.19 $\pm$ 4.74	13.41 $\pm$ 3.31	26.74 $\pm$ 3.23	158.37 $\pm$ 10.37

IL: Interleukin; TNF- α : Tumor necrosis factor-alpha; sICAM-1: Soluble Interleukin Adhesion Molecule-1; LTD4: Leukotriene D4.**Table 4 Comparison of adverse reactions between the two groups of patients (mean $\pm$ SD)**

Group	Dry nose	Dry throat discomfort	Bitter	Minor erosive osmosis of nasal mucosa	Adverse reaction rate
Control group (40)	1	3	4	3	11 (27.5)
Observation group (40)	0	2	1	0	3 (7.5)
<i>t</i> value					5.541
<i>P</i> value					0.019

In this study, the comparison of serum inflammatory indexes between the two groups of patients before treatment was not statistically significant ($P > 0.05$), while the IL-6, IL-10, TNF- α , sICAM-1, and LTD4 were outstandingly different after treatment. Compared with the control group, the curative effect of the observation group is better. The therapeutic effect of glucocorticoid and antihistamine is obvious, and the serum inflammatory index level is reduced. IL-6 can induce inflammatory reactions, and its source range is very wide. A large number of experimental data show that IL-6 participates in a variety of inflammatory reactions and diseases in the human body and is an important pro-inflammatory factor[19]. IL-6 can promote the proliferation and differentiation of immune cells, and improve their immune function. IL-6 can activate T lymphocytes, induce B lymphocytes to differentiate, and finally secrete immunoglobulin. Therefore, the expression level of IL-6 will show a certain downward trend with the improvement of human health level, and gradually tend to the conventional expression level. In addition, IL-10 is a multifunctional cytokine derived from multiple cells. It is an important human anti-inflammatory factor. It can regulate immune response, regulate cell growth and differentiation, and participate in anti-inflammatory reaction[20]. In addition, IL-10 is also an important immune system regulator in human body, and it is currently recognized as an immunosuppressive factor in the medical field.

As a multifunctional inflammatory cytokine, TNF- α has anti-infection and anti-tumor effects[21]. Its reaction mechanism is to stimulate eosinophils and lymphocytes to produce IL-6 and other factors by mediating the accumulation of eosinophils and lymphocytes at the inflammatory site, thus indirectly playing an anti-tumor and anti-inflammatory role. Aggregation is activated, and the vascular endothelium synthesizes fibroblasts to synthesize colony stimulating factors, thereby causing the body to produce a persistent inflammatory response[22]. TNF- α can stimulate the proliferation of B cells and promote Ig secretion. The level of TNF- α can affect the body's anti-infection ability. Macrophages release a lot of TNF- α factors, which activate neutrophils and endothelial cells to adhere to inflammatory receptors and promote the release of inflammatory factors. Therefore, in the inflammatory reaction, the more serious the virus or bacterial infection, the higher the level of TNF- α [23].

Under normal circumstances, the severity of the condition can be judged according to the TNF-level in the patient's body, which has positive significance. The main role of sICAM-1 is to mediate the interaction between cells, cells and

extracellular matrix, and play an important role in the development of physiology and pathology[24]. LTD4 is a strong active leukotriene D, which promotes the contraction of bronchial smooth muscle, increases capillary permeability and promotes mucus secretion[25]. When the body has an allergic reaction or an inflammatory reaction, phospholipase acts on the phospholipid bilayer of the cell membrane to promote the production of 5-HPETE. Leukotriene receptor antagonists inhibit the release of inflammatory mediators and cytokines by down-regulating the expression of adhesion molecules induced by sICAM-1. Down-regulating the expression of LTD4, inhibiting the release of LTD4 in the airway, and reducing the proliferation of inflammatory cells, leukotriene receptor antagonists induce apoptosis by activating the 5-HPETE tetraenoic acid pathway.

CONCLUSION

Although the idea of this study is novel, there are some shortcomings. Although this study discussed the clinical efficacy of glucocorticoid, antihistamine and leukotriene receptor antagonist in patients with AR, there was a lack of research on its specific long-term mechanism. The research object of this study comes from the same hospital, and the sample source is single and representative. In addition, the exclusion and inclusion criteria of this study are subjectively set according to the experimental direction, which may lead to biased results. In summary, glucocorticoids and antihistamines have obvious therapeutic effects, reduce serum inflammatory index levels, relieve symptoms and signs of patients, promote patient recovery, and provide a certain reference for clinical treatment of AR.

ARTICLE HIGHLIGHTS

Research background

Glucocorticoids, antihistamines and leukotriene receptor antagonists are excellent therapeutic agents for allergic rhinitis (AR) at present, but the existing research lacks the comprehensive clinical effect comparison and analysis of the three.

Research motivation

To explore the clinical and anti-inflammatory effects of glucocorticoid, antihistamine and leukotriene receptor antagonist on AR.

Research objectives

To systematically evaluate the therapeutic effect of glucocorticoid, antihistamine and leukotriene receptor antagonist on AR, and evaluate its anti-inflammatory level to guide the follow-up clinical treatment.

Research methods

To evaluate the clinical efficacy, anti-inflammatory reaction and adverse reactions before and after treatment and between groups, and to comprehensively evaluate the efficacy of leukotriene receptor antagonists, glucocorticoids and antihistamines.

Research results

All three drugs have certain therapeutic effects, and the therapeutic effect of glucocorticoid combined with antihistamine is better than that of leukotriene receptor antagonist. Glucocorticoid combined with antihistamine has better anti-inflammatory effect and lower adverse reactions.

Research conclusions

Glucocorticoids and antihistamines are more effective in the treatment of AR, and it is better to reduce the level of serum inflammatory indicators and have lower adverse reactions.

Research perspectives

Summarize the common drugs used in the treatment of AR, and comprehensively analyze the clinical efficacy, anti-inflammatory level and adverse reactions of these drugs to serve clinical treatment.

FOOTNOTES

Author contributions: Qiu C and Feng D contributed equally to this work; Feng D conceived and designed the experiments; Qiu C and Feng D selected the literature, extracted data and analyzed it; Feng D and Qiu C wrote the manuscript.

Institutional review board statement: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent statement: This study is a retrospective study, therefore informed consent forms are exempted.

Conflict-of-interest statement: The authors declare no competing financial interest.

Data sharing statement: The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Chen Qiu 0009-0001-7145-0231.

S-Editor: Fan JR

L-Editor: A

P-Editor: Cai YX

REFERENCES

- Kalmarzi RN, Atace P, Fathollahpour A, Behzadifar M, Moradi M, Sharifian F, Kalani N, Kooti W. The Prevalence of Allergic Rhinitis among Iranian Children: A Systematic Review and Meta-Analysis. *Endocr Metab Immune Disord Drug Targets* 2020; **20**: 189-197 [PMID: 31113353 DOI: 10.2174/1871530319666190515100735]
- Field K, Blaiss MS. Sublingual Versus Subcutaneous Immunotherapy for Allergic Rhinitis: What Are the Important Therapeutic and Real-World Considerations? *Curr Allergy Asthma Rep* 2020; **20**: 45 [PMID: 32548677 DOI: 10.1007/s11882-020-00934-4]
- Leonardi A, Castegnaro A, Valerio AL, Lazzarini D. Epidemiology of allergic conjunctivitis: clinical appearance and treatment patterns in a population-based study. *Curr Opin Allergy Clin Immunol* 2015; **15**: 482-488 [PMID: 26258920 DOI: 10.1097/ACI.0000000000000204]
- Ahmed OG, Lane AP. Is Sublingual Immunotherapy an Effective Therapy for Allergic Rhinitis? *Laryngoscope* 2020; **130**: 2301-2302 [PMID: 32068887 DOI: 10.1002/lary.28574]
- Scadding GK, Kariyawasam HH, Scadding G, Mirakian R, Buckley RJ, Dixon T, Durham SR, Farooque S, Jones N, Leech S, Nasser SM, Powell R, Roberts G, Rotiroti G, Simpson A, Smith H, Clark AT. BSACI guideline for the diagnosis and management of allergic and non-allergic rhinitis (Revised Edition 2017; First edition 2007). *Clin Exp Allergy* 2017; **47**: 856-889 [PMID: 30239057 DOI: 10.1111/cea.12953]
- Di Salvo E, Patella V, Casciaro M, Gangemi S. The leukotriene receptor antagonist Montelukast can induce adverse skin reactions in asthmatic patients. *Pulm Pharmacol Ther* 2020; **60**: 101875 [PMID: 31837440 DOI: 10.1016/j.pupt.2019.101875]
- Roditi RE, Shin JJ. The Influence of Age on the Relationship Between Allergic Rhinitis and Otitis Media. *Curr Allergy Asthma Rep* 2018; **18**: 68 [PMID: 30343453 DOI: 10.1007/s11882-018-0826-2]
- Gómez F, Rondón C, Salas M, Campo P. Local allergic rhinitis: mechanisms, diagnosis and relevance for occupational rhinitis. *Curr Opin Allergy Clin Immunol* 2015; **15**: 111-116 [PMID: 25961385 DOI: 10.1097/ACI.0000000000000150]
- China Allergic Rhinitis Research Cooperative Group. [Expert consensus on subcutaneous immunotherapy for allergic rhinitis 2015]. *Zhongguo Er Bi Yanhou Tuijin Jiehe* 2015; **22**: 379-404 [DOI: 10.16066/j.1672-7002.2015.08.001]
- Bei ZP, Shu H, Zhou L. Diagnostic criteria for ophthalmology, ear, nose, and throat diseases M. Beijing: Science Press, 2001: 295-296. Available from: <https://baike.baidu.com/reference/7679045/216aMCCWI-ohmvi4Axp-rQc7nUkJsCiLsvZ1wDqJ5h26AtpEp6aba34FKaR332hmQZMrBCgCW7T7BDnQs8YUnsXHpp3C>
- Licari A, Castagnoli R, Brambilla I, Tosca MA, De Filippo M, Marseglia G, Ciprandi G. Biomarkers of immunotherapy response in patients with allergic rhinitis. *Expert Rev Clin Immunol* 2018; **14**: 657-663 [PMID: 30039714 DOI: 10.1080/1744666X.2018.1504679]
- Madison S, Brown EA, Franklin R, Wickersham EA, McCarthy LH. Clinical Question: Nasal saline or intranasal corticosteroids to treat allergic rhinitis in children. *J Okla State Med Assoc* 2016; **109**: 152-153 [PMID: 27328556]
- Wallace DV, Dykewicz MS. Seasonal Allergic Rhinitis: A focused systematic review and practice parameter update. *Curr Opin Allergy Clin Immunol* 2017; **17**: 286-294 [PMID: 28658067 DOI: 10.1097/ACI.0000000000000375]
- Singewald N, Schmuckermair C, Whittle N, Holmes A, Ressler KJ. Pharmacology of cognitive enhancers for exposure-based therapy of fear, anxiety and trauma-related disorders. *Pharmacol Ther* 2015; **149**: 150-190 [PMID: 25550231 DOI: 10.1016/j.pharmthera.2014.12.004]
- Manson ML, Säfholm J, James A, Johnsson AK, Bergman P, Al-Ameri M, Orre AC, Kärman-Mårdh C, Dahlén SE, Adner M. IL-13 and IL-4, but not IL-5 nor IL-17A, induce hyperresponsiveness in isolated human small airways. *J Allergy Clin Immunol* 2020; **145**: 808-817.e2 [PMID: 31805312 DOI: 10.1016/j.jaci.2019.10.037]
- Nunez J, Santillanes G. Anaphylaxis in pediatric patients: early recognition and treatment are critical for best outcomes. *Pediatr Emerg Med Pract* 2019; **16**: 1-24 [PMID: 31124642]
- Curto-Barredo L, Giménez-Arnau AM. Treatment of chronic spontaneous urticaria with an inadequate response to H1-antihistamine. *G Ital Dermatol Venereol* 2019; **154**: 444-456 [PMID: 30717573 DOI: 10.23736/S0392-0488.19.06274-6]
- Castells M, Butterfield J. Mast Cell Activation Syndrome and Mastocytosis: Initial Treatment Options and Long-Term Management. *J Allergy Clin Immunol Pract* 2019; **7**: 1097-1106 [PMID: 30961835 DOI: 10.1016/j.jaip.2019.02.002]
- Mihara M, Hashizume M, Yoshida H, Suzuki M, Shiina M. IL-6/IL-6 receptor system and its role in physiological and pathological conditions. *Clin Sci (Lond)* 2012; **122**: 143-159 [PMID: 22029668 DOI: 10.1042/CS20110340]
- Ouyang W, Rutz S, Crellin NK, Valdez PA, Hymowitz SG. Regulation and functions of the IL-10 family of cytokines in inflammation and

- disease. *Annu Rev Immunol* 2011; **29**: 71-109 [PMID: [21166540](#) DOI: [10.1146/annurev-immunol-031210-101312](#)]
- 21 **Ma K**, Zhang H, Baloch Z. Pathogenetic and Therapeutic Applications of Tumor Necrosis Factor- α (TNF- α) in Major Depressive Disorder: A Systematic Review. *Int J Mol Sci* 2016; **17** [PMID: [27187381](#) DOI: [10.3390/ijms17050733](#)]
- 22 **Cruceriu D**, Baldasici O, Balacescu O, Berindan-Neagoe I. The dual role of tumor necrosis factor-alpha (TNF- α) in breast cancer: molecular insights and therapeutic approaches. *Cell Oncol (Dordr)* 2020; **43**: 1-18 [PMID: [31900901](#) DOI: [10.1007/s13402-019-00489-1](#)]
- 23 **Idriss HT**, Naismith JH. TNF alpha and the TNF receptor superfamily: structure-function relationship(s). *Microsc Res Tech* 2000; **50**: 184-195 [PMID: [10891884](#) DOI: [10.1002/1097-0029\(20000801\)50:3<184::AID-JEMT2>3.0.CO;2-H](#)]
- 24 **Gregory MA**, Manuel-Apolinar L, Sánchez-García S, Villa Romero AR, de Jesús Iuit Rivera J, Basurto Acevedo L, Grijalva-Otero I, Cuadros-Moreno J, García-de la Torre P, Guerrero Cantera J, García Domínguez JA, Martínez Gallardo S, Vega García S, Mejía Alonso LA, Sánchez-Arenas R. Soluble Intercellular Adhesion Molecule-1 (sICAM-1) as a Biomarker of Vascular Cognitive Impairment in Older Adults. *Dement Geriatr Cogn Disord* 2019; **47**: 243-253 [PMID: [31408858](#) DOI: [10.1159/000500068](#)]
- 25 **Hoxha M**, Rovati GE, Cavanillas AB. The leukotriene receptor antagonist montelukast and its possible role in the cardiovascular field. *Eur J Clin Pharmacol* 2017; **73**: 799-809 [PMID: [28374082](#) DOI: [10.1007/s00228-017-2242-2](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

