

World Journal of *Clinical Cases*

World J Clin Cases 2023 November 16; 11(32): 7741-7939



EDITORIAL

- 7741 Efficacy of probiotics supplementation in amelioration of celiac disease symptoms and enhancement of immune system

Moawad MHE, Alkhaldeh IM, Naswhan AJ

ORIGINAL ARTICLE**Retrospective Cohort Study**

- 7745 Effect of enhanced recovery after surgery with multidisciplinary collaboration on nursing outcomes after total knee arthroplasty

Liu J, Zheng QQ, Wu YT

Retrospective Study

- 7753 Appropriate leucine-rich α -2 glycoprotein cut-off value for Japanese patients with ulcerative colitis

Yamazato M, Yanai S, Oizumi T, Eizuka M, Yamada S, Toya Y, Uesugi N, Sugai T, Matsumoto T

- 7761 Association between depressive mood and body image and menopausal symptoms and sexual function in perimenopausal women

Ling J, Wang YH

- 7770 Clinical study of NFNC in the treatment of acute exacerbation chronic obstructive pulmonary disease patients with respiratory failure

Chen X, Dai L, Ma JZ, Chu XX, Dai L, Liu JM, Guo SW, Ru XW, Zhuang XS

- 7778 Mortal condition in an unusual localization, analysis of isolated tongue and tongue base abscesses

Bal KK, Gür H, Demir I, Ismi O, Vayisoglu Y, Gorur K, Ozcan C, Unal M

- 7785 Adult-onset hypophosphatemic osteomalacia as a cause of widespread musculoskeletal pain: A retrospective case series of single center experience

Kim S, Kim SW, Lee BC, Kim DH, Sung DH

Prospective Study

- 7795 Efficacy and safety of laparoscopic *vs* open gastrectomy after neoadjuvant therapy for locally advanced gastric cancer

Yu CD, Zhang K

Randomized Controlled Trial

- 7806 Effect of anesthesia induction with butorphanol on postoperative nausea and vomiting: A randomized controlled trial

Xie F, Sun DF, Yang L, Sun ZL

Randomized Clinical Trial

- 7814** Efficacy and safety of aspirin antiplatelet therapy within 48 h of symptom onset in patients with acute stroke
Zhang JQ, Pan ZB

META-ANALYSIS

- 7822** Analysis of the effectiveness of cognitive rehabilitation for patients with chronic mental illness: A meta-analysis
Jang JS, Oh S, Kim G, Lee N, Song H, Park J, Lee Y, Kim M, Kwon M

SCIENTOMETRICS

- 7833** Emerging trends and hotspots of Nuclear factor erythroid 2-related factor 2 in nervous system diseases
Chang XQ, Xu L, Zuo YX, Liu YG, Li J, Chi HT

CASE REPORT

- 7852** Transcatheter embolization for hemorrhage from aberrant testicular artery after partial nephrectomy: A case report
Youm J, Choi MJ, Kim BM, Seo Y
- 7858** Persistent left superior vena cava in right hemiarch replacement under deep hypothermic circulatory arrest: A case report
Mi ZY, He G, Gao HL, Li C
- 7865** Type II Abernethy malformation with cystic fibrosis in a 12-year-old girl: A case report
Zhang LJ, Liu XY, Chen TF, Xu ZY, Yin HJ
- 7872** Glucocorticoid reduction induced chorea in pediatric-onset systemic lupus erythematosus: A case report
Xu YQ, Wang M, Zhang Y
- 7876** Primary pulmonary lymphoepithelioma-like carcinoma misdiagnosed as lung squamous cell carcinoma: A case report
Yin CJ, Wang GJ, Su XM, Li D
- 7881** Median arcuate ligament syndrome complicated with gallbladder stones: A case report
Dang JQ, Wang QQ, Yang YL, Shang L, Bian QT, Xiang HJ
- 7888** Uterine rupture due to adenomyosis in an adolescent: A case report and review of literature
Kim NI, Lee JS, Nam JH
- 7895** Multiple therapies relieve long-term tardive dyskinesia in a patient with chronic schizophrenia: A case report
Lv L, Guo P, Feng M, Fang Y, Wang SK, Chen HX
- 7900** Lung ultrasound for the early diagnosis of acute lung injury: A case report
Zheng X, Liu N

- 7905** New treatment for gastric duplication cyst: Endoscopic ultrasonography-guided fine-needle aspiration combined with lauromacrogol sclerotherapy: A case report
Bu YW, Han RQ, Ma WQ, Wang GN, Er LM
- 7911** Immunotherapy in SMARCB1 (INI-1)-deficient sinonasal carcinoma: Two case reports
Zhang L, Gao AX, He YL, Xu MJ, Lu HJ
- 7920** Interstitial pneumonia combined with nocardia cyriacigeorgica infection: A case report
Qi DD, Zhuang Y, Chen Y, Guo JJ, Zhang Z, Gu Y
- 7926** Intracranial infection accompanied sweet's syndrome in a patient with anti-interferon- γ autoantibodies: A case report
Zheng JH, Wu D, Guo XY

LETTER TO THE EDITOR

- 7935** Potential and limitations of ChatGPT and generative artificial intelligence in medical safety education
Wang X, Liu XQ

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Guo-Xin Ni, MD, PhD, Chief Doctor, Professor, School of Sport Medicine and Rehabilitation, Beijing Sport University, Beijing 100084, China. guoxinni@bjmu.edu.cn

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJCC* as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Hua-Ge Yan*, Production Department Director: *Xu Guo*, Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

November 16, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Randomized Controlled Trial

Effect of anesthesia induction with butorphanol on postoperative nausea and vomiting: A randomized controlled trial

Fang Xie, De-Feng Sun, Lin Yang, Zhong-Liang Sun

Specialty type: Medicine, research and experimental**Provenance and peer review:** Unsolicited article; Externally peer reviewed.**Peer-review model:** Single blind**Peer-review report's scientific quality classification**Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0**P-Reviewer:** Higa K, Japan**Received:** August 10, 2023**Peer-review started:** August 10, 2023**First decision:** August 30, 2023**Revised:** October 18, 2023**Accepted:** November 9, 2023**Article in press:** November 9, 2023**Published online:** November 16, 2023**Fang Xie, De-Feng Sun, Zhong-Liang Sun**, Department of Anesthesiology, The First Affiliated Hospital of Dalian Medical University, Dalian 116011, Liaoning Province, China**Lin Yang**, Department of Neuroelectrophysiology, The First Affiliated Hospital of Dalian Medical University, Dalian 116011, Liaoning Province, China**Corresponding author:** De-Feng Sun, MS, Professor, Department of Anesthesiology, The First Affiliated Hospital of Dalian Medical University, No. 5 Longbin Road, Dalian 116011, Liaoning Province, China. sundefengyl@163.com**Abstract****BACKGROUND**

Postoperative nausea and vomiting (PONV) are common complications that affect the recovery and well-being of elderly patients undergoing gastrointestinal laparoscopic surgery.

AIM

To investigate the effect of butorphanol on PONV in this patient population.

METHODS

A total of 110 elderly patients (≥ 65 years old) who underwent gastrointestinal laparoscopic surgery were randomly assigned to receive butorphanol (40 $\mu\text{g}/\text{kg}$) or sufentanil (0.3 $\mu\text{g}/\text{kg}$) during anesthesia induction in a 1:1 ratio. The measured outcomes included the incidence of PONV at 48 h after surgery, intraoperative dose of propofol and remifentanil, Bruggmann Comfort Scale score in the postanesthesia care unit (PACU), number of compressions for postoperative patient-controlled intravenous analgesia (PCIA), and time to first flatulence after surgery.

RESULTS

The results revealed a noteworthy reduction in the occurrence of PONV at 24 h after surgery in the butorphanol group, when compared to the sufentanil group (T1: 23.64% vs 5.45%, T2: 43.64% vs 20.00%, $P < 0.05$). However, no significant variations were observed between the two groups, in terms of the clinical characteristics, such as the PONV or motion sickness history, intraoperative and postoperative 48-h total infusion volume and hemodynamic parameters, intraoperative dose of propofol and remifentanil, number of postoperative PCIA compressions, time until the first occurrence of postoperative flatulence, and incidence of PONV at 48 h post-surgery (all, $P > 0.05$). Furthermore, patients in

the butorphanol group were more comfortable, when compared to patients in the sufentanil group in the PACU.

CONCLUSION

The present study revealed that butorphanol can be an efficacious substitute for sufentanil during anesthesia induction to diminish PONV within 24 h following gastrointestinal laparoscopic surgery in the elderly, simultaneously improving patient comfort in the PACU.

Key Words: Butorphanol; Sufentanil; Enhanced recovery after surgery; Anesthesiology; Gastrointestinal surgery; Postoperative nausea and vomiting

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: In this study, butorphanol was used for anesthesia induction, and it was found that the incidence of postoperative nausea and vomiting was significantly lower at 24 h after surgery in the butorphanol group, when compared to the sufentanil group. In addition, the Bruggmann Comfort Scale scores in the postanesthesia care unit were significantly better in the butorphanol group, when compared to the sufentanil group.

Citation: Xie F, Sun DF, Yang L, Sun ZL. Effect of anesthesia induction with butorphanol on postoperative nausea and vomiting: A randomized controlled trial. *World J Clin Cases* 2023; 11(32): 7806-7813

URL: <https://www.wjgnet.com/2307-8960/full/v11/i32/7806.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i32.7806>

INTRODUCTION

Postoperative nausea and vomiting (PONV) is the second most common postoperative adverse reaction after pain, which has an estimated incidence of 30% in the general surgical population, and an incidence that can reach as high as 80% in high-risk patients[1]. Aspiration pneumonia caused by PONV is a severe risk, particularly in elderly patients with poor pharyngeal reflex recovery after general anesthesia. In addition, PONV may cause electrolyte imbalance, poor incision healing, insufficient blood volume, and delayed discharge from the hospital.

The concept of enhanced rehabilitation after surgery emphasizes the role of minimizing adverse reactions after surgery, in order to improve the quality and pace of recovery[2]. The high-risk types of surgery with PONV include laparoscopic, bariatric, and gynecological surgery. The mechanism of PONV induced by the laparoscopic surgery remains unclear. Recent clinical studies have suggested that this may be correlated to the decrease in pain threshold of patients undergoing laparoscopic surgery, the stimulation of residual postoperative carbon dioxide in the abdominal cavity, and the pulling state of the peritoneum, which can result in increased demand for postoperative analgesia, such as opioids, leading to an increased likelihood of PONV[3]. For patients undergoing gastrointestinal laparoscopic surgery, nausea and vomiting are more likely to occur after surgery. Therefore, the balance between analgesia and PONV remains as a major challenge for anesthesiologists.

Traditional opioids produce an analgesic effect by exciting the μ (μ_1 and μ_2) receptors. However, the excitation of μ_2 receptors can enhance the sensitivity to vestibule stimulation, affect the chemoreceptor triggering area, and delay gastric emptying, thereby triggering PONV[4]. In contrast, butorphanol, which is a synthetic opioid receptor agonist-antagonist with 5-8 times the analgesic potency of morphine, exhibits low activity to δ receptors, while stimulating the κ and μ_1 receptors, and antagonizing μ_2 receptors[5]. Through its antagonistic effect on μ_2 receptors, butorphanol significantly reduces the incidence of PONV caused by traditional opioids. Furthermore, clinical studies have revealed that butorphanol has a good analgesic effect on patients with chronic visceral pain through the activation of κ receptors[6,7]. At present, few studies have compared butorphanol and sufentanil in the incidence of PONV during general anesthesia. Therefore, the present study aimed to investigate the effect of butorphanol on PONV in elderly patients who underwent gastrointestinal laparoscopic surgery.

MATERIALS AND METHODS

General information

The present study was approved by the Ethics Committee of the First Affiliated Hospital of Dalian Medical University (PJ-KS-KY-2020-161 [X]), and registered in the China Clinical Trial Center (ChiCTR2100045860). Patients ≥ 65 years old, who underwent gastrointestinal laparoscopic surgery from February 2020 to February 2021, were enrolled for the present study. Using the computer statistics software, these patients were randomly allocated into two groups in a 1:1 ratio: Sufentanil and butorphanol groups.

Based on preliminary experiments and previous studies[8], the sample size was calculated according to the incidence of PONV. The preliminary experiment results indicated that the incidence of PONV was approximately 35% in the sufentanil group, and 13% in the butorphanol group. In order to ensure adequate statistical power with 85% power at 5% level of significance, at least 49 patients were required for each group. Accounting for the potential 10% dropout rate, a total of 110 patients were included for the present study.

Inclusion and exclusion criteria

Inclusion criteria: Patients ≥ 65 years old, who underwent gastrointestinal laparoscopic surgery, and provided a written informed consent. Exclusion criteria: Hypersensitivity to butorphanol and sufentanil, serious respiratory complications, severe obstructive sleep apnea-hypopnea syndrome or obesity [body mass index (BMI) ≥ 28 kg/m²], opioid dependence, significant abnormalities in liver or kidney function, and severe visual or auditory impairment.

Anesthesia monitoring

The following basic clinical information were recorded at one day prior to surgery: Age, gender, height, weight, BMI, American Society of Anesthesiologists (ASA) classification, smoking status, and history of PONV and motion sickness. These patients were required to fast for six hours, and have water deprivation for two hours before the surgery, with no preoperative drugs administered. Upon entering the operation room, the electrocardiogram, heart rate, oxygen saturation (SpO₂), non-invasive blood pressure, bispectral index, and oral and sublingual temperature were monitored. In addition, invasive arterial blood pressure was monitored *via* radial artery catheterization and internal jugular vein catheterization, in order to detect any hemodynamic changes, and facilitate the administration of fluids and medications, when necessary.

Anesthetic method

Anesthesia induction was administered to patients in the sufentanil group at a dose of 0.3 μ g/kg of sufentanil, while patients in the butorphanol group were given 40 μ g/kg of butorphanol, based on the analgesic titer ratio. During the anesthesia induction, the intravenous administration of 1-2 mg/kg of propofol and 0.3 mg/kg of benzenesulfonate atracurium was performed, while remifentanyl was pumped at a rate of 5-10 μ g/kg/h. Then, tracheal intubation was performed under visual laryngoscopy after the muscle relaxant took effect. The anesthesia maintenance during the operation consisted of the intravenous infusion of 4-6 mg/kg/h of propofol, 5-10 μ g/kg/h of remifentanyl, and 0.10-0.15 mg/kg/h of benzenesulfonate atracurium. When the surgery was completed, the infusion of benzenesulfonate atracurium, propofol and remifentanyl were stopped, while 0.1 mg/kg of butorphanol was given for patient-controlled intravenous analgesia (PCIA). Then, these patients were transferred to the postanesthesia care unit (PACU), and vital signs monitoring was continued for 48 h. The study assistants were responsible for the preparation and administration of the studied medications. The other assistants were responsible for the monitoring and recording of the results during data collection. All assistants were blinded to the study.

Outcome measures

The primary outcome of the study was the incidence of PONV, which was evaluated using the PONV grading scale in the PACU (T1), and at 24 h (T2) and 48 h (T3) after surgery (Table 1). The other observed parameters were, as follows: Intra-operative dose of propofol and remifentanyl, total infusion volume (at intraoperative and postoperative 24 and 48 h), operation time, the agitation[9] and Bruggmann Comfort Scale (BCS)[10] scores in the PACU, the number of compressions for PCIA within 48 h after surgery, and the time to first postoperative flatulence. The cumulative dose of propofol and remifentanyl administered through a micropump infusion device, both during the induction and maintenance phases of anesthesia, was calculated using the following formula: Dose = infusion rate (mg/kg/min) \times patient weight (kg) \times duration of surgery (min). The BCS scores were utilized to assess the level of patient comfort in the two groups: 0, indicates continuous pain; 1, represents no pain at rest, but with severe pain during deep breathing or coughing; 2, indicates no pain while lying at rest, and slight pain during deep breathing or coughing; 3, represents no pain during deep breathing; 4, represents no pain during coughing[11].

Statistical methods

For normally distributed measurement data, mean \pm SD was used for the statistical description, and independent sample *t*-test was performed to determine the statistical difference. For non-normally distributed measurement data, median (M) and interquartile range were used for the statistical description, and the Mann-Whitney *U*-test was performed to determine the statistical difference. χ^2 test was used to analyze the difference between groups for the enumeration data. Frequency (rate) was used to describe the ordinal data, and this was analyzed using the Wilcoxon rank sum test. SPSS 26.0 was used for the statistical analysis. A *P* value of < 0.05 was considered statistically significant.


RESULTS

A total of 168 elderly patients, who underwent gastrointestinal laparoscopic surgery from February 2020 to February 2021, were screened in the present study. Among these patients, 35 patients did not agree to participate, and 18 patients were excluded based on the exclusion criteria. During the trial, five patients were excluded due to the following reasons: Rejection and loss to follow-up. Finally, a total of 110 patients (66 male and 44 female patients) were included for the present study (Figure 1).

Table 1 Postoperative nausea and vomiting grading scale

PONV grade	Patient response
0	Without PONV
I	Nausea without vomiting
II	Nausea with vomiting (< 3 times/d)
III	Vomiting ≥ 3 times/d

PONV: Postoperative nausea and vomiting.



DOI: 10.12998/wjcc.v11.i32.7806 Copyright ©The Author(s) 2023.

Figure 1 The patient inclusion, randomization, and follow-up flowchart.

Comparison of baseline characteristics

No significant differences were observed between the two groups, in terms of age, BMI, gender, ASA grade, smoking history, PONV or motion sickness history, intraoperative and postoperative 48-h total infusion volume (Table 2), and hemodynamic parameters (Table 3) ($P > 0.05$).

Comparison of PONVs at postoperative 48 h

As shown in Table 4, there was a significant difference in the occurrence of PONV at T1 ($P = 0.005$) and T2 ($P = 0.001$), while there was no statistical difference at T3 ($P = 0.169$), between the sufentanil and butorphanol groups (Table 4).

Comparison of intraoperative propofol and remifentanyl

There was no significant difference in the total dose of intraoperative propofol ($P = 0.893$) and remifentanyl ($P = 0.438$) between the sufentanil and butorphanol groups (Table 5).

Comparison of agitation and BCS scores

The BCS scores were significantly better in the butorphanol group, when compared to the sufentanil group ($P = 0.028$), although there was no significant difference in agitation scores in the PACU between the two groups ($P = 0.439$) (Table 5).

Comparison of PCIA effective compressions and time to first postoperative flatulence

There were no statistically significant differences observed between the two groups, in terms of the number of PCIA effective compressions at postoperative 48 h ($P = 0.881$), and at the time to first postoperative flatulence ($P = 0.822$) (Table 5).

Table 2 Comparison of baseline characteristics between the sufentanil and butorphanol groups

	Sufentanil group (n = 55)	Butorphanol group (n = 55)	P value
Age	71.0 ± 5.7	69.6 ± 5.7	0.199
Gender (male/female)	32/23	34/21	0.698
ASA (I/II/III)	0/33/22	0/30/25	0.847
Weight (kg)	63.6 ± 10.3	68.6 ± 11.3	0.058
BMI (kg/m ²)	21.1 ± 1.8	20.81 ± 1.7	0.403
Smoking (yes/no)	25/30	26/29	0.703
PONV or motion sickness history (yes/no)	14/41	20/35	0.218
Operation time (h)	3.41 ± 1.30	3.25 ± 1.07	0.484
Intraoperative infusion volume (mL)	1290.9 ± 404.7	1243.6 ± 316.8	0.497
Postoperative 24-h infusion volume (mL)	2380.9 ± 137.6	2342.7 ± 133.8	0.143
Postoperative 48-h infusion volume (mL)	2152.7 ± 128.9	2125.5 ± 117.4	0.249

ASA: American Society of Anesthesiologists; BMI: Body mass index; PONV: Postoperative nausea and vomiting.

Table 3 Comparison of hemodynamics between the sufentanil and butorphanol groups

	HR (BPM)			SBP (mmHg)			DBP (mmHg)		
	Sufentanil	Butorphanol	P value	Sufentanil	Butorphanol	P value	Sufentanil	Butorphanol	P value
Pre-operation	69.2 ± 9.0	67.9 ± 7.1	0.400	145.2 ± 15.2	146.1 ± 12.0	0.748	67.8 ± 6.2	67.51 ± 5.1	0.828
One minute before induction	70.2 ± 9.5	68.8 ± 6.5	0.348	149.5 ± 17.9	149.3 ± 12.4	0.966	71.2 ± 8.8	68.4 ± 5.4	0.056
One minute after tracheal intubation	69.8 ± 9.2	68.8 ± 6.1	0.480	144.3 ± 18.4	147.8 ± 10.5	0.233	69.5 ± 8.7	67.6 ± 4.3	0.156
Intraoperative maintenance	67.3 ± 8.1	68.5 ± 6.3	0.388	145.2 ± 13.9	149.1 ± 9.8	0.098	68.3 ± 7.6	68.1 ± 4.5	0.867

HR: Heart rate; SBP: Systolic blood pressure; DBP: Diastolic blood pressure.

Table 4 Comparison of postoperative nausea and vomiting within postoperative 48 h between the sufentanil and butorphanol groups

	Sufentanil group (n = 55)	Butorphanol group (n = 55)	Z	P value
T1 PONV (0/1/II/III)	42/6/6/1	52/3/0/0	-2.786	0.005 ^a
T2 PONV (0/1/II/III)	21/11/11/2	44/4/7/0	-3.188	0.001 ^a
T3 PONV (0/1/II/III)	50/3/2/0	54/1/0/0	-1.375	0.169

^aThere is significant statistical difference between the two groups, with $P < 0.05$.

T1: During the postanesthesia care unit period; T2: Return to the ward for 24 h; T3: Return to the ward for 24-48 h. PONV: Postoperative nausea and vomiting.

DISCUSSION

The present study compared the effects of sufentanil and butorphanol on the incidence of PONV in elderly patients who underwent gastrointestinal laparoscopic surgery. The results revealed that the incidence of PONV was lower in the PACU, and at 24 h after surgery in the butorphanol group, when compared to the sufentanil group, although there was no statistical difference observed at 48 h after surgery between the two groups.

The complex mechanism of PONV involves the following risk factors: Female gender, smoking, history of PONV or motion sickness, and opioids[12,13]. Several studies have revealed that traditional opioids that are commonly used for pain management, such as μ agonists, have been associated with nausea and vomiting, while providing analgesic efficacy [14,15]. Traditional opioids produce an analgesic effect by exciting the μ (μ_1 and μ_2) receptors. However, the excitation of

Table 5 Comparison of actual doses of propofol and remifentanyl, agitation scores, Bruggmann comfort scale scores, and effective compressions of patient-controlled intravenous analgesia pump

	Sufentanil group (n = 55)	Butorphanol group (n = 55)	Z	P value
Propofol (mg)	1027.2 ± 461.6	1016.4 ± 379.4	-	0.893
Remifentanyl (mg)	1.3 ± 0.6	1.2 ± 0.5	-	0.438
Agitation score (0/1/2/3)	42/8/5/0	47/6/2/0	-0.774	0.439
BCS score (0/1/2/3/4)	6/5/30/14/0	1/4/24/25/1	-2.195	0.028 ^a
Effective compressions of PCIA	1.00 (2.00-1.00)	1.00 (2.00-1.00)	-	0.881

^aThere is significant statistical difference between the two groups, with $P < 0.05$.

BCS: Bruggmann comfort scale; PCIA: Patient-controlled intravenous analgesia pump.

$\mu 2$ receptors can enhance the sensitivity to vestibule stimulation, affect the chemoreceptor triggering area, and delay gastric emptying, thereby triggering PONV[4].

Butorphanol, which is a synthetic opioid receptor agonist-antagonist with 5-8 times the analgesic potency of morphine, exhibits low activity to δ receptors, while stimulating the κ and $\mu 1$ receptors, and antagonizing $\mu 2$ receptors[5]. Sufentanil has a long clearance half-life in elderly patients, and its effect on opioid receptors can persist for several hours after surgery, increasing the incidence and duration of PONV. Since sufentanil undergoes metabolism and clearance over time, its effect on opioid receptors decreases, which may explain the different effects of sufentanil and butorphanol on PONV at different time points.

Recent studies have revealed that butorphanol can effectively inhibit the hemodynamic fluctuations caused by tracheal intubation during anesthesia induction, which is consistent with the results of the hemodynamic parameter analysis in a previous study[16]. In the present study, there was no significant difference in hemodynamic fluctuations before and after endotracheal intubation between the sufentanil and butorphanol groups, and both drugs effectively inhibited the circulation fluctuations caused by the endotracheal intubation. Furthermore, there was no significant difference in intraoperative remifentanyl dose, PACU agitation score, or the number of effective compressions for postoperative PCIA between the sufentanil and butorphanol groups. Thus, it was considered that the induction of anesthesia with butorphanol can produce similar and relatively complete analgesic effects as sufentanil. More importantly, butorphanol can activate the κ receptors, and exert sedative effects. Although there was no statistical difference in intraoperative propofol dose between the two groups in the present study, the BCS scores were higher in the butorphanol group, indicating that the postoperative comfort level of patients induced by butorphanol was higher.

Previous studies have reported that intravenous butorphanol can promote the recovery of postoperative gastrointestinal function, and shorten the time to first postoperative flatulence in elderly patients undergoing radical laparoscopic nephrectomy[17]. However, there was no statistical difference in the time to first postoperative flatulence between the sufentanil and butorphanol groups, which was possibly due to the following factors: Postoperative ambulation time, postoperative dietary recovery, and the use of glycerine enema. Therefore, further comprehensive analyses are required to verify this conclusion.

The limitations of the present study should be acknowledged. Merely the occurrence of nausea and vomiting within 48 h after surgery were observed, and the PDNV was not followed up. Furthermore, the present study merely included elderly patients ≥ 65 years old, who underwent gastrointestinal laparoscopic surgery. Thus, patients in other age groups, especially young women, needs to be investigated. Moreover, the specific operation methods of gastrointestinal surgery were not statistically analyzed in the present study. In addition, other high-risk surgeries, such as pelvic surgery, thyroid surgery, strabismus repair, and middle ear surgery, were not included in the present study[18,19]. Therefore, the conclusions need to be supported by further evidence and more information.

CONCLUSION

In summary, the administration of butorphanol has shown potential in significantly reducing the occurrence of PONV within 24 h after gastrointestinal surgery in elderly patients, and improving the comfort of patients in the PACU. Therefore, the present study contributes valuable evidence that support strategies targeted at mitigating PONV during the perioperative period.

ARTICLE HIGHLIGHTS

Research background

Postoperative nausea and vomiting (PONV) are common complications after surgery, seriously affects the prognosis of elderly patients for laparoscopic gastrointestinal surgery.

Research motivation

This prospective, double-blind randomized controlled trial aimed to investigate the effect of butorphanol on PONV in this patient population.

Research objectives

Elderly patients (≥ 65 years old) who underwent gastrointestinal laparoscopic surgery.

Research methods

Patients were randomly assigned to receive butorphanol (40 $\mu\text{g}/\text{kg}$) or sufentanil (0.3 $\mu\text{g}/\text{kg}$) during anesthesia induction in a 1:1 ratio. The measured outcomes included the incidence of PONV at 48 h after surgery, intraoperative dose of propofol and remifentanyl, Bruggmann Comfort Scale (BCS) score in the postanesthesia care unit (PACU), number of compressions for postoperative patient-controlled intravenous analgesia (PCIA), and time to first flatulence after surgery.

Research results

The results revealed a noteworthy reduction in the occurrence of PONV at 24 h after surgery in the butorphanol group, when compared to the sufentanil group. However, no significant variations were observed between the two groups, in terms of the clinical characteristics, such as the PONV or motion sickness history, intraoperative and postoperative 48-h total infusion volume and hemodynamic parameters, intraoperative dose of propofol and remifentanyl, number of postoperative PCIA compressions, time until the first occurrence of postoperative flatulence, and incidence of PONV at 48 h post-surgery. Furthermore, patients in the butorphanol group were more comfortable, when compared to patients in the sufentanil group in the PACU.

Research conclusions

The administration of butorphanol has shown potential in significantly reducing the occurrence of PONV within 24 h after gastrointestinal surgery in elderly patients, and improving the comfort of patients in the PACU.

Research perspectives

Anesthesia induction with butorphanol may reduce the incidence of PONV, especially for some patients with a high risk of PONV (young women, no-smoking, PONV or motion sickness history, high-risk surgeries, such as pelvic surgery, thyroid surgery, strabismus repair, and middle ear surgery).

FOOTNOTES

Co-first authors: Fang Xie and Lin Yang.

Author contributions: Xie F drafted the manuscript and critically revised the manuscript for important intellectual content; Sun DF approved the final version to be published; Yang L agreement to be accountable for all aspects of the work, ensuring that the questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved; Sun ZL substantial contribution to the conception and design of the study; and all authors read and approved the final manuscript.

Institutional review board statement: The study was approved by the Ethics Committee of the First Affiliated Hospital of Dalian Medical University (PJ-KS-KY-2020-161 [X]).

Clinical trial registration statement: The study is registered in the China Clinical Trial Center (ChiCTR2100045860, 25/04/2021).

Informed consent statement: All participants provided a signed informed consent.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: The datasets generated and/or analyzed in the study are not publicly available due to the limitations of ethical approval, which involve the patient data and anonymity. However, these are available from the corresponding author on reasonable request.

CONSORT 2010 statement: The authors have read the CONSORT 2010 Statement, and the manuscript was prepared and revised according to the CONSORT 2010 Statement.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Fang Xie 0009-0005-1109-7375; De-Feng Sun 0000-0002-5147-2409; Lin Yang 0000-0003-4232-6052; Zhong-Liang Sun 0000-0002-3657-6963.

S-Editor: Wang JJ

L-Editor: A

P-Editor: Zhao S

REFERENCES

- 1 **Gan TJ**, Belani KG, Bergese S, Chung F, Diemunsch P, Habib AS, Jin Z, Kovac AL, Meyer TA, Urman RD, Apfel CC, Ayad S, Beagley L, Candiotti K, Englesakis M, Hedrick TL, Kranke P, Lee S, Lipman D, Minkowitz HS, Morton J, Philip BK. Fourth Consensus Guidelines for the Management of Postoperative Nausea and Vomiting. *Anesth Analg* 2020; **131**: 411-448 [PMID: 32467512 DOI: 10.1213/ANE.0000000000004833]
- 2 **Seyfried S**, Herrle F, Schröter M, Hardt J, Betzler A, Rahbari NN, Reißfelder C. [Initial experiences with the implementation of the enhanced recovery after surgery (ERAS®) protocol]. *Chirurg* 2021; **92**: 428-433 [PMID: 33471183 DOI: 10.1007/s00104-020-01341-1]
- 3 **Stoops S**, Kovac A. New insights into the pathophysiology and risk factors for PONV. *Best Pract Res Clin Anaesthesiol* 2020; **34**: 667-679 [PMID: 33288117 DOI: 10.1016/j.bpa.2020.06.001]
- 4 **Sande TA**, Laird BJA, Fallon MT. The Management of Opioid-Induced Nausea and Vomiting in Patients with Cancer: A Systematic Review. *J Palliat Med* 2019; **22**: 90-97 [PMID: 30239277 DOI: 10.1089/jpm.2018.0260]
- 5 **Laffey DA**, Kay NH. Premedication with butorphanol. A comparison with morphine. *Br J Anaesth* 1984; **56**: 363-367 [PMID: 6712850 DOI: 10.1093/bja/56.4.363]
- 6 **Deflers H**, Gandar F, Bolen G, Dettleux J, Sandersen C, Marlier D. Effects of a Single Opioid Dose on Gastrointestinal Motility in Rabbits (*Oryctolagus cuniculus*): Comparisons among Morphine, Butorphanol, and Tramadol. *Vet Sci* 2022; **9** [PMID: 35051113 DOI: 10.3390/vetsci9010028]
- 7 **Tang W**, Luo L, Hu B, Zheng M. Butorphanol alleviates lipopolysaccharide-induced inflammation and apoptosis of cardiomyocytes via activation of the κ -opioid receptor. *Exp Ther Med* 2021; **22**: 1248 [PMID: 34539844 DOI: 10.3892/etm.2021.10683]
- 8 **Kranke P**, Meybohm P, Diemunsch P, Eberhart L. Risk-adapted strategy or universal multimodal approach for PONV prophylaxis? *Best Pract Res Clin Anaesthesiol* 2020; **34**: 721-734 [PMID: 33288122 DOI: 10.1016/j.bpa.2020.05.003]
- 9 **Zhao Y**, Qin F, Liu Y, Dai Y, Cen X. The Safety of Propofol Versus Sevoflurane for General Anesthesia in Children: A Meta-Analysis of Randomized Controlled Trials. *Front Surg* 2022; **9**: 924647 [PMID: 35813045 DOI: 10.3389/fsurg.2022.924647]
- 10 **Zong S**, Du J, Chen Y, Tao H. Application effect of dexmedetomidine combined with flurbiprofen axetil and flurbiprofen axetil monotherapy in radical operation of lung cancer and evaluation of the immune function. *J BUON* 2021; **26**: 1432-1439 [PMID: 34565001]
- 11 **Ren C**, Xu H, Xu G, Liu L, Liu G, Zhang Z, Cao JL. Effect of intraoperative infusion of dexmedetomidine on postoperative recovery in patients undergoing endovascular interventional therapies: A prospective, randomized, controlled trial. *Brain Behav* 2019; **9**: e01317 [PMID: 31099992 DOI: 10.1002/brb3.1317]
- 12 **Huynh P**, Villaluz J, Bhandal H, Alem N, Dayal R. Long-Term Opioid Therapy: The Burden of Adverse Effects. *Pain Med* 2021; **22**: 2128-2130 [PMID: 33674861 DOI: 10.1093/pm/pnab071]
- 13 **Azzam AAH**, McDonald J, Lambert DG. Hot topics in opioid pharmacology: mixed and biased opioids. *Br J Anaesth* 2019; **122**: e136-e145 [PMID: 31010646 DOI: 10.1016/j.bja.2019.03.006]
- 14 **Mercadante S**. Opioid Analgesics Adverse Effects: The Other Side of the Coin. *Curr Pharm Des* 2019; **25**: 3197-3202 [PMID: 31333113 DOI: 10.2174/1381612825666190717152226]
- 15 **Mercadante S**, Arcuri E, Santoni A. Opioid-Induced Tolerance and Hyperalgesia. *CNS Drugs* 2019; **33**: 943-955 [PMID: 31578704 DOI: 10.1007/s40263-019-00660-0]
- 16 **Kaur J**, Srilata M, Padmaja D, Gopinath R, Bajwa SJ, Kenneth DJ, Kumar PS, Nitish C, Reddy WS. Dose sparing of induction dose of propofol by fentanyl and butorphanol: A comparison based on entropy analysis. *Saudi J Anaesth* 2013; **7**: 128-133 [PMID: 23956709 DOI: 10.4103/1658-354X.114052]
- 17 **Lyu SG**, Lu XH, Yang TJ, Sun YL, Li XT, Miao CH. [Effects of patient-controlled intravenous analgesia with butorphanol versus sufentanil on early postoperative rehabilitation following radical laparoscopic nephrectomy]. *Zhonghua Yi Xue Za Zhi* 2020; **100**: 2947-2951 [PMID: 32993256 DOI: 10.3760/cma.j.cn112137-20200616-01866]
- 18 **Aydin A**, Kaçmaz M, Boyacı A. Comparison of ondansetron, tropisetron, and palonosetron for the prevention of postoperative nausea and vomiting after middle ear surgery. *Curr Ther Res Clin Exp* 2019; **91**: 17-21 [PMID: 31384338 DOI: 10.1016/j.curtheres.2019.06.002]
- 19 **Künzli BM**, Walensi M, Wilimsky J, Bucher C, Bühner T, Kull C, Zuse A, Maurer CA. Impact of drains on nausea and vomiting after thyroid and parathyroid surgery: a randomized controlled trial. *Langenbecks Arch Surg* 2019; **404**: 693-701 [PMID: 31243574 DOI: 10.1007/s00423-019-01799-6]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

