

CONSORT 2010 checklist of information to include when reporting a randomised trial*

Blinding	Implementation	mechanism	Allocation	generation	Sequence	Randomisation:		Sample size			Outcomes		Interventions		Participants		Trial design	Methods	objectives	Background and	Introduction			Title and abstract	Section/Topic
11a	10		9	86	8a		7b	7a	6b		6a		51	4b	4a	<u>3</u> b	3a		2b	2a		1 b	ā		Item No
If done, who was blinded after assignment to interventions (for example, participants, care providers, those	Who generated the random allocation sequence, who enrolled participants, and who assigned participants to interventions	describing any steps taken to conceal the sequence until interventions were assigned	Mechanism used to implement the random allocation sequence (such as sequentially numbered containers),	Type of randomisation; details of any restriction (such as blocking and block size)	Method used to generate the random allocation sequence		When applicable, explanation of any interim analyses and stopping guidelines	How sample size was determined	Any changes to trial outcomes after the trial commenced, with reasons		Completely defined pre-specified primary and secondary outcome measures, including how and when they	actually administered	The interventions for each group with sufficient details to allow replication, including how and when they were	Settings and locations where the data were collected		Important changes to methods after trial commencement (such as eligibility criteria), with reasons	Description of trial design (such as parallel, factorial) including allocation ratio		Specific objectives or hypotheses	Scientific background and explanation of rationale		Structured summary of trial design, methods, results, and conclusions (for specific guidance see CONSORT for abstracts)	Identification as a randomised trial in the title		Checklist item
2 7	7			D .	2	1	E 2	ZIA	2	ノー% *					ما	LA .	7-5	7		7		4)4	E/A		Reported on page No

	Sources of funding and other support (such as supply of drugs), role of funders	25	Funding
	Where the full trial protocol can be accessed, if available	24	Protocol
	Registration number and name of trial registry	23	Registration
			Other information
idence	Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence	22	Interpretation
	Generalisability (external validity, applicability) of the trial findings	21	Generalisability
ses	Trial limitations, addressing sources of potential bias, imprecision, and, if relevant, multiplicity of analyses	20	Limitations
			Discussion
	All important harms or unintended effects in each group (for specific guidance see CONSORT for harms)	19	Harms
	pre-specified from exploratory		
ishing	Results of any other analyses performed, including subgroup analyses and adjusted analyses, distinguishing	18	Ancillary analyses
	For binary outcomes, presentation of both absolute and relative effect sizes is recommended	17b	
	precision (such as 95% confidence interval)		estimation
	For each primary and secondary outcome, results for each group, and the estimated effect size and its	17a	Outcomes and
	by original assigned groups		
sis was	For each group, number of participants (denominator) included in each analysis and whether the analysis was	16	Numbers analysed
	A table showing baseline demographic and clinical characteristics for each group	15	Baseline data
	Why the trial ended or was stopped	14b	
	Dates defining the periods of recruitment and follow-up	14a	Recruitment
	For each group, losses and exclusions after randomisation, together with reasons	13b	recommended)
	were analysed for the primary outcome		diagram is strongly
t, and	For each group, the numbers of participants who were randomly assigned, received intended treatment, and	13a	Participant flow (a
			Results
	Methods for additional analyses, such as subgroup analyses and adjusted analyses	12b	
	Statistical methods used to compare groups for primary and secondary outcomes	12a	Statistical methods
	If relevant, description of the similarity of interventions	11b	
	assessing outcomes) and how		

recommend reading CONSORT extensions for cluster randomised trials, non-inferiority and equivalence trials, non-pharmacological treatments, herbal interventions, and pragmatic trials. Additional extensions are forthcoming: for those and for up to date references relevant to this checklist, see www.consort-statement.org. *We strongly recommend reading this statement in conjunction with the CONSORT 2010 Explanation and Elaboration for important clarifications on all the items. If relevant, we also