

7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA **Telephone:** +1-925-399-1568 **E-mail:** office@baishideng.com https://www.wjgnet.com

PEER-REVIEW REPORT

Name of journal: World Journal of Clinical Cases

Manuscript NO: 88638

Title: Efficacy and safety of Nafamostat mesylate in patients with end-stage renal failure

Provenance and peer review: Unsolicited Manuscript; Externally peer reviewed

Peer-review model: Single blind

Reviewer's code: 06099458

Position: Peer Reviewer

Academic degree: MD, PhD

Professional title: Assistant Professor

Reviewer's Country/Territory: Canada

Author's Country/Territory: China

Manuscript submission date: 2023-11-10

Reviewer chosen by: AI Technique

Reviewer accepted review: 2023-11-13 09:28

Reviewer performed review: 2023-11-21 09:13

Review time: 7 Days and 23 Hours

	[] Grade A: Excellent [] Grade B: Very good [Y] Grade C:
Scientific quality	Good
	[] Grade D: Fair [] Grade E: Do not publish
Novelty of this manuscript	[] Grade A: Excellent [] Grade B: Good [Y] Grade C: Fair [] Grade D: No novelty
Creativity or innovation of this manuscript	 [] Grade A: Excellent [Y] Grade B: Good [] Grade C: Fair [] Grade D: No creativity or innovation



7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-399-1568 E-mail: office@baishideng.com https://www.wjgnet.com

Scientific significance of the conclusion in this manuscript	 [] Grade A: Excellent [Y] Grade B: Good [] Grade C: Fair [] Grade D: No scientific significance
Language quality	[] Grade A: Priority publishing [Y] Grade B: Minor language polishing [] Grade C: A great deal of language polishing [] Grade D: Rejection
Conclusion	[] Accept (High priority) [] Accept (General priority) [Y] Minor revision [] Major revision [] Rejection
Re-review	[Y]Yes []No
Peer-reviewer statements	Peer-Review: [Y] Anonymous [] Onymous Conflicts-of-Interest: [] Yes [Y] No

SPECIFIC COMMENTS TO AUTHORS

In this study, the authors evaluated the efficacy and safety of naproxat mesylate in patients with end-stage renal failure. The results show that the post-treatment prothrombin time, activated partial thromboplastin time, thrombin time, and International Normalized Ratio values in the control group were higher than those in the observation group, while the fibrinogen level in the control group was lower than that in the observation group. Minor comments: 1. The manuscript should be edited. Some minor language polishing should be revised. 2. Please move the tables to the end of the manuscript. 3. Please update the references with PMID and Doi numbers.